Citation for published version:

Belham, P, Hill, MS & Kociok-Kohn, G 2015, 'Alkali metal-mediated Dehydrocoupling of $\text{Me}_2\text{NH.BH}_3$ ', *Dalton Transactions*, vol. 44, no. 27, pp. 12078-12081. <https://doi.org/10.1039/C5DT00178A>

DOI:

[10.1039/C5DT00178A](https://doi.org/10.1039/C5DT00178A)

Publication date:

2015

Document Version

Peer reviewed version

[Link to publication](#)

Publisher Rights

CC BY-ND

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1

Alkali metal-mediated dehydrocoupling of $\text{Me}_2\text{NH}\cdot\text{BH}_3$

Peter Bellham, Michael S. Hill* and
Gabriele Kociok-Köhn

Bis(trimethylsilyl)amide derivatives of the group 1 elements (Li, Na, K) are competent pre-catalysts for the dehydrocoupling of $\text{Me}_2\text{NH}\cdot\text{BH}_3$ via the formation of intermediates containing $[\text{H}_3\text{BNMe}_2\text{BH}_2\text{Me}_2\text{N}]^-$ anions.

Please check this proof carefully. **Our staff will not read it in detail after you have returned it.**

Translation errors between word-processor files and typesetting systems can occur so the whole proof needs to be read. Please pay particular attention to: tabulated material; equations; numerical data; figures and graphics; and references. If you have not already indicated the corresponding author(s) please mark their name(s) with an asterisk. Please e-mail a list of corrections or the PDF with electronic notes attached – do not change the text within the PDF file or send a revised manuscript. Corrections at this stage should be minor and not involve extensive changes. All corrections must be sent at the same time.

Please bear in mind that minor layout improvements, e.g. in line breaking, table widths and graphic placement, are routinely applied to the final version.

We will publish articles on the web as soon as possible after receiving your corrections; **no late corrections will be made.**

Please return your **final** corrections, where possible within **48 hours** of receipt, by e-mail to: dalton@rsc.org

Queries for the attention of the authors

Journal: **Dalton Transactions**

Paper: **c5dt00178a**

Title: **Alkali metal-mediated dehydrocoupling of Me₂NH·BH₃**

Editor's queries are marked like this [Q1, Q2, ...], and for your convenience line numbers are indicated like this [5, 10, 15, ...].

Please ensure that all queries are answered when returning your proof corrections so that publication of your article is not delayed.

Query Reference	Query	Remarks
Q1	For your information: You can cite this article before you receive notification of the page numbers by using the following format: (authors), Dalton Trans., (year), DOI: 10.1039/c5dt00178a.	
Q2	Please carefully check the spelling of all author names. This is important for the correct indexing and future citation of your article. No late corrections can be made.	
Q3	Please check that the inserted CCDC number is correct.	
Q4	Ref. 4 <i>b</i> : Can this reference be updated?	
Q5	Please indicate where ref. 19 should be cited in the text.	

Peter Bellham, Michael S. Hill* and Gabriele Kociok-Köhn

www.rsc.org/dalton

of groups 3–5 and aluminium and gallium in group 13.^{5,6} Our own work has concentrated on the reactivity of similarly d⁰ complexes of magnesium and calcium amides and alkyls with secondary amine boranes, R₂NH·BH₃.⁷ In such cases the formation of the ultimate products of dehydrocoupling, cyclic borazanes of the form [R₂NBH₂]₂, are found to be produced through the formation of isolable compounds containing co-ordinated [H₃BNR₂BH₂R₂N][−] anions (for example, compounds **1** and **2**).

To account for these observations we have suggested that this process is predominantly metal-mediated and dependent upon the mechanism illustrated in Scheme 1,⁷ for which some supporting evidence has recently emerged from computational DFT studies.⁸ Under this regime, the primary B–N bond forming reaction requires a sequence of metallated amidoborane β -hydride elimination and polarised $R_2N=BH_2$ insertion and δ -hydride elimination steps. We have proposed that this latter reaction occurs through the intermediacy of species similar to **1** and **2** and that the kinetic facility for this reaction

† Electronic supplementary information (ESI) available. CCDC 1043568. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5dt00178a

is dictated by the charge density and resultant ability of the M^{2+} centre to polarise and activate a B–H bond of the ligated $[H_3BNR_2BH_2R_2N]^-$ anion, such that the dehydrocoupling activity describes a reactivity trend $Mg > Ca > Sr > Ba$.⁷ Although the parent sodium amidoborane was reported as early as 1938 and more recent research has investigated $[LiNH_2BH_3]$ as a hydrogen storage material,⁹ we are not aware of any report of use of an alkali metal compound to mediate the *catalytic* dehydrocoupling of a representative amine borane. Notably, Sneddon has reported $[Na\{N(SiMe_3)_2\}]$ as a convenient soluble reagent for the formation of five-membered, linear aminoborane anions $Na^+[BH_3N(R)HBH_2N(R)HBH_3]^-$.¹⁰ In this contribution, therefore, we extend our studies to the group 1 (Li, Na and K) bis(trimethylsilyl)amides and provide an initial appraisal of the ability of this series of M^+ cations to mediate the dehydrocoupling of the secondary amine borane, $Me_2NH \cdot BH_3$.

More recent interest in hydrogen storage is pre-dated by over 30 years by Keller's investigation of the reactivity of group 1 reagents with a variety of borane molecules, including dimethylamine borane.^{11–17} The synthesis and solid-state structures of $[Li(NMe_2BH_3)]$ (3) and $[K(NMe_2BH_3)]$ (4) have also recently been described through syntheses employing stoichiometric reactions of $Me_2NH \cdot BH_3$ with *n*-butyl-lithium or the appropriate metal hydride.^{17,18} Although compound 3 has also been utilised as an intermediate in the synthesis of amine borane derivatives of less electropositive metals, the synthesis of these compounds is commonly frustrated by their apparent instability in ethereal solvents. Isolated samples of both 3 and 4 have been described to include up to 40% of an impurity, which is observed as unassigned triplet and quartet resonances in the ^{11}B NMR spectra.¹⁸ Although such species have not been definitively identified, Keller has also reported that the formation of 4, through the reaction of KH with dimethylamine borane, takes place *via* a metallated intermediate $[NMe_2BH_2NMe_2BH_3]^-$ anion which is unstable in the presence of excess metal hydride.^{12,14}

In an effort to resolve these observations we, thus, carried out a study of the reactivity of dimethylamine borane with the group 1 bis(trimethylsilyl)amides, $MN(SiMe_3)_2$ ($M = Li, Na, K$). These reagents were selected in preference to the group 1 hydrides due to their greater solubility in organic media and their greater purity which ensured that accurate stoichiometric quantities of the alkali metal reagent could be reacted with the amine borane. An initial reaction between $[K\{N(SiMe_3)_2\}]$ and two equivalents of $Me_2NH \cdot BH_3$ at room temperature was, thus, undertaken and found to result in the clean formation of Keller's previously described $[K\{NMe_2BH_2NMe_2BH_3\}]$ derivative (5) (eqn (1)).

The ^{11}B NMR spectrum of 5 at room temperature comprised two broad, unresolved resonances at $\delta = -12.3$ ppm and $\delta = 2.6$ ppm, assigned to the BH_3 and BH_2 units of a potassium co-

ordinated $[NMe_2BH_2NMe_2BH_3]^-$ anion respectively, which did not display any resolvable $^1J_{BH}$ coupling even at reduced temperature. Although this ^{11}B NMR chemical shift data is comparable with those of the homoleptic magnesium complex $[(THF)Mg\{NMe_2BH_2NMe_2BH_3\}_2]$ ($\delta = -15.3, 4.0$ ppm)^{7a} and the scandium species $[Sc(NMe_2BH_2NMe_2BH_3)_2\{N(SiMe_3)_2\}]$ ($\delta = -12.9, 4.0$ ppm),⁵ these latter signals were notably better resolved, as quartet and triplet resonances respectively, than those observed for compound 5; rationalised as a consequence of the greater lability of the group 1 element.

Single crystals of compound 5 suitable for single crystal X-ray diffraction analysis were isolated from a concentrated THF solution at $-30^\circ C$.[†] The results of this analysis are shown in Fig. 1 and selected bond length and angle data are provided in the figure caption. The solid-state structure of 5 may be considered as a dimeric molecule, which assembles as a linear coordination polymer of individual $[K\{NMe_2BH_2NMe_2BH_3\} \cdot THF_2]$ and $[K\{NMe_2BH_2NMe_2BH_3\} \cdot THF]$ units which are propagated by intermolecular $BH_2 \cdots K$ and $BH_3 \cdots K$ multi-centre-bonding interactions. The potassium centres are co-ordinated by the nitrogen atom and further multi-centre-bonding interactions with two boron-bound hydrides of the associated $[NMe_2BH_2NMe_2BH_3]^-$ anion and multi-centre-bonding interactions with boron-bound hydrides of a second $[NMe_2BH_2NMe_2BH_3]^-$ unit, augmented by molecules of THF. The linear coordination polymer which propagates through intermolecular interactions of the BH_3 units with the K atoms of adjacent dimeric units is illustrated in Fig. 2.

Whereas the solid-state structure of compound 5 exhibits some variation in the K–O bond distances, the K1–N1 and K2–N3

Fig. 1 ORTEP representation of the asymmetric unit of compound 5. Thermal ellipsoids set at 25% probability. Hydrogen atoms other than the boron-bound hydrides removed for clarity. Selected bond lengths (Å) and angles ($^\circ$); K1–N1 2.8726(16), K2–N3 2.8679(17), K1–O1 2.7781(15), K1–O2 2.7448(15), K2–O3 2.7328(16), N1–B1 1.496(3), B1–N2 1.635(2), N2–B2 1.588(3), N3–B3 1.509(3), B3–N4 1.643(3), N4–B4 1.592(3), O2–K1–O2 172.76(5), K1–N1–B1 86.39(10), N1–B1–N2 116.87(16), B1–N2–B2 111.73(14), O3–K2–N3 98.79(5), K2–N3–B3 86.58(11), N3–B3–N4 116.85(17), B3–N4–B4 112.16(15).

Fig. 2 Propagation of the linear coordination polymer formed by compound 5.

bond lengths are identical within experimental error. The bond lengths of the potassium-bound N–B units [N1–B1 1.496(3), N3–B3 1.509(3) Å] within compound 5 are, however, significantly shorter than those of any previously reported metallated $[\text{NMe}_2\text{BH}_2\text{NMe}_2\text{BH}_3]^-$ anion (typically 1.56 Å), whilst the remaining B–N bonds are longer than the corresponding linkages within any previously reported comparable group 2 or group 3 compounds.^{5,7} Although these variations may be rationalised by the adjustment to the identity of the d^0 metal centre, it is likely that the formation of a coordination polymer propagated through multi-centre-bonding interactions between the potassium centres and hydrides of both boron atoms also exerts a significant influence over the precise metrical parameters within the individual $[\text{NMe}_2\text{BH}_2\text{NMe}_2\text{BH}_3]^-$ anions.

Identical reactions performed between $[\text{Li}\{\text{N}(\text{SiMe}_3)_2\}]$ and $[\text{Na}\{\text{N}(\text{SiMe}_3)_2\}]$ with two equivalents of $\text{Me}_2\text{NH}\cdot\text{BH}_3$ at room temperature provided access to the corresponding lithium (6) and sodium (7) $[\text{NMe}_2\text{BH}_2\text{NMe}_2\text{BH}_3]^-$ derivatives, characterised as quartet and triplet resonances (6: $\delta = -14.7$ ppm $^1J_{\text{BH}} = 89$ Hz and 3.5 ppm $^1J_{\text{BH}} = 98$ Hz; 7: $\delta = -14.7$ ppm $^1J_{\text{BH}} = 87$ Hz and 1.9 ppm $^1J_{\text{BH}} = 98$ Hz) in the respective ^{11}B NMR spectra. Both reactions, however, proceeded markedly more slowly than for the synthesis of compound 5 and neither 6 nor 7 could be isolated in pure form. The formation of the sodium derivative was also accompanied by $[\text{Na}\{\text{NMe}_2\text{BH}_3\}]$, which appeared as a quartet resonance coincidental with the BH_3 unit of 7 in the ^{11}B NMR spectrum.

The similarity of the outcome the stoichiometric reactivity employed in the formation of compounds 5–7 to the species identified as intermediates during the group 2- and group 3-catalysed dehydrocoupling of $\text{Me}_2\text{NH}\cdot\text{BH}_3$ (e.g. 1 and 2)^{5,7} prompted us to assess the activity of the lithium, sodium and potassium bis(trimethylsilyl)amides under catalytic conditions. An initial assessment of the catalytic dehydrocoupling of $\text{Me}_2\text{NH}\cdot\text{BH}_3$ was, thus, undertaken using 5 mol% of each group 1 reagent in toluene solution. Dehydrocoupling was observed to take place at 80 °C albeit very slowly over the course of 124 hours. In each case the formation of compounds 5–7, comprising ca. 5–10% of the total boron-containing species, was clearly observed in the relevant ^{11}B NMR spectra and the ultimate products of the catalysis were observed to be the cyclic borazane, $[\text{Me}_2\text{NBH}_2]_2$ and a minor quantity of the diaminoborane, $[\text{HB}(\text{NMe}_2)_2]$. We have previously suggested that the production of the latter species is a consequence of a

Table 1 Conversions (%) deduced by integration of ^{11}B NMR spectra for the dehydrocoupling of $\text{Me}_2\text{NH}\cdot\text{BH}_3$ by 5 mol% of the group 1 bis(trimethylsilyl)amides after heating at 80 °C for ca. 124 hours. (mass balance provided by unreacted $\text{Me}_2\text{NH}\cdot\text{BH}_3$)

	$[\text{Li}\{\text{N}(\text{SiMe}_3)_2\}]$	$[\text{Na}\{\text{N}(\text{SiMe}_3)_2\}]$	$[\text{K}\{\text{N}(\text{SiMe}_3)_2\}]$
$[\text{Me}_2\text{NBH}_2]_2$	72	43	43
$[\text{HB}(\text{NMe}_2)_2]$	5	3	6

β -hydride elimination pathway which is competitive with the δ -hydride elimination required for production of the cyclic borazane (Scheme 1). It is notable, however, that both processes result in the production of the reactive group 1 hydride required for onward reaction with $\text{Me}_2\text{NH}\cdot\text{BH}_3$ within the catalytic manifold. The conversions observed in this qualitative study, summarised in Table 1, indicated that a greater efficacy for the dehydrocoupling reaction is displayed by the smaller lithium centre. Notably the sodium and potassium-based reactions were observed to slow markedly at more advanced reaction times.

To establish the veracity of this apparent trend in reactivity, the reaction kinetics were interrogated by examination of the initial rates of the three reactions between 0.5 mmol $\text{Me}_2\text{NH}\cdot\text{BH}_3$ and 5 mol% of the relevant bis(trimethylsilyl)amide at 50 °C, with the consumption of $\text{Me}_2\text{NH}\cdot\text{BH}_3$ monitored by ^{11}B NMR spectroscopy. The results of these experiments revealed that the initial dehydrocoupling catalysed by the lithium and sodium reagents proceeded at comparable rates, which were intermediate between those previously observed magnesium- and calcium-based catalyses. In contrast the reaction between $[\text{K}\{\text{N}(\text{SiMe}_3)_2\}]$ and $\text{Me}_2\text{NH}\cdot\text{BH}_3$ progressed more slowly and at a rate commensurate with that observed in our previous study of $[\text{Ca}\{\text{N}(\text{SiMe}_3)_2\}_2]$.⁷ Although these initial rate data, thus, suggest a trend in reactivity $\text{Mg} > \text{Li} \approx \text{Na} > \text{K} \approx \text{Ca}$, they must be treated with caution as they take no account of other factors at play during more advanced stages of the catalytic reactions. We tentatively ascribe, for example, the lower conversion for the sodium-based reaction (Table 1), which was observed to slow markedly at more advanced stages of the reaction, as a likely consequence of the aggregation of a hydrocarbon-insoluble sodium hydride species.

Conclusions

Group 1 bis(trimethylsilyl)amides act as viable pre-catalysts for the dehydrocoupling of $\text{Me}_2\text{NH}\cdot\text{BH}_3$. The catalytic reactivity displays similar features to previous reports of dehydrocoupling processes catalysed by d^0 metal centres. The reactions proceed through intermediates containing $[\text{H}_3\text{BNMe}_2\text{BH}_2\text{Me}_2\text{N}]^-$ anions and display a notable kinetic dependence on the identity of the catalytic group 1 reagent, which we suggest is a consequence of variation in charge density of the d^0 centre.

Acknowledgements

We thank the University of Bath for a University Research Scholarship (PB).

Notes and references

†X-Ray diffraction data for 5. $\text{C}_{20}\text{H}_{58}\text{B}_4\text{K}_2\text{N}_4\text{O}_{34}$, $M = 524.14$, monoclinic, $P2_1/c$, $a = 12.5404(3)$ Å, $b = 17.0035(4)$ Å, $c = 15.9352(4)$ Å, $\beta = 106.6308(12)^\circ$, $V = 3255.74(14)$ Å³, $Z = 4$, $\rho = 1.069$ g cm³, temperature 150(2) K, $R_1 [I > 2\sigma(I)] = 0.0424$, $wR_2 [I > 2\sigma(I)] = 0.0962$, $R_1 [\text{all data}] = 0.0699$, $wR_2 [\text{all data}] = 0.1096$, measured reflections = 61 939, unique reflections = 6385, $R_{\text{int}} = 0.0977$. Crystallographic data for compound 5 are available as CCDC 1043568.

- 1 T. Autrey, M. Bowden and A. Karkamkar, *Faraday Discuss.*, 2011, **151**, 157–169.
- 2 Z. T. Xiong, C. K. Yong, G. T. Wu, P. Chen, W. Shaw, A. Karkamkar, T. Autrey, M. O. Jones, S. R. Johnson, P. P. Edwards and W. I. F. David, *Nat. Mater.*, 2008, **7**, 138–141.
- 3 See, for example: (a) Y. S. Chua, H. Wu, W. Zhou, T. J. Udovic, G. Wu, Z. Xiong, M. W. Wong and P. Chen, *Inorg. Chem.*, 2012, **51**, 1599–1603; (b) K. Shimoda, Y. Zhang, T. Ichikawa, H. Miyaoka and Y. Kojima, *J. Mater. Chem.*, 2011, **21**, 2609–2615.
- 4 For reviews, see: (a) F. H. Stephens, V. Pons and T. R. Baker, *Dalton Trans.*, 2007, 2613–2626; (b) H. C. Johnson, T. N. Hooper and A. S. Weller, *Top. Organomet. Chem.*, 2015, DOI: 10.1007/978-3-319-13054-5_6, in press.
- 5 M. S. Hill, G. Kociok-Köhn and T. P. Robinson, *Chem. Commun.*, 2010, **46**, 7587–7589.
- 6 (a) H. J. Cowley, M. S. Holt, R. L. Melen, J. M. Rawson and D. S. Wright, *Chem. Commun.*, 2011, **47**, 2682–2684; (b) R. J. Less, R. L. Melen and D. S. Wright, *RSC Adv.*, 2012, **2**, 2191–2199; (c) M. M. Hansmann, R. L. Melen and D. S. Wright, *Chem. Sci.*, 2011, **2**, 1554–1559.
- 7 (a) D. J. Liptrot, M. S. Hill, M. F. Mahon and D. J. MacDougall, *Chem. – Eur. J.*, 2010, **16**, 8508–8515; (b) M. S. Hill, M. Hodgson, D. J. Liptrot and M. F. Mahon, *Dalton Trans.*, 2011, **40**, 7783–7790.
- 8 V. Butera, N. Russo and E. Sicilia, *Chem. – Eur. J.*, 2014, **20**, 5967–5976.
- 9 H. I. Schlesinger and A. B. Burg, *J. Am. Chem. Soc.*, 1938, **60**, 290.
- 10 W. C. Ewing, P. J. Carroll and L. G. Sneddon, *Inorg. Chem.*, 2013, **52**, 10690–10697.
- 11 C. R. Balulescu and P. C. Keller, *Inorg. Chem.*, 1978, **17**, 3707–3708.
- 12 P. C. Keller, *Inorg. Chem.*, 1975, **14**, 438–440.
- 13 P. C. Keller, *Inorg. Chem.*, 1975, **14**, 440–440.
- 14 P. C. Keller, *J. Am. Chem. Soc.*, 1974, **96**, 3078–3085.
- 15 L. D. Schwartz and P. C. Keller, *Inorg. Chem.*, 1973, **12**, 947–949.
- 16 P. C. Keller, *Inorg. Chem.*, 1971, **10**, 2256–2259.
- 17 L. D. Schwartz and P. C. Keller, *Inorg. Chem.*, 1971, **10**, 645–649.
- 18 D. J. Wolstenholme, J. Flogeras, F. N. Che, A. Decken and G. S. McGrady, *J. Am. Chem. Soc.*, 2013, **135**, 2439–2442.
- 19 C. A. Jaska, K. Temple, A. J. Lough and I. Manners, *J. Am. Chem. Soc.*, 2003, **125**, 9424–9434.