Citation for published version:

Keogh, E 2015, 'Men, masculinity and pain', *Pain*, vol. 156, no. 12, pp. 2408-2412.
<https://doi.org/10.1097/j.pain.0000000000000328>

DOI:

[10.1097/j.pain.0000000000000328](https://doi.org/10.1097/j.pain.0000000000000328)

Publication date:

2015

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Men, masculinity and pain

Edmund Keogh

University of Bath

(In press, PAIN)

Correspondence: Edmund Keogh, Department of Psychology & Centre for Pain Research, University of Bath, Claverton Down, Bath, BA2 7AY, United Kingdom, Tel: +44(0)1225 383671; Fax: +44(0)1225 386752; E-mail: e.m.keogh@bath.ac.uk; URL: <http://www.bath.ac.uk/pain>

Type of manuscript: Topical Review

Number of pages: 23

Number of Figures: 1

Word count: 1,997 (2,000 max)

Running head: Men, masculinity and pain

Keywords: sex; gender; men; masculinity; gender-roles; pain.

Disclosures: The author has received unrelated research funding support from Reckitt Benckiser Healthcare (UK) Limited, and Engineering and Physical Sciences Research Council.

1. Introduction

Men's pain is not well understood, despite clear evidence that sex and gender play important roles in pain and disability [35]. Outside pain, men's health has emerged as an area of interest, and not only identified issues specific to men, but helped us better understand male-female differences as well. By taking a men's health approach to pain, we can view existing evidence in a different way, and identify potential gaps in our understanding of men (and women's) pain. The purpose of this selective review is therefore to identify key themes from the men's health literature and consider them within the context of men's pain. Whilst this review will mostly highlight psychosocial factors, it is acknowledged that biological factors contribute to men's pain. The review will focus on three core areas - morbidity/mortality, health behaviours, and masculinity – as well as the relationship between them (see Figure 1). The final section considers future directions, including implications for clinical research and practice.

Figure 1

2. Gender differences in mortality and morbidity

One key theme that dominates men's health literature is the general, yet paradoxical, gender difference in mortality and morbidity rates. Typically, males have lower life expectancy [41; 75; 113] – males are more vulnerable to common life-limiting conditions [65; 75], and more likely to die from accidental and non-accident injury [11; 29; 82], including suicide [78]. Females, however, seemed to have worse overall health i.e., higher morbidity [22; 79; 92].

This health-survival difference is a good starting point for the current review. It forces us to step back, identify general themes, and ask question that might not have yet been asked within the context of pain. For example, chronic pain is viewed as long-term condition, although the life-limiting aspects of pain are starting to emerge [99; 105; 118]. However, few studies investigate the impact of gender on the relationship between pain and mortality beyond statistical control. In one of the few known studies, Docking et al. [30] found a greater mortality risk in older women with lower back pain, but not in men. It is not known whether this bias occurs in other age groups or painful conditions. Given the increased risk of suicide in both chronic pain and men [52; 100], it is possible that males might show greater pain-related suicide. One way to examine this could be to look at non-accidental opioid-related deaths. Whilst there is a general trend for women to use poisoning/overdose as a method of suicide [5; 78], opioid-related overdoses can be higher in men [84; 85]. Taking a men's health perspective highlights the need to consider outcomes such as mortality, suicide etc.

Despite ambiguities around pain-related mortality, gender results in a clear and well recognised pattern of pain morbidity. Prevalence of pain is generally higher in women than men [35; 58; 63]. However, these studies also show that pain still occurs to a high degree in men – and the rate is not far behind women. Men are not immune from pain and suffering. There are also common male-specific conditions where pain plays a significant role e.g., testicular/prostate cancer, prostatitis. Although fewer in number, there are shared pain conditions that show a male bias

e.g., cluster headache [37]. What this tells us is that men suffer from pain, and this should not be overlooked when considering gender-related issues.

3. Men's health behaviours

3.1 Health behaviour and utilization

Explanations for these gender differences can also be found in the men's health literature. Males might have poorer survival rates because they engage in riskier behaviours [66; 87]. Men are more likely to smoke, use excessive alcohol, have poor diets etc [28; 96; 110]. Men are less likely to engage with healthcare services, and have fewer health checks or physician consultations [2; 34; 43; 68; 111]. Given the importance of catching life-threatening conditions early, delays negatively affect survival rates. When men do engage, health behaviours can be good e.g., some studies find medication adherence to be better in men [48; 71].

An important priority is to understand why men continue to engage in riskier behaviours, and seem resistant to health promotion messages. Barriers men face around help seeking include ineffective communication skills, embarrassment, and anxiety [116]. Lack of support has been related to poorer health [47]. Men tend to have a narrower range of support networks [7], engaging more with family members than those outside the family [10; 25; 97]. Levels of available support can get worse for men with increasing age [36]. Health campaigns help men overcome embarrassment, adopt better health behaviours, and improve social support networks. For example, the 'men's sheds' initiative provide male friendly environments that help reduce isolation and improve health awareness [114].

3.2 Pain care utilisation and pain coping behaviours

These general explanations can be applied to pain. We know men have a lower take up of pain services [24; 33; 55] and are less likely to report using analgesics [9; 44; 83]. This could be because women suffer more pain [77]. Alternatively, health seeking behaviours may help explain lower pain service use by men. However, few studies consider gender [45], and results are mixed. Access to back pain services do not seem to show male-female differences in help-seeking [1; 51] - it is unknown whether men avoid seeking help across other painful conditions. It would also be interesting to know whether there are expectations around men's 'need' for analgesia, and if asking for pain relief is seen by some as a 'signal of weakness'.

Few studies take a men's health approach to pain behaviours, and so relatively little is known about men's self-management of pain [38]. Men seem to use a narrower range of pain coping strategies [32; 107]. They report using distraction-type approaches [61], even though they may benefit from focusing on pain [62]. Men are more likely to use alcohol to cope with pain [19; 94], and more males present at emergency rooms following combined alcohol and opioid use [54]. Males reported using 'adaptive' coping [32], but also avoidance [90; 91], and the usefulness of such approaches depend on context. Fear avoidance may contribute to pain chronicity [109], and the relationship between pain anxiety and pain can be stronger in men [31; 40] - although conflicting reports have been found [74; 102]. Men who perceive pain as more threatening have also been found to adopt passive coping responses [53]. Together, it seems that male coping patterns are worth further consideration, especially if related to pain avoidance.

4. Masculinity, health and pain

4.1 Masculinity and health

At the heart of men's health is masculinity, which refers to attitudes and beliefs about what it is to be 'male'. Masculinity is culturally defined, and so we need to be mindful that much of the research and measures used stems from North America and Europe [103]. Masculine behaviours are exemplified as being stoic and less emotional, independent and more dominant or aggressive. These views shape an individual's gender identity and influence behaviour, including health [2; 26; 69; 70; 98]. Identification with strength and independence explains why some men are less willing to express health concerns and seek help [101]. Gender beliefs are thought to be acquired in childhood (friends, family), then shaped and reinforced throughout life [20; 72]. Media representations of male health often reflect strength [8], and violation of male gender norms can result in negative responses [27; 50].

The relationship between masculinity and health is reciprocal - poor health can be perceived as a threat to masculinity and result in a loss of male identity [27; 42; 117]. This loss can lead to gender role strain [64], increased depression, and may even be linked to suicide [49; 73; 80]. Some interventions attempt to challenge traditional masculine beliefs in order to improve wellbeing and promote healthy behaviours [69].

4.2 Masculinity, gender identity and pain

Although masculinity has not been extensively considered within the context of pain, but there are signs it is important [16; 38]. Greater male identification (high masculinity/low femininity) is associated with higher pain thresholds in laboratory studies [6; 39]. Gender identity is also embedded in how men describe clinical pain

[3; 4; 106]. Stereotypical beliefs exist around male pain - the typical male is viewed as less willing to report pain, and more likely to use distraction [60; 95]. Female pain is also perceived to be wider ranging and related to reproductive status [18]. Gender-role beliefs explain male-female variation in laboratory-induced pain, and affect how pain signals are interpreted [88; 115]. Unfortunately, the link between masculinity and pain behaviours has not been examined from a developmental perspective, and it is unclear whether they co-develop during childhood [58; 76; 108].

The expression of pain could challenge masculine identity, and so pain can have a demasculinising effect [17; 86]. Chronic pain can affect sexuality and sexual functioning [81; 89], which can also impact on gender identity. Interestingly, opioids can affect testosterone [12; 14], and reduced testosterone is implicated in some common health problems [56, 57]. It is unclear whether psychosocial pain interventions that seek to improve pain and disability affect masculine identity in any way. It is unknown whether pain-related threats to masculine identity contribute to mortality, including suicide. Campaigns that raise awareness around effects of chronic pain on men and masculinity have yet to be developed and tested.

5. Next steps/future directions

- There is an urgent need to determine whether a gender pain-survival paradox exists. We know there is an increased female vulnerability to pain, but not whether males have higher chronic pain-related deaths. There is a related need to consider gender differences in suicide amongst those in pain.
- The general absence of a men's health focus has led to calls for specific health policies and targeted campaigns that focus on men [13; 15; 34; 93;

112]. What are the healthcare needs and policy implications for the management of men's pain? There is a general need to improve interventions for chronic pain, with a move towards greater personalisation of treatment. Should there be specific interventions that recognise the impact of pain on men? Can a men's health approach be used to help improve treatments for both men and women in pain? For example, if women are more open to discussing issues and using social support, how should this be used to inform how to best deliver group-based interventions?

- Little is known about the structure, nature and impact of masculinity in pain, beyond simple conceptualisations [103]. To what extent are masculine gender roles specific to pain, or are they part of a more general health behaviour to consider? If they are general health behaviours, are they still relevant to pain? There are also multiple masculinities (e.g., identity, stereotypes, attributions [104]), and dispositional/situational aspects – how is this relevant to pain? Whilst masculinity is most often examined in men, it is relevant for women also. There is little pain research into the impact of masculine behaviours in women [46], and feminine behaviours in men.
- Greater understanding of the reciprocal relationship between masculinity and pain is required. What effect does masculine gender identity have on pain, pain reporting and pain behaviours. To what extent do masculine beliefs affect empathy, and the identification of pain in others [59]? Similarly, little is known about the consequences of pain-related changes to male gender identity e.g., social relationships, quality of life. We do not know whether this relationship changes across the lifespan, from childhood, through work, and into older life.

6. Conclusions

Men experience a high level of pain, but are generally less likely to engage with health care services. We know little about how pain specifically affects men, and whether we should take a male focused approach to pain management for men. Advances in men's health, and our understanding of constructs such as masculinity, are emerging as important for pain as well. This review points to areas for immediate attention, such as morbidity/mortality differences, health behaviours, and health regulation/promotion. It also highlights the great potential in fully considering the gendered context in which pain operates.

7. Acknowledgements/Conflict of interest statement

The author has no known conflicts of interest associated with this work.

The author has received unrelated research funding support from Reckitt Benckiser Healthcare (UK) Limited, and Engineering and Physical Sciences Research Council.

9. References

- [1] Adamson J, Hunt K, Nazareth I. The influence of socio-demographic characteristics on consultation for back pain-a review of the literature. *Fam Pract* 2011;28(2):163-171.
- [2] Addis ME, Mahalik JR. Men, masculinity, and the contexts of help seeking. *Am Psychol* 2003;58(1):5-14.
- [3] Ahlsen B, Bondevik H, Mengshoel AM, Solbraekke KN. (Un)doing gender in a rehabilitation context: a narrative analysis of gender and self in stories of chronic muscle pain. *Disabil Rehabil* 2014;36(5):359-366.
- [4] Ahlsen B, Mengshoel AM, Solbraekke KN. Troubled bodies - troubled men: a narrative analysis of men's stories of chronic muscle pain. *Disabil Rehabil* 2012;34(21):1765-1773.
- [5] Ajdacic-Gross V, Weiss MG, Ring M, Hepp U, Bopp M, Gutzwiller F, Rossler W. Methods of suicide: international suicide patterns derived from the WHO mortality database. *B World Health Organ* 2008;86(9):726-732.
- [6] Alabas OA, Tashani OA, Tabasam G, Johnson MI. Gender role affects experimental pain responses: a systematic review with meta-analysis. *Eur J Pain* 2012;16(9):1211-1223.
- [7] Almquist YB, Ostberg V, Rostila M, Edling C, Rydgren J. Friendship network characteristics and psychological well-being in late adolescence: exploring differences by gender and gender composition. *Scand J Public Health* 2014;42(2):146-154.
- [8] Anstiss D, Lyons A. From men to the media and back again: help-seeking in popular men's magazines. *J Health Psychol* 2014;19(11):1358-1370.

- [9] Antonov KI, Isacson DG. Prescription and nonprescription analgesic use in Sweden. *Ann Pharmacother* 1998;32(4):485-494.
- [10] Antonucci TC, Akiyama H. An examination of sex-differences in social support among older men and women. *Sex Roles* 1987;17(11-12):737-749.
- [11] Archer J. Sex differences in aggression in real-world settings: a meta-analytic review. *Rev Gen Psychol* 2004;8(4):291-322.
- [12] Basaria S, Travison TG, Alford D, Knapp PE, Teeter K, Cahalan C, Eder R, Lakshman K, Bachman E, Mensing G, Martel MO, Le D, Stroh H, Bhasin S, Wasan AD, Edwards RR. Effects of testosterone replacement in men with opioid-induced androgen deficiency: a randomized controlled trial. *Pain* 2015;156(2):280-288.
- [13] Bates LM, Hankivsky O, Springer KW. Gender and health inequities: a comment on the final report of the WHO Commission on the social determinants of health. *Soc Sci Med* 2009;69(7):1002-1004.
- [14] Bawor M, Bami H, Dennis BB, Plater C, Worster A, Varenbut M, Daiter J, Marsh DC, Steiner M, Anglin R, Coote M, Pare G, Thabane L, Samaan Z. Testosterone suppression in opioid users: a systematic review and meta-analysis. *Drug Alcohol Depend* in press.
- [15] Bell DL, Breland DJ, Ott MA. Adolescent and young adult male health: a review. *Pediatrics* 2013;132(3):535-546.
- [16] Bernardes SF, Keogh E, Lima ML. Bridging the gap between pain and gender research: a selective literature review. *Eur J Pain* 2008;12(4):427-440.
- [17] Bernardes SF, Lima ML. Being less of a man or less of a woman: Perceptions of chronic pain patients' gender identities. *Eur J Pain* 2010;14(2):194-199.

- [18] Bernardes SF, Silva SA, Carvalho H, Costa M, Pereira S. Is it a (fe) male pain? Portuguese nurses' and laypeople's gendered representations of common pains. *Eur J Pain* 2014;18(4):530-539.
- [19] Brennan PL, Schutte KK, SooHoo S, Moos RH. Painful medical conditions and alcohol use: a prospective study among older adults. *Pain Med* 2011;12(7):1049-1059.
- [20] Bussey K, Bandura A. Social cognitive theory of gender development and differentiation. *Psychol Rev* 1999;106(4):676-713.
- [21] Campbell CI, Weisner C, LeResche L, Ray GT, Saunders K, Sullivan MD, Banta-Green CJ, Merrill JO, Silverberg MJ, Boudreau D, Satre DD, Von Korff M. Age and gender trends in long-term opioid analgesic use for noncancer pain. *Am J Public Health* 2010;100(12):2541-2547.
- [22] Case A, Paxson C. Sex differences in morbidity and mortality. *Demography* 2005;42(2):189-214.
- [23] Cicero TJ, Wong G, Tian Y, Lynskey M, Todorov A, Isenberg K. Co-morbidity and utilization of medical services by pain patients receiving opioid medications: data from an insurance claims database. *Pain* 2009;144(1-2):20-27.
- [24] Cornally N, McCarthy G. Chronic pain: the help-seeking behavior, attitudes, and beliefs of older adults living in the community. *Pain Manag Nurs* 2011;12(4):206-217.
- [25] Cornwell B. Independence through social networks: bridging potential among older women and men. *J Gerontol B-Psychol* 2011;66(6):782-794.
- [26] Courtenay WH. Constructions of masculinity and their influence on men's well-being: a theory of gender and health. *Soc Sci Med* 2000;50(10):1385-1401.

- [27] Courtenay WH. Key determinants of the health and well-being of men and boys. *Int J Mens Health* 2003;2(1):1-30.
- [28] Craig R, Mindell J. Health Survey for England 2011: health, social care and lifestyles. Leeds: Health and Social Care Information Centre, 2012.
- [29] de Almeida RM, Cabral JC, Narvaes R. Behavioural, hormonal and neurobiological mechanisms of aggressive behaviour in human and nonhuman primates. *Physiol Behav* 2015;143:121-135.
- [30] Docking RE, Fleming J, Brayne C, Zhao J, Macfarlane GJ, Jones GT, Cambridge City over-75s Cohort study collaboration. The relationship between back pain and mortality in older adults varies with disability and gender: results from the Cambridge City over-75s Cohort (CC75C) study. *Eur J Pain* 2015;19(4):466-472.
- [31] Edwards RR, Augustson EM, Fillingim RB. Sex-specific effects of pain-related anxiety on adjustment to chronic pain. *Clin J Pain* 2000;16(1):46-53.
- [32] El-Shormilisy N, Strong J, Meredith PJ. Associations between gender, coping patterns and functioning for individuals with chronic pain: a systematic review. *Pain Res Manag* 2015;20(1):48-55.
- [33] Eriksen J, Sjogren P, Ekholm O, Rasmussen NK. Health care utilisation among individuals reporting long-term pain: an epidemiological study based on Danish National Health Surveys. *Eur J Pain* 2004;8(6):517-523.
- [34] European Commission. The state of men's health in Europe - extended report. Brussels: European Union, 2011.
- [35] Fillingim RB, King CD, Ribeiro-Dasilva MC, Rahim-Williams B, Riley JL, 3rd. Sex, gender, and pain: a review of recent clinical and experimental findings. *J Pain* 2009;10(5):447-485.

- [36] Fischer CS, Beresford L. Changes in support networks in late middle age: the extension of gender and educational differences. *J Gerontol B Psychol Sci Soc Sci* 2015;70(1):123-131.
- [37] Fischera M, Marziniak M, Gralow I, Evers S. The incidence and prevalence of cluster headache: a meta-analysis of population-based studies. *Cephalalgia* 2008;28(6):614-618.
- [38] Flurey CA, Hewlett S, Rodham K, White A, Noddings R, Kirwan J. Men, rheumatoid arthritis, psychosocial impact and self-management: A narrative review. *J Health Psychol* in press.
- [39] Fowler SL, Rasinski HM, Geers AL, Helfer SG, France CR. Concept priming and pain: an experimental approach to understanding gender roles in sex-related pain differences. *J Behav Med* 2011;34(2):139-147.
- [40] Frot M, Feine JS, Bushnell MC. Sex differences in pain perception and anxiety. A psychophysical study with topical capsaicin. *Pain* 2004;108(3):230-236.
- [41] G. B. D. Mortality and Causes of Death Collaborators. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;385(9963):117-171.
- [42] Garfield CF, Isacco A, Rogers TE. A review of men's health and masculinity. *Am J Lifestyle Med* 2008;2(6):474-487.
- [43] Green CA, Pope CR. Gender, psychosocial factors and the use of medical services: a longitudinal analysis. *Soc Sci Med* 1999;48(10):1363-1372.
- [44] Hargreave M, Andersen TV, Nielsen A, Munk C, Liaw KL, Kjaer SK. Factors associated with a continuous regular analgesic use-a population-based study

of more than 45 000 Danish women and men 18-45 years of age.

Pharmacoeconom Dr S 2010;19(1):65-74.

[45] Henderson EM, Keogh E, Rosser BA, Eccleston C. Searching the Internet for help with pain: adolescent search, coping, and medication behaviour. Br J Health Psychol 2013;18(1):218-232.

[46] Himmelstein MS, Sanchez DT. Masculinity impediments: internalized masculinity contributes to healthcare avoidance in men and women. J Health Psychol 2014.

[47] Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: a meta-analytic review. Plos Med 2010;7(7): e1000316.

[48] Holt E, Joyce C, Dornelles A, Morisky D, Webber LS, Muntner P, Krousel-Wood M. Sex differences in barriers to antihypertensive medication adherence: findings from the cohort study of medication adherence among older adults. J Am Geriatr Soc 2013;61(4):558-564.

[49] Houle J, Mishara BL, Chagnon F. An empirical test of a mediation model of the impact of the traditional male gender role on suicidal behavior in men. J Affect Disorders 2008;107(1-3):37-43.

[50] Hoyt MA, Stanton AL, Irwin MR, Thomas KS. Cancer-related masculine threat, emotional approach coping, and physical functioning following treatment for prostate cancer. Health Psychol 2013;32(1):66-74.

[51] Hunt K, Adamson J, Hewitt C, Nazareth I. Do women consult more than men? A review of gender and consultation for back pain and headache. J Health Serv Res Po 2011;16(2):108-117.

- [52] Ilgen MA, Kleinberg F, Ignacio RV, Bohnert ASB, Valenstein M, McCarthy JF, Blow FC, Katz IR. Noncancer pain conditions and risk of suicide. *JAMA Psychiat* 2013;70(7):692-697.
- [53] Jackson T, Wang YL, Wang Y, Fan HY. Self-efficacy and chronic pain outcomes: a meta-analytic review. *J Pain* 2014;15(8):800-814.
- [54] Jones CM, Paulozzi LJ, Mack KA, Centers for Disease C, Prevention. Alcohol involvement in opioid pain reliever and benzodiazepine drug abuse-related emergency department visits and drug-related deaths - United States, 2010. *MMWR Morbidity and mortality weekly report* 2014;63(40):881-885.
- [55] Kaur S, Stechuchak KM, Coffman CJ, Allen KD, Bastian LA. Gender differences in health care utilization among veterans with chronic pain. *J Gen Intern Med* 2007;22(2):228-233.
- [56] Kelly DM, Jones TH. Testosterone: a metabolic hormone in health and disease. *J Endocrinol* 2013;217(3):R25-R45.
- [57] Kelly DM, Jones TH. Testosterone: a vascular hormone in health and disease. *J Endocrinol* 2013;217(3):R47-R71.
- [58] Keogh E. Sex differences in pain across the life course. In: RA Moore, editor. *Handbook of pain and palliative care: biobehavioral approaches for the life course*. New York: Springer, 2012. pp. 347-366.
- [59] Keogh E. Gender differences in the nonverbal communication of pain: a new direction for sex, gender, and pain research? *Pain* 2014;155(10):1927-1931.
- [60] Keogh E, Denford S. Sex differences in perceptions of pain coping strategy usage. *Eur J Pain* 2009;13(6):629-634.
- [61] Keogh E, Eccleston C. Sex differences in adolescent chronic pain and pain-related coping. *Pain* 2006;123(3):275-284.

- [62] Keogh E, Herdenfeldt M. Gender, coping and the perception of pain. *Pain* 2002;97(3):195-201.
- [63] LeResche L. Epidemiology of pain conditions with higher prevalence in women. In: C Chin, RB Fillingim, TJ Ness, editors. *Pain in women*. Oxford: Oxford University Press, 2013. pp. 3-15.
- [64] Levant RF. Research in the psychology of men and masculinity using the gender role strain paradigm as a framework. *Am Psychol* 2011;66(8):765-776.
- [65] Lipsky MS, Cannon M, Lutfiyya MN. Gender and health disparities: the case of male gender. *Dis Mon* 2014;60(4):138-144.
- [66] Loefer M, Walach H. The combined effects of healthy lifestyle behaviors on all cause mortality: a systematic review and meta-analysis. *Prev Med* 2012;55(3):163-170.
- [67] Luy M, Minagawa Y. Gender gaps - life expectancy and proportion of life in poor health. *Health Rep* 2014;25(12):12-19.
- [68] Mackenzie CS, Gekoski WL, Knox VJ. Age, gender, and the underutilization of mental health services: the influence of help-seeking attitudes. *Aging Ment Health* 2006;10(6):574-582.
- [69] Mahalik JR, Burns SM, Syzdek M. Masculinity and perceived normative health behaviors as predictors of men's health behaviors. *Soc Sci Med* 2007;64(11):2201-2209.
- [70] Manierre MJ. Gaps in knowledge: tracking and explaining gender differences in health information seeking. *Soc Sci Med* 2015;128:151-158.
- [71] Manteuffel M, Williams S, Chen W, Verbrugge RR, Pittman DG, Steinkellner A. Influence of patient sex and gender on medication use, adherence, and prescribing alignment with guidelines. *J Womens Health* 2014;23(2):112-119.

- [72] Martin CL, Ruble DN. Patterns of gender development. *Annu Rev Psychol* 2010;61:353-381.
- [73] Moller-Leimkuhler AM. The gender gap in suicide and premature death or: why are men so vulnerable? *Eur Arch Psy Clin N* 2003;253(1):1-8.
- [74] Moore DJ, Eccleston C, Keogh E. Does sex moderate the relationship between anxiety and pain? *Psychol Health* 2013;28(7):746-764.
- [75] Murphy SL, Xu J, Kochanek KD. Deaths: final data for 2010. *Natl Vital Stat Rep* 2013;61(4):1-117.
- [76] Myers CD, Tsao JCI, Glover DA, Kim SC, Turk N, Zeltzer LK. Sex, gender, and age: contributions to laboratory pain responding in children and adolescents. *J Pain* 2006;7(8):556-564.
- [77] Niesters M, Dahan A, Kest B, Zacny J, Stijnen T, Aarts L, Sarton E. Do sex differences exist in opioid analgesia? A systematic review and meta-analysis of human experimental and clinical studies. *Pain* 2010.
- [78] Office of National Statistics. Suicides in the United Kingdom, 2013 registrations 2015.
- [79] Oksuzyan A, Juel K, Vaupel JW, Christensen K. Men: good health and high mortality. Sex differences in health and aging. *Aging Clin Exp Res* 2008;20(2):91-102.
- [80] Oliffe JL, Han CSE, Ogrodniczuk JS, Phillips JC, Roy P. Suicide from the perspectives of older men who experience depression: a gender analysis. *Am J Mens Health* 2011;5(5):444-454.
- [81] Pakpour AH, Nikoobakht M, Campbell P. Association of pain and depression in those with chronic low back pain the mediation effect of patient sexual functioning. *Clin J Pain* 2015;31(1):44-51.

- [82] Parks SE, Johnson LL, McDaniel DD, Gladden M, Centers for Disease C, Prevention. Surveillance for violent deaths - National Violent Death Reporting System, 16 states, 2010. *MMWR Surveill Summ* 2014;63(1):1-33.
- [83] Paulose-Ram R, Hirsch R, Dillon C, Losonczy K, Cooper M, Ostchega Y. Prescription and non-prescription analgesic use among the US adult population: Results from the third National Health and Nutrition Examination Survey (NHANES III). *Pharmacoepidemiol Drug Saf* 2003;12(4):315-326.
- [84] Paulozzi LJ. Prescription drug overdoses: a review. *J Safety Res* 2012;43(4):283-289.
- [85] Paulozzi LJ, Kilbourne EM, Shah NG, Nolte KB, Desai HA, Landen MG, Harvey W, Loring LD. A history of being prescribed controlled substances and risk of drug overdose death. *Pain Med* 2012;13(1):87-95.
- [86] Paulson M, Danielson E, Norberg A. Nurses' and physicians' narratives about long-term non-malignant pain among men. *J Adv Nurs* 1999;30(5):1097-1105.
- [87] Pinkhasov RM, Wong J, Kashanian J, Lee M, Samadi DB, Pinkhasov MM, Shabsigh R. Are men shortchanged on health? Perspective on health care utilization and health risk behavior in men and women in the United States. *Int J Clin Pract* 2010;64(4):475-487.
- [88] Pronina I, Rule NO. Inducing bias modulates sensitivity to nonverbal cues of others' pain. *Eur J Pain* 2014;18(10):1452-1457.
- [89] Puhse G, Wachsmuth JU, Kemper S, Husstedt IW, Evers S, Kliesch S. Chronic pain has a negative impact on sexuality in testis cancer survivors. *J Androl* 2012;33(5):886-893.

- [90] Racine M, Castarlenas E, de la Vega R, Tome-Pires C, Sole E, Miro J, Jensen MP, Moulin DE, Nielson WR. Sex differences in psychological response to pain in patients with fibromyalgia syndrome. *Clin J Pain* 2015;31(5):425-432.
- [91] Ramirez-Maestre C, Esteve R. The role of sex/gender in the experience of pain: resilience, fear, and acceptance as central variables in the adjustment of men and women with chronic pain. *J Pain* 2014;15(6):608-618.
- [92] Read JG, Gorman BK. Gender and health inequality. *Annu Rev Sociol* 2010;36:371-386.
- [93] Richardson N, Smith JA. National men's health policies in Ireland and Australia: what are the challenges associated with transitioning from development to implementation? *Public Health* 2011;125(7):424-432.
- [94] Riley JL, King C. Self-report of alcohol use for pain in a multi-ethnic community sample. *J Pain* 2009;10(9):944-952.
- [95] Robinson ME, Riley JL, Myers CD, Papas RK, Wise EA, Waxenberg LB, Fillingim RB. Gender role expectations of pain: relationship to sex differences in pain. *J Pain* 2001;2(5):251-257.
- [96] Ryu SY, Park J, Choi SW, Han MA. Associations between socio-demographic characteristics and healthy lifestyles in Korean Adults: the result of the 2010 Community Health Survey. *J Prev Med Public Health* 2014;47(2):113-123.
- [97] Shye D, Mullooly JP, Freeborn DK, Pope CR. Gender differences in the relationship between social network support and mortality - a longitudinal study of an elderly cohort. *Soc Sci Med* 1995;41(7):935-947.
- [98] Sloan C, Conner M, Gough B. How does masculinity impact on health? A quantitative study of masculinity and health behavior in a sample of UK men and women. *Psychol Men Masc* 2015;16(2):206-217.

- [99] Smith D, Wilkie R, Uthman O, Jordan JL, McBeth J. Chronic pain and mortality: a systematic review. *Plos One* 2014;9(6).
- [100] Tang NKY, Crane C. Suicidality in chronic pain: a review of the prevalence, risk factors and psychological links. *Psychol Med* 2006;36(5):575-586.
- [101] Tannenbaum C, Frank B. Masculinity and health in late life men. *Am J Mens Health* 2011;5(3):243-254.
- [102] Thibodeau MA, Welch PG, Katz J, Asmundson GJG. Pain-related anxiety influences pain perception differently in men and women: a quantitative sensory test across thermal pain modalities. *Pain* 2013;154(3):419-426.
- [103] Thompson EH, Bennett KM. Measurement of masculinity ideologies: a (critical) review. *Psychol Men Masculin* 2015;16(2):115-133.
- [104] Tobin DD, Menon M, Menon M, Spatta BC, Hodges EVE, Perry DG. The intrapsychics of gender: a model of self-socialization. *Psychol Rev* 2010;117(2):601-622.
- [105] Torrance N, Elliott AM, Lee AJ, Smith BH. Severe chronic pain is associated with increased 10 year mortality. A cohort record linkage study. *Eur J Pain* 2010;14(4):380-386.
- [106] Trudeau KJ, Danoff-Burg S, Revenson TA, Paget SA. Agency and communion in people with rheumatoid arthritis. *Sex Roles* 2003;49(7-8):303-311.
- [107] Unruh A, Ritchie J, Merskey H. Does gender affect appraisal of pain and pain coping strategies? *Clin J Pain* 1999;15(1):31-40.
- [108] Vierhaus M, Lohaus A, Schmitz AK. Sex, gender, coping, and self-efficacy: mediation of sex differences in pain perception in children and adolescents. *Eur J Pain* 2011;15(6):621.e1-621.e8.

- [109] Vlaeyen JWS, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. 2000;85(3):317-332.
- [110] Wagner FA, Anthony JC. Male-female differences in the risk of progression from first use to dependence upon cannabis, cocaine, and alcohol. *Drug Alcohol Depen* 2007;86(2-3):191-198.
- [111] Wang YY, Hunt K, Nazareth I, Freemantle N, Petersen I. Do men consult less than women? An analysis of routinely collected UK general practice data. *BMJ Open* 2013;3(8).
- [112] White A, De Sousa B, De Visser R, Hogston R, Madsen SA, Makara P, McKee M, Raine G, Richardson N, Clarke N, Zatonski W. Men's health in Europe. *J Mens Health* 2011;8(3):192-201.
- [113] White A, McKee M, de Sousa B, de Visser R, Hogston R, Madsen SA, Makara P, Richardson N, Zatonski W, Raine G. An examination of the association between premature mortality and life expectancy among men in Europe. *Eur J Public Health* 2014;24(4):673-679.
- [114] Wilson NJ, Cordier R. A narrative review of Men's Sheds literature: reducing social isolation and promoting men's health and well-being. *Health Soc Care Comm* 2013;21(5):451-463.
- [115] Wise EA, Price DD, Myers CD, Heft MW, Robinson ME. Gender role expectations of pain: Relationship to experimental pain perception. *Pain* 2002;96(3):335-342.
- [116] Yousaf O, Grunfeld EA, Hunter MS. A systematic review of the factors associated with delays in medical and psychological help-seeking among men. *Health Psychol Rev* in press.

- [117] Zaider T, Manne S, Nelson C, Mulhall J, Kissane D. Loss of masculine identity, marital affection, and sexual bother in men with localized prostate cancer. *J Sex Med* 2012;9(10):2724-2732.
- [118] Zhu K, Devine A, Dick IM, Prince RL. Association of back pain frequency with mortality, coronary heart events, mobility, and quality of life in elderly women. *Spine* 2007;32(18):2012-2018.

Figure 1: Possible relationship between the key areas covered within the review.