Citation for published version:

Gonzalez, JT 2016, 'Appetite effect of breakfast omission - constrained to morning?', *Nutrition*, vol. 32, no. 2, pp. 287-288. <https://doi.org/10.1016/j.nut.2015.08.003>

DOI:

[10.1016/j.nut.2015.08.003](https://doi.org/10.1016/j.nut.2015.08.003)

Publication date:

2016

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Title: Appetite effects of breakfast – restrained to the morning?

Author: Javier T. Gonzalez, BSc., MRes., PhD.

Email: J.T.Gonzalez@Bath.ac.uk

Author list for indexing: Gonzalez

Author disclosure: JT Gonzalez has no conflicts of interest.

Word count: 457 words.

Address for correspondence:

Department for Health,
University of Bath,
Bath,
BA2 7AY,
UK.

Tel: 0(+44)1225 38 5518

Recommendations to regularly consume breakfast in order to control or lose body weight are prevalent in guidelines from charities [1] and public health authorities [2, 3]. Yet, when the evidence base is scrutinized, the vast majority of research is observational in nature [4]. Whilst observational studies are useful for establishing a potential link between a behavior and an outcome, randomized controlled trials are essential in order to make definitive causal statements. More recent studies using free-living designs over periods of weeks and months [5-8], combined with evidence from more tightly-controlled laboratory studies conducted within a single day [9-11] are beginning to address the causality of breakfast consumption, body weight regulation and health, whilst providing mechanistic insight into any causal effects.

When attempting to control or lose body weight, exercise can be a powerful tool to improve body composition. Exercise also potentially improves metabolic health, even when the energy deficit that exercise usually creates is removed [12]. As exercise drastically alters the metabolic responses to breakfast consumption [9, 13], it is important to study the effects of breakfast consumption/omission in the context of exercise.

In this edition of *Nutrition*, Clayton *et al.* [14] add to this growing body of literature by examining the subjective appetite, metabolic and hormonal responses to breakfast consumption compared to extended overnight fasting, with the particularly novel component being an exercise bout performed in the afternoon (as opposed to the morning), between a standardized lunch and dinner. This design not only allows the findings to be generalized to those who perform exercise towards the end of the day, but also provides information as to whether breakfast consumption results in any carry-over effects on exercise metabolism, even after lunch has been consumed [15]. Subjects reported feeling less hungry after breakfast consumption (compared to extended fasting), up until the standardized lunch was consumed. However, no significant differences were found in hunger (nor in any other subjective

appetite ratings) for the rest of the day, suggesting that any effects of breakfast consumption on subjective appetite ratings are transient in nature and do not carry-over after a standard lunch is consumed. Similarly, the metabolic and hormonal responses were not significantly different following lunch, although a higher frequency of sampling may be needed to detect some metabolic responses [9, 13].

By fixing energy intake at lunch and dinner, information about within-day compensation cannot be determined, however, it can be argued that when individuals are attempting to control or lose body weight, then *ad libitum* feeding is not necessarily the default situation, and energy intake is in many cases, prescribed to some extent. This study provides information about appetite responses under prescribed feeding and exercise, a crucial component to study given the benefits of exercise for body composition and health.

Acknowledgements

The author has sole responsibility for all aspects of this manuscript. J.T.G is currently receiving funding from the European Society for Clinical Nutrition and Metabolism, and the Rank Prize Funds. The author declares no conflicts of interest in relation to this manuscript.

References

1. British Heart Foundation: **Facts not fads - Your simple guide to healthy weight loss**. vol. M2. pp. 36-37 2015:36-37.
2. **Why does eating a healthy breakfast help control weight?** [<http://www.mayoclinic.org/food-and-nutrition/expert-answers/FAQ-20058449>]
3. National Institute for Health and Care Excellence: **Obesity guidance on the prevention, assessment and management of overweight and obesity in adults and children (NICE clinical guideline 43)**. (Agency HD ed. London 2006.
4. Brown AW, Bohan Brown MM, Allison DB: **Belief beyond the evidence: using the proposed effect of breakfast on obesity to show 2 practices that distort scientific evidence**. *Am J Clin Nutr* 2013, **98**:1298-1308.
5. Betts JA, Richardson JD, Chowdhury EA, Holman GD, Tsintzas K, Thompson D: **The causal role of breakfast in energy balance and health: a randomized controlled trial in lean adults**. *Am J Clin Nutr* 2014, **100**:539-547.
6. Dhurandhar EJ, Dawson J, Alcorn A, Larsen LH, Thomas EA, Cardel M, Bourland AC, Astrup A, St-Onge MP, Hill JO, et al: **The effectiveness of breakfast recommendations on weight loss: a randomized controlled trial**. *Am J Clin Nutr* 2014, **100**:507-513.
7. Geliebter A, Astbury NM, Aviram-Friedman R, Yahav E, Hashim S: **Skipping breakfast leads to weight loss but also elevated cholesterol compared with consuming daily breakfasts of oat porridge or frosted cornflakes in overweight individuals: a randomised controlled trial**. *J Nutr Sci* 2014, **3**:e56.

8. Farshchi HR, Taylor MA, Macdonald IA: **Deleterious effects of omitting breakfast on insulin sensitivity and fasting lipid profiles in healthy lean women.** *Am J Clin Nutr* 2005, **81**:388-396.
9. Gonzalez JT, Veasey RC, Rumbold PL, Stevenson EJ: **Breakfast and exercise contingently affect postprandial metabolism and energy balance in physically active males.** *Br J Nutr* 2013, **110**:721-732.
10. Chowdhury EA, Richardson JD, Tsintzas K, Thompson D, Betts JA: **Carbohydrate-rich breakfast attenuates glycaemic, insulinaemic and ghrelin response to ad libitum lunch relative to morning fasting in lean adults.** *Br J Nutr* 2015:1-10.
11. Astbury NM, Taylor MA, Macdonald IA: **Breakfast consumption affects appetite, energy intake, and the metabolic and endocrine responses to foods consumed later in the day in male habitual breakfast eaters.** *J Nutr* 2011, **141**:1381-1389.
12. Walhin JP, Richardson JD, Betts JA, Thompson D: **Exercise counteracts the effects of short-term overfeeding and reduced physical activity independent of energy imbalance in healthy young men.** *J Physiol* 2013, **591**:6231-6243.
13. Gonzalez JT: **Paradoxical second-meal phenomenon in the acute post-exercise period.** *Nutrition* 2014, **30**:961-967.
14. Clayton DJ, Stensel DJ, James LJ: **Effect of breakfast omission on subjective appetite, metabolism, acylated ghrelin and GLP-1(7-36) during rest and exercise.** *Nutrition* Ahead of print.
15. Clayton DJ, Barutcu A, Machin C, Stensel DJ, James LJ: **Effect of Breakfast Omission on Energy Intake and Evening Exercise Performance.** *Med Sci Sports Exerc* 2015.