Citation for published version:

Ross, NP, Hamann, F, Zakamska, NL, Richards, GT, Villforth, C, Strauss, MA, Greene, JE, Alexandroff, R, Brandt, WN, Liu, G, Myers, AD, Paris, I & Schneider, DP 2015, 'Extremely red quasars from SDSS, BOSS and WISE: classification of optical spectra', *Monthly Notices of the Royal Astronomical Society*, vol. 453, no. 4, pp. 3932-3952. <https://doi.org/10.1093/mnras/stv1710>

DOI:

[10.1093/mnras/stv1710](https://doi.org/10.1093/mnras/stv1710)

Publication date:

2015

Document Version

Early version, also known as pre-print

[Link to publication](#)

This article has been accepted for publication in *Monthly Notices of the Royal Astronomical Society*, Published by Oxford University Press on behalf of the Royal Astronomical Society.

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Extremely Red Quasars from SDSS, BOSS and WISE: Classification of Optical Spectra

Nicholas P. Ross^{1,2*}, Fred Hamann³, Nadia L. Zakamska⁴, Gordon T. Richards^{1,5},
Carolyn Villforth^{3,6}, Michael A. Strauss⁷, Jenny E. Greene^{7,8}, Rachael Alexandroff⁴,
W. Niel Brandt^{9,10}, Guilin Liu⁴, Adam D. Myers^{11,5}, Isabelle Pâris¹²,
Donald P. Schneider^{9,10}

¹*Department of Physics, Drexel University, 3141 Chestnut Street, Philadelphia, PA 19104, USA*

²*Institute for Astronomy, SUPA†, University of Edinburgh, Royal Observatory, Edinburgh, EH9 3HJ, United Kingdom*

³*Department of Astronomy, University of Florida, Gainesville, FL 32611-2055, USA*

⁴*Department of Physics & Astronomy, Johns Hopkins University, 3400 N. Charles St., Baltimore, MD 21218, USA*

⁵*Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany*

⁶*SUPA, School of Physics and Astronomy, University of St Andrews, North Haugh, St Andrews, KY16 9SS, UK*

⁷*Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA*

⁸*Alfred P. Sloan Fellow*

⁹*Department of Astronomy and Astrophysics, The Pennsylvania State University, University Park, PA 16802, USA*

¹⁰*Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, PA 16802, USA*

¹¹*Department of Physics and Astronomy, University of Wyoming, Laramie WY 82071, USA*

¹²*INAF - Osservatorio Astronomico di Trieste, Via G. B. Tiepolo 11, I-34131 Trieste, Italy*

24 July 2015

ABSTRACT

Quasars with extremely red infrared-to-optical colours are an interesting population that can test ideas about quasar evolution as well as orientation, obscuration and geometric effects in the so-called AGN unified model. To identify such a population we match the quasar catalogues of the Sloan Digital Sky Survey (SDSS), the Baryon Oscillation Spectroscopic Survey (BOSS) to the Wide-Field Infrared Survey Explorer (WISE) to identify quasars with extremely high infrared-to-optical ratios. We identify 65 objects with $r_{\text{AB}} - W4_{\text{Vega}} > 14$ mag (i.e., $F_{\nu}(22\mu\text{m})/F_{\nu}(r) \gtrsim 1000$). This sample spans a redshift range of $0.28 < z < 4.36$ and has a bimodal distribution, with peaks at $z \sim 0.8$ and $z \sim 2.5$. It includes three $z > 2.6$ objects that are detected in the $W4$ -band but not $W1$ or $W2$ (i.e., “ $W1W2$ -dropouts”). The SDSS/BOSS spectra show that the majority of the objects are reddened Type 1 quasars, Type 2 quasars (both at low and high redshift) or objects with deep low-ionization broad absorption lines (BALs) that suppress the observed r -band flux. In addition, we identify a class of Type 1 permitted broad-emission line objects at $z \simeq 2 - 3$ which are characterized by emission line rest-frame equivalent widths (REWs) of $\gtrsim 150\text{\AA}$, much larger than those of typical quasars. In particular, 55% (45%) of the non-BAL Type 1s with measurable CIV in our sample have $\text{REW}(\text{CIV}) > 100$ (150) $\text{\AA}$, compared to only 5.8% (1.3%) for non-BAL quasars in BOSS. These objects often also have unusual line ratios, such as very high $\text{N V}/\text{Ly}\alpha$ ratios. These large REWs might be caused by suppressed continuum emission analogous to Type 2 quasars; however, there is no obvious mechanism in standard Unified Models to suppress the continuum without also obscuring the broad emission lines.

Key words: Astronomical data bases: surveys – Quasars: general – galaxies: evolution – galaxies: infrared.

1 INTRODUCTION

Quasars are the most luminous non-transient objects in the Universe, with L_{bol} reaching $\sim 10^{47-48}$ erg s^{−1}, powered by

* email: npross@physics.drexel.edu

accretion of matter onto supermassive black holes. To explain how quasars are fueled and become active, models (e.g., Sanders et al. 1988; Hopkins et al. 2005, 2006) suggest that mergers between gas-rich galaxies produce starbursts and drive gas to the inner galactic nuclear regions. This activity fuels the growth of the supermassive black holes, which actively accrete matter at this stage, with much of the black hole growth happening in this ultra-violet (UV) and optically obscured phase. Feedback, potentially from a radiation-driven quasar wind, leads to removal of gas and dust from the central region, allowing the quasar to be seen in the UV and optical. These major-merger models are invoked to explain the obscuration/reddening observations noted in e.g., Urrutia et al. (2008), Glikman et al. (2012), and predict that red/obscured quasars should appear preferentially during an early stage of quasar evolution. However, the exact nature of the AGN triggering mechanism is not straightforward, as observations find that levels of disturbance in AGN hosts are consistent with those of inactive control samples (e.g. Kocevski et al. 2012; Villforth et al. 2014), or only slightly elevated at low- z (Rosario et al. 2015).

Quasars appear with a wide range of observational properties. Intrinsically luminous quasars may or may not appear bright at optical, ultra-violet and X-ray wavelengths. Much of this variance can be explained in the context of the geometry-based ‘unification models’ (Antonucci 1993). If gas and dust are present near the active nucleus but do not completely surround it, then some lines of sight to the nucleus are clear, whereas others are blocked. In the former case, the observer can directly view the emission from the accretion disk, with the quasar appearing bright at X-ray, ultra-violet and optical wavelengths; these objects are termed unobscured active galactic nuclei. Often these objects display broad (several thousand km s^{-1}) emission lines in the optical spectra and are classified as ‘Type 1’ quasars (Khachikian & Weedman 1974). Less than complete blockage of the quasar continuum by dust will redden the spectrum (e.g., Richards et al. 2003; Krawczyk et al. 2015).

This contrasts with the situation where the observer’s line of sight is blocked by circumnuclear clouds; in this case X-rays are absorbed by the intervening gas and ultra-violet and optical photons are scattered and absorbed by the intervening dust. At optical wavelengths, often the only signature of nuclear activity in this case is strong, narrow emission lines produced in the material illuminated by the quasar along unobscured directions. These objects with weak ultra-violet and optical continua are obscured AGN, and if narrow UV/optical emission lines are present, are designated Type 2 sources (Antonucci & Miller 1985; Smith et al. 2002; Zakamska et al. 2003; Brandt & Hasinger 2005; Reyes et al. 2008). Thus, obscuration, as part of an evolutionary phase or an orientation effect, is critical in determining the observed properties of quasars. However, because of their faintness at rest-frame optical and ultra-violet wavelengths, identifying obscured objects, at the peak of quasar activity $z \sim 2$, remains challenging (Stern et al. 2002; Norman et al. 2002; Alexandroff et al. 2013; Greene et al. 2014).

The circumnuclear clouds of gas and dust absorb X-ray, ultra-violet and optical radiation from the quasar and re-emit this energy thermally at infrared wavelengths. This is why unobscured quasars have similar luminosities at

near- and mid-infrared wavelengths ($\sim 2 - 30\mu\text{m}$ in the rest-frame) as they do in the optical and in the ultra-violet (Elvis et al. 1994; Richards et al. 2006; Polletta et al. 2008; Elvis 2010). Extremely obscured and dusty objects, in which optical emission is partially extinguished or completely blocked, are therefore expected to show much higher infrared-to-optical ratios than unobscured quasars. Indeed, a host of previous studies have used a K -band excess selection (Chiu et al. 2007; Maddox et al. 2008; Jurek et al. 2008; Nakos et al. 2009; Souchay et al. 2009; Wu & Jia 2010; Peth et al. 2011; Wu et al. 2011; Maddox et al. 2012; Fynbo et al. 2013; Wu et al. 2013), a near-infrared+radio selection (Glikman et al. 2004, 2007, 2012, 2013) or a mid-infrared selection (Lacy et al. 2004; Stern et al. 2005; Martínez-Sansigre et al. 2006; Richards et al. 2009; Donley et al. 2012; Stern et al. 2012; Banerji et al. 2013; Assef et al. 2013) to identify obscured (as well as unobscured) AGN.

In this paper, we describe a search for quasars with extremely red colours in order to study their range of spectral properties. We use optical photometry and spectroscopy from the Sloan Digital Sky Survey (SDSS; York et al. 2000) and the SDSS-III (Eisenstein et al. 2011) Baryon Oscillation Spectroscopic Survey (BOSS; Dawson et al. 2013) as well as mid-infrared photometry from the Wide-Field Infrared Survey Explorer (WISE; Wright et al. 2010). A companion paper (Hamann et al. 2015, in prep.) presents a detailed investigation into a new class of objects, the extreme rest-frame equivalent width quasars, which are introduced here.

This paper is organized as follows. In Section 2, we describe our datasets and sample selection, and in Section 3 we present the basis sample properties of the extremely red quasars. In Section 4 we discuss the optical spectra of the extremely red quasars, classifying objects along the traditional lines of broad-line Type 1s, narrow-line Type 2s and those with interesting absorption features. In Section 4.7, we introduce the new class of extreme equivalent width quasars. In Section 5 we discuss the selection of the extremely red quasars and place these objects in a broader physical and evolutionary context. We conclude in Section 6.

2 DATA AND SAMPLE SELECTION

2.1 Parent dataset

Our starting point is the spectroscopic quasar catalogues of the SDSS Seventh Data Release (DR7Q; Schneider et al. 2010; Shen et al. 2011) and the SDSS-III Tenth Data Release (DR10Q; Pâris et al. 2014). SDSS quasar targets with $i \leq 19.1$ were selected if the colours were consistent with being at redshift $z \lesssim 3$, and $i \leq 20.2$ objects were selected if $z \gtrsim 3$, as outlined in Richards et al. (2002). BOSS quasar targets are selected to a magnitude limit of $g \leq 22.0$ or $r \leq 21.85$, with the primary goal to select quasars in the redshift range $2.2 \leq z \leq 3.5$ as described by Ross et al. (2012, and references therein). In both SDSS and BOSS, quasar targets are also selected if they are matched within $2''$ ($1''$ in the case of BOSS) of an object in the Faint Images of the Radio Sky at Twenty-cm (FIRST) catalogue of radio sources (Becker et al. 1995). Both the SDSS DR7Q and BOSS DR10Q include quasars that were selected by algorithms

Figure 1. *Top:* $r - W4$ colour of the $W4$ -detected SDSS/BOSS quasars (black dots) as a function of $W4$ (i.e., $22\mu\text{m}$) magnitude. The quasars with $r - W4 > 14$ are shown as large red circles. *Bottom:* The r_{AB} -band magnitude versus $W4$. As a guide, $W4_{\text{Vega}} = 8.0$ mag corresponds to $W4_{\text{AB}} = 14.6$, or a flux density of $F_{\nu} \approx 6\text{mJy}$ (Wright et al. 2010).

other than the main quasar selections; these sources appear in the catalogue due to being targeted by the respective galaxy selections, being a ‘serendipitous’ (Stoughton et al. 2002) or ‘special’ (Adelman-McCarthy et al. 2006) target in SDSS, or an ‘ancillary’ target in BOSS (Dawson et al. 2013; Alam et al. 2015).

Luminous distant quasars are expected to be unresolved in optical ground-based observations, so we use point-spread function (PSF) magnitudes reported in the SDSS DR7 and BOSS DR10 quasar catalogues, corrected for Galactic extinction (Schlegel et al. 1998). Using the 2.5m Sloan telescope (Gunn et al. 2006), imaging data (Gunn et al. 1998) and spectra were obtained with the double-armed SDSS/BOSS spectrographs which have $R = \lambda/\text{FWHM} \sim 2000$ (Smee et al. 2013). Redshifts are measured from the spectra using the methods described in Bolton et al. (2012) and the data products are detailed in the SDSS DR7 (Abazajian et al. 2009) and DR10 (Ahn et al. 2014) papers.

The SDSS DR7 and BOSS DR10 quasar catalogues include 105,783 objects across 9380 deg^2 , and 166,583 objects across $6,370\text{ deg}^2$, respectively, with 16,420 objects in common to both catalogues. Thus our superset of optical data has 255,946 objects, and we use the BOSS spectrum (which has higher S/N and a larger wavelength coverage) when there are duplicate spectra between the two catalogues.

WISE mapped the sky in four filters centered at 3.4, 4.6, 12, and $22\mu\text{m}$ ($W1$, $W2$, $W3$, and $W4$ bands), achieving 5σ point-source sensitivities better than 0.08, 0.11, 1, and 6 mJy, respectively. The WISE Explanatory Supplement¹ provides further details about the astrometry and photometry

in the source catalogue. We retrieved photometric quantities for each of the four WISE bands. Hereafter, we use the notation SNR_{Wx} for the signal-to-noise ratio and Wx for the Vega-based WISE magnitudes, in band $x = 1, 2, 3$ and 4.

Because of established conventions, we report Sloan Digital Sky Survey (SDSS) *ugriz* magnitudes on the AB zero-point system (Oke & Gunn 1983; Fukugita et al. 1996), while the Wide-Field Infrared Survey Explorer (WISE) $W1 - 4$ magnitudes are calibrated on the Vega system (Wright et al. 2010). For WISE bands, $m_{\text{AB}} = m_{\text{Vega}} + \Delta m$ where $\Delta m = (2.699, 3.339, 5.174, 6.620)$ for $W1$, $W2$, $W3$ and $W4$, respectively (Cutri et al. 2011). We make use of the Explanatory Supplement to the WISE All-Sky Data Release, as well as the WISE AllWISE Data Release Products online.

The WISE team has released two all sky catalogues: the 9 month cryogenic phase of the mission led to the WISE All-Sky (“AllSky”) Data Release; while the WISE AllWISE Data Release (“AllWISE”) combines the AllSky data with the NEOWISE program (Mainzer et al. 2011). The resulting AllWISE dataset is deeper in the two shorter WISE-bands (5σ point-source sensitivities now 0.054 and 0.071 mJy), and the data processing algorithms were improved in all four W -bands. Our analysis began with AllSky, moved to AllWISE when it became available, but we use both since we had already performed visual inspections using AllSky.

We match the sample of the 255,946 unique quasars in the SDSS DR7Q and BOSS DR10Q, and use both the AllSky and the AllWISE data available at the NASA/IPAC Infrared Science Archive (IRSA)², with a matching radius of $2''$. For the combined DR7Q+DR10Q, we find 203,680 matches; 102,083 objects (96.5%) of the DR7Q and 111,779 (67%) of the DR10Q are matched in one or more WISE bands. The difference in the percentage of matches is expected since BOSS is fainter than SDSS. Matching with DR10Q positions offset in Right Ascension and Declination by $6''$, $12''$ and $18''$ (~ 1 , 2 and 3 times the WISE angular resolution in $W1/2/3$) return, respectively, 508, 893 and 873 matches within $2''$, suggesting our false-positive matching rate is $\lesssim 1\%$ (see also Krawczyk et al. 2013).

With a matching radius of $2''$, 15,843 of the SDSS DR7 quasars, and 2,979 of the (optically fainter) BOSS DR10 quasars have a good match in the WISE $W4$ band, i.e., $\text{SNR}_{W4} \geq 3$, $W4 < 8.00$ (close to the nominal 5σ point source sensitivity of $W4 = 7.9$ mags, Wright et al. 2010) and the WISE contamination and confusion flag, `cc_flags` set to “0000”, suggesting the source is unaffected by known artifacts in all four bands. Table 1 presents an overview of the numbers of SDSS/BOSS quasars in the WISE-matched dataset.

2.2 Extremely red quasar criterion

Dey et al. (2008) discovered a population of $z \sim 2$ dust-obscured galaxies, revealed by the *Spitzer* Space Telescope. In addition to having AGN activity, these objects are strongly star forming and dusty which contributes to their red colours (Brand et al. 2008; Soifer et al. 2008; Pope et al. 2008; Busmann et al. 2009a,b; Melbourne et al.

¹ wise2.ipac.caltech.edu/docs/release/allwise/expsup/index.html

² http://irsa.ipac.caltech.edu/

Description	Unique objects	from SDSS	from BOSS
DR7Q and DR10Q quasars ^a	255 946	105 783	166 583
+ matched to WISE	203 680	102 083	111 779
+ SNR _{W4} ≥ 3	52 873	41 922	10 951
+ W4 − σ _{W4} ≤ 8.00	28 513	24 041	4 472
+ W4 ≤ 8.00	18 615	16 035	2 580
+ cc.flags ^b	17 744	15 300	2 444
and r − W4 > 13.0	286	104	182
and r − W4 > 13.5	136	34	102
and r − W4 > 14.0	65	19	48

Table 1. Overview of the numbers of SDSS/BOSS quasars in the WISE-matched dataset. The third row gives the number of quasars which satisfy $W4$ SNR ≥ 3; the fourth row is the number of quasars having a magnitude of $W4 - \sigma_{W4} \leq 8.00$, while the 5th row is the number having $W4 \leq 8.00$. Numbers of objects passing the $r_{AB} - W4_{\text{Vega}}$ colour cut is given. ^aThere are objects common to both SDSS DR7Q and BOSS DR10Q that have WISE matches. Hence, the sum of the “From SDSS” and “From BOSS” columns is greater than the given total. ^b“Good” $W4$ detections have SNR_{W4} ≥ 3 and $W4 < 8.00$ and cc.flags equal to “0000”.

2009, 2012). Motivated by this finding, we apply similar selection criteria to the SDSS/BOSS+WISE matched quasars; see Figure 1. At high redshift our WISE $W4$ i.e., $22\mu\text{m}$ -based selection, is sensitive to hot dust since $W4$ probes $\lambda \simeq 7.3\mu\text{m}$ at $z \approx 2$ (Draine & Li 2007; Elitzur 2008; Diamond-Stanic & Rieke 2010). This is of particular interest because at these redshifts shorter wavebands (e.g. $W1$) may cease to trace dust emission, i.e. $W1$ probes $\lambda \simeq 1\mu\text{m}$, where dust does not emit strongly.

In particular, we require the spectroscopic quasars to have a reliable detection in the WISE $W4$ -band:

$$\text{SNR}_{W4} \geq 3, \quad (1)$$

a colour of

$$r_{AB} - W4_{\text{Vega}} > 14.0 \quad (2)$$

i.e., $r_{AB} - W4_{AB} \gtrsim 7.5$, which corresponds to $F_{\nu}(22\mu\text{m})/F_{\nu}(r) \gtrsim 1000$, and

$$W4_{\text{Vega}} \leq 8.00. \quad (3)$$

We further include objects with $W4 > 8.00$ if the object has SNR_{W4} ≥ 3 and $W4 - \sigma_{W4} < 8.00$. In Table 1 we outline the stages in the selection and the resulting number of quasars after every step.

The $r - W4 > 14$ selection has been shown to select Type 2 quasars with $z \sim 2$ (Brand et al. 2006, 2007) and while both Dey et al. (2008) and our selection have a colour cut of $r_{AB} - W4_{\text{Vega}} > 14.0$, we note that Dey et al. (2008) went to much fainter magnitudes, basing their initial selection in the IR, matched to optical catalogues, and then obtained additional spectra of their candidates. Also, while Dey et al. (2008) start from the IR selection, we start with UV/visible selected quasars in SDSS/BOSS and then look at $W4$ to find the reddest cases.

Our selection applied to both the AllSky and AllWISE catalogues results in very similar samples, and any object that passes the criteria in equations (1)-(3) in either of these data releases is included. There are 65 quasars that have spectra from SDSS/BOSS and satisfy the extremely red criterion from the union between the two WISE datasets.

Figure 2. The minimal amount of dust extinction required to produce high infrared-to-optical ratios consistent with our selection criteria, as a function of the redshift of the quasar. The Richards et al. (2006) spectral energy distribution for bright quasars is assumed to represent an unextinguished mean quasar. We then apply varying amounts of SMC-like extinction (from Weingartner & Draine 2001) and convolve with the respective filter curves. We assume $R_V \equiv \frac{A_V}{E(B-V)} = 2.87$ (Gordon & Clayton 1998). The three curves show the A_V values required to generate colours of $r - W4 > 13$ (yellow) $r - W4 > 13.5$ (orange) and $r - W4 > 14$ (red).

The $W4$ PSF has a full width at half maximum (FWHM) of $12''$, much larger than the optical PSF, so a nearby galaxy and a quasar cannot be deblended in $W4$ observations if they are only a few arcseconds apart. To check this possibility, we visually inspected the SDSS and WISE images of the 65 objects using the SDSS Image List Tool and the IRSA WISE Image Service, and found that there is no other obvious optical source (down to the 5σ -depth of the SDSS photometric survey of $r = 22.7$) within $6''$ of any of our targets. In only two cases there is potential concern that another optical object boosts the $W4$ flux - we return to this in Section 4.4. We do note, however, that dusty star-forming galaxies (particularly at high redshift) are very faint optically, so there is still a possibility that there is an additional contaminating galaxy; inspecting the relatively shallow SDSS images only eliminates the possibility of optically bright contaminating galaxies.

Also, although they are bright in $W1$ and $W2$, brown dwarfs tend to be significantly fainter than 8th magnitude in $W4$ (Kirkpatrick et al. 2011), so we do not suspect contamination from these objects. Thus we consider all 65 $r - W4 > 14$ quasars to be good matches.

2.3 Effects of dust extinction

Before we discuss in detail the observed spectral properties of our sample of extremely red quasars, we describe our expectations for how much extinction would be required to redden the overall spectral energy distribution of a quasar in order for it to be picked up by our colour selection. We acknowledge that the extremely red colours of our objects may not be all due to dust, and thus this exercise is for general guidance.

We simulate the colours of reddened quasars at different redshifts using the Richards et al. (2006) spectral energy distribution of bright quasars, assuming that the spectral energy distributions (SEDs) of all unextincted quasars are the same. We then apply varying amounts of SMC-like extinction (parametrized by the amount of extinction in the V-band, A_V , in magnitudes), place this object at different redshifts and record its $r - W4$ colour.

The result is shown in Figure 2, where the amount of extinction as a function of redshift is shown to generate colours of $r - W4 > 13$, $r - W4 > 13.5$ and $r - W4 > 14$ (the yellow, orange and red curves, respectively). The amount of extinction is very sensitive to the redshift of the quasar, and here we are only concerned with the dust reddening of a Type 1 SED. If there is any optical emission that is not affected by the circumnuclear dust extinction – for example, the extended emission from the host galaxy – then it would make a reddened quasar appear less red in the $r - W4$ colour. Thus the values of A_V plotted in Figure 2 should be regarded as *lower* limits on the amount of quasar extinction to produce the requisite colours. At low redshifts, the observed r -band probes rest-frame optical emission of the quasar, so the amount of extinction necessary to produce a very red optical-to-infrared colour is considerable. However, at high redshifts (where BOSS observations probe the rest-frame ultra-violet emission), an A_V of just a few tenths is sufficient to strongly suppress the ultra-violet emission while having a small effect on the infrared; this produces a very red $r - W4$ colour. Vardanyan et al. (2014) discuss the rest $0.25\mu\text{m}/7.8\mu\text{m}$ luminosities and properties of $1.4 < z < 5$ quasars in the SDSS, and define ‘obscured’ quasars based on $\gtrsim 5$ mags of UV extinction; considerably larger than we consider for our extremely red quasars over the same redshift range.

Due to the shallow nature of the WISE $W4$ band, at high redshift, any $W4$ -detected galaxy that does not show nuclear activity will have $L_{\text{IR}} > 10^{12} L_{\odot}$ and star-formation rates $> 100 M_{\odot} \text{ yr}^{-1}$. Indeed, many of the non-AGN WISE $W4$ detections will have substantially higher infrared luminosities and SFRs.

We have also checked the $R - W4$ colours of galaxies from the Canada-France-Hawaii Telescope Legacy Survey³ and find that they again have a range of colours, $\sim 9 < R - W4 \lesssim 15$.

The optical colours of our sample are inconsistent with simple dust extinction of a standard quasar SED (e.g. Richards et al. 2003; Krawczyk et al. 2015) and this is a current area of investigation. We discuss possible solutions to this in our companion papers, Hamann et al. (2015,

in prep.) and Zakamska et al. (2015, in prep); in particular, contribution from scattered light is a possibility (e.g., Zakamska et al. 2005, 2006) and so is patchy obscuration Veilleux et al. (2013). In particular, we investigate further the nature of the obscuration from the SEDs, which is not consistent with simple reddening from our analysis. As such, we are very hesitant to draw any further conclusions on dust reddening and A_V values in this paper, and leave detailed discussions to Hamann et al. (2015, in prep).

³ <http://www.cfht.hawaii.edu/Science/CFHLS/>.

Table 2: The 65 quasars from SDSS and BOSS that have $r - W4 > 14$, in order of increasing redshift. The r -band AB magnitude is from either Schneider et al. (2010) or Pâris et al. (2014), corrected for Galactic extinction (Schlegel et al. 1998), while the WISE W4 Vega magnitude is from the AllWISE Data Release catalogue. The Radio fluxes are the integrated flux densities (F_{int}) measured in mJy at 20cm from the FIRST survey (Becker et al. 1995). The Rest-frame Equivalent Widths (REWs) in Ångströms are given for MgII at $z \leq 1.5$ and CIV for objects at redshift $z > 1.5$. The Full-Width, Half-Maximum (FWHM) of the relevant emission line is also quoted. Objects with similar optical spectral properties are grouped together, using the bitwise values described in Table 3, converted here to binary (with a space gap for easier inspection).

Name (SDSS J)	R.A. (J2000)	Decl. (J2000)	z	r -mag (AB)	W4 mag (Vega)	$r - W4$	Radio flux/mJy	REWs / Å	FWHM / km s ⁻¹	Notes	Spectral Classification
101034.27+372514.7	152.64283	37.42077	0.282	18.39±0.03	4.26±0.02	14.14	-	-	-	Type 1; Strong narrow [OII], starburst	0000 0101
0941000.8+143614.4	145.25339	14.60402	0.384	19.43±0.02	5.39±0.04	14.04	7.6	-	-	Type 1; Strong narrow [OII], starburst	0001 0110
105327.92+375804.2	163.36636	37.96786	0.448	19.64±0.03	5.59±0.04	14.05	2.8	-	-	Type 1; Weak [OII], strong [NeIII] and [OIII], post-starburst	0001 1010
112657.76+163912.0	171.74069	16.65334	0.464	19.20±0.02	4.79±0.03	14.41	2.4	-	-	Type 1; Strong narrow [OIII], post-starburst	0001 1010
102834.03-023659.6	157.14182	-2.61657	0.470	18.90±0.02	4.80±0.03	14.10	68.5	-	-	Type 1 based on MgII, blazar candidate	0001 0001
110550.54+112702.0	166.46059	11.45058	0.498	20.65±0.05	6.17±0.06	14.48	2.9	-	-	Potential Type 2 candidate, but broad H α	0001 0010
092501.78+274607.9	141.25744	27.76888	0.531	19.43±0.02	4.47±0.02	14.97	26.7	-	-	Type 2 in Reyes et al. (2008)	0001 0010
091157.55+014327.5	137.98980	1.72432	0.603	21.48±0.08	6.60±0.09	14.88	4.6	-	-	Type 2 in Reyes et al. (2008)	0001 0010
225612.16-010508.0	344.05071	-1.08556	0.651	21.53±0.07	7.16±0.13	14.37	3.0	27±8	1870±730	Type 2 in Reyes et al. (2008)	0001 0010
161558.92+252736.1	243.99552	25.46004	0.676	22.04±0.09	7.26±0.09	14.78	4.2	69±8	918±174	Type 2	0001 0010
111354.66+124439.0	168.47776	12.74418	0.680	19.86±0.03	5.20±0.04	14.66	4.2	-	-	Red Type 1 (broad H β , strong [OII], Fe II)	0001 0101
124836.10+424259.3	192.15043	42.71650	0.682	21.61±0.07	7.32±0.10	14.29	3.4	22±11	3103±1138	Type 2	0001 0010
101303.09+072731.9	153.26288	7.45887	0.697	20.99±0.05	6.91±0.10	14.08	-	35±5	1793±294	Type 2	0000 0010
154215.25+161648.9	235.56354	16.28028	0.714	21.70±0.06	7.54±0.10	14.17	1.5	23±6	1972±507	Type 2	0001 0010
144642.28+011303.0	221.67618	1.21751	0.726	21.02±0.05	6.08±0.05	14.94	5.8	-	-	Type 2 in Reyes et al. (2008)	0001 0010
005009.81-003900.6	12.54091	-0.65017	0.728	20.11±0.03	5.87±0.06	14.24	4.4	-	-	Type 2	0001 0010
142240.99+383710.4	215.67083	38.61957	0.758	20.57±0.03	6.53±0.05	14.04	9.8	-	-	Type 2 in Reyes et al. (2008)	0001 0010
140744.00+360109.5	211.93333	36.01931	0.783	21.29±0.05	6.85±0.07	14.44	5.3	112±5	(4000)	Type 1 with O[II]	0001 0101
124231.04+023307.0	190.62933	2.55195	0.785	21.16±0.05	6.81±0.08	14.35	-	-	-	Red Type 1	0000 0001
091501.71+241812.1	138.75714	24.30336	0.844	20.39±0.03	4.82±0.03	15.57	10.1	-	-	Type 1; Very bright in W4	0001 0001
152504.74+123401.7	231.26977	12.56715	0.851	21.14±0.04	6.92±0.07	14.22	-	43±3	1984±167	Type 2	0000 0010
151354.48+145125.2	228.47702	14.85702	0.881	21.00±0.04	6.37±0.05	14.63	7.6	20±2	2212±209	Type 2	0001 0010
144855.23+213622.0	222.23016	21.60612	0.882	21.70±0.07	7.72±0.12	13.98	5.2	101±17	7868±1347	Type 2	0001 0010
210712.77+005439.4	316.80322	0.91096	0.924	20.56±0.03	6.31±0.06	14.25	0.5	-	-	Type 1, strong FeII, LoBAL MgII absorption	0011 0001
075257.36+351834.6	118.23901	35.30962	0.926	23.23±0.30	7.57±0.19	15.66	-	52±4	2979±198	Type 1	0000 0001
141702.32+364839.5	214.25969	36.81098	1.004	21.58±0.06	6.89±0.06	14.69	7.8	181±33	21698±3680	Type 2	0001 0010
142131.51+210719.4	215.38131	21.12207	1.061	21.78±0.08	7.72±0.14	14.06	1.0	37±4	1425±137	Type 2	0001 0010
092356.70+230030.9	140.98627	23.00860	1.095	22.34±0.12	7.68±0.20	14.66	-	30±8	2112±410	Type 2, strong O[II]	0000 0110
091103.50+444630.3	137.76458	44.77511	1.302	20.88±0.06	6.58±0.06	14.30	-	-	-	Type 1; Strong FeII emission, LoBAL MgII absorption	0010 0001
111004.78+375236.7	167.51994	37.87686	1.308	20.34±0.04	5.61±0.04	14.74	7.1	-	-	Red Type 1	0001 0001
165258.51+390249.7	253.24380	39.04716	1.308	21.21±0.04	6.79±0.06	14.42	249.3	7±1	2966±304	Type 1 based on MgII, blazar candidate	0001 0001
102425.41+033239.8	156.10589	3.54441	1.477	22.40±0.17	7.11±0.17	15.29	-	35±8	1166±377	Type 2 based on v. narrow CIV	0000 0010
144929.61+394824.2	222.37341	39.80674	1.491	21.74±0.06	7.66±0.11	14.08	24.3	57±9	10075±1405	Type 2 based on v. narrow CIV and HeII	0001 0010
153542.41+090341.1	233.92671	9.06143	1.533	21.33±0.04	6.65±0.07	14.68	8.1	137±14	1556±140	Type 1; Very strong FeII emission in UV2/3, UV1, and strong AlIII with weak/absent CIII] and SiIII]; see Appendix A1.	0011 0001
080304.76+532627.0	120.76987	53.44084	1.573	20.85±0.05	6.60±0.07	14.26	2.9	123±7	3196±138	Type 1; LoBAL	0011 0001
122453.20+084303.2	186.22167	8.71757	1.960	23.55±0.73	8.01±0.25	15.54	-	0±0	0±0	Red Type 1	0000 0001
211329.61+001841.7	318.37340	0.31161	1.996	23.40±0.28	8.14±0.29	15.26	-	192±7	1536±49	Type 2, "Extreme" REW (EREW)	1000 0010
163313.29+401338.9	248.30541	40.22749	2.010	21.18±0.05	7.12±0.08	14.07	10.5	20±10	11936±99	Type 1; Blazar candidate	0001 0001
173049.10+585059.5	262.70459	58.84988	2.035	21.39±0.07	7.37±0.09	14.02	-	-	-	Type 1; FeLoBAL	0010 0001
133611.79+404522.9	204.04913	40.75637	2.071	21.07±0.06	6.38±0.05	14.69	3.1	20±3	6244±688	Type 1	0001 0001
013435.66-093102.9	23.64861	-9.51748	2.220	21.30±0.06	6.58±0.07	14.71	960	-	-	Gravitational Lens, Red Type 1, see Appendix A2	0001 0001
084447.66+462338.7	131.19862	46.39411	2.224	21.48±0.08	7.48±0.13	14.00	-	175±4	1744±34	Type 2, with narrow CIV	0000 0010
094317.59+541705.1	145.82333	54.28475	2.230	21.41±0.07	7.33±0.09	14.08	2.2	-	-	Type 1; FeLoBAL	0011 0001
222646.53+005211.2	336.69390	0.86980	2.247	21.54±0.07	7.54±0.15	14.00	653	65±4	2663±212	Red Type 1, strong associated absorbtion	0001 0001
000610.67+121501.2	1.54449	12.25035	2.309	22.18±0.13	6.56±0.07	15.61	-	108±6	4589±202	Type 1, EREW	1000 0001
232326.17-010033.1	350.85906	-1.00920	2.355	21.83±0.08	7.76±0.22	14.07	-	256±5	3977±62	Type 1, N v/Ly α > 1, EREW	1000 0001
123241.73+091209.3	188.17391	9.20261	2.385	21.09±0.05	6.78±0.09	14.31	-	230±3	4873±53	Type 1, N v/Ly α > 1, EREW	1000 0001
083413.90+511214.6	128.55793	51.20407	2.391	19.77±0.04	5.43±0.04	14.33	1.9	-	-	Type 1, red slope, z from absorption at red end	0001 0001

Name (SDSS J)	R.A. (J2000)	Decl. (J2000)	z	r-mag (AB)	W4 mag (Vega)	r - W4	Radio flux/mJy	REWs / Å	FWHM / km s ⁻¹	Notes	Spectral Classification
0832000.2+1615000.3	128.00085	16.25011	2.423	21.82±0.08	7.51±0.16	14.31	1.1	-	-	Type 1, EREW, strong associated absorption lines	1001 0001
011110.83+204543.8	17.79517	20.76218	2.432	22.03±0.09	7.96±0.21	14.06	-	0±0	0±0	Type 1; Ly-α emitter QSO	0000 0001
090630.73+082837.3	136.62806	8.47705	2.437	21.77±0.09	7.76±0.20	14.00	22.8	3±1	340±159	Type 1 (from Ly-α width)	0001 0001
093638.41+101930.3	144.16006	10.32510	2.458	21.72±0.06	7.61±0.17	14.11	-	169±4	1273±20	Type 2, but with EREW	1000 0010
134026.99+083427.2	205.11248	8.57424	2.490	22.31±0.13	8.10±0.21	14.21	-	338±11	2088±51	Type 2, but with EREW	1000 0010
221524.00-005643.8	333.85003	-0.94550	2.493	22.29±0.12	7.91±0.24	14.38	-	152±5	4226±112	Type 1; EREW	1000 0001
111017.13+193012.5	167.57139	19.50347	2.497	20.61±0.03	6.59±0.07	14.01	-	-	-	Type 1; EREW	1000 0010
155102.79+084401.1	237.76163	8.73366	2.520	20.66±0.03	6.61±0.06	14.05	2.9	10±3	2467±426	Type 1; FeLoBAL	0011 0001
083448.48+015921.1	128.70202	1.98921	2.594	21.19±0.05	6.88±0.09	14.31	-	214±7	2864±73	EREW, Type 1	1000 0001
220337.79+121955.3	330.90749	12.33204	2.626	21.57±0.05	7.45±0.14	14.12	-	262±3	1072±10	EREW, Type 2	1000 0010
085124.78+314855.7	132.85329	31.81548	2.638	21.63±0.07	6.79±0.08	14.84	-	76±5	3584±181	Type 1, W1W2-dropout	0100 0001
100424.88+122922.2	151.10370	12.48952	2.640	22.15±0.12	7.87±0.19	14.28	12.3	-	-	Type FeLoBAL	0011 0001
104611.50+024351.6	161.54796	2.73103	2.772	21.51±0.07	7.11±0.11	14.40	-	203±7	5181±144	Type 1, N v/Lyα > 1, EREW	1000 0001
155434.17+110950.6	238.64238	11.16407	2.936	21.19±0.05	6.63±0.08	14.56	-	-	-	Type 1, BAL, probably FeLoBAL	0010 0001
135959.73+052512.3	209.99888	5.42008	3.055	22.08±0.11	7.27±0.09	14.81	-	70±10	7250±833	Type 1, W1W2-dropout	0100 0001
022052.11+013711.1	35.21715	1.61976	3.138	21.82±0.08	7.08±0.09	14.74	-	341±19	2843±167	EREW, Type 1, W1W2-dropout	0100 0001
101439.51+413830.6	153.66466	41.64183	4.360	21.84±0.08	7.74±0.16	14.10	-	118±8	13568±684	Type 1, (Fe)LoBAL	0010 0001

Figure 3. The redshift distribution of the 17,744 SDSS/BOSS quasars with good W4 matches is given by the black (dashed) histogram. The redshift distribution of extremely red quasars (with $r - W4 > 14$) is given by the solid (red) line. It has a binwidth of $\Delta z = 0.20$ and is renormalized to peak at 300. This maximum bin, at $z \approx 0.70$, actually has 12 objects.

General Description	Bitwise Value	Number in Sample
Type 1 Quasar	0000 0001	37
Type 2 Quasar	0000 0010	28
Starburst	0000 0100	5
post-starburst	0000 1000	2
Radio Detected	0001 0000	37
(Fe)LoBAL	0010 0000	10
W1W2-dropouts	0100 0000	3
“Extreme” REWs	1000 0000	12

Table 3. The classifications of the optical spectra of the extremely red quasar sample, the bitwise value associated with each classification and the number in the sample. Aside from the Type 1/Type 2 dichotomy, the categories are not mutually exclusive. Quasars are flagged as ‘Radio Detected’ if they are detected in the FIRST survey.

3 KEY PROPERTIES OF THE EXTREMELY RED QUASARS

3.1 Magnitudes and redshift distribution

The key properties of the extremely red quasar sample are provided in Table 2: the catalogue name of the object, its Right Ascension, Declination, redshift, and magnitude and colour properties. The Radio fluxes are the integrated flux densities (F_{int}) measured in mJy at 20cm from the FIRST survey (Becker et al. 1995). The Rest-frame Equivalent Widths (REWs) are given for MgII at $z \leq 1.5$ and C IV for objects at redshift $z > 1.5$. The Full-Width, Half-Maximum (FWHM) of the relevant emission line is also quoted, with the REW and FWHM measurements coming from fits to the line profiles, as described in Hamann et al. (in prep.).

We give each object a classification based on optical spectral properties, indicated by the bitwise values described in Table 3. These classifications were all performed by visual inspection, with objects assigned to the classes given in Table 3 depending on the presence (or absence) of broad/narrow emission lines, broad absorption features, indications of ongoing or recently ceased star-formation, or objects with large equivalent width in the C IV emission line. These visual inspections are generally qualitative but as discussed in Section 4, aid us to understand the sample and physical types of quasars that the $r - W4 > 14$ colour selects. We also present a short note on each object in Table 2, qualitative assessments based on visual inspection of the optical spectra, and use these notes to make the link between the classifications given in Table 3 and the discussions presented in Section 4. We call a quasar “red” when its spectral slope is shallower than the Vanden Berk et al. (2001) template. The acronym EREW stands for Extreme Rest-frame Equivalent Width, referring to a group of objects we introduce and describe in Section 4.7.

The redshift range for the extremely red quasars is $0.282 < z < 4.36$, and half the sample is fainter than $r=21.4$. Figure 3 shows the redshift distribution of the sample compared to the full redshift distribution of the 17,744 SDSS/BOSS quasars with good W4 matches. The redshift distribution of the $r - W4 > 14$ sample is bimodal, with peaks around $z \sim 0.8$ and $z \sim 2.5$. This bimodality is the result of the two different SDSS quasar selections. The $z \sim 2.5$ redshift peak is due to the increased depth of the BOSS selection, as seen in Figure 4.

In the left panel of Figure 4 the redshifts and i -band magnitudes are displayed for the $r - W4 > 14$ objects, and compared to the distributions from the full SDSS DR7 and BOSS DR10 quasar catalogues. SDSS quasars use the i -band for selection limits, thus one can identify the $i = 19.1$ magnitude limit contour for the SDSS $z \lesssim 2.5$ quasars, as well as the redshift histogram peaks at $z \sim 0.6 - 1.0$ and $z \sim 2.2 - 2.7$ seen in the BOSS quasar distribution (e.g. Fig. 2 in Pâris et al. 2014). In the right panel of Fig. 4, the extremely red quasars are compared to the SDSS/BOSS+WISE catalogue, for the fully matched (i.e. any WISE band) sample given by the black contours, and for the W4-matched objects only, given by the orange contours. The redshift- i magnitude distribution of the $\approx 18,000$ objects with good W4 matches is given by the orange contours and points. These W4 objects are at the bright end in the optical, and are generally at $z < 2$.

Since we are selecting such extreme objects it is important to see how the selection affects the redshift distribution. In Figure 5 we show how the number of objects selected, and the sample redshift, depend on $r - W4$ colour. The dip in mean redshift at $r - W4 \sim 10 - 11$ comes from the fact that fainter BOSS quasars, generally at $z \gtrsim 2$, are lost quicker than $z \lesssim 1$ quasars. The upturn at $r - W4 > 11.5$ is due to the fact that the W4-bright BOSS quasars that are very red remain in the sample even to the reddest colours.

Figure 6 presents the magnitude-redshift and colour-redshift relations for the SDSS/BOSS+WISE matched catalogue highlighting the extremely red sample. This plot can be directly compared with Yan et al. (2013) (and their Figure 20). Yan et al. (2013) have also merged the WISE photometry with the SDSS (though not BOSS), and describe

Figure 4. The redshift versus i -band magnitude distribution for the extremely red quasars satisfying the $r - W4 > 14$ selection. *Left:* The blue and black contours show the distributions for the SDSS DR7 and the BOSS DR10 quasar catalogues, respectively. *Right:* The black contours show the distribution of all SDSS/BOSS quasars that have a match in any of the WISE bands, while the orange contours are SDSS/BOSS quasars with good $22\mu\text{m}$ W4-band detections.

Figure 5. The number of quasars selected, and their mean redshift, as a function of $r_{\text{AB}} - W4_{\text{Vega}}$ colour. The dashed (red) lines give the 1σ -spread on the mean redshift value. In total, 17,744 quasars have $r - W4 > 8.0$. Note that the extremely red quasar sample at $r - W4 \geq 14.0$ is the tail end of the W4-matched quasar sample.

the spectroscopic observations of eight objects that have $r - W4 > 14$ (see their Table 1 for details). Although we have an order of magnitude more $r - W4 > 14$ objects at $z > 2$, the Yan et al. objects are significantly fainter in the optical than our SDSS/BOSS detected objects. Based on the spectral properties of the Yan et al. (2013) sample, 3 (out of 8) objects are classified as Type-1 AGN, 3 objects are Type-2 AGN, and the remaining two objects have fea-

tures consistent with either a Type-2 AGN, or a star-forming galaxy, broadly in line with the composition given in Table 3. Interestingly, Yan et al. (2013) motivate the use of the $r - W4 > 14$ colour selection in order to select ultraluminous infrared galaxy (ULIRG) candidates, not red quasars (they use the shorter WISE bands to select reddened quasars), and indeed find this colour selection identifies IR luminous galaxies over a wide range of redshift, $z \sim 0.7 - 3$. However, the relative contribution to the W4 luminosity from AGN and star formation warrants investigation, especially at high redshift where the rest-frame $22\mu\text{m}$ can probe warm (AGN) dust, as well as PAH features.

3.2 Mid-Infrared Colours of the ERQs

The mid-infrared colour-colour distributions for the SDSS/BOSS quasars that are detected in WISE are shown in Figure 7. Contours are given for the full “good” W4 sample, with objects that have redshifts $z < 2$ and $z > 2$ given by the blue and red contours, respectively. The colours of the extremely red quasar sample are given by the large solid circles. We also split the extremely red quasar sample into three broad classes, ‘Type 1’ (section 4.1), ‘Type 2’ (sections 4.2 and 4.3) and the Extreme Rest-frame Equivalent Width (EREW) objects (section 4.7).

Fig. 7 shows that the Type 2 and EREW quasar populations are redder in $(W2 - W3)$ than the general W4-detected population. Upon visual inspection of the W4-detected objects with very red, $(W2 - W3) \gtrsim 3.8$ colours, many were found to lie along the Ecliptic and did not have secure W4 photometry due to Mars, Jupiter and Saturn producing a significant number of spurious mid-infrared detections⁴.

⁴ wise2.ipac.caltech.edu/docs/release/allsky/expsup/sec5_3.html#planets

Figure 6. The magnitude-redshift and colour-redshift diagrams for the objects with a WISE detection (black contours) and objects with a good $W4$ match (orange contours). The $r - W4 > 14$ extremely red quasar population is shown by the red circles. *Top left:* Redshift vs. $r - W4$ colour; *bottom left:* Redshift vs. r -band AB magnitude and *bottom right:* Redshift vs. $W4$ Vega magnitude. The data from Yan et al. (2013) are indicated by the blue squares.

These detections are not plotted in Fig. 7 and does not affect any of our $r - W4 > 14$ extremely red quasars.

In $(W1 - W2)$, the extremely red quasar population is perhaps slightly more representative of the $W4$ -detected population; none of the red quasars are bluer than $(W1 - W2) \approx 0.5$. The $W1W2W3$ colour-colour diagram has become a key diagnostic for WISE objects, and its power in separating different classes of objects is seen in Wright et al. (2010); Eisenhardt et al. (2012) and Yan et al. (2013). In particular, Eisenhardt et al. (2012) use a colour selection based on $W1W2W3$ colours to select “W1W2-dropouts”. Three of our quasars satisfy this criterion, signified by the stars in Fig 7. We discuss these objects in Section 4.6.

3.3 Radio Properties

Table 2 also gives the radio fluxes of the $r_{AB} - W4_{Vega} > 14$ sample. 37 (56%) of the full extremely red quasar sample is detected in the FIRST survey (Becker et al. 1995), which has a flux limit of ≈ 1 mJy at 20 cm, with 22 out of 28 objects at $z < 1.1$ (79%) being detected. The magnitude of radio flux at 20cm ranges from 0.55 to 960 mJy.

While the radio loudness is typically defined by the ratio of the 5 GHz radio to 4400Å flux densities (e.g., Kellermann et al. 1989; Padovani et al. 2003), with the advent of the FIRST and SDSS surveys, much of the recent analysis of radio loudness has been done in terms of the 20cm to i -band ratio, $\mathcal{R} = f_{\text{radio}}/f_{\text{optical}}$, e.g., Ivezić et al. (2002), White et al. (2007) and Kratzer & Richards (2015). Typically no k -correction is applied as the mean spectral index in the radio and optical are similar (though this assumption may not hold true for these extremely red, and unusual ob-

Figure 7. The mid-infrared “W1W2W3” colour-colour distributions for objects in the SDSS and BOSS Quasar catalogues, that have a match in WISE and have a $W4$ -band $SNR \geq 3$. Objects with redshifts $z < 2$ and $z > 2$ are given by the blue and red contours, respectively, with the contouring level the same for the subsets of the distributions. The insert gives the redshift distribution for the $W4$ -matched WISE sample which have good $W1W2W3$ detections (dashed histogram) and which pass the “good $W4$ ” criteria (solid histogram). The large coloured points indicated the $r - W4 > 14$ quasars. We split this sample into three broad classes. The EREW sample generally lies on the edge of the overall $W4$ -matched population, with roughly half the sample having very red colours i.e., $W2 - W3 \gtrsim 4.5$. The stars signify objects that pass the “W1W2-dropout” selection of Eisenhardt et al. (2012) (see text for details).

jects). While there is no true gap in the distribution of $\mathcal{R}$, at $\mathcal{R} \sim 10$ there is a subtle change in the distribution (e.g., White et al. 2007; Kratzer & Richards 2015) and it is common to refer to objects with $\mathcal{R} \geq 10$ as “radio-loud”. The value of $\mathcal{R}$ can be calculated straight from Table 2 for each of our objects, and we find that all 37 of our radio-detected objects are nominally radio-loud. However, without knowing the extent of reddening from dust in our sources, these values should be considered as an upper limit to the true value of $\mathcal{R}$. We also find that all of the 37 radio-detected objects are unresolved in FIRST.

A total of 30 objects (46%) in our sample have a radio target selection flag set indicating that these objects were detected in the FIRST survey and passed the SDSS/BOSS quasar selection to be observed as quasar targets. Of these 30 objects, 10 are from SDSS and 20 from BOSS. 17 objects (26%) are selected for spectroscopy *solely* due to their radio properties, all of which are from BOSS.

Yan et al. (2013) discovered ULIRGs at $z \sim 2$ with extremely red colours, which suggests that star-formation will lead to an enhancement of the $W4$ emission we detect from our sample. While the sample definition of $r - W4 > 14$ selects quasars that are red due to dust obscuration (i.e., the targeted obscured quasar population), there is the possibility it will also select systems with particularly bright IR emission due to star formation. The large fraction of radio-

Figure 8. The ratio between the observed $W4$ flux of our sample and the $W4$ flux that would be expected given the observed radio luminosities and the conversions between radio and star formation rates reported in Bell (2003). A ratio of > 1 means that the $W4$ emission is likely AGN dominated. For objects undetected in the radio (FIRST) survey, we only calculate an upper limit on the star formation rate and thus a lower limit on the observed-to-predicted flux ratio, given by the red arrows.

detected quasars in the resulting sample, and the known relations between radio and IR emission from star formation, suggest that this may be an aspect of the extremely red quasar selection.

To investigate this further, we calculate rest-frame 1.4 GHz radio luminosity from the observed fluxes using a radio spectral index of $\alpha = -0.7$. In the first instance, we assume that all of the radio emission is due to star formation, which will result in upper limits on the SFR derived from the radio and MIR. We use the conversions between radio and star formation rates reported in Bell (2003) to calculate the necessary SFRs and total IR luminosity of star formation $L_{\text{SFR}, 8-1000\mu\text{m}}$. Then we scale seven star-formation templates from Kirkpatrick et al. (2012) and Mullaney et al. (2011) to match the SFRs necessary to explain the observed radio emission. We then redshift them to the redshift of the target, convolved with the $W4$ filter from Jarrett et al. (2011) and calculate the predicted IR flux. Figure 8 shows the ratio between the observed and the predicted $W4$ fluxes; those are shown as black points, where the error bars reflect the spread amongst templates (and therefore provide a measure of the systematic error). For objects undetected in the radio, we can only calculate an upper limit on the SFR rate and thus a lower limit on the observed-to-predicted flux ratio; those are shown as red upward arrows, and those are conservative (i.e., the lowest lower limit of the seven templates). A ratio of > 1 means the MIR emission is likely AGN dominated, while ratios < 1 mean the predicted SFR is less than expected, given the radio flux.

There are 37 objects detected in the radio; if all the radio flux is associated with star-formation, which is unlikely, then for these objects the median luminosity due to star formation would need to be $L_{8-1000\mu\text{m}} = 10^{13.6} L_{\odot}$ in order to explain the median radio luminosity of $\nu L_{\nu}[1.4\text{GHz}] =$

$10^{41.6} \text{ erg sec}^{-1}$. In other words, these objects would need to be forming stars at the median rate of $\approx 7700 M_{\odot} \text{ yr}^{-1}$ in order to explain the observed radio emission, and this rate is $\sim 4\text{--}10\times$ higher than the e.g. $850\mu\text{m}$ -selected submillimeter galaxy (SMG) population at redshifts $z \sim 2 - 3$ (Wardlow et al. 2013; Fu et al. 2013; Casey et al. 2014). However, this SFR is an upper limit, as the AGN is almost certainly contributing significant flux in all objects. It is more likely that both the radio emission and the $W4$ emission in these objects are due to the AGN and star formation, and thus the SFRs are more in-line with those of SMGs and WISE-selected luminous infrared galaxies (Wu et al. 2012; Jones et al. 2014, 2015; Tsai et al. 2015).

The two points with the lowest $W4_{\text{obs}}/W4_{\text{SFR,predicted}}$ ratios are potentially beamed sources with an apparent extremely high radio luminosity $\nu L_{\nu}[1.4\text{GHz}] \simeq 10^{44.5} \text{ erg s}^{-1}$ which has nothing to do with star formation in the host galaxy (though our data are insufficient to determine the origin of the $W4$ emission in these sources). For the less extreme sources at $z < 1$, we find that the required median SFR necessary to explain the observed radio and $W4$ emission is $2180 M_{\odot} \text{ yr}^{-1}$, which is still very high, but could potentially be the result of an e.g. (major) merger. While we do not have direct observations (e.g., far-infrared) which would rule out such star formation rates, various authors find it unlikely that star formation dominates mid-infrared and radio emission of luminous Type 2 quasars at $z < 1$ (e.g., Harrison et al. 2014; Zakamska & Greene 2014; Zakamska et al. 2015, in prep.).

The remaining objects are undetected in the radio and therefore we can place only upper limits on star formation rates. For this population we find the median lower limit $W4_{\text{obs}}/W4_{\text{SFR,predicted}}$ of 2.4, meaning that the star formation is inadequate by at least this factor in explaining the observed $W4$ flux. Thus we conclude that the $W4$ fluxes in these objects must be dominated by the AGN as well. However, it seems that ultimately, we currently do not have enough data to determine whether the radio emission is due to AGN or star formation. New FIR/submm data is required, with further SED decomposition, and indeed this is where future analysis will be concentrated (Hamann et al., 2015, in prep.; Zakamska et al., 2015, in prep.).

4 CLASSIFICATIONS FROM OPTICAL SPECTROSCOPY

The spectra of the extremely red quasar sample are publicly available at the SDSS-III Science Archive Server⁵. We broadly classify the objects that satisfy the $r - W4 > 14$ selection into: (i): unobscured (but reddened) Type 1 quasars possessing broad lines (bitwise value 1 in Table 3); (ii): objects that show evidence for being narrow-line Type 2 quasars from their optical spectra (bitwise value 2 in Table 3); (iii): objects that suggest ongoing, or recently ceased star formation (bitwise values 4 and 8); (iv): radio-detected quasars (in the FIRST survey, bitwise value 16); (v): quasars with blue-shifted absorption features (bitwise value 32); (vi): objects that are not detected in the WISE $W1/2$ -bands

⁵ <http://dr10.sdss3.org>

(bitwise 64) and (vii): a class of Type 1 sources with extreme rest equivalent widths in their UV broad lines, namely, $\text{REW} > 150\text{\AA}$ in either C IV or Mg II (bitwise 128). These classification groups are not mutually exclusive; several objects are assigned to multiple groups. However, we class all of the sources as either Type 1 (37/65; 57%) or Type 2 (28/65; 43%). These sources are indicated by the bitwise values in the last column of Table 2 and are described in Table 3. Example spectra are presented in what follows.

4.1 Type 1 Quasars

Figure 9 presents the spectra of six examples of extremely red quasars, classified as Type 1 based on broad emission lines seen in their optical spectroscopy. Where measured, the mean line widths are $\text{FWHM}(\text{Mg II}) \sim 6440 \text{ km s}^{-1}$ and $\text{FWHM}(\text{C IV}) \sim 4900 \text{ km s}^{-1}$. Figure 9 lists the SDSS name, redshift, classification (from Table 3) and $r - W4$ colour of each object.

The spectra in Figure 9 are heterogeneous; some show blue continua, while some are red. The first object (with the lowest redshift in our sample) has unusually broad [O III] with non-Gaussian structure (Liu et al. 2010, see also section 4.4). Many of the examples of Type 1 quasars from our selection appear to be significantly reddened (Richards et al. 2003) often with a host galaxy component.

Glikman et al. (2013) identify dust-reddened quasars by matching the FIRST radio catalogue to the UKIRT Infrared Deep Sky Survey (UKIDSS; Lawrence et al. 2007). This study identified 14 reddened quasars with $E(B - V) > 0.1$, including three at $z > 2$. However, Glikman et al. (2013) find no heavily reddened ($E(B - V) \gtrsim 0.5$) quasars at high redshifts ($z > 2$). The object ULASJ1234+0907 ($z = 2.503$) discovered by Banerji et al. (2014) is currently the reddest broad-line Type 1 quasar known, with $(i - K)_{\text{AB}} > 7.1$. Since all our extremely red quasars have $i \lesssim 22$ mag and a range of $1.62 < (i - K)_{\text{AB}} < 4.73$, none of our objects have optical-to-near infrared colours as red as ULASJ1234+0907.

We note the object J165258+390249 has a relatively blue continuum and is clearly not suffering from dust obscuration. This object may be optically variable, and at the time of the optical and IR photometry had a large enough $r - W4$ to satisfy our colour cut. We also note it is a radio source, and has features consistent with an optical blazar spectrum. This further highlights the importance of understanding the radio properties (and potentially the synchrotron radiation contribution) of these quasars and how it affects the IR and the resulting sample of objects.

4.2 Type 2 Quasar candidates at low redshift

We define Type 2 quasars as objects with narrow permitted emission lines that do not have an underlying broad component, and with line ratios characteristic of non-stellar ionizing radiation (Zakamska et al. 2003; Reyes et al. 2008). Specifically, in Type 2 quasars the broad-line region is completely blocked from the observer by strong obscuration ($A_V > 10$), and nuclear activity is inferred indirectly via the emission line ratios of ionized gas which are characteristic of photo-ionization by the quasar (Baldwin et al. 1981; Veilleux & Osterbrock 1987). At low redshifts, where $\text{H}\beta$

and [O III] are accessible in the optical spectrum, we can define Type 2 objects as those in which the kinematic structure of the permitted emission lines is similar to that of the forbidden lines; the permitted lines should not display an additional broad ($\text{FWHM} \gtrsim 2000 \text{ km s}^{-1}$) component. We also require $[\text{O III}] / \text{H}\beta > 3$.

Figure 10 presents examples of Type 2 quasar candidates at $z \lesssim 1$. No line-fitting has been performed, but all objects in this class are seen by visual inspection to have similar same kinematics in the permitted $\text{H}\beta$ and forbidden [O III] lines. In the extremely red quasar sample, the quasar nature of the Type 2 candidates is not in doubt: the typical values of $[\text{O III}] / \text{H}\beta$ are 10 or greater, and high-ionization species such as [Ne V] are detected in most cases. However, $\text{H}\beta$ can be weak and there may be broad components in some $\text{H}\alpha$ lines (Zakamska et al. in prep) which are redshifted out of the BOSS wavelength coverage.

Just over half (60%) of our Type 2s are at $z < 1$. This is due to the fact that we are using the spectroscopic quasar catalogues as our initial sample, before matching to the WISE photometry. Thus, from the requirement of having a relatively bright detection in the optical (cf. the Brand et al. studies) there will be a selection bias towards lower redshift objects.

Among the 28 objects with $z < 1.1$, 20 (71%) are classified as Type 2 candidates from their optical spectra; see Figure 10. This high fraction of Type 2 objects is completely uncharacteristic of the parent sample of the SDSS/BOSS quasars, making it clear that the selection based on high infrared-to-optical colours can indeed recover obscured quasars. All seven Type 2 quasars in our sample that were observed before July 2006 are found in the catalogue of Reyes et al. (2008).

4.3 Type 2 Quasar candidates at high redshift

We find objects that have narrow emission lines, weak continua and satisfy the definition of Type 2 candidates, at high redshift ($z > 1.5$) as well; see Figure 11. The classifications are performed by visual inspection of the optical spectra and all objects in this class with $z > 1.5$ have $\text{FWHM}(\text{C IV}) \lesssim 2000 \text{ km s}^{-1}$.

These objects have similar optical spectra to those described in Yan et al. (2013) and Alexandroff et al. (2013), although there are no objects in common between our sample and that presented in Alexandroff et al. (2013). Those authors present a sample of 145 candidate Type 2 quasars from BOSS at redshifts between 2 and 4.3, using data from DR9 (Ahn et al. 2012). These objects are characterized by weak continuum in the rest-frame ultraviolet with a typical continuum magnitude of $i \approx 22$ (i.e., $M_i(z = 2.5) \approx -25$) and strong lines of C IV and $\text{Ly}\alpha$, with $\text{FWHM} \leq 2000 \text{ km s}^{-1}$. Typical ($\sim L^*$) galaxies are ≥ 1 magnitude fainter at these redshifts (Marchesini et al. 2007, 2012), suggesting host-galaxy light is not sufficient to explain the continuum luminosity and some level of AGN light is getting through/around the absorber.

The reddest quasar in Alexandroff et al. (2013) has $r - W4 = 13.30$. In general, our selection of Type 2 candidates is not as strict as that of Alexandroff et al. (2013), and we find no overlap between the sample here and the DR9-based sample in Alexandroff et al. (2013). While this class of

Figure 9. Examples of extremely red quasars classified as Type 1 quasars based on their optical spectroscopy. The SDSS name, redshift, classification and $r - W4$ colour are given in each panel. The red part of the spectrum marks the contribution to the observed r -band flux, while the blue line is the estimated error per pixel (Lee et al. 2013). The spectra are smoothed using an 8-pixel boxcar average. Prominent emission lines often seen in AGN spectra; Ly β 1025.72, Ly α 1215.67, N v 1240.14, Si iv 1393.76, 1402.77, C iv 1548.19, 1550.76 C iii] 1908.73, Mg ii 2798.75 Ne v 3346.82, 3426.84 [O ii] 3728.48 [Ne iii] 3869.85, H δ 4102.89, H γ 4341.68, H β 4862.68, [O iii] 4960.30, 5008.24 and H α 6564.61 Å, are marked by the vertical dashed lines, where we use vacuum wavelengths for identifying emission lines. The feature at 5575 Å in the lower two panels is the [O i] bright sky line. Spectra that cut off at 3800 Å and 9200 Å are SDSS spectra, while BOSS spectra extend to 3600 Å and 10,400 Å. Objects with starred classifications were targeted due to their radio properties. Note that the y -axis scales are different in each panel.

Figure 10. Same as for Fig 9, except showing examples of extremely red quasars classified as Type 2 quasars based on their optical spectroscopy, at $z \lesssim 1.1$.

Figure 11. Same as for Fig 9, except showing examples of extremely red quasars classified as Type 2 quasars based on their optical spectroscopy, at $z \gtrsim 1.5$.

object shows only narrow emission lines in the rest-frame UV spectra, Greene et al. (2014) obtained near-infrared spectroscopy for a subset of the Alexandroff et al. (2013) sample, demonstrating that the H α emission line consistently requires a broad component. This result implies that the typical extinction in these objects is $A_V \lesssim$ a few, sufficient to block the rest-frame UV continuum and broad lines but not the rest-frame optical - consistent with the (relatively) bright continuum.

4.4 Objects with starburst and post-starburst signatures

Our sample includes quasars that have starburst or post-starburst signatures in their optical spectra. Examples are given in Figure 12.

The [OII]/[OIII] ratio can be used as a starburst indicator in AGN and galaxies (e.g., Kauffmann et al. 2003; Groves et al. 2006). Objects from our sample are placed into the starburst class if they exhibit evidence of obvious [OII] emission (Kennicutt 1998; Kewley & Dopita 2002; Kewley et al. 2004; Ho 2005; Moustakas et al. 2006; Mostek et al. 2012), though we acknowledge that [OII] emission can also be associated with the narrow-line region of the AGN.

The post-starburst objects are objects which have either a Balmer break, and/or an obvious Balmer series, originating from an A-star population that is not dominated by blue light from OB-stellar populations. Quasars with post-starbursting signatures have been identified previously, (e.g., Brotherton et al. 1999, 2002; Cales et al. 2013, 2014) and Matsuoka et al. (2015) derive physical properties from optical spectra while Wei et al. (2013) study the MIR spectral properties of post-starburst quasars.

Both post-starburst objects in our sample are at relatively low redshift, and not all the host galaxy light is likely to be captured by the 2" spectroscopic fiber. However, the quasar J112657.76+163912.0 (hereafter J1126+1639) has a pronounced Balmer break and we estimate that the fraction of e.g A-star host galaxy flux required is at least that of the AGN. J1126+1639 may have other optical objects within the W4-beam, as we mentioned in Section 2.2. J1126+1639 appears to be at the center of a group/small cluster of other optically red galaxies. However, the other group members are *blue* in (W1 – W2) colour, suggesting that these objects are not bright in W4. Our likely candidate appears disturbed in the SDSS optical image, suggesting that it hosts an on-going merger.

4.5 Broad Absorption Line Quasars

Our $r - W4 > 14$ quasar selection also finds broad absorption line (BAL) objects (Figure 13). Although we do not use it for our classifications, the CIV balnicity index (BI; Weymann et al. 1991) for our extremely red BAL quasars as reported in the DR10Q catalogue is generally $\log_{10}(\text{BI}_{\text{CIV}}) \gtrsim 3$. In particular, we find examples of BALs that exhibit absorption from low-ionization species, i.e. LoBALs, and objects that have Fe II absorption, i.e. FeLoBALs. At $z = 4.36$, J101439.51+413830.6 is the highest redshift object in the full sample, but due to this high redshift, any Fe II

Figure 13. Same as for Fig 9, except showing examples of the Broad Absorption Line quasars. These three objects are examples of the rare subclass of FeLoBALs. At $z = 4.36$, J101439.51+413830.6 is the highest redshift object in our full sample.

absorption will be outside the optical wavelength coverage of the SDSS/BOSS spectra.

It has been known for some time that BALs can be redder (in optical-to-near infrared colour) than the general quasar population e.g., Hall et al. (1997). Hall et al. (2002b) report on a number of heavily-reddened, extreme BAL quasars discovered in SDSS, while Trump et al. (2006), Gibson et al. (2009) and Allen et al. (2011) explore this reddening-BAL relation. However, we note that the presence of the BAL objects in our sample is not due just to dust reddening; in these objects, the BAL troughs remove a large fraction of the continuum that would otherwise contribute to the r -band flux, which makes the $r - W4$ colour even redder. We find FeLoBALs in our sample not because they have red continua, but because the BALs have wiped out the r -band flux; in this sense these are in the sample for different reasons than for the other very red quasars.

4.6 W1W2-dropout Quasars

Objects that are selected to be bright at 12 or 22 μm , but are undetected by WISE at 3.4 and 4.6 μm are “W1W2-dropouts”. Eisenhardt et al. (2012) find ~ 1000 such objects over the whole sky, including the $z = 2.452$ source WISE J181417.29+341224.9, a hyper-luminous infrared galaxy (HyLIRG) with $L_{\text{IR}} > 10^{13} L_{\odot}$. WISE 1814+3412 has a W4 flux density of $14.38 \pm 0.87 \text{ mJy}$; a 350 μm detection (Wu et al. 2012) implies a minimum bolometric luminosity of $3.7 \times 10^{13} L_{\odot}$ suggesting that the W1W2-dropouts are extreme cases of luminous, hot (60-120K) dust-obscured galaxies possibly representing a short evolutionary phase during galaxy merging and evolution. Due to these dust temper-

Figure 12. Same as for Fig 9, except showing examples of extremely red quasars that display signatures of an ongoing, or post, star-bursting phase. In particular we identify the Balmer edge in J105327.92+375804.2 and J112657.76+163912.0.

Figure 14. Same as for Fig 9, except showing quasars that are “W1W2-dropouts”. The feature at 5575 Å in the lower two panels is the [O I] bright sky line.

atures, the W1W2-dropouts are also known as hot, dust-obscured galaxies (“Hot DOGs”).

Three of our 65 objects ($\sim 5\%$) satisfy the W1W2-dropout selection criteria. Their optical spectra are shown in Figure 14. These three quasars were all discovered by BOSS, have r -band fluxes of $\approx 5 - 8 \mu\text{Jy}$ (cf., $2.25 \pm 0.11 \mu\text{Jy}$ for WISE 1814+3412) and redshifts $z = 2.638 - 3.138$, placing them in the HyLIRG regime.

4.7 Extreme Equivalent Width Objects

The objects that we present in Figure 15 all have C IV FWHM $> 2000 \text{ km s}^{-1}$ (Table 2) and are thus classified as Type 1 quasars. These six examples are a set of objects at $z \simeq 2 - 3$ which are characterized by extreme rest-frame equivalent widths (EREWs) of $\gtrsim 150 \text{ Å}$ (the measured REWs from the C IV line are given in Fig. 15). Four sources have $\text{REW}(\text{C IV}) > 200 \text{ Å}$. For comparison, typical quasars have $\text{REW}(\text{C IV}) \approx 25\text{--}50 \text{ Å}$. Some of the extreme REW sources also have unusual line properties, including high N V/Ly α , enhanced Si IV+O IV] and weak He II and C III]. Some also have strong Ly β and unusual Ly α profiles.

To quantify how unusual these objects are, we use emission line data from the DR10Q catalogue to select sources that have: $2.0 < z < 3.4$, the visual inspection flag for BALs set to 0 (i.e. no BALs), $\text{FWHM}(\text{C IV}) > 2000 \text{ km s}^{-1}$, and the ratio of the C IV rest-frame equivalent width to its uncertainty > 3 . A total of 98807 quasars satisfy these criteria and in the cumulative distribution, 5.8% of objects have $\text{REW}(\text{C IV}) > 100 \text{ Å}$ with 1.3% having $\text{REW}(\text{C IV}) > 150 \text{ Å}$. This is to be compared to 55% (45%) of the non-BAL Type 1s with measurable C IV in our sample having $\text{REW}(\text{C IV}) > 100$ (150) Å.

The quasar J1535+0903 (see Appendix A1) may also be a member of this class. We measure $\text{REW}(\text{C IV}) = 136$

Figure 16. The optical $(g-r)$ colour of the W4-matched quasars (orange points) as a function of redshift. The mean colour of the W4-matched sample is given by the solid blue line. At $z > 1.9$ the colour distribution of our extremely red quasars is impressively broad, demonstrating the heterogeneity of our sample. The EREW population is *bluer* than the overall population in $(g-r)$ at $z \approx 2.5$. All optical magnitudes are AB.

Å, and $\text{REW}(\text{Mg II}) \sim 280 \text{ Å}$, the precise value depending on uncertain continuum placement. Like the N V/Ly $\alpha > 1$ objects, J1535+0903 has very strong Al III $\lambda 1860$.

The large REWs in these objects might be caused by suppressed continuum emission analogous to Type 2 quasars in the Unified Model. However, the large line widths (at least compared to typical narrow line regions; Liu et al. 2013) and in some cases very high electron densities (Section A1; Hamann et al., in prep.) suggest that the line-emitting regions are close to the central continuum source, as in typical broad line regions. It therefore seems difficult to have dust obscuring the continuum source without also obscuring the broad emission lines. We investigate these issues further in Hamann et al. (in prep.).

5 COLOURS AND SELECTION OF THE EXTREMELY RED QUASARS

The optical SDSS/BOSS spectra of the extremely red quasars have been presented in Section 4. The heterogeneity of the optical spectra is immediately obvious. Upon visual inspection of the optical spectra, we have classified the extremely red quasar sample into various groups given in Tables 2 and 3. In this section, we explore the colours and selection of the extremely red quasars, in order to understand the relationship of the sub-classes to each other, and with respect to the general quasar population.

5.1 Colours of the extremely red quasar groups

Figure 16 presents the optical $(g-r)$ colours for the full W4 matched sample as a function of redshift. The mean $(g-r)$ colour of all quasars with a W4-detection is given by the

Figure 15. Same as for Fig 9, except showing examples of the new set of “Extreme Rest-frame Equivalent Width” (EREW) objects. These objects possess large REWs of e.g. the CIV line (values given in Table 2), unusual profiles of the Ly α to NV complex (and the ratio of Ly α :NV) and flat continua in all examples. The quasar J022052.11+013711.1 (bottom panel) is also a W1W2-dropout (Fig. 14).

Figure 17. The optical-midinfrared colour for our W4-matched sample (orange points), with $r - W1$, $r - W2$, $r - W3$ and $r - W4$ shown from top to bottom. The mean colour of the W4-matched sample is given by the solid blue line. As the optical-MIR wavelength arm extends from W1 ($3.4\mu\text{m}$) to W4 ($22\mu\text{m}$) the selected population moves further away from the mean W4-matched sample. All optical magnitudes are AB, all WISE magnitudes are Vega.

blue line. As in Figure 6, the three groups are indicated separately: Type 1s (dark blue circles), Type 2s (purple) and EREW objects (dark green). The extremely red quasars are generally redder than the mean W4 detected population in $(g - r)$ colour. However, the EREW objects are the exception; they are bluer than the full W4 population at all redshifts that they are found. This suggests that we are perhaps not sensitive to all EREWs, but just those objects with redshifts such that there are a number of very strong lines in the g -band.

Fig. 17 presents the optical to infrared colours, $(r - W1)$, $(r - W2)$, $(r - W3)$ and $(r - W4)$, for the full W4 matched sample as a function of redshift. The quasars detected at $22\mu\text{m}$ generally have a $r - W1$ colour of 5 or more for redshifts $z = 0 - 3$, and again are redder than the general population of quasars. However, some of the Type 2 objects do have $(r - W1)$ and $(r - W2)$ colours consistent with the general W4 detected population. However, all the extremely red quasars are considerably redder in $(r - W3)$ which samples the ratio of rest-frame UV/blue to $\sim 6 - 7\mu\text{m}$ light at redshift $z \sim 0.8$ and rest-frame far UV to $\sim 3.4\mu\text{m}$ colour at redshift $z \sim 2.5$. By design, our sample is much redder than the general W4 detected quasar population in $(r - W4)$ colour, the latter of which has $10 \lesssim (r - W4) \lesssim 12$ for redshifts $z = 0 - 3$.

Figure 18. The optical colour-colour properties of SDSS/BOSS+WISE matched sample and the extremely red quasars: (top left): $(u - g)$ vs. $(g - r)$; (top right): $(g - r)$ vs. $(r - i)$; (bottom left): $(r - i)$ vs. $(i - z)$; (bottom right): $(g - i)$ vs. i -band PSF magnitude. The distribution of all the SDSS/BOSS quasars with a good match in any WISE band is given by the black contours, while those objects with good W4-matches, and their colour distributions are given by the solid orange contours. We also show the location in colour-colour space of the three broad groups of our extremely red quasars – Type 1s (dark blue circles), Type 2s (purple) and EREW objects (dark green). The EREWs occupy a particular place in gri -colour-colour space, being very blue in $(g - i)$ colour.

Figure 18 shows the optical colour-colour distribution of the full SDSS/BOSS WISE matched sample (black contours), and those that have good W4 detections (orange contours). The extremely red quasar population tends to avoid the $(u - g) \leq 0.4$ and $(g - r) \leq 0.3$ region. The EREWs occupy a particular location in gri colour-colour and $(g - i)$ - i colour-magnitude space. The EREWs have a very blue colour of $(g - i) \sim 0$, in contrast to the overall extremely red quasar population, which tend to have redder $(g - i)$ colours than the general W4-matched sample. Further investigations into the colours and selection properties of the extremely red quasars and the EREWs are presented in Hamann et al. (in prep.).

Note that $z \approx 1.5$ is the redshift where C IV begins to contribute to the flux in the SDSS g -band. Thus, Type 2 quasars at $z \lesssim 1.5$ may have colours inconsistent with the SDSS/BOSS optical quasar selection criteria and thus enter the SDSS/BOSS sample via their radio properties. On the other hand, Type 2 quasars at $z \gtrsim 1.5$ with C IV and Ly α emission really have blue continua.

6 SUMMARY AND CONCLUSIONS

We have matched the quasar catalogues of the SDSS and BOSS with WISE to identify quasars with extremely high

infrared-to-optical ratios $r_{AB} - W4_{\text{Vega}} > 14$ mag (i.e., $F_{\nu}(22\mu\text{m})/F_{\nu}(r) \gtrsim 1000$). We identify 65 objects and note the following findings and conclusions:

- This sample spans a redshift range of $0.28 < z < 4.36$ and has a bimodal distribution, with peaks at $z \sim 0.8$ and $z \sim 2.5$.
- We recover a wide range of quasar spectra in this selection. The majority of the objects have spectra of reddened Type 1 quasars, Type 2 quasars (both at low and high redshift) and objects with strong absorption features.
- There is a relatively high fraction of Type 2 objects at low redshift, suggesting that a high optical-to-infrared colour can be an efficient selection of narrow-line quasars.
- There are three objects that are detected in the $W4$ -band but not $W1$ or $W2$ (i.e., “ $W1W2$ -dropouts”), all of which are at $z > 2.6$.
- We identify an intriguing class of objects at $z \simeq 2 - 3$ which are characterized by equivalent widths of $\text{REW}(\text{CIV}) \gtrsim 150\text{\AA}$. These objects often also have unusual line properties. We speculate that the large REWs may be caused by suppressed continuum emission analogous to Type 2 quasars in the Unified Model. However, there is no obvious mechanism in the Unified Model to suppress the continuum without also suppressing the broad emission lines, thus potentially providing an interesting challenge to quasar models.

There are numerous avenues that the current dataset can contribute to understanding quasar–host galaxy evolution and quasar orientation. Hamann et al. (in prep.) investigate the nature and origins of extreme broad emission line REWs in BOSS quasars. This will involve the detailed examination of emission line properties for a subset of extreme REW quasars to gain a better understanding of the physical conditions and possible transient nature of this phenomenon. With the addition of $\sim 18,000$ spectra of $22\mu\text{m}$ detected quasars, one can significantly update Roseboom et al. (2013) where those authors investigate the range of covering factors, determined from the ratio of IR to UV/optical luminosity, seen in luminous Type 1 quasars. However, the inhomogeneous selection of our dataset makes computing the completeness challenging. One can also measure the clustering of the red-selected quasars, (e.g., Donoso et al. 2014; DiPompeo et al. 2014), using spectroscopic redshifts to enable a more robust 3D-clustering measurement. These investigations into environmental, and host galaxy, e.g. morphology, star-formation, properties of the very red optical–IR colour-selected population will be able to elucidate on the nature of the “extremely red quasar” population.

ACKNOWLEDGEMENTS

We thank R. M. Cutri and the IPAC team for the Explanatory Supplement to the WISE All-Sky and AllWISE Data Release Products resource. Peregrine McGehee and the IRSA HelpDesk were also very useful. N.P.R. thanks Nathan Bourne for useful discussions on infrared-to-radio flux ratios. AllWISE makes use of data from WISE, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, and NEOWISE, which is a project of the

Figure A1. The spectrum of SDSS J153542.41+090341.1. This object has exceptionally strong FeII emission in the UV1 (2600Å), UV2/3 (2400Å) and 2740Å complexes with weak emission from other FeII features.

Jet Propulsion Laboratory/California Institute of Technology. WISE and NEOWISE are funded by the National Aeronautics and Space Administration. N.P.R. acknowledges that funding for part of this project was supplied by NASA and the *Hubble* Grant Number: HST-GO-13014.06. FH acknowledges support from the USA National Science Foundation grant AST-1009628.

Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the U.S. Department of Energy. The SDSS-III web site is <http://www.sdss3.org/>. SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration including the University of Arizona, the Brazilian Participation Group, Brookhaven National Laboratory, University of Cambridge, University of Florida, the French Participation Group, the German Participation Group, the Instituto de Astrofísica de Canarias, the Michigan State/Notre Dame/JINA Participation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, New Mexico State University, New York University, Ohio State University, Pennsylvania State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of Utah, Vanderbilt University, University of Virginia, University of Washington, and Yale University.

Facilities: SDSS, WISE

APPENDIX A: NOTES ON INDIVIDUAL OBJECTS

A1 J153542.41+090341.1

Figure A1 displays the spectrum of SDSS J153542.41+090341.1, hereafter J1535+0903, a remarkable object with a unique pattern of emission lines. We classify this quasar as a Type 2 candidate based on line widths in both CIV (Table 2) and Mg II ($\text{FWHM} = 1600 \pm 60 \text{ km s}^{-1}$). However, the exceptionally strong emission in Al III 1860 and in Fe II at $\sim 2300 - 2770\text{\AA}$ point to high densities indicative of a broad line region and thus a Type 1 classification. In particular, photoionization models by Baldwin et al. (1996) demonstrate that large ratios of Al III 1860 compared

to the inter-combination lines with similar ionizations, C III] 1909 and especially Si III] 1892, require densities $n_e \gtrsim 10^{12} \text{ cm}^{-3}$. Similarly, Baldwin et al. (2004) showed that this peculiar pattern of FeII emission lines, dominated by a few spikes that include the resonance multiplets UV1 and UV 2,3, also requires densities $\gtrsim 10^{12} \text{ cm}^{-3}$ in a gas with very low ionization parameter. Hamann et al. (in prep.) discuss this object further.

A2 J013435.66-093102.9

SDSS J013435.66-093102.9, hereafter J0134-0931, is a gravitational lens. J0134-0931 has $r_{\text{psf}} = 21.33$, $W4 = 6.59$ and a redshift $z = 2.22$ and is also detected in FIRST at 0.96 Jy (integrated flux). J013435 has previously been reported and studied in detail in Gregg et al. (2002); Hall et al. (2002a); Keeton (2001); Keeton & Winn (2003); Winn et al. (2002) and Winn et al. (2003). The flux is the linear sum of the lensing galaxy and the (boosted) quasar and at least one of the galaxy or quasar is exceedingly red. If there is dust in the foreground galaxy, it will extinct the quasar, and cause the $r - W4$ colour to be red. However, the foreground galaxy, at $z = 0.76$, is faint enough to not contribute significantly to the overall flux, so the background quasar is extremely red on its own.

REFERENCES

- Abazajian K. N., et al., 2009, ApJS, 182, 543
 Adelman-McCarthy J. K., et al., 2006, ApJS, 162, 38
 Ahn C. P., et al., 2012, ApJS, 203, 21
 Ahn C. P., et al., 2014, ApJS, 211, 17
 Alam S., et al., 2015, ArXiv e-prints
 Alexandroff R., et al., 2013, MNRAS, 435, 3306
 Allen J. T., Hewett P. C., Maddox N., Richards G. T., Belokurov V., 2011, MNRAS, 410, 860
 Antonucci R., 1993, ARA&A, 31, 473
 Antonucci R. R. J., Miller J. S., 1985, ApJ, 297, 621
 Assef R. J., et al., 2013, ApJ, 772, 26
 Baldwin J. A., Ferland G. J., Korista K. T., Carswell R. F., Hamann F., Phillips M. M., Verner D., Wilkes B. J., Williams R. E., 1996, ApJ, 461, 664
 Baldwin J. A., Ferland G. J., Korista K. T., Hamann F., LaCluyzé A., 2004, ApJ, 615, 610
 Baldwin J. A., Phillips M. M., Terlevich R., 1981, PASP, 93, 5
 Banerji M., Fabian A. C., McMahon R. G., 2014, MNRAS
 Banerji M., McMahon R. G., Hewett P. C., Gonzalez-Solares E., Koposov S. E., 2013, MNRAS, 429, L55
 Becker R. H., White R. L., Helfand D. J., 1995, ApJ, 450, 559
 Bell E. F., 2003, ApJ, 586, 794
 Bolton A. S., et al., 2012, AJ, 144, 144
 Brand K., et al., 2006, ApJ, 644, 143
 Brand K., et al., 2007, ApJ, 663, 204
 Brand K., et al., 2008, ApJ, 680, 119
 Brandt W. N., Hasinger G., 2005, ARA&A, 43, 827
 Brotherton M. S., et al., 1999, ApJ Lett., 520, L87
 Brotherton M. S., Grabelsky M., Canalizo G., van Breugel W., Filippenko A. V., Croom S., Boyle B., Shanks T., 2002, PASP, 114, 593
 Bussmann R. S., et al., 2009a, ApJ, 693, 750
 Bussmann R. S., et al., 2009b, ApJ, 705, 184
 Cales S., Brotherton M. S., Shang Z., Bennert V., Canalizo G., Diamond-Stanic A. M., 2014 Vol. 223 of AAS, #115.06
 Cales S. L., et al., 2013, ApJ, 762, 90
 Casey C. M., Narayanan D., Cooray A., 2014, Phys. Rep., 541, 45
 Chiu K., Richards G. T., Hewett P. C., Maddox N., 2007, MNRAS, 375, 1180
 Cutri R. M., et al., 2011, Technical report, Explanatory Supplement to the WISE Preliminary Data Release Products
 Dawson K., et al., 2013, AJ, 145, 10
 Dey A., et al., 2008, ApJ, 677, 943
 Diamond-Stanic A. M., Rieke G. H., 2010, ApJ, 724, 140
 DiPompeo M. A., Myers A. D., Hickox R. C., Geach J. E., Hainline K. N., 2014, MNRAS, 442, 3443
 Donley J. L., et al., 2012, ApJ, 748, 142
 Donoso E., Yan L., Stern D., Assef R. J., 2014, ApJ, 789, 44
 Draine B. T., Li A., 2007, ApJ, 657, 810
 Eisenhardt P. R. M., et al., 2012, ApJ, 755, 173
 Eisenstein D. J., Weinberg D. H., et al., 2011, AJ, 142, 72
 Elitzur M., 2008, NewAR, 52, 274
 Elvis M., 2010, in IAU Symposium Vol. 267, The Quasar Continuum. pp 55–64
 Elvis M., et al., 1994, ApJS, 95, 1
 Fu H., et al., 2013, Nat, 498, 338
 Fukugita M., Ichikawa T., Gunn J. E., Doi M., Shimasaku K., Schneider D. P., 1996, AJ, 111, 1748
 Fynbo J. P. U., Krogager J.-K., Venemans B., Noterdaeme P., Vestergaard M., Møller P., Ledoux C., Geier S., 2013, ApJS, 204, 6
 Gibson R. R., et al., 2009, ApJ, 692, 758
 Glikman E., et al., 2012, ApJ, 757, 51
 Glikman E., et al., 2013, ApJ, 778, 127
 Glikman E., Gregg M. D., Lacy M., Helfand D. J., Becker R. H., White R. L., 2004, ApJ, 607, 60
 Glikman E., Helfand D. J., White R. L., Becker R. H., Gregg M. D., Lacy M., 2007, ApJ, 667, 673
 Gordon K. D., Clayton G. C., 1998, ApJ, 500, 816
 Greene J. E., et al., 2014, ApJ, 788, 91
 Gregg M. D., Lacy M., White R. L., Glikman E., Helfand D., Becker R. H., Brotherton M. S., 2002, ApJ, 564, 133
 Groves B. A., Heckman T. M., Kauffmann G., 2006, MNRAS, 371, 1559
 Gunn J. E., et al., 1998, AJ, 116, 3040
 Gunn J. E., et al., 2006, AJ, 131, 2332
 Hall P. B., et al., 2002a, ApJ Lett., 575, L51
 Hall P. B., et al., 2002b, ApJS, 141, 267
 Hall P. B., Martini P., Depoy D. L., Gatley I., 1997, ApJ Lett., 484, L17
 Harrison C. M., Alexander D. M., Mullaney J. R., Swinbank A. M., 2014, MNRAS, 441, 3306
 Ho L. C., 2005, ApJ, 629, 680
 Hopkins P. F., et al., 2005, ApJ, 630, 705
 Hopkins P. F., et al., 2006, ApJS, 163, 1
 Ivezić Ž., et al., 2002, AJ, 124, 2364
 Jarrett T. H., et al., 2011, ApJ, 735, 112
 Jones S. F., et al., 2014, MNRAS, 443, 146
 Jones S. F., et al., 2015, MNRAS, 448, 3325

- Jurek R. J., Drinkwater M. J., Francis P. J., Pimblett K. A., 2008, *MNRAS*, 383, 673
- Kauffmann G., et al., 2003, *MNRAS*, 346, 1055
- Keeton C. R., 2001, *ArXiv Astrophysics e-prints*
- Keeton C. R., Winn J. N., 2003, *ApJ*, 590, 39
- Kellermann K. I., Sramek R., Schmidt M., Shaffer D. B., Green R., 1989, *AJ*, 98, 1195
- Kennicutt Jr. R. C., 1998, *ARA&A*, 36, 189
- Kewley L. J., Dopita M. A., 2002, *ApJS*, 142, 35
- Kewley L. J., Geller M. J., Jansen R. A., 2004, *AJ*, 127, 2002
- Khachikian E. Y., Weedman D. W., 1974, *ApJ*, 192, 581
- Kirkpatrick A., et al., 2012, *ApJ*, 759, 139
- Kirkpatrick J. D., et al., 2011, *ApJS*, 197, 19
- Kocevski D. D., et al., 2012, *ApJ*, 744, 148
- Kratzer R. M., Richards G. T., 2015, *AJ*, 149, 61
- Krawczyk C. M., Richards G. T., Mehta S. S., Vogeley M. S., Gallagher S. C., Leighly K. M., Ross N. P., Schneider D. P., 2013, *ApJS*, 206, 4
- Krawczyk C. M., Richards G. T., Mehta S. S., Vogeley M. S., Gallagher S. C., Leighly K. M., Ross N. P., Schneider D. P., 2015, *ApJS*, 999, 4
- Lacy M., et al., 2004, *ApJS*, 154, 166
- Lawrence A., et al., 2007, *MNRAS*, 379, 1599
- Lee K.-G., et al., 2013, *AJ*, 145, 69
- Liu G., Zakamska N. L., Greene J. E., Nesvadba N. P. H., Liu X., 2013, *MNRAS*, 436, 2576
- Liu X., Shen Y., Strauss M. A., Greene J. E., 2010, *ApJ*, 708, 427
- Maddox N., Hewett P. C., Péroux C., Nestor D. B., Wisotzki L., 2012, *MNRAS*, 424, 2876
- Maddox N., Hewett P. C., Warren S. J., Croom S. M., 2008, *MNRAS*, 386, 1605
- Mainzer A., et al., 2011, *ApJ*, 731, 55
- Marchesini D., et al., 2007, *ApJ*, 656, 42
- Marchesini D., Stefanon M., Brammer G. B., Whitaker K. E., 2012, *ApJ*, 748, 126
- Martínez-Sansigre A., Rawlings S., Lacy M., Fadda D., Jarvis M. J., Marleau F. R., Simpson C., Willott C. J., 2006, *MNRAS*, 370, 1479
- Matsuoka Y., et al., 2015, *ArXiv e-prints*
- Melbourne J., et al., 2009, *AJ*, 137, 4854
- Melbourne J., et al., 2012, *AJ*, 143, 125
- Mostek N., Coil A. L., Moustakas J., Salim S., Weiner B. J., 2012, *ApJ*, 746, 124
- Moustakas J., Kennicutt Jr. R. C., Tremonti C. A., 2006, *ApJ*, 642, 775
- Mullaney J. R., Alexander D. M., Goulding A. D., Hickox R. C., 2011, *MNRAS*, 414, 1082
- Nakos T., et al., 2009, *Astron. & Astrophys.*, 494, 579
- Norman C., et al., 2002, *ApJ*, 571, 218
- Oke J. B., Gunn J. E., 1983, *ApJ*, 266, 713
- Padovani P., Perlman E. S., Landt H., Giommi P., Perri M., 2003, *ApJ*, 588, 128
- Pâris I., et al., 2014, *Astron. & Astrophys.*, 563, A54
- Peth M. A., Ross N. P., Schneider D. P., 2011, *AJ*, 141, 105
- Polletta A., et al., 2008, *ApJ*, 675, 960
- Pope A., et al., 2008, *ApJ*, 689, 127
- Reyes R., et al., 2008, *AJ*, 136, 2373
- Richards G. T., et al., 2002, *AJ*, 123, 2945
- Richards G. T., et al., 2003, *AJ*, 126, 1131
- Richards G. T., et al., 2006, *ApJS*, 166, 470
- Richards G. T., et al., 2009, *AJ*, 137, 3884
- Rosario D. J., et al., 2015, *Astron. & Astrophys.*, 573, A85
- Roseboom I. G., Lawrence A., Elvis M., Petty S., Shen Y., Hao H., 2013, *MNRAS*, 429, 1494
- Ross N. P., et al., 2012, *ApJS*, 199, 3
- Sanders D. B., Soifer B. T., Elias J. H., Madore B. F., Matthews K., Neugebauer G., Scoville N. Z., 1988, *ApJ*, 325, 74
- Schlegel D. J., Finkbeiner D. P., Davis M., 1998, *ApJ*, 500, 525
- Schneider D. P., et al., 2010, *AJ*, 139, 2360
- Shen Y., et al., 2011, *ApJS*, 194, 45
- Smee S. A., et al., 2013, *AJ*, 146, 32
- Smith J. A., et al., 2002, *AJ*, 123, 2121
- Soifer B. T., Helou G., Werner M., 2008, *ARA&A*, 46, 201
- Souchay J., et al., 2009, *Astron. & Astrophys.*, 494, 799
- Stern D., et al., 2002, *ApJ*, 568, 71
- Stern D., et al., 2005, *ApJ*, 631, 163
- Stern D., et al., 2012, *ApJ*, 753, 30
- Stoughton C., et al., 2002, *AJ*, 123, 485
- Trump J. R., et al., 2006, *ApJS*, 165, 1
- Tsai C.-W., et al., 2015, *ApJ*, 805, 90
- Urrutia T., Lacy M., Becker R. H., 2008, *ApJ*, 674, 80
- Vanden Berk D. E., et al., 2001, *AJ*, 122, 549
- Vardanyan V., Weedman D., Sargsyan L., 2014, *ApJ*, 790, 88
- Veilleux S., et al., 2013, *ApJ*, 764, 15
- Veilleux S., Osterbrock D. E., 1987, *ApJS*, 63, 295
- Villforth C., et al., 2014, *MNRAS*, 439, 3342
- Wardlow J. L., et al., 2013, *ApJ*, 762, 59
- Wei P., Shang Z., Brotherton M. S., Cales S. L., Hines D. C., Dale D. A., Ganguly R., Canalizo G., 2013, *ApJ*, 772, 28
- Weingartner J. C., Draine B. T., 2001, *ApJ*, 548, 296
- Weymann R. J., Morris S. L., Foltz C. B., Hewett P. C., 1991, *ApJ*, 373, 23
- White M., Zheng Z., Brown M. J. I., Dey A., Jannuzi B. T., 2007, *ApJ Lett.*, 655, L69
- White R. L., Helfand D. J., Becker R. H., Glikman E., de Vries W., 2007, *ApJ*, 654, 99
- Winn J. N., Kochanek C. S., Keeton C. R., Lovell J. E. J., 2003, *ApJ*, 590, 26
- Winn J. N., Lovell J. E. J., Chen H.-W., Fletcher A. B., Hewitt J. N., Patnaik A. R., Schechter P. L., 2002, *ApJ*, 564, 143
- Wright E. L., et al., 2010, *AJ*, 140, 1868
- Wu J., et al., 2012, *ApJ*, 756, 96
- Wu X., Jia Z., 2010, *MNRAS*, 406, 1583
- Wu X.-B., Wang R., Schmidt K. B., Bian F., Jiang L., Fan X., 2011, *AJ*, 142, 78
- Wu X.-B., Zuo W., Yang J., Yang Q., Wang F., 2013, *AJ*, 146, 100
- Yan L., et al., 2013, *AJ*, 145, 55
- York D. G., et al., 2000, *AJ*, 120, 1579
- Zakamska N. L., et al., 2003, *AJ*, 126, 2125
- Zakamska N. L., et al., 2005, *AJ*, 129, 1212
- Zakamska N. L., et al., 2006, *AJ*, 132, 1496
- Zakamska N. L., et al., 2015
- Zakamska N. L., Greene J. E., 2014, *MNRAS*, 442, 784