Citation for published version:

Arrowsmith, M, Hill, MS, Johnson, AL, Kociok-Kohn, G & Mahon, MF 2015, 'Attenuated organo-magnesium activation of white phosphorus', *Angewandte Chemie-International Edition*, vol. 54, no. 27, pp. 7882-7885.
<https://doi.org/10.1002/anie.201503065>

DOI:

[10.1002/anie.201503065](https://doi.org/10.1002/anie.201503065)

Publication date:

2015

Document Version

Early version, also known as pre-print

[Link to publication](#)

This is the peer reviewed version of the following article: Arrowsmith, M. , Hill, M. S., Johnson, A. L., Kociok Köhn, G. and Mahon, M. F. (2015), Attenuated Organomagnesium Activation of White Phosphorus. *Angew. Chem. Int. Ed.*, 54: 7882-7885. doi:10.1002/anie.201503065, which has been published in final form at <https://doi.org/10.1002/anie.201503065>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Attenuated Organo-magnesium Activation of White Phosphorus

Dr Merle Arrowsmith, Prof. Michael S. Hill,* Dr Andrew L. Johnson, Dr Gabriele Kociok-Köhn and Dr Mary F. Mahon*

Department of Chemistry, University of Bath, Claverton Down, Bath, BA2 7AY, UK

Email: msh27@bath.ac.uk

*Sequential reactions between a 2,6-di-iso-propylphenyl-substituted β -diketiminato magnesium *n*-butyl derivative and P_4 allow the highly discriminating synthesis of unusual $[nBu_2P_4]^{2-}$ and $[nBu_2P_8]^{2-}$ cluster dianions.*

Keywords: cluster compounds · magnesium · P_4 activation · phosphorus · structure elucidation

White phosphorus is the starting point in the synthesis of many organophosphanes and phosphorus-containing molecules. As a consequence, the controlled activation of P_4 has attracted considerable attention.^[1] While notable metal-free examples of P_4 activation are provided by Bertrand's use of nucleophilic *N*-heterocyclic carbenes for sequential P-P activation of the tetrahedral molecule,^[2] a majority of approaches have utilized either reactive transition metal or main group reagents. Typical transition metal-based activation can induce not only the cleavage of bonds within P_4 but also the aggregation of smaller fragments to form P_n ($n > 4$) cluster anions. Cummins, for example, has shown that disproportionation of a niobium(IV) tris-enolate in the presence of P_4 provides the $\{P_8\}$ cluster compound **1**,^[3] while in a very recent report, Bergman and Arnold have described a trinuclear β -diketiminato μ - P_{12} niobium complex, which was isolated as one of multiple minor by-products during the reaction of the Nb(V) complex $(^{Dipp}BDI)Nb(N^tBu)Me_2$ ($^{Dipp}BDI = HC\{C(Me)_2N(2,6-i-Pr_2C_6H_3)\}_2$) with H_2 in the presence of P_4 .^[4] Notably, this latter reaction was reported to yield a compound containing a *cyclo*- P_4 dianion as its predominant reaction product and, in some respects, the irrationality of the formation both of these compounds is typical of the synthesis of high nuclearity $\{P_n\}$ clusters.^[1f]

1 (Ad = adamantylidene; Mes = 2,4,6-Me₃C₆H₂)

2 (Dipp = 2,6-*i*-Pr₂C₆H₃)

3 (Dipp = 2,6-*i*-Pr₂C₆H₃)

4

5 (Mes* = 2,4,6-*t*-Bu₃C₆H₂)

6 (a Ar = Mes*
b Ar = 2,6-Mes₂C₆H₃)

Although reductive chemistry to yield *cyclo*-P₄ anions has also been described through the formal oxidative addition of P₄ to low oxidation state Group 13 and 14 derivatives (for example, compounds **2** and **3**)^[1c,5,6] and there is precedent for the activation of P₄ through its interaction with transition metal hydrides and alkyls,^[7] well defined reactions with anionic Group 14 derivatives of elements from Groups 1 and 2 are uncommon and dominated by the use of highly sterically demanding carbon- or silicon-reagents. Wright and co-workers have described the synthesis of the [R₂P₈]²⁻ cluster dianion (**4**) derived from a {P₇}³⁻ Zintl ion core through the reaction of the very bulky hypersilyl potassium reagent with P₄.^[8] The disubstituted butterfly compound **5** (Mes* = 2,4,6-*t*-Bu₃C₆H₂) and the diphosphene Mes*P=P Mes* were similarly isolated from the reaction of Mes*Li and P₄ in the presence of Mes*Br.^[9] Production of compound **5** presumably occurs through initial addition of Mes*Li across a P-P bond within the P₄ unit and the intermediacy of the consequent organopolyposphorus anion. The existence of species of this latter type has very recently been confirmed by Lammertsma and co-workers who reported that the B(C₆F₅)₃-stabilized lithium bicyclo[1.1.0]tetraphosphabutane species **6a** and **6b** could be isolated from reactions of P₄ and sterically demanding lithium aryls in the presence of the borane Lewis acid.^[10] Although similar but indiscriminate reactivity was posited some 50 years ago during reactions with simple Grignard reagents,^[11] the activation of white phosphorus by nucleophilic Group 2 reagents is even less widespread as that by the alkali metals. Our own research interests lie in the development of a defined bond activation and homogeneous catalytic chemistry for the Earth-abundant and environmentally benign alkaline earth

elements.^[12] As a first step toward the incorporation of white phosphorus in Group 2-derived catalysis, we present in this contribution the results of our initial studies which indicate that remarkable kinetic control during P₄ activation is achievable with organomagnesium reagents (Scheme 1). Key to the success of this chemistry is the application of the sterically demanding ^{Dipp}BDI supporting ligand employed in the syntheses of Arnold and Bergman's aforementioned niobium systems,^[4] compounds **2** and **3** and very recently described binuclear cobalt(I) and copper(I) species containing a P₄-derived tetraphospacyclobutadiene and an unactivated P₄ ligand respectively.^[13]

Scheme 1: Synthesis of the complexes described herein.

An initial reaction between an equimolar quantity of P₄ and the β -diketiminato *n*-butylmagnesium complex, [(^{Dipp}BDI)Mg*n*-Bu] (**7**), resulted in the smooth consumption of the magnesium starting material to provide a single new species (**8**) characterized by two mutually coupled doublet resonances at δ –25.6 and –195.5 ppm (¹J_{P-P} = 99.6 Hz) in the resultant ³¹P{¹H} NMR spectrum. Notably, although compound **8** formed with only 50% consumption of the P₄, some evidence of further onward reaction was apparent on continued monitoring (*vide infra*). Repetition of this reaction with the reagent quantities adjusted to reflect the apparent 2:1 stoichiometry resulted in the complete consumption of both P₄ and **7** and the sole production of compound **8**, which was also observed to comprise a single set of temperature invariant (from –90 to +90 °C) *n*-butyl and (^{Dipp}BDI) ligand environments in the corresponding ¹H and ¹³C NMR spectra. The origin of these observations was resolved through the isolation of colorless single crystals of compound **8** suitable for X-ray diffraction analysis from *n*-hexane solution at low temperature, the results of which are displayed in

Figure 1 with selected bond length and angle data presented in the figure caption. The molecular structure of **8** comprises a dinuclear magnesium complex of an unprecedented non-planar $[n\text{Bu}_2\text{P}_4]^{2-}$ dianion which may be considered as arising from the formal addition of two magnesium *n*-butyl fragments to, and the consequent activation of two P-P single bonds within, the P_4 tetrahedron. The P-C bonds to the *n*-butyl chains are located on the adjacent P(1) and P(2) atoms while the magnesium centers each coordinate to one of these alkylated phosphorus atoms and the phosphorus atom located at the opposite diagonal corners of the tetraphosphorus unit. The adoption of a square array is prevented by a distinct pyramidalization at each phosphorus center such that the P-P-P angles are compressed to ca. 80° . The P(3)-P(4) bond length [2.2647(7) Å] is significantly elongated in comparison to the interatomic separation between the alkylated phosphorus atoms [P(1)-P(2) 2.1762(7) Å] but is comparable to the P-P distances observed in molecules containing the $[\text{P}_4]^{4-}$ anion such as compound **2** [mean 2.29 Å].^[5] The Mg-P distances to P(3) and P(4) [Mg(1)-P(3) 2.5625(7), Mg(2)-P(4) 2.5592(7) Å] are also shorter than those to P(1) and P(2) [Mg(1)-P(1) 2.6336(7), Mg(2)-P(2) 2.6652(7) Å] and we interpret both of these structural features to indicate that the distribution of negative charge is polarized toward the non-alkylated P(3) and P(4) centers and that the $[n\text{Bu}_2\text{P}_4]^{2-}$ unit as a whole may be considered as derived from the formal addition of two *n*-butyl carbenium ions to the $[\text{P}_4]^{4-}$ constituent anion of compound **2**.

Figure 1: ORTEP representation (30% probability ellipsoids) of compound **8**. Hydrogen atoms and *iso*-propyl groups are removed for clarity. Selected bond lengths (Å) and angles (°): Mg(1)-N(1) 2.0462(15), Mg(1)-N(2) 2.0465(15), Mg(1)-P(3) 2.5625(7), Mg(1)-P(1) 2.6336(7), Mg(1)-P(4) 3.1607(7), Mg(2)-N(4) 2.0424(15), Mg(2)-N(3) 2.0466(15), Mg(2)-P(4) 2.5592(7), Mg(2)-P(2) 2.6652(7), P(1)-C(59) 1.8584(19), P(1)-P(2) 2.1762(7), P(1)-P(4) 2.2127(7), P(2)-C(63) 1.8575(19), P(2)-P(3) 2.2108(7), P(3)-P(4) 2.2647(7), N(1)-Mg(1)-P(3) 126.57(5), N(2)-Mg(1)-P(3) 122.72(5), N(1)-Mg(1)-P(1) 127.76(5), N(2)-Mg(1)-P(1) 122.16(5), P(3)-Mg(1)-P(1) 67.14(2), N(1)-Mg(1)-P(4) 167.48(5), N(2)-Mg(1)-P(4) 99.06(5), P(3)-Mg(1)-P(4) 45.131(17), P(1)-Mg(1)-P(4) 43.731(16), N(4)-Mg(2)-N(3) 93.52(6), N(4)-Mg(2)-P(4) 126.28(5), N(3)-Mg(2)-P(4) 121.83(5), N(4)-Mg(2)-P(2) 121.40(5), N(3)-Mg(2)-P(2) 130.79(5), P(4)-Mg(2)-P(2) 66.130(19).

The previously noted onward reaction of compound **8** and additional P₄ (*vide supra*) prompted us to carry out a further reaction of an equimolar quantity of P₄ and compound **8**. Reaction in benzene solution at room temperature provided the slow and simultaneous consumption of the starting materials and direct conversion to a single new compound (**9**) characterized by the emergence of eight mutually coupled signals in the ³¹P NMR spectrum. The corresponding ¹H and ¹³C NMR spectra indicated that this new compound comprised two differentiated groups of resonances indicative of two separate *n*-butyl and (^{Dipp}BDI) ligand environments. Compound **9** could also be synthesized directly through the slow (7 days) reaction of compound **7** with an equimolar amount of P₄ at room temperature or by heating of

a similar reaction at 60°C for 2 days. In this latter case compound **9** crystallized as colorless single crystals suitable for a further X-ray diffraction analysis, allowing the resolution of the solution NMR spectroscopic observations. The result of this analysis is shown in Figure 2 and highlights that the disposition of the phosphorus and magnesium centers within compound **9** is consistent with the coupling pattern observed in the solution-state ^{31}P NMR spectrum and the multiplicity of signals arising in the ^1H and ^{13}C NMR data. The structure comprises two ($^{\text{Dipp}}\text{BDI}$) magnesium units bonded to a $[\text{n-Bu}_2\text{P}_8]^{2-}$ cluster dianion derived from the formal reductive coupling of two neutral P_4 units. Although the rationality of this $\{\text{P}_8\}$ cluster synthesis from the combination of two $\{\text{P}_4\}$ fragments appears to be unique, its structure may be considered as a heptaphosphanorbornane fused to an *exo*-oriented cyclotriphosphane ring and a constitutional isomer of the $\{\text{P}_8\}$ cage within the niobium(V) species $[\text{Ph}_2\text{CP}_8\text{Nb}(\text{OC}[\text{}^2\text{Ad}]\text{Mes})_3]$, which was derived through treatment of compound **1** with benzophenone.^[14] The single *tris*-enolato niobium dication and the $\{\text{Ph}_2\text{C}\}$ unit of this previously described species is replaced in the present case by two ($^{\text{Dipp}}\text{BDI}$) magnesium monocations and two *n*-butyl substituents. The magnesium centers contact the $\{\text{P}_8\}$ cage structure in *endo* [Mg(1)] and *exo* [Mg(2)] configurations with respect to the P(4) bridgehead and provide Mg-P distances of 2.575(3) Å [P(4)-Mg(2)], 2.625(3) Å [P(8)-Mg(2)], 2.594(3) Å [P(2)-Mg(1)] and 2.831(3) Å [P(7)-Mg(1)]. Although these latter values indicate some variability in the charge donation across the cage structure, it should be noted that the Mg-P contacts are all within the range expected [2.45 to 2.985 Å] from a search of the Cambridge structural database.^[15] All of the P-P distances within this new cluster are also consistent with P-P single bonds and range between 2.096(3) Å [P(3)-P(4)] and 2.262(4) Å [P(5)-P(6)].

Figure 2: ORTEP representation (30% probability ellipsoids) of compound **9**. Hydrogen atoms, *iso*-propyl methyl groups and *n*-butyl groups aside from the P-bonded C(59) and C(63) atoms are removed for clarity. Selected bond lengths (Å) and angles (°): P(1)-P(5) 2.195(3), P(2)-P(3) 2.160(2), P(2)-Mg(1) 2.594(3), P(3)-P(4) 2.096(3), P(3)-P(7) 2.186(3), P(4)-P(5) 2.222(3), P(4)-Mg(2) 2.575(3), P(5)-P(6) 2.262(4), P(6)-P(7) 2.170(4), P(6)-P(8) 2.185(3), P(7)-P(8) 2.153(3), P(7)-Mg(1) 2.831(3), P(8)-Mg(2) 2.625(3) Mg(1)-N(2) 2.046(5), P(2)-P(1)-P(5) 105.23(12), P(1)-P(2)-P(3) 96.79(11), P(1)-P(2)-Mg(1) 90.77(11) P(3)-P(2)-Mg(1) 97.47(9), P(4)-P(3)-P(7) 113.74(12), P(2)-P(3)-P(7) 97.95(10), P(3)-P(4)-P(5) 88.12(10), P(3)-P(4)-Mg(2) 101.38(10), P(5)-P(4)-Mg(2) 91.12(11), P(1)-P(5)-P(4) 102.66(14), P(1)-P(5)-P(6) 92.00(12), P(4)-P(5)-P(6), 105.80(12), P(7)-P(6)-P(8) 59.27(11), P(7)-P(6)-P(5) 106.69(12).

Subsequent preliminary investigations indicated that the kinetic control exerted during the syntheses of compounds **8** and **9** is an apparent consequence of the steric demands of both the supporting (^{Dipp}BDI) platform and the reactive *n*-butyl co-ligand. Analysis by ³¹P NMR spectroscopy of a reaction between [(^{Dipp}BDI)MgH]₂ and half a molar equivalent of P₄ provided a mixture of several phosphorus-containing species. Although a minor component comprised broadened resonances at δ −3.5 and −159 ppm reminiscent of those observed for compound **8**, the major product of this reaction was characterized by a triplet signal observed at δ −284.8 ppm (¹J_{P-H} = 45 Hz). This latter species was identified as the trimeric primary magnesium phosphide, compound **10**, through a further X-ray analysis performed on single crystals produced by fractional crystallization of the reaction mixture (Figure 3a). Analytically pure samples of compound **10**, notable as the first primary phosphide of a Group 2 element to have been structurally characterized, were found to be unstable once redissolved

in toluene- d_8 . Monitoring by ^1H and ^{31}P NMR spectroscopy at room temperature evidenced the onset of a redistributive process to as yet unidentified magnesium-containing species and the production of PH_3 , which was observed as a binomial quartet at $\delta -245.0$ ppm in the ^{31}P NMR spectrum.

In a similar manner, monitoring by NMR spectroscopy of a reaction analogous to that employed in the synthesis of compound **8** but with replacement of the $N\text{-}\{2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3\}$ groups of the ($^{\text{Dipp}}\text{BDI}$) ligand of **7** by less sterically demanding $N\text{-}\{2,4,6\text{-Me}_3\text{C}_6\text{H}_2\}$ substituents ($^{\text{Mes}}\text{BDI}$), indicated less discriminating behavior and reactivity which again resulted in the formation of multiple products. A further reaction performed between $[(^{\text{Mes}}\text{BDI})\text{Mg}n\text{-Bu}]$ and P_4 in a 2:3 molar ratio again provided a complex mixture containing at least three new phosphorus-containing compounds. Fractional crystallization from this solution and a resultant single crystal X-ray diffraction analysis of compound **11** (Figure 3b) provided some insight into the nature of these processes. Compound **11** may be viewed as a $[\text{P}_7]^{3-}$ Zintl ion cage reminiscent of compound **4** but decorated with three β -diketiminato magnesium units. We suggest, therefore, that the formation of this phosphorus cage constitutes a thermodynamic sink during the solution comproportionation of magnesium polyphosphorus species of varying nuclearity.^[16]

Figure 3: ORTEP representations (30% probability ellipsoids) of (a) compound **10** with hydrogen atoms except those attached to P(1), P(2) and P(3) and *iso*-propyl methyl groups removed for clarity; (b) compound **11** with hydrogen atoms and mesityl methyl groups removed for clarity.

In conclusion we have shown that the controlled, sequential organomagnesium-mediated activation of white phosphorus may be achieved through selection of an appropriate combination of supporting and reactive co-ligands. We are continuing to explore the parameters of this reactivity and will report our findings in future publications.

Methods

Details of the synthesis, characterization data and the crystallographic protocols employed in this study are given in the Supporting Information. CCDC 1057657 - 1057660 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Center via www.ccdc.ca.ac.uk/data_request/cif.

Acknowledgement

We thank the EPSRC (UK) for funding.

References

1. For reviews, see (a) M. Scheer, G. Balázs, A. Seitz, *Chem. Rev.* **2010**, *110*, 4236-4256; (b) B. M. Cossairt, N. A. Piro, C. C. Cummins, *Chem. Rev.* **2010**, *110*, 4164-4177; (c) S. Khan, S. S. Sen, H. W. Roesky, *Chem. Commun.* **2012**, *48*, 2169-2178; (d) M. Serrano-Ruiz, A. Romerosa, P. Lorenzo-Luis, *Eur. J. Inorg. Chem.* **2014**, 1587-1598; (e) M. Caporali, L. Gonsalvi, A. Rossin, M. Peruzzini, *Chem. Rev.* **2010**, *110*, 4178-4235; (f) M. Ehses, A. Romerosa, M. Peruzzini, *Top. Curr. Chem.* **2002**, *220*, 107-140.
- [2] (a) J. D. Masuda, W. W. Schoeller, B. Donnadieu, G. Bertrand, *Angew. Chem. Int. Ed.* **2007**, *46*, 7052-7055; *Angew. Chem.* **2007**, *119*, 7182-7185; (b) O. Back, G. Kuchenbeiser, B. Donnadieu, G. Bertrand, *Angew. Chem. Int. Ed.* **2009**, *48*, 5530-5533; *Angew. Chem.* **2009**, *121*, 5638-5641; (c) C. D. Martin, C. M. Weinstein, C. E. Moore, A. L. Rheingold, G. Bertrand, *Chem. Commun.* **2013**, *49*, 4486-4488.
- [3] (a) B. M. Cossairt, C. C. Cummins, *Angew. Chem. Int. Ed.* **2008**, *47*, 169-172; *Angew. Chem.* **2008**, *120*, 175-178; (b) B. M. Cossairt, C. C. Cummins, *Angew. Chem. Int. Ed.* **2008**, *47*, 8863-8866; *Angew. Chem.* **2008**, *120*, 8995-8998.
- [4] C. Camp, L. Maron, R. G. Bergman, J. Arnold, *J. Am. Chem. Soc.* **2014**, *136*, 17652-17661.
- [5] Y. Peng, H. Fan, H. Zhu, H. W. Roesky, J. Magull, C. Hughes, *Angew. Chem. Int. Ed.* **2004**, *43*, 3443-3445; *Angew. Chem.* **2004**, *116*, 3525-3527.

- [6] Y. Xiong, S. Yau, M. Brym, M. Driess, *Angew. Chem. Int. Ed.* **2008**, *47*, 8863-8866; *Angew. Chem.* **2007**, *119*, 4595–4597.
- [7] See, for example; P. Barbaro, A. Ienco, C. Mealli, M. Peruzzini, O. J. Scherer, G. Schmitt, F. Vizza, G. Wolmershäuser, *Chem. Eur. J.* **2003**, *9*, 5195-5210.
- [8] W. T. K. Chan, F. García, A. D. Hopkins, L. C. Martin, M. McPartlin, D. S. Wright, *Angew. Chem. Int. Ed.* **2007**, *46*, 3084–3086; *Angew. Chem.* **2007**, *119*, 3144–3146.
- [9] R. Riedel, H.-D. Hausen, E. Fluck, *Angew. Chem. Int. Ed. Engl.* **1985**, *24*, 1056-1057; *Angew. Chem.* **1985**, *97*, 1050-1050.
- [10] J. E. Borger, A. W. Ehlers, M. Lutz, J. C. Slootweg, K. Lammertsma, *Angew. Chem. Int. Ed. Engl.* **2014**, *53*, 12836-12839; *Angew. Chem.* **2014**, *126*, 13050–13053.
- [11] (a) M. M. Rahut, A. M. Semsel, *J. Org. Chem.* **1963**, *28*, 471-473; (b)) M. M. Rahut, A. M. Semsel, *J. Org. Chem.* **1963**, *28*, 473-478.
- [12] For reviews, see; (a) M. Arrowsmith, M. S. Hill, ‘*Alkaline Earth Chemistry: Applications in Catalysis*’, *Comprehensive Inorganic Chemistry II*. ed. T. Chivers, Elsevier, **2013**, *1*, 1189; (b) A. G. M. Barrett, M. R. Crimmin, M. S. Hill, P. A. Procopiou, *Proc. R. Soc. A*, **2010**, *466*, 927; (c) M. R. Crimmin, M. S. Hill, *Topics in Organometallic Chemistry*, ed. S. Harder, **2013**, *45*, 191; (d) S. Harder, *Chem. Rev.*, **2010**, *110*, 3852.
- [13] (a) S. Yao, N. Lindenmaier, Y. Xiong, S. Inoue, T. Szilvási, M. Adelhardt, J. Sutter, K. Meyer, M. Driess, *Angew. Chem. Int. Ed. Engl.* **2015**, *54*, 1250-1254; *Angew. Chem.* **2015**, *127*, 1266–1270; (b) F. Spitzer, M. Sierka, M. Latronico, A. V. Virovets, M. Scheer, *Angew. Chem. Int. Ed.* **2015**, *54*, 4392–4396; *Angew. Chem.* **2015**, *127*, 4467–4472.
- [14] B. M. Cossairt, C. C. Cummins, *Inorg. Chem.* **2008**, *47*, 9363-9371.
- [15] CSD search performed 31/3/15.
- [16] R. S. P. Turbervill, J. M. Goicoechea, *Chem. Rev.* **2014**, *114*, 10807-10828.

For Table of Contents:

Taking the P: Sequential reactions between a 2,6-di-iso-propylphenyl-substituted β-diketiminato magnesium n-butyl derivative and P₄ allow the highly discriminating synthesis of unusual [nBu₂P₄]²⁻ and [nBu₂P₈]²⁻ cluster dianions.