Citation for published version:

Chapman, R, Lawrence, R, Williams, J & Bull, S 2017, 'Formyloxyacetoxypheylmethane as an *N*-Formylating Reagent for Amines, Amino Acids, and Peptides', pp. 4908-4911. <https://doi.org/10.1021/acs.orglett.7b02382>

DOI:

[10.1021/acs.orglett.7b02382](https://doi.org/10.1021/acs.orglett.7b02382)

Publication date:

2017

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Formyloxyacetoxyphenylmethane as an *N*-Formylating Reagent for Amines, Amino Acids and Peptides

Robert. S. L. Chapman,[‡] Ruth Lawrence,[‡] Jonathan M. J. Williams and Steven. D. Bull*

Department of Chemistry, University of Bath, Bath, BA2 7AY, UK.

ABSTRACT: Formyloxyacetoxyphenylmethane is a stable, water-tolerant, *N*-formylating reagent for primary and secondary amines that can be used under solvent-free conditions at room temperature to prepare a range of *N*-formamides, *N*-formyl-anilines, *N*-formyl- α -amino-acids, *N*-formyl-peptides and an isocyanide.

N-Formylation reactions of primary and secondary amines are important transformations in organic chemistry, because they provide direct access to the highly versatile formamide group.¹ Highlighting their synthetic utility, *N*-formyl derivatives can be used as precursors for the synthesis of formamidines,² ureas,³ carbamates,³ aryl amides,⁴ isocyanates,⁵ nitriles⁶ and isocyanides.⁷ Consequently, they have often been used as intermediates for the synthesis of pharmaceuticals.⁸ Alternatively, the *N*-formyl group may be used as a protecting group for peptide synthesis,⁹ whilst many naturally occurring formamides exhibit important medicinal and biological activities.¹⁰ The formamide moiety is also present in pharmaceuticals, including Leucovorin (chemotherapeutic),^{11a} Formoterol (asthma)^{11b} and Orlistat (anti-obesity).^{11c}

Due to their importance, a number of *N*-formylating reagents for primary and secondary amines have been developed,¹ including the use of formic acid,^{12a} formic acid in the presence of coupling reagents (e.g. DCC),^{12b} acetic formic anhydride,^{12c} formate esters,^{12d} formamides,^{12e} trialkylorthoformates,^{12f} enol formates,^{12g} ammonium formate,^{12h} and *N*-formyl-benzotriazole.¹²ⁱ However, many of these reagents are only moderately reactive towards sterically hindered or electronically deactivated amines, often requiring elevated temperatures, extended reaction times or the use of excess reagent for their *N*-formylation reactions to proceed to completion.¹ A number of these formylating reagents are also susceptible to hydrolysis, whilst some reagents can decompose to liberate hazardous by-products (e.g. C=O).¹ Given these limitations, a number of catalytic *N*-formylation protocols have also been developed that employ *N*-formyl donors in combination with acid catalysts, Lewis acids, organocatalysts, metal catalysts and biocatalysts.¹³

Given the limitations of using many *N*-formylating reagents under catalyst free conditions, we now report herein that formyloxyacetoxyphenylmethane (FAPM) **5** can be used as a bench-stable formylating reagent for primary/secondary amines under solvent-free conditions at room temperature. Importantly, the stability of FAPM **5** towards hydrolysis, enables it to be used

as a catalyst free *N*-formylating reagent for α -amino acids and peptides in aqueous based solvent systems.

As part of investigations into the Baeyer-Villiger oxidation reactions of α,β -unsaturated ketones, we found that treatment of benzylidene acetone **1** with 4 equiv. of *meta*-chloroperoxybenzoic acid (*m*CPBA) in toluene for 24 h gave FAPM **5** in 84% isolated yield (Scheme 1a).¹⁴ In this remarkable one-pot 4-step reaction, benzylidene acetone **1** first undergoes an *m*CPBA mediated Baeyer-Villiger reaction to afford enol acetate **2**, whose electron-rich alkene bond is then epoxidised by a second equivalent of *m*CPBA to afford epoxy-acetate **3**. This unstable epoxy-acetate **3** then undergoes intramolecular rearrangement to afford aldehyde **4**, which then undergoes a further *m*CPBA mediated Baeyer-Villiger reaction to afford FAPM **5**. The formation of intermediates **2** and **4** in this oxidation reaction were confirmed by ¹H NMR spectroscopic analysis of a series of control reactions that were carried out using fewer equivalents of *m*CPBA over shorter periods of time (see SI for details).¹⁵

A second route to FAPM **5** was also developed using a variant of a protocol previously developed by Raphaeli and coworkers for the synthesis of non-symmetric acyloxyalkyl esters.¹⁶ This involved treatment of benzaldehyde with 1.5 equiv. of acetyl chloride in the presence of a catalytic amount of *para*-toluene sulfonic acid to afford α -chlorobenzyl acetate **6**,¹⁷ that was subsequently reacted with formic acid and Et₃N in acetone to afford FAPM **5** in 76% yield (Scheme 1b).¹⁸

FAPM **5** proved to be a bench-stable liquid that could be stored at room temperature under nitrogen for up to 6 months without decomposition. We reasoned that FAPM **5** might act as a selective *N*-formyl donor for amines, which was confirmed by reacting 1.5 equiv. of FAPM **5** with 1 equiv. of benzylamine **7a** in THF at rt for 4.5 h to give *N*-formyl-benzylamine **8a** and imine **9in** 90% and 10% conversion respectively (Table 1, entry 1). Importantly, ¹H NMR spectroscopic analysis revealed no evidence of any *N*-acetyl-benzylamide having been formed *via* competing *N*-acyl transfer from FAPM **5**. Mechanistically, we propose that

benzylamine **10a** attacks the reactive formyl group of FAPM to irreversibly afford formamide **8a**,

Scheme 1. Syntheses of formyloxyacetoxyphenylmethane **5** via: (a) *m*CPBA mediated oxidation reaction of benzylidene acetone **1**. (b) Nucleophilic attack of formic acid on α -chlorobenzyl acetate **6**.

with benzaldehyde and acetic acid being generated as by-products. Unwanted formation of imine **9** then occurs through acid catalyzed reaction of benzylamine **7a** with the benzaldehyde by-product. A solvent screen was carried out with the aim of identifying conditions that would suppress formation of imine **9**. Use of ethanol and 1,4-dioxane (Table 1, entries 2-3) also afforded significant amounts of the unwanted imine **9** by-product (8-12%). Carrying out the *N*-formylation reaction in MeCN (Table 1, entry 4) gave reduced amounts of imine (4%), whilst CH_2Cl_2 , toluene and EtOAc (Table 1, entries 5-7) gave excellent conversion to *N*-formyl-benzylamine **8a** with only trace amounts of imine **9** ($\leq 2\%$) being formed. Since imine formation is potentially reversible, a series of *N*-formylation reactions were carried out in mixed aqueous solvents, in the hope that the presence of bulk water would perturb the reaction equilibrium away from imine formation. Gratifyingly, use of 1:1 mixtures of THF : H_2O , MeCN : H_2O and 1,4-dioxane : H_2O (Table 1, entries 8-10) gave good conversions to *N*-formyl-benzylamine **8a**, with $\leq 1\%$ of imine **9** being formed in each case. The tolerance of FAPM **5** towards hydrolysis in aqueous systems was confirmed by carrying out an *N*-formylation reaction in water,¹⁹ with 1.5 equiv. of FAPM **5** resulting in complete conversion of benzylamine to *N*-formyl-benzylamine **8a** (Table 1, entry 11). Finally, FAPM **5** could also be used under solvent free conditions (Table 1, entry 12), resulting in clean *N*-formylation of benzylamine, with only 2% of imine **9** being produced.²⁰

¹H NMR spectroscopic analysis of the *N*-formylation reaction of benzylamine in d_8 -toluene revealed that complete consumption of benzylamine **7a** occurred after 1 h. Analysis after 15 min revealed the presence of a 87 : 13 mixture of formamide **8a** : imine **9**, which subsequently equilibrated to a 99 : 1 mixture of formamide **8a** : imine **9** after 1 hr. With this information in hand, we employed 1.5 equiv. of FAPM to *N*-formylate a range of sixteen primary and secondary amines **7a**- to afford a series of *N*-formamides **8a-p** in 53-96% yield after chromatographic purification (Scheme 2). Most of these *N*-formylation reactions were

carried out under solvent-free conditions by dissolving the amine in neat FAPM and stirring at rt, followed by purification by silica chromatography. In those cases where the parent amine was insoluble in FAPM **5**, then EtOAc was employed as a cosolvent to afford a homogeneous reaction mixture. *N*-Formylation reactions of primary aliphatic amines proved particularly facile; typically affording their corresponding formamides **8a-e** in good to excellent yields (81-95%). Pleasingly, amines containing acid sensitive functionalities, such as furan and dimethyl acetal were well tolerated, affording formamides **8b** and

Table 1. Solvent screen to optimize the *N*-formylation reaction of benzylamine with FAPM **5**

Entry	Solvent	Formamide 8a	Imine 9
1	THF	90	10
2	EtOH	88	12
3	Dioxane	92	8
4	MeCN	96	4
5	CH_2Cl_2	100	-
6	Toluene	100	-
7	EtOAc	98	2
8	THF : H_2O (1 : 1)	99	1
9	MeCN : H_2O (1 : 1)	100	-
10	Dioxane : H_2O (1 : 1)	99	1
11	H_2O	100	-
12	Neat	98	2

Product ratios determined from integrals of diagnostic resonances for **8a** and **9** in ¹H NMR spectra of crude reaction products.

8e in 95% and 85% yields respectively. FAPM **5** reacted exclusively with the primary amine functionality of tryptamine to afford formamide **8d** in 90% yield. Whilst the formamide **8f** of sterically congested benzhydrylamine was formed in good 83% yield, formylation of *tert*-butylamine gave *N*-formamide **8g** in only 53% isolated yield, arising from losses during reaction work-up due to its volatility. (*S*)-Valine methyl ester afforded *N*-formamide **8h** in an excellent 93% yield, with no evidence of any racemization having occurred. Acyclic and cyclic secondary amines were also formylated using FAPM **5**, with moderate to excellent 58-96% yields of *N*-formamides **8i-k** being isolated. Pleasingly, electron-deficient aniline and its secondary *N*-methyl derivative gave their corresponding formamides **8l-m** in excellent 90% and 91% yields respectively. However, formylation of anilines containing deactivating halogen or ester groups required extended reaction times of 24 h for their corresponding *N*-formamides **8o-p** to be formed in 87-95% yields respectively.

Alternative work-up procedures were then developed to allow for these *N*-formylation reactions to be carried out on-scale, without the need for chromatographic purification. When the desired formamide product was crystalline, or a high boiling oil, then the acetic acid and benzaldehyde by-products could be removed by fractional distillation *in vacuo*. For more volatile formamides, purification could be achieved by first extracting the crude reaction product with $\text{NaHCO}_3(\text{aq})$ to remove acetic acid, followed by extraction with $\text{NaHSO}_3(\text{aq})$ to remove benzaldehyde. This affords a crude product containing the formamide that is then purified by recrystallisation or fractional distillation as required (See SI for details).

Although several reagents have been developed for the *N*-formylation of amino ester derivatives,²¹ extension of these

methods to unprotected α -amino acids has not been widely reported. This is due to the instability of many formylating reagents towards hydrolysis in aqueous media and the insolubility of α -amino acids in non-aqueous solvents. Currently, *N*-formyl- α -amino acids are most commonly prepared via treatment of the parent α -amino acids with a large excess of Ac_2O and formic acid (via formation of formyl acetate *in situ*), which often requires forcing conditions and/or extended reaction times.²² *N*-formylation of α -amino acids using formic acid in water has been reported, however this protocol requires Oxyma as a stoichiometric coupling reagent.²² Given the observed stability of FAPM in aqueous solvents, a range of six L-amino acids **10a-f** were treated with 1.5 equiv. of FAPM in water containing 5 equiv. of NaHCO_3 as a base, which resulted in 80–85% conversion to their corresponding formamides **11a-f** after 16 h. However, we subsequently found that modifying these formylation reactions, by incorporating addition of a second portion of FAPM **5** (1.5 equiv.) after 8 h, resulted in complete consumption of each α -amino acid, giving *N*-formyl-amino acids **11a-f** in good to excellent 71–89% isolated yields, with no racemization having occurred (Scheme 3).

Scheme 2: FAPM **5** as an *N*-formylating reagent for the synthesis of *N*-formamides **8a-p**

(a) Reaction conditions: 2 mmol of amine, 1.5 equiv. FAPM **5**, 1 h, rt. (b) Reaction conditions: 2 mmol of amine, 1.5 equiv. FAPM **5**, EtOAc, 1 h, rt. (c) Reaction conditions: 2 mmol of amine, 1.5 equiv. FAPM **5**, EtOAc, rt, 24 h.

N-Formylmethionine-leucyl-phenylalanine (f-MLP) **14** is a chemotactic *N*-formyl-tripeptide that is a potent chemoattractant for leukocytes and a macrophage activator that is involved in triggering the innate immune response towards bacterial pathogens,²⁴ that was chosen as a suitable target to test the utility of FAPM for the *N*-formylation of peptidic substrates. The parent tripeptide ester **12** was treated with 1.5 equiv. of FAPM in EtOAc for 5 h to afford formamide **13** in 87% yield, which was then hydrolysed using NaOH/MeOH to afford f-MLP **14** in 93% yield (Scheme 4a). Similarly, a decapeptide with the sequence Ac-

ADGIVNGVKA- NH_2 **15** containing *N*-acetamide and C-primary amide termini, was reacted with 5 equiv. of FAPM in a mixture of $\text{MeCN}/\text{H}_2\text{O}$ (pH 8.0) containing a few drops of DMSO. This resulted in *N*-formylation of the free ω -amino group of its lysine residue affording *N*-formyl-decapeptide **16**, as confirmed from HRMS analysis (Scheme 4b).

Formamides are often employed for the synthesis of isocyanides,⁷ and so we investigated whether FAPM **5** could be used to develop a stepwise 'one-pot' protocol to directly convert an amine into its corresponding isocyanide. 4-Bromoaniline **7n** and FAPM **5** (1.5 equiv.) were stirred in CH_2Cl_2 at rt for 24 h to afford its corresponding formamide **8n** *in situ* (confirmed by ^1H NMR spectroscopic analysis). Sequential dropwise addition of diisopropylamine (4 equiv.) and POCl_3 (1.2 equiv.) to this solution of formamide **8n** in CH_2Cl_2 at 0 °C, followed by stirring at 0 °C for 2 h and rt for 4 h, gave 1-bromo-4-isocyanobenzene **17** in 74% yield over two steps (Scheme 5).

Scheme 3: FAPM **5** as an *N*-formylating reagent for the synthesis of *N*-formyl- α -amino acids **11a-f**.

Scheme 4: FAPM **5** for the synthesis of: (a) *N*-formylmethionyl-leucyl-phenylalanine (f-MLP) **14**; (b) *N*-formyl-decapeptide **16**

Scheme 5: FAPM **5** for the one-pot synthesis of isocyanide **17**

In conclusion, two practical syntheses of formyloxycetoxyphe-nylmethane **5** have been developed and its use as a water toler-

ant *N*-formylating reagent for primary and secondary amines under solvent-free conditions demonstrated. Its potential as a reagent for the synthesis of *N*-formyl- α -amino acids and *N*-formyl-peptides in aqueous systems has been established and it has also been used as an *N*-formylating reagent in a one pot protocol to convert an aniline into its corresponding isocyanide.

ASSOCIATED CONTENT

Complete experimental procedure and relevant spectra (^{13}C and ^1H NMR spectra) for all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>

AUTHOR INFORMATION

Corresponding Author

*E.mail:s.d.bull@bath.ac.uk

Author Contributions

‡These authors contributed equally.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We thank the University of Bath, EPSRC and the Centre for Doctoral Training in Sustainable Chemical Technologies (EP/L016354/1) for funding. We would like to thank Dr Jody Mason for donating a sample of decapeptide **15** that was prepared using conventional solid-phase peptide synthesis.

REFERENCES

- Gerack, C. J.; McElwee-White, L. *Molecules* **2014**, *19*, 7689-7713.
- Han, Y.; Cai, L. S. *Tetrahedron Lett.* **1997**, *38*, 5423-5426.
- Faraj, M. K. *Synthesis of isocyanate precursors from primary formamides*, U.S. Patent 5,686,645, **1997**.
- Jo, Y.; Ju, J.; Choe, J.; Song, K. H.; Lee, S. J. *Org. Chem.* **2009**, *74*, 6358-6361.
- Fu, P. P.; Boyer, J. H. *J. Chem. Soc. Perkin Trans 1* **1974**, 2246-2250.
- Arlt, D.; Klein, G. *Preparation of nitriles from formamides*, U.S. Patent 4419297, **1983**.
- Obrecht, R.; Herrmann, R.; Ugi, I. *Synthesis* **1985**, 400-402.
- Jackson, A.; Meth-Cohn, O. *J. Chem. Soc. Chem. Commun.* **1995**, 1319.
- Yamashiro, D.; Li, C. H. *J. Org. Chem.* **1973**, *38*, 2594-2597.
- Wright, A. D.; Lang-Unnasch, N. *J. Nat. Prod.* **2009**, *72*, 492-495.
- (a) Machover, D. *Cancer* **1997**, *80*, 1179-1187. (b) Hett, R.; Fang, Q. K.; Gao, Y.; Wald, S. A.; Senanayake, C. H. *Org. Process Res. Dev.* **1998**, *2*, 96-99. (c) Ma, G.; Zancanella, M.; Oyola, Y.; Richardson, R. D.; Smith, J. W.; Romo, D. *Org. Lett.* **2006**, *8*, 4497-4500.
- (a) Jung, S. H.; Ahn, J. H.; Park, S. K.; Choi, J. K. *Bull. Korean Chem. Soc.* **2002**, *23*, 149-150. (b) Tornesello, A. L.; Sanseverino, M.; Buonaguro, F. M. *Molecules* **2016**, *21*, 736-742; (c) Strazzolini, P.; Giumanini, A. G.; Cauci, S. *Tetrahedron* **1990**, *46*, 1081-1081. (d) Christensen, S. B.; Hansen, A. M.; Franzky, H. J. *Peptide Sci.* **2017**, *23*, 410-415. (e) Cochet, T.; Bellosta, V.; Greiner, A.; Roche, D.; Cossy, J. *Synlett* **2011**, 1920-1922. (f) Kaboudin, B.; Khodamorady, M. *Synlett* **2010**, 2905-2907. (g) Neveux, M.; Bruneau, C.; Digneuf, P. H. *J. Chem. Soc. Perkin Trans 1* **1991**, 1197-1199. (h) Reddy, P. G.; Kumar, G. D. K.; Baskaran, S. *Tetrahedron Lett.* **2000**, *41*, 9149-9151. (i) Katritzky, A. R.; Chang, H. X.; Yang, B. Z. *Synthesis* **1995**, 503-505.
- (a) Brahmachari, G.; Laskar, S. *Tetrahedron Lett.* **2010**, *51*, 2319-2322. (b) Nguyen, T. B.; Sorres, J.; Tran, M. Q.; Ermolenko, L.; Al-Mourabit, A. *Org. Lett.* **2012**, *14*, 3202-3205. (c) Lei, M.; Ma, L.; Hu, L. *Tetrahedron Lett.* **2010**, *51*, 4186-4188. (d) Habibi, D.; Nasrollahzadeh, M.; Shebekhtari, H. *J. Mol. Catal. A* **2013**, *378*, 148-155. (e) Kim, J. G.; Jang, D. O. *Synlett* **2010**, 2093-2096. (f) Deutsch, J.; Eckelt, R.; Kockritz, A.; Martin, A. *Tetrahedron* **2009**, *65*, 10365-10369. (f) Sonawane, R. B.; Rasal, N. K.; Jagtap, S. V. *Org. Lett.* **2017**, *19*, 2078-2081. (g) Patre, R. E.; Mal, S.; Nilkanth, P. R.; Ghorai, S. K.; Deshpande, S. H.; El Qacemi, M.; Smejkal, T.; Pal, S.; Manjunath, B. N. *Chem. Commun.* **2017**, 53, 2382-2385. (h) Hosseini-Sarvari, M.; Sharghi, H. *J. Org. Chem.* **2006**, *71*, 6652-6654. (h) Allen, C. L.; Atkinson, B. N.; Williams, J. M. J. *Angew. Chem. Int. Ed. Engl.* **2012**, *51*, 1383-1386. (i) Mamani, L.; Sheykhan, M.; Heydari, A.; Faraji, M.; Yamini, Y. *Appl. Cat. A* **2010**, *377*, 64-69.
- For previous reports where peracid mediated Baeyer-Villiger reactions of α,β -unsaturated ketones gave mixtures of products containing formyl-acylals **18-20**, see: (a) Formyl-acylal **18**: Baures, P. W.; Eggleston, D. S.; Flisak, J. R.; Gombatz, K.; Lantos, I.; Mendelson, W.; Remich, J. J. *Tetrahedron Lett.* **1990**, *31*, 6501-6504. (b) Formyl-acylal **19**: Gorodetsky, M.; Danieli, N.; Mazur, Y. *J. Org. Chem.* **1967**, *32*, 760-764. (c) Formyl-acylal **20**: DeBoer, A.; Ellwanger, R. E. *J. Org. Chem.* **1974**, *39*, 77-83.

- A similar mechanistic rationale was proposed to explain formation of mixed acylals **19** and **20**; See refs 14b-c for details.
- Nudelman, A.; Levovich, I.; Cutts, S. M.; Phillips, D. R.; Rephaeli, A. *J. Med. Chem.* **2005**, *48*, 1042-1054.
- Su, W.; Can, J. *J. Chem. Res.* **2005**, 88-90.
- Formyl-acylals have been prepared previously in moderate 44-57% yields via reaction of aldehydes with an acid anhydride and formic acid in the presence of phosphorus pentoxide, see: Scheeren, J. W.; Tax, W. J. M.; Schijf, R. *Synthesis* **1973**, 151-153.
- For other *N*-formylation reactions in water see: (a) Habibi, D.; Sahebkhietari, H.; Nasrollahzadeh, M.; Taghipour, A. *Lett. Org. Chem.* **2013**, *10*, 209-212. (b) Kaboudin, B.; Khodamorady, M. *Synlett* **2010**, 2905-2907.
- For other non-catalyzed solvent-free *N*-formylation reactions, see: (a) Rahman, M.; Kundu, D.; Hajra, A.; Majee, A. *Tetrahedron Lett.* **2010**, *51*, 2896-2899. (b) Habibi, D.; Nasrollahzadeh, M. *Compt. Rend. Chim.* **2013**, *16*, 1008-1016. (c) Dhake, K. P.; Tambade, P. J.; Singhal, R. S.; Bhanage, B. M. *Green Chem. Lett. Rev.* **2011**, *4*, 151-157. (d) Lebleu, T.; Kotsuki, H.; Maddaluno, J.; Legros, J. *Tetrahedron Lett.*, **2014**, *55*, 362-364. (e) Batuta, S.; Begum, N. A. *Synth. Comm.* **2017**, *47*, 137-147.
- (a) Nishikawa, Y.; Nakamura, H.; Ukai, N.; Adachi, W.; Hara, O. *Tetrahedron Lett.* **2017**, *58*, 860-863. (b) El-Dine, T. M.; Evans, D.; Rouden, J.; Blanchet, J. *Chem. Eur. J.* **2016**, *22*, 5894-5898. (c) Hill, D. R.; Hsiao, C. N.; Kurukuluriya, R.; Wittenberger, S. J. *Org. Lett.* **2002**, *4*, 111-113. (d) Suchy, M.; Elmehriki, A. A. H.; Hudson, R. H. E. *Org. Lett.* **2011**, *13*, 3952-3955. (e) Ducek, W.; Deutsch, J.; Vieth, S.; Niclas, H. J. *Synthesis* **1996**, 37-38. (f) Kisfaludy, L.; Otvos, L. *Synthesis* **1987**, 510. (g) Kim, J.-G.; Jang, D. O. *Synlett* **2010**, 1231-1234. (h) Chancellor, T.; Morton, C. *Synthesis* **1994**, 1023-1025.
- (a) Fayol, A.; Housseman, C.; Sun, X. W.; Janvier, P.; Bienaymé, H.; Zhu, J. P. *Synthesis* **2005**, 161-165. (b) Yang, P.-Y.; Liu, K.; Ngai, M.-H.; Lear, M. J.; Wenk, M. R.; Yao, S. Q. *J. Am. Chem. Soc.* **2010**, *132*, 656-666. (c) Fukuda, T.; Wagatsuma, H.; Kominami, Y.; Nogata, Y.; Yoshimura, E.; Chiba, K.; Kitano, Y.

- Chem. Biodiv.* **2016**, *13*, 1502-1510. (d) Soriano-Maldonado, P.; Rodriguez-Alonso, M. J.; Hernandez-Cervantes, C.; Rodriguez-Garcia, I.; Clemente-Jimenez, J. M.; Rodriguez-Vico, F.; Martinez-Rodriguez, S.; Las Heras-Vazquez, F. J. *Process Biochem.* **2014**, *49*, 1281-1287.
23. Aleiwi, B. A.; Mitachi, K.; Kurosu, M. *Tetrahedron Lett.* **2013**, *54*, 2077-2081.
24. Yang, K. H.; Fang, H.; Ye, J. S.; Gong, J. Z.; Wang, J. T.; Xu, W. F. *Pharmazie* **2008**, *63*, 779-783.