Citation for published version:

Bennett, HM, Williamson, SM, Walsh, TK, Woods, DJ & Wolstenholme, AJ 2012, 'ACR-26: A novel nicotinic receptor subunit of parasitic nematodes', *Molecular and Biochemical Parasitology*, vol. 183, no. 2, pp. 151-157. <https://doi.org/10.1016/j.molbiopara.2012.02.010>

DOI:

[10.1016/j.molbiopara.2012.02.010](https://doi.org/10.1016/j.molbiopara.2012.02.010)

Publication date:

2012

Document Version

Peer reviewed version

[Link to publication](#)

NOTICE: this is the author's version of a work that was accepted for publication in *Molecular and Biochemical Parasitology*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Molecular and Biochemical Parasitology* vol 183, issue 2, 2012, DOI 10.1016/j.molbiopara.2012.02.010

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ACR-26: a novel nicotinic receptor subunit of parasitic nematodes

Hayley M. Bennett^{a,b}, Sally M. Williamson^{a,b,1}, Thomas K. Walsh^{b,2}, Debra J. Woods^c, Adrian J. Wolstenholme^{a,b*}

^aDept of Infectious Diseases & Center for Tropical and Emerging Global Diseases, University of Georgia, Athens, GA 30602, USA; ^bDept of Biology & Biochemistry, University of Bath, Bath BA2 7AY, UK; ^cVeterinary Medicine Discovery Parasitology, Pfizer Animal Health, 7000 Portage Rd, Kalamazoo MI 49001 USA.

*Author for Correspondence: Dr Adrian Wolstenholme, Dept of Infectious Diseases, University of Georgia, 501 DW Brooks Drive, Athens, GA 30602 USA. Email adrianw@uga.edu, Tel 1 706 542 2404.

¹Current address: Institute of Neuroscience, Henry Wellcome Building, Newcastle University, Newcastle Upon Tyne NE1 7RU, United Kingdom.

²Current address: CSIRO, Clunies Ross Street, Black Mountain, ACTON ACT 2601, Australia

Abstract

Nematode nicotinic acetylcholine receptors are the targets for many effective anthelmintics, including those recently introduced into the market. We have identified a novel nicotinic receptor subunit sequence, *acr-26*, that is expressed in all the animal parasitic nematodes we examined from clades III, IV and V, but is not present in the genomes of *Trichinella spiralis*, *Caenorhabditis elegans*, *Pristionchus pacificus* and *Meloidogyne spp.* In *Ascaris suum*, ACR-26 is expressed on muscle cells isolated from the head, but not from the mid-body region. Sequence comparisons with other vertebrate and nematode subunits suggested that ACR-26 may be capable of forming a functional homomeric receptor; when *acr-26* cRNA was injected into *Xenopus* oocytes along with *X. laevis ric-3* cRNA we occasionally observed the formation of acetylcholine- and nicotine-sensitive channels. The unreliable expression of ACR-26 *in vitro* may suggest that additional subunits or chaperones may be required for efficient formation of the functional receptors. ACR-26 may represent a novel target for the development of cholinergic anthelmintics specific for animal parasites.

Keywords: anthelmintic, filaria, immunofluorescence, nicotinic receptor

1. Introduction

Control of parasitic nematode infections continues to rely on the use of chemical anthelmintics. Many of these compounds act at ion channels, including the nicotinic acetylcholine receptors (nAChRs) found at the nematode neuromuscular junction, on pharyngeal muscle and within the central nervous system [1,2,3,4]. Indeed, the two most recently introduced veterinary anthelmintics, monepantel [5] and derquantel [6], as well as tribendimidine, proposed for use against human infections [7], all act at nicotinic receptors [8,9,10]. Nematodes possess many nAChR genes, and these vary considerably between species, with several parasites having fewer than the model organism, *Caenorhabditis elegans* [11]. From *in vitro* studies that have reconstituted levamisole-sensitive receptors [12,13,14] it is clear that the subunit composition and pharmacology of the neuromuscular nAChR are different between parasites and *C. elegans*, and this has led us to search for novel nAChR sequences in the parasitic nematode, *Ascaris suum*. Such novel receptors may have potential as drug targets. This large worm was selected for these studies because the physiology and pharmacology of many of its nAChR can be studied *ex vivo* [15,16], and because the distribution of the receptor subunits can be studied in dissected tissues and on disassociated muscle cells.

We report here the identification and expression of a nAChR subunit sequence, *acr-26*, that is present in *A. suum* and orthologues of which are widely distributed in animal parasitic nematodes, but are absent from the genomes of several free-living and plant parasitic species.

2 Materials and Methods

2.1 Parasite Material

Adult *A. suum* were a kind gift of Prof A. Maule (Queen's University, Belfast) and Dr Richard Martin (Iowa State University). *Haemonchus contortus* L3 larvae of the drug sensitive ISE isolate were kindly supplied by Dr Philip Skuce (Moredun Institute).

2.2 Molecular cloning

An EST sequence (Accession number FE918510), derived from *A. suum* and showing significant identity to known nAChR subunit cDNAs, was identified in the database. 5' and 3' RACE reactions (primers; 5' RACE, SL1 (GGTTTAATTACCCAAGTTTGAG) and ACR26-RV3 (AACGTTTATCGTCAACACCTG); 3' RACE, anchor (GACCACGCGTATCGATGTCGAC) and ACR26-FW2 (TAATTATGTTGTGTCGGGTG) were carried out as described previously [12,17] to amplify the rest of the cDNA sequence. These partial products were cloned into pGEM-Teasy and sequenced. Specific primers (forward – ATGATGGCAACTCGTCGG; reverse – TTAATGCAGACCATATAAAGAC) were used to amplify a full-length sequence from *A. suum* cDNA (made from RNA extracted from the head region); this sequence was also cloned into pGEM-Teasy and sequenced. The sequence was deposited in the database under the Accession Number GU135625. An essentially identical procedure was used to amplify a full-length cDNA from *Haemonchus contortus*, which was deposited under Accession number EU006791.

In order to search for related sequences in cDNA from other species (*Cooperia oncophora*, *Ostertagia ostertagi* and *Teladorsagia circumcincta*) degenerate oligonucleotide primers were

designed based on the aligned sequences from *A. suum* and *H. contortus* and used to amplify partial sequences from the target organisms.

2.3 Immunofluorescence

A specific goat antiserum (Sigma-Genosys, USA) was raised against a synthetic multiple antigenic peptide, EIDGTATDEQKLLHLL, (Alta Biociences, UK) corresponding to the N-terminus of the mature ACR-26 polypeptide, essentially as described [12]. IgG was isolated from the serum by affinity chromatography over a CPG column to which the antigenic peptide had been immobilised, and the purified antibody used in immunofluorescence experiments in dissociated muscle cells isolated from the body wall and head regions. Adult *A. suum* were kindly provided by Prof, A. Maule (Queen's University, Belfast, UK) and were shipped and stored in Ascaris Ringer Solution (4mM NaCl, 5.9mM CaCl₂, 4.9mM MgCl₂, 5mM Tris-HCl pH7.4, 125mM sodium acetate, 24.5mM KCl). They were used within 24hrs of their arrival. The worms were pinned out on a dissection tray and injected at 3cm intervals with 5mg/ml collagenase 1A in ARS. After 2 hrs at 37° the cuticle was cut longitudinally and pinned flat at the head end. Disassociated head and muscle cells were removed independently with a Pasteur pipette and fixed in 5% (v/v) formaldehyde in ARS for 9hrs at 4°. The cells were washed three times in 0.1% (v/v) Triton X-100 in phosphate-buffered saline (PBS). The affinity-purified anti-ACR-26 was applied at a 1:200 dilution in PBS and the cells incubated with gentle agitation for 40hrs at 4°. Control cells were incubated with purified control goat IgG under the same conditions. The cells were washed three times in Triton X-100/PBS as before and then an FITC-conjugated rabbit anti-goat IgG (Sigma, Poole, UK, catalogue number F7367), diluted 1:200 in PBS, added for 4hrs at 4°. The cells were washed three times in Triton X-100/PBS before being

103 mounted in Mowiol 4-88 reagent (Polysciences, Inc, USA) and observed under a Zeiss LSM510
104 confocal microscope.

106 ***2.4 Functional Expression***

107 The *acr-26* cDNA was subcloned into the BglII and SpeI sites of the pT7TS vector, which was
108 linearised and transcribed into cRNA using the mMessage mMachine T7 kit (Ambion). The
109 cRNA was injected into defolliculated *Xenopus* oocytes along with cRNA encoding the *X. laevis*
110 orthologue of RIC-3 (Bennett et al., unpublished), which were screened for acetylcholine-gated
111 channels as described [12].

3 Results

3.1 Cloning of *acr-26* cDNA from parasitic nematodes

We successfully amplified a full-length cDNA encoding the novel nAChR subunit from *A. suum*, extending the EST sequence (Accession number FE918510) that had already been deposited in the database. We compared the sequence of the *A. suum* subunit with the other nAChR subunits from both *A. suum* and *C. elegans* (Figure 1). The results showed that the new subunit was not orthologous to any of those from *C. elegans* and we therefore named the new sequence *acr-26* to distinguish it from them. In order to determine whether *acr-26* was confined to *A. suum* or was present in other parasitic nematodes, we searched the partial genome sequence of *H. contortus* for similar sequences and used the results of that search to amplify a full-length orthologous sequence from this clade V parasite. Alignments of the Asu-ACR-26 and Hco-ACR-26 sequences (Figure 2) showed that they were very similar, and would almost overlap if both were plotted on the tree shown in Figure 1. Both subunits shared key amino-acid residues in loops that form the agonist binding sites with vertebrate $\alpha 7$ subunits, especially in the complementary loop D normally provided by β -subunits in heteromeric receptors, suggesting that ACR-26 may be able to form a homomeric nAChR [18]. Further BLAST searches (Table 1) revealed that *acr-26* like sequences are present in the filarial parasites *Brugia malayi*, *Dirofilaria immitis*, *Loa loa* and *Wuchereria bancrofti*, and *Strongyloides ratti*, but not in *Trichinella spiralis*, the plant parasitic *Meloidogyne spp.* or in the free living *Pristionchus pacificus*. When we searched other invertebrate phyla for ACR-26-like sequences, the best hits were with the nAChR subunits G and D from *Lymnaea stagnalis* [19], which shared 67% and 61% amino-acid identity with Asu-ACR-26, respectively. In order to investigate whether or not the gene is present in other trichostrongylid nematodes of economic importance, we amplified and sequenced partial *acr-26*

cDNAs from *C. oncophora*, *O. ostertagi* and *T. circumcincta*. The partial clones were sequenced and proved to possess high levels of identity to Hco-ACR-26, showing that this subunit is also expressed in these parasites (Figure 3). They are deposited in the sequence database under Accession numbers JN966888, JN966889 and JN966890.

3.2 Distribution of ACR-26 in *Ascaris suum*

In order to determine where in the parasite ACR-26 was expressed, we raised an antiserum against a synthetic peptide corresponding to the predicted N-terminal sequence of the mature polypeptide. Antibodies purified from this antiserum recognized an HA-tagged version of ACR-26 when this was expressed in mammalian cells, and when examined under confocal microscopy the anti-ACR-26 immunofluorescence completely overlapped with that produced by an anti-HA antibody (data not shown). We applied the purified anti-ACR-26 antibody to isolated muscle cells, derived both from the mid-body region and from the head. No specific staining of the body wall muscles was observed (Fig 4), but immunoreactivity was detected on the surface of the head muscle cells. This was consistent with the cloning of the *acr-26* cDNA from RNA isolated from the head region of the worm. No fluorescence was observed when preparations were treated with a control goat IgG.

3.3 ACR-26 forms a functional nicotinic receptor

The amino-acid sequence of ACR-26, specifically the conservation of residues in the loops forming the ligand-binding site with those present in vertebrate $\alpha 7$ subunits (Fig 1), suggested that it may be able to form a functional nAChR when expressed as a homomer. We therefore injected *Xenopus* oocytes with *Asu-acr-26* or *Hco-acr-26* cRNA and attempted to detect the

160 formation of Ach- and nicotine-sensitive channels. Expression of Asu-ACR-26 nAChRs was
161 sporadic and unreliable, but on occasion channels were detected in response to the application of
162 Ach and nicotine (Fig 5). These channels were extremely sensitive to Ach, with concentrations
163 $>100\mu\text{M}$ producing maximal responses; the unreliable expression of this receptor makes an
164 accurate estimate of the EC_{50} for Ach very difficult but it was between 10 and 100nM. For
165 nicotine the EC_{50} was $25\mu\text{M}$ (95% confidence limits 15-42 μM), with a Hill coefficient of 1.66
166 ± 1.29 . Since ACR-26 is expressed on some muscle cells (Fig 4), it is possible that it co-
167 assembles with other muscle nAChR subunits *in vivo*, but attempts to improve the reliability and
168 reproducibility of *in vitro* ACR-26 expression by co-expression with *Asu-unc-29* or *Asu-unc-38*
169 cRNAs were unsuccessful. No functional channels were detected in oocytes injected with *Hco-*
170 *acr-26* cRNA.

4 Discussion

Nematodes encode a rich variety of nAChR and these continue to be exploited as effective targets for the development of new anthelmintic drugs [5,6,7]. We report here that many animal parasitic species possess a new gene, *acr-26*, that is not present in *C. elegans*, several other free living species or the *Meloidogyne* genus of plant parasites. In *A. suum*, the ACR-26 subunit is expressed in head, but not body-wall, muscle cells, and is capable of forming a homomeric receptor – though expression of this receptor is unreliable, which might indicate that further subunits are required for full activity *in vivo*. Attempts to express ACR-26 homomers from a second parasitic species, *H. contortus*, were unsuccessful. It would be interesting to add ACR-26 to the reconstituted *H. contortus* nAChR recently reported by Boulin et al. [13]. The pharmacology of the homomeric receptor, if it reflects that of native ACR-26 containing nAChR, would appear to be distinct from those previously reported for reconstituted levamisole receptors [12,13,14]. The ACR-26 channels were extremely sensitive to Ach, and the EC₅₀ of between 10 and 100nM was 1-2 orders of magnitude less than that of the *A. suum* UNC-29/UNC-38 receptor (~1μM) [12] – compare the responses to 10 and 100nm Ach with those to 10 and 30μM nicotine in Figure 5. If this reflects the pharmacology of native ACR-26 containing receptors on head muscle cells, it suggests that they may mediate responses to lower levels of cholinergic signaling than the previously characterized levamisole-sensitive receptors [12-14]. This novel pharmacology might enable it to be developed as a target for compounds that are effective against nematode parasites, but are less dangerous to free-living species in the environment.

The expression of ACR-26 on *Ascaris* head but not body-wall, muscles, is distinct from that of other nicotinic subunits, such as UNC-29 and UNC-38, that are found on both muscle types [12]

and suggests that it may have a specific function there. It is tempting to relate the function of ACR-26 to the more complex movements in the nematode head - body-wall muscle permits only dorsal-ventral bends, whereas the head can also move laterally – but it is difficult to explain why the subunit would only be present in parasitic and not free-living species. We have as yet no information on the distribution of ACR-26 in nematodes other than *Ascaris*. The evolutionary history of this gene is interesting; it is conserved in animal parasitic species of multiple clades [19], though not clade I, implying that it pre-dates their appearance, but is absent from free-living species of clade V and plant parasitic species of clade IV. Analysis of nematode phylogenetics has led to the conclusion that animal parasitism evolved multiple times [20,21,22,23], which is on the face of it hard to reconcile with a gene that is specifically associated with parasitic species, even if, as suggested by van Megen et al [23], convergent evolution seems to be a feature of the Nematoda. It is probably more likely that *acr-26* has been lost in the free-living and plant parasitic species; this, together with its specific expression in head muscles, raises interesting questions about its likely function. Further developments in functional genetics methods for parasitic nematodes [24,25,26] may allow us to understand that function better. The high level of amino-acid identity between ACR-26 and the molluscan D and G subunits [19] is interesting; these subunits form a small out-group on the phylogenetic tree of mollusc nAChR and are expressed only at low levels in the CNS. No functional expression of either could be detected in *Xenopus* oocytes [27]. ACR-26 may thus be a member of a small group of invertebrate nAChR subunits whose function has yet to be determined.

Acknowledgements

216 We would like to thank Aaron Maule (Queen's University, Belfast) and Richard Martin (Iowa
217 State University) for their generosity in supplying adult *Ascaris suum*. This work was funded by
218 an Industrial Partnership Grant from BBSRC (UK) and Pfizer Animal Health.

219

References

- [1] Jones AK, Sattelle DB. Functional genomics of the nicotinic acetylcholine receptor gene family of the nematode, *Caenorhabditis elegans*. *Bioessays* 2004;26:39-49.
- [2] Martin RJ. Modes of action of anthelmintic drugs. *Vet J* 2004;154:11-34.
- [3] Sattelle DB. Invertebrate nicotinic acetylcholine receptors-targets for chemicals and drugs important in agriculture, veterinary medicine and human health. *J Pestic Sci* 2009;34:233-40.
- [4] Wolstenholme AJ. Ion channels and receptors as targets for the control of parasitic nematodes. *Int J Parasit - Drugs Drug Resist* 2011;1:In Press.
- [5] Kaminsky R, Ducray P, Jung M, et al. A new class of anthelmintics effective against drug-resistant nematodes. *Nature* 2008;452:176-80.
- [6] Little PR, Hodge A, Maeder SJ, et al. Efficacy of a combined oral formulation of derquantel-abamectin against the adult and larval stages of nematodes in sheep, including anthelmintic-resistant strains. *Vet Parasitol* 2011;181:180-93.
- [7] Steinmann P, Zhou XN, Du ZW, et al. (2008) Tribendimidine and albendazole for treating soil-transmitted helminths, *Strongyloides stercoralis* and *Taenia* spp.: open-label randomized trial. *PLoS Negl Trop Dis* 2008;2:e322.
- [8] Rufener L, Baur R, Kaminsky R, Maser P, Sigel E. Monepantel allosterically activates DEG-3/DES-2 channels of the gastrointestinal nematode *Haemonchus contortus*. *Mol Pharmacol* 2010;78:895-902.
- [9]. Robertson AP, Clark CL, Burns TA, et al. Paraherquamide and 2-deoxy-paraherquamide distinguish cholinergic receptor subtypes in *Ascaris* muscle. *J Pharmacol Exp Therapeut* 2002;302:853-60.
- [10] Hu Y, Xiao SH, Aroian RV. The new anthelmintic tribendimidine is an L-type (levamisole and pyrantel) nicotinic acetylcholine receptor agonist. *PLoS Negl Trop Dis* 2009;3:e499.
- [11] Williamson SM, Walsh TK, Wolstenholme AJ. The cyst-loop ligand-gated ion channel gene family of *Brugia malayi* and *Trichinella spiralis*: a comparison with *Caenorhabditis elegans*. *Invert Neurosci* 2007;7:219-26.
- [12] Williamson SM, Robertson AP, Brown L, et al. The nicotinic acetylcholine receptors of the parasitic nematode *Ascaris suum*: formation of two distinct drug targets by varying the relative expression levels of two subunits. *PLoS Pathog* 2009;5:e1000517.
- [13] Boulin T, Fauvin A, Charvet CL, et al. Functional reconstitution of *Haemonchus contortus* acetylcholine receptors in *Xenopus* oocytes provides mechanistic insights into levamisole resistance. *Br J Pharmacol* 2011;164:1421-32.
- [14] Boulin T, Gielen M, Richmond JE, et al. Eight genes are required for functional reconstitution of the *Caenorhabditis elegans* levamisole-sensitive acetylcholine receptor. *Proc Nat'l Acad Sci USA* 2008;105: 18590-95.
- [15] Martin RJ, Pennington AJ, Duittoz AH, Robertson S, Kusel JR. The physiology and pharmacology of neuromuscular transmission in the nematode parasite, *Ascaris suum*. *Parasitology* 1991;102:S41-S58.
- [16] Qian H, Martin RJ, Robertson AP. Pharmacology of N-, L-, and B-subtypes of nematode nAChR resolved at the single-channel level in *Ascaris suum*. *FASEB Journal* 2006;20:2606.
- [17] Laughton DL, Amar M, Thomas P, et al. Cloning of a putative inhibitory amino acid receptor subunit from the nematode *Haemonchus contortus*. *Recept Channels* 1994;2:155-63.

- [18] Brejc K, van Dijk WJ, Klaassen RV, et al. Crystal structure of an ACh-binding protein reveals the ligand-binding domain of nicotinic receptors. *Nature* 2001;411:269-76.
- [19] van Nierop P, Keramidas A, Bertrand S, et al. Identification of molluscan nicotinic acetylcholine receptor (nAChR) subunits involved in formation of cation- and anion-selective nAChRs. *J Neurosci* 2005;25:10617-26.
- [20] Blaxter ML, De ley P, Garey JR, et al. A molecular evolution framework for the phylum Nematoda. *Nature* 1998;392:71-5.
- [21] Holterman M, van der Wurff A, van den Elsen S, et al. Phylum-wide analysis of SSU rDNA reveals deep phylogenetic relationships among nematodes and accelerated evolution towards crown clades. *Mol Biol Evol* 2006;23:1792-800.
- [22] Meldal BHM, Debenham NJ, De Ley P, et al. An improved molecular phylogeny of the Nematoda with special emphasis on marine taxa. *Mol Phylogenet Evol* 2007;42:622-36.
- [23] van Megen H, van den Elsen S, Holterman M, et al. A phylogenetic tree of nematodes based on about 1200 full-length small subunit ribosomal DNA sequences. *Nematol* 2009;11:927-54.
- [24] Xu S, Liu C, Tzertzinis G, et al. In vivo transfection of developmentally competent *Brugia malayi* infective larvae. *Int J Parasitol* 2011;41:355-62.
- [25] Lok JB, Artis D. Transgenesis and neuronal ablation in parasitic nematodes: revolutionary new tools to dissect host-parasite interactions. *Parasite Immunol* 2008;30:203-14.
- [26] Li X, Shao H, Junio A, et al. Transgenesis in the parasitic nematode *Strongyloides ratti*. *Mol Biochem Parasitol* 2011;179:114-9.
- [27] van Nierop P, Bertrand S, Munno DW, et al. (2006) Identification and functional expression of a family of nicotinic acetylcholine receptor subunits in the central nervous system of the mollusc *Lymnaea stagnalis*. *J Biol Chem* 2006;281:1680-91.
- [28] Mitreva M, Jasmer DP, Zarlenga DS, et al. The draft genome of the parasitic nematode *Trichinella spiralis*. *Nature Genetics* 2011;43:228-74.
- [29] Wang JB, Czech B, Crunk A, et al. Deep small RNA sequencing from the nematode *Ascaris* reveals conservation, functional diversification, and novel developmental profiles. *Genome Res* 2011;21:1462-77.
- [30] Ghedin E, Wang S, Spiro D, et al. Draft genome of the filarial nematode parasite *Brugia malayi*. *Science* 2007;317:1756-60.
- [31] Elsworth B, Wasmuth J, Blaxter M. NEMBASE4: The nematode transcriptome resource. *Int J Parasitol* 2011;41:881-94.
- [32] Godel C, Kumar S, Koutsovoulos G, et al. The genome of the heartworm, *Dirofilaria immitis*. *Proc Nat'l Acad Sci. USA*; submitted.
- [33] Abad P, Gouzy J, Aury JM, et al. Genome sequence of the metazoan plant-parasitic nematode *Meloidogyne incognita*. *Nature Biotech* 2008;26:909-15.
- [34] Opperman CH, Bird DM, Williamson VM, et al. Sequence and genetic map of *Meloidogyne hapla*: A compact nematode genome for plant parasitism. *Proc Nat'l Acad Sci USA* 2008;105:14802-7.
- [35] *C. elegans* Sequence Consortium. Genome sequence of the nematode *C. elegans*: A platform for investigating biology. *Science* 1998;282:2012-18.
- [36] Dieterich C, Clifton SW, Schuster LN, et al. The *Pristionchus pacificus* genome provides a unique perspective on nematode lifestyle and parasitism. *Nature Genetics* 2008;40:1193-8.

Figure Legends

Figure 1. Tree of *C. elegans* and *A. suum* nAChR subunits. A maximum likelihood neighbour joining bootstrapped tree of translated nAChR sequences drawn with Geneious Pro 5.4. Sequences of *C. elegans* nAChR subunits are shown in black. *A. suum* nAChR subunit sequences identified in the transcriptome [29] are shown in red, and are named after their *C. elegans* orthologue, with the exception of ACR-26 (red box), which has no orthologue. All orthologue pairings gave bootstrap values of 100.

Figure 2. Alignment of nematode ACR-26 sequences with vertebrate $\alpha 7$. An alignment of the ACR-26 polypeptides from *A. suum* and *H. contortus* with the murine $\alpha 7$ nAChR subunit was made with Clustal (<http://www.ebi.ac.uk/Tools/msa/clustalw2/>). Amino-acid residues conserved between the two ACR-26 subunits, and $\alpha 7$, are shown in bold. The yellow shading indicates the residues predicted to form the loops of the ligand-binding site, and the grey shading shows the predicted membrane-spanning regions. The underlined residues are those that were used to raise an antibody against the *A. suum* subunit.

Figure 3. Partial *acr-26* sequences from nematodes of medical and agricultural importance. Full-length amino-acid sequences of ACR-26 from *A. suum*, *B. malayi* and *H. contortus* were aligned with the translated partial sequences from *O. ostertagi*, *C. oncophora* and *T. circumcincta* obtained from PCR reactions using degenerate primers designed to conserved regions.

Figure 4. Immunostaining of *A. suum* muscle cells with anti-ACR-26 antibody. Muscle cells were isolated from either *A. suum* body wall muscle or from the 2 cm most anterior region (in front of the nerve ring), fixed and stained. A) A representative head muscle cell in a negative control condition, treated with control goat IgG and anti-goat IgG FITC. In the experimental condition cells were treated with affinity purified anti-ACR-26 and anti-goat IgG FITC. B) A representative body wall muscle cell treated with the anti-ACR-26 antibody. C) A representative head muscle cell demonstrating a strong positive signal. D). Close-up of the muscle arm after anti-ACR-26 staining on head muscle. Confocal image of head muscle stained with affinity-purified anti-ACR-26 antibody. Part of the arm is out of the plane of the image.

Figure 5. Asu-ACR-26 is capable of forming a functional nAChR in *Xenopus* oocytes. A) Example dose-dependent responses to applied ACh. B) Dose response curve for nicotine at the ACR-26 receptor. C) Example dose-dependent responses to applied nicotine.

Table 1. Distribution of *acr-26* sequences in nematode species.

Clade[19]	Trophic ecology	Species	<i>Acr-26</i> ?	Length of known coding sequence and reference numbers	Source
I	Vertebrate parasite	<i>Trichinella spiralis</i>	No	-	BLAST completed genome project [268]
III	Vertebrate parasite	<i>Ascaris suum</i>	Yes	Full-length (1533 bp), GenBank: GU135625	EST [279], this manuscript.
III	Vertebrate parasite	<i>Brugia malayi</i>	Yes	Full-length (1542 bp), NCBI: XM_001901191	BLAST completed genome project[2830]
III	Vertebrate parasite	<i>Loa loa</i>	Yes	Full-length (1428 bp), NCBI: XM_003140235 (LOAG_04698)	BLAST completed genome project (http://www.broadinstitute.org/annotation/genome/filarial_worms/Info.html)
III	Vertebrate parasite	<i>Wuchereria bancrofti</i>	Yes	Partial (630 bp), Broad Institute: WUBG_13499	BLAST incomplete genome project (http://www.broadinstitute.org/annotation/genome/filarial_worms/Info.html)
III	Vertebrate parasite	<i>Dirofilaria immitis</i>	Yes	Partial (577 bp), EST cluster: DIC00454	EST and completed genome project [29,30,31,32]
IV	Phytoparasite	<i>Meloidogyne incognita</i>	No	-	BLAST completed genome project [343]
IV	Phytoparasite	<i>Meloidogyne hapla</i>	No	-	BLAST completed genome project [324]
IV	Vertebrate parasite	<i>Strongyloides ratti</i>	Yes	Sanger Institute, pathogen_RATTI_Contig 74886	BLAST genome project (http://www.sanger.ac.uk/resources/downloads/helminths/strongyloides-ratti.html)
V	Free-living bacterivore	<i>Caenorhabditis elegans</i>	No	-	BLAST completed genome project [335]
V	Free-living omnivore	<i>Pristionchus pacificus</i>	No	-	BLAST completed genome project [346]
V	Vertebrate parasite	<i>Cooperia oncophora</i>	Yes	Partial (639 bp) GenBank: JN966889	Cloning – this manuscript
V	Vertebrate parasite	<i>Haemonchus contortus</i>	Yes	Full-length (1573 bp), GenBank: EU006791	BLAST incomplete genome project (http://www.sanger.ac.uk/resources/downloads/helminths/haemonchus-contortus.html), Cloning – this manuscript
V	Vertebrate parasite	<i>Ostertagia ostertagi</i>	Yes	Partial (597 bp) GenBank: JN966890	Cloning – this manuscript
V	Vertebrate parasite	<i>Teladorsagia circumcincta</i>	Yes	Partial (708 bp) GenBank: JN966888	Cloning – this manuscript

Asu-ACR-26	MMATRRELILLIFIHGVVDVVLADSSITDLSEIDGTATDEQKLLYHL	50
Hco-ACR-26	-MLIFCSIMSLVNAIPLNVSEAPTVPPTEDHAGGRRYTDEQRLLYHLMK	49
Mus α7	-----MCGRRGGIWLALAAALLHVS LQG-----EFQRRLYKELVK	35
	Loop D	
Asu-ACR-26	QYEKAVRPVRNASH T V T V K L G M T M T N I F E M D E K N Q V L T I N V W L D Q E W K D E	100
Hco-ACR-26	DYERSVRPVRNASH T V T V R L G M T M T N I F D M D E R N Q V P T I N V W L D Q E W N D E	99
Mus α7	NYNPLERPVANDS Q P L T V Y F S L S L L Q I M D V D E K N Q V L T T N I W L Q M S W T D H	85
	Loop A	Loop E
Asu-ACR-26	LLRWDPKEFGGIQSIRIPCDLIWLPDIVLYNNADDYTAGYMRSRAMV FYD	150
Hco-ACR-26	LLRWNPDDFNGIQSLRIPCDLIWLPDIVLYNNADDYTAGYMRSRAMVLYT	149
Mus α7	YLQWNMSEYPGVKNVRF PDGQIWKPDILLYNSADERFDATEHTNVLVNAS	135
	Loop B	
Asu-ACR-26	GTVFWPPP T Q L R S T C K I D V T Y F P F D S Q H C A L K F G S W T Y H G F Q V D I T N R S D	200
Hco-ACR-26	GTVFWPPP K Q L R S T C K V D V S L F P F D E Q R C S L K F G S W T Y H G F Q V D I T N R S E	199
Mus α7	GHCQYLP P G I F K S S C Y I D V R W F P F D V Q Q C K L K F G S W S Y G G W S L D L Q --MQ	183
	Loop F	Loop C
Asu-ACR-26	NVDLSNYVVSGEFDLVRVHQKRRVVKYTCCEPEPYDVTFFFIHIRRKTLYY	250
Hco-ACR-26	NIDL TNYPVSGEFDLVKVYQKRRVVKYTCCEPEPYDITFFFIYIRRKTLYY	249
Mus α7	EADISSYIPNGEWDLMGIPGKRNEKFYECCEPEPYDVTYTVTMRRTLYY	233
Asu-ACR-26	LYNVVFPCLMMSVLTLLVFLLPDPSGEKIALGITVLLAFSVFVLAIAEKM	300
Hco-ACR-26	LYNIVFPCLMMSVLTLLVFLVPDPSGEKIALGITVLLAFSVFVLAIAEKM	299
Mus α7	GLNLLIPCVLISALALLVFLLPADPSGEKISLGITVLLSLTVFMLLVAEIM	283
Asu-ACR-26	PETSDSMPLIGIYLTVVMAMTSVSVVMTVMVLNFHHRGPFNQAVPKWVHR	350
Hco-ACR-26	PETSDSMPLIGIYLTVVMSMTSVSVVMTVMVLNFHHRGPFNEPVPNWARV	349
Mus α7	PATSDSVPLIAQYFASTMIIVGLSVVTVIVLRYHHHD PDGGKMPKWTRI	333
Asu-ACR-26	LVLNRLRRALCMRLPYTGWKDNGFCSANGMTKTI SIGLAMDDFN-----	394
Hco-ACR-26	LVLDRLLRLLRMKLSTRGESARVSVCPNSMRRMSVRVAMDDMRKEIISV	399
Mus α7	ILLNWCAWFLRMKRPGEDKVRPACQHKPRCSLASVELSAGAGPPTSNGN	383
Asu-ACR-26	-----VDSFEELD N Q F L Q M Q E T E L V N V D D A A L R K K K C R P N E L H	433
Hco-ACR-26	LGPGLLTGGNGVAESARLESLLLDVPQTEFS-VEQNGVVRRPRVGD DLQ	448
Mus α7	LLYIGFRGLEGMHCAPT P D S G V V C G R L A C S P T H D E H L M H G T H P S D G D P D L	433
Asu-ACR-26	SRLLKTLQVLIRQEMEDLYQTLANEWRQVAQVIDRLLFWVFLVCTVIIT	483
Hco-ACR-26	AKLLRTLQVLVKRQENEDASERVANEWRHVAQVIDRLLLWIFL FATTAIT	498
Mus α7	AKILEEVRYIANRFRCQDESEVICSEWKFAACVDRCLMAFSVF TIICT	483
Asu-ACR-26	LILLIIIPAVHRSMESDVFDESLYGLH	510
Hco-ACR-26	FVLLVLIPSLPYTYNYEDPE-----	519
Mus α7	IGILMSAPNFVEAVSKDFA-----	502

		Loop A	Loop E	
Tci	WLDQEWKDELLRWDPKEFGGIQSIRIPCDLI	WLPDIVLYNNADDYTAGY	MRSRAMVFPYDG	60
Asu	WLDQEWKDELLRWDPKEFGGIQSIRIPCDLI	WLPDIVLYNNADDYTAGY	MRSRAMVFPYDG	151
Oos	-----	WLPDIVLYNNADDYTAGY	MRSRAMVLYTG	29
Hco	WLDQEWNDELLRWNPDDFNGIQSLRIPCDLI	WLPDIVLYNNADDYTAGY	MRSRAMVLYTG	150
Con	-----	WLPDIVLYNNADDYTAGY	MRSRAMVLYTG	29
Bma	WLDQEWKDELLVWDPKKFGGIKSVRVPCDLI	WLPDIVLYNSADDYTVGS	MHSRAILFYDG	159
		*****	*:****::*	

	Loop E	Loop B	Loop F	
Tci	TVFWPPPTQLRSTCKIDVTYPPFDSQH	CALKFGSWTYHGFQVDITN	RSNDNVDLSNYVVS	120
Asu	TVFWPPPTQLRSTCKIDVTYPPFDSQH	CALKFGSWTYHGFQVDITN	RSNDNVDLSNYVVS	211
Oos	TVFWPPPTQLRSTCKVDVSLPPFDEQRC	SLRFGSWTYHGFQVDITN	RSNDNIDLINYPVS	89
Hco	TVFWPPPKQLRSTCKVDVSLPPFDEQRC	SLKFGSWTYHGFQVDITN	RSENIDLINYPVS	210
Con	TVFWPPPTQLRSTCKVDVSLPPFDEQRC	SLKFGSWTYHGFQVDITN	RSGNIDLINYPVS	89
Bma	TVFWPPPTQLRSTCKTDVTYPPFDSQH	CSIKFGSWTYHGLQVDITN	RSINVDLSNYVES	220
	*****	:*****	*:****::*	

	Loop C	TM1	
Tci	EFDLVRVHQKRGVVKY-TCCPEPYPDVTFFIHVRRKTL	YYLYNVVFPCTMMSVLTLLVFL	180
Asu	EFDLVRVHQKRRVVKY-TCCPEPYPDVTFFIHRRKTL	YYLYNVVFPCTMMSVLTLLVFL	271
Oos	EFDLVKVYRKRRVVKY-TCCPEPYPDITFFIYIRKTL	YYLYNIVFPCLMMSVLTLLVFI	149
Hco	EFDLVKVYRKRRVVKY-TCCPEPYPDITFFIYIRKTL	YYLYNIVFPCLMMSVLTLLVFL	270
Con	EFDLVKVYRKRRVVKY-TCCPEPYPDITFFIHIRKTL	YYLYNIVFPCLMMSVLTLLVFI	149
Bma	EFDLVRVFCRRVVKY-TCCLEPYPDVTFFIHIRKTL	YYLYNVVFPCLMMSVLTLLVFL	280
	*****	:*****	*:****::*

	TM2	
Tci	PPDSGEKIALGITVLLAFSVFVLAIAADKMPET	212
Asu	PPDSGEKIALGITVLLAFSVFVLAIAEKMPET	303
Oos	PPDSGEKIALGITVLLAFSVFVLAIAEKMPET	181
Hco	PPDSGEKIALGITVLLAFSVFVLAIAEKMPET	302
Con	PPDSGEKIALGITVLLASSVFVLAIAEKMPET	181
Bma	PPDSNEKIALGITVLLAFSVSVLAIAEKMPET	312
	****	:*****

A

B

C

D