



Citation for published version:

Gerasimovičs, A, Hocquet, A & Nilssen, T 2021, 'Non-autonomous rough semilinear PDEs and the multiplicative Sewing lemma', *Journal of Functional Analysis*, vol. 281, no. 10, 109200.
<https://doi.org/10.1016/j.jfa.2021.109200>

DOI:

[10.1016/j.jfa.2021.109200](https://doi.org/10.1016/j.jfa.2021.109200)

Publication date:

2021

Document Version

Peer reviewed version

[Link to publication](#)

Publisher Rights

CC BY-NC-ND

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Non-autonomous rough semilinear PDEs and the multiplicative Sewing Lemma

Andris Gerasimovičs¹, Antoine Hocquet², and Torstein Nilssen³

¹University of Bath, Bath, United Kingdom;
Email: ag2616@bath.ac.uk

²Technische Universität Berlin, Berlin, Germany;
Email: antoine.hocquet86@gmail.com

³University of Agder, Kristiansand, Norway;
Email: torstein.nilssen@uia.no

August 9, 2020

Abstract

We investigate existence, uniqueness and regularity for local solutions of rough parabolic equations with subcritical noise of the form $du_t - L_t u_t dt = N(u_t) dt + \sum_{i=1}^d F_i(u_t) d\mathbf{X}_t^i$ where $(L_t)_{t \in [0, T]}$ is a time-dependent family of unbounded operators acting on some scale of Banach spaces, while $\mathbf{X} \equiv (X, \mathbb{X})$ is a two-step (non-necessarily geometric) rough path of Hölder regularity $\gamma > 1/3$. Besides dealing with non-autonomous evolution equations, our results also allow for unbounded operations in the noise term (up to some critical loss of regularity depending on that of the rough path \mathbf{X}). As a technical tool, we introduce a version of the multiplicative sewing lemma, which allows to construct the so called product integrals in infinite dimensions. We later use it to construct a semigroup analogue for the non-autonomous linear PDEs as well as show how to deduce the semigroup version of the usual sewing lemma from it.

Keywords: Rough path, rough partial differential equations, semilinear equations, multiplicative Sewing Lemma, propagator.

Mathematics Subject Classification (2010): 60L50, 60H15, 35K58, 32A70

Contents

1	Introduction	2
1.1	Motivations	2
1.2	General notation	7
2	Settings and main results	8
2.1	Functional analytic framework	8
2.2	Hölder spaces and controls	9
2.3	Assumptions and main results	11
2.4	An illustrating example: non-autonomous stochastic reaction-diffusion equations	16

3	A multiplicative sewing Lemma and some by-products	18
3.1	The main result	18
3.2	A new proof of Tanabe/Sobolevskii’s Theorem (proof of Theorem 2.10)	22
3.3	Lie-Trotter product formula	26
4	Controlled Path according to a monotone family of interpolation spaces	30
4.1	Affine Sewing Lemma	30
4.2	Controlled rough paths	34
5	Equations with subcritical multiplicative noise: proof of Theorem 2.15	40
5.1	Solutions to subcritical RPDEs	40
5.2	Continuity of the solution map and smoothing away from zero	43
5.3	Weak solutions. Proofs of Theorems 2.18 and 2.19	44
6	Further examples	49
6.1	Stochastic Navier-Stokes equation	49
6.2	Cahn-Hilliard equation	50
6.3	Parabolic equations with transport noise	50

1 Introduction

1.1 Motivations

We are interested in non-autonomous semilinear evolution equations of the form

$$\begin{cases} du_t = (L_t u_t + N(u_t))dt + \sum_{i=1}^d F_i(u_t) d\mathbf{X}_t^i, & t \in [0, T], \\ u_0 = x \in \mathcal{B}, \end{cases} \quad (1.1)$$

where the unknown u is continuous with values in some Banach space \mathcal{B} , $(L_t)_{t \in [0, T]}$ is a continuous in time family of unbounded operators satisfying a suitable sector condition, $X: [0, T] \rightarrow \mathbf{R}^d$ is a path of Hölder regularity $\gamma > 1/3$, while N and F denote some non-linearities.

Parabolic equations like (1.1) appear in a stochastic context (e.g. in filtering theory), where X denotes for instance a finite-dimensional Brownian motion. They were first investigated in the late 70’s through the work of Pardoux, Krylov and Rozovskii [41, 51], using an appropriate functional setting in which Itô calculus can be used, together with monotonicity arguments. In the autonomous case, the so called ‘mild approach’ was largely developed by Da Prato and his school, which culminated in the monograph [16].

In a ‘rough path’ context, the treatment of equations similar to (1.1) originates in the mild approach by Gubinelli, Lejay and Tindel [32] for Young paths, later extended to rough paths by Gubinelli and Tindel in [31] (see also the works of Deya [18, 17, 19]). Parallel to that, a viscosity formulation following ideas from Lions and Souganidis [46, 47] was proposed by Caruana, Friz, Oberhauser [12, 24]. Recently, a variational approach to evolutionary rough PDEs with transport noise was introduced by Gubinelli, Deya, Hofmanová and Tindel [20] (see also Gubinelli and Bailleul [4]). Modelled on Sobolev spaces, their notion of solution is ‘intrinsic’, in the sense that they work directly at the level of a rough evolution equation, avoiding the use of flow transforms. The mild approach gained some renewed interest in [27], where it was used to prove a Hörmander type theorem for degenerate SPDEs.

When $X = W$ is a Brownian Motion in \mathbf{R}^d (the case of a Wiener Process defined on some abstract Wiener space could be carried out by letting $d = \infty$ in the sum below), and L_t is a deterministic family of operators, one defines a mild solution by the Duhamel Formula

$$u_t - S_{t,0}x = \int_0^t S_{t,r}N(u_r)dr + \sum_{i=1}^d \int_0^t S_{t,r}F_i(u_r)dW_r^i, \quad t \in [0, T], \quad (1.2)$$

where the stochastic integral is understood in the Itô sense, $S_{t,s} \in \mathcal{L}(\mathcal{B}, \mathcal{B})$ for $0 \leq s \leq t \leq T$ is the propagator associated to (L_t) , that is, $S_{t,s}x$ denotes the solution v evaluated at time t (if it exists and is unique), of the linear equation

$$\partial_t v = L_t v_t \quad \text{on } (s, T], \quad v_s := x \in \mathcal{B}. \quad (1.3)$$

As natural as it may look, the formula (1.2) is meaningful only when L_t is a deterministic family. Even assuming adaptedness of L_t does not guarantee that formula (1.2) is meaningful, since in that case the stochastic integrand $S_{t,r}F_i(u_r)$ is only \mathcal{F}_t -measurable (and not \mathcal{F}_r -measurable, as required in standard Itô calculus). This anticipative nature causes major technical difficulties in stochastic contexts where a representation like (1.2) is needed. The situation of random L_t is even more problematic since a solution constructed using Skorohod integral (which would be a natural candidate) is not, generally speaking, a weak solution of (1.1) (see [44]). A solution was nonetheless provided by Leon and Nualart in [45], relying on the formalism of ‘forward integral’ introduced earlier by Russo and Vallois in [54]. Despite being quite successful in the non-autonomous, semilinear scenario, their approach does not seem to easily extend to broader contexts such as quasilinear equations.

The situation when the operator L_t is random arises naturally, for instance in the context of non-degenerate quasilinear SPDEs of the form

$$du_t = A(u_t)u_t dt + \sigma(u_t)dW_t, \quad t \in (0, T], \quad u_0 \in \mathcal{B}, \quad (1.4)$$

where to prove existence and uniqueness one is led to investigate a fixed point for the map

$$v = (v_t(\omega, x)) \mapsto \left(t \mapsto S_{t,0}^v u_0 + \int_0^t S_{t,s}^v \sigma(u_s) dW_s \right).$$

Here $S_{t,s}^v$ denotes the random propagator generated by the elliptic part $L_t = A(v_t)$, and is therefore only \mathcal{F}_t -measurable. In a deterministic setting, this functional analytic approach is due to Amann [2]. Based on an alternative (non-anticipative) formulation for (1.4), we note that similar quasilinear stochastic equations were investigated in the recent work [42], though it remains unclear how to translate Amann’s method to a stochastic context.

Another motivation for considering random generators lies in the study of infinite-dimensional Kolmogorov equations. A concrete example is given by the weak error analysis for the following SPDE

$$dX_t^x = AX_t^x dt + \sigma(X_t^x), \quad X_0^x = x \in H = L^2(0, 1), \quad (1.5)$$

seen as an evolution equation in the Hilbert space H , where the superscript in X^x emphasizes the dependency of the solution w.r.t. the initial datum. Here A is a uniformly elliptic operator (for instance the Dirichlet Laplacian) and σ denotes a Lipschitz continuous Nemytskii operator. In [9] the authors use that the directional derivative $\langle Du_t^x, h \rangle$ of the functional

$u(t, x) = \mathbb{E}[\varphi(X_t^x)]$ for any bounded and continuously differentiable $\varphi: H \rightarrow \mathbf{R}$, is given formally for each $h \in H$ by the anticipative integral

$$\langle Du_t^x, h \rangle = \int_0^t \langle S_{t,s} \sigma'(X_t^x), e^{sA} h dW_s \rangle \quad (1.6)$$

where this time $S_{t,s}$ is the propagator associated with the formal generator ' $L_t := A + \sigma(X_t) \frac{dW_t}{dt}$ '. Otherwise said, $S_{t,s}: H \rightarrow H$ is the solution map $z \mapsto Z_t^s$ for the linear equation

$$dZ_t^s = AZ_t^s dt + \sigma'(X_t^x) Z_t^s dW_t, \quad \text{on } (s, t], \quad Z_s^s = z.$$

The representation (1.6) is a crucial tool in obtaining estimates on the solutions of Kolmogorov equations (see [9, Sec. 5]). These estimates are then needed to address the weak order analysis of an explicit Euler scheme for (1.5). We remark that the authors of the previous paper cannot use (1.6) and have to hinge their analysis on a discretized version of (1.5). Though it is not our intention here to discuss the meaning of the ill-defined product in the corresponding formal generator, we note that the tools developed in Section 4 allow us to make perfect sense of the integral (1.6), and to obtain estimates for it.

Our main objective in this paper is to build quite a broad framework for a pathwise mild solution theory to semilinear SPDEs of the form (1.1). We do this through the theory of rough paths introduced by Lyons [46], in a spirit that is similar to Gubinelli's controlled rough path approach [29]. In keeping with this view, we will look for solutions u of (1.1) that 'locally look like X '. Loosely speaking, (u, u') will be said to be controlled by X if

$$u_t - u_s = u'_s \cdot (X_t - X_s) + R_{t,s},$$

where $R_{t,s} = O(|t - s|^{2\gamma})$ and $\gamma > 1/3$ is the Hölder regularity of X . The main difference with [29] is that here the above control happens not in the space \mathcal{B} where the initial datum lies, but in some larger space. Indeed, as seen from (1.2) one cannot expect u_t to be Hölder continuous in that space, since even the solution to the corresponding linear equation is not. One could potentially work in spaces with a weight at time 0 (see [59] for the attempt in that direction, though authors can only consider F_i that improves spatial regularity), but going to the larger space allows sticking very closely to the classical theory of controlled rough paths. We nevertheless point out that, using the smoothing properties of the propagator $S_{t,s}$, we will show that the solution u is indeed continuous in time with values in the original space \mathcal{B} .

One of the main advantages of the pathwise approach as it was numerous shown in finite dimensions, is that it allows to show that the solution depends continuously on the noise and the initial condition. This is in strong contrast to the approach using Itô calculus, where in general only measurability of the solution map is available. Another advantage is that the notion of the rough integral that we will use is deterministic in its nature and therefore does not rely on the adaptedness of the integrands. This, in fact, allows to treat equation (1.1) with random L_t in a less technical way than the one of Leon and Nualart. We will also show that in the case of adapted L_t , our mild solutions are also weak Itô solutions when X is the Wiener process. Finally, the rough path approach allows to consider equation (1.1) with a Gaussian noise which is not a martingale, in particular allowing to treat the case of a fractional Brownian motion with Hurst parameter less than $1/2$.

Our main result in this part is to state sufficient conditions on N and F under which existence, uniqueness and continuity of the solution map holds for (1.1). It can be loosely formulated as follows (precise statements will be given in Theorems 2.15, 2.18, 5.1, 5.4 and 5.9).

Metatheorem. *Let $(t \mapsto L_t)$ be a Hölder continuous, sectorial family of linear operators acting on a scale of Banach spaces $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$, in such a way that $L_t : \mathcal{B}_\alpha \rightarrow \mathcal{B}_{\alpha-1}$ for any t and α . Fix some initial datum $x \in \mathcal{B}_0$. Consider nonlinearities N and F such that N is polynomial, while F is three times continuously differentiable and subcritical. There exists a unique, local in time, mild solution to (1.1) such that u is controlled by X with Gubinelli derivative $F(u)$. The solution map depends continuously on the initial condition, as well as on the rough path \mathbf{X} . It is also a weak solution in the usual sense.*

Roughly speaking, the above ‘subcriticality’ assumption means that F , seen as a non-linear operator acting on the scale (\mathcal{B}_α) does not cause any loss of regularity greater than the Hölder time-regularity of the rough path \mathbf{X} . This condition illustrates the subtle interplay between space and time regularity that occurs in problems of the form (1.1). An elementary but illustrative example that fits within this solvability theory is the linear equation

$$du_t = \Delta u_t dt + (-\Delta)^\sigma u_t d\mathbf{X}_t, \quad x \in \mathcal{B}_0 := L^p(\mathbf{R}^n), \quad p \in (1, \infty), \quad (1.7)$$

under the subcritical assumption $\sigma < \gamma$, where (\mathcal{B}_α) is the Bessel potential scale (see below). In the case of X being a Brownian motion, we see that σ can be arbitrary close to $1/2$, which in strength agrees to the well-known similar results obtained using mild formulation and Itô calculus (see for instance [10] and the references therein). Note that, in the ‘super-critical’ case $\gamma < 1/2 = \sigma$, there is at least one type of equations that still possess a unique solution, which consists in transport noise of the form $F(u_t) = \sigma \cdot \nabla u_t$. As observed by Deya, Gubinelli, and Tindel [20] (see also [36]), it is possible in this case to prove a priori estimates, which in turn allow inferring existence and uniqueness of solutions. As pointed out in [36, Remark 2.4], in this case, the assumption that the rough path X be *geometric* is essential, in contrast with the subcritical case dealt with in the present work. This limitation constitutes another motivation for us to introduce an alternative formulation.

A secondary objective of our manuscript is to introduce a new version of the multiplicative Sewing Lemma, which allows constructing product integrals in a rather general class of metric spaces (\mathcal{M}, d) equipped with a product operation. Product integrals are going to be the limits of the form

$$\varphi_{t,s} = \lim_{|\pi| \rightarrow 0} \prod_{[v,u] \in \pi} \mu_{v,u}, \quad (1.8)$$

where the limit is taken over arbitrary partitions π of $[s, t]$ with their mesh size going to zero. The novelty of our version is that it will apply to various cases where the limit lives in an infinite dimensional space. The main difference from the usual additive sewing lemma is that the product here is non-commutative in general. Note that a version of multiplicative sewing lemma for the non-commutative products has already been introduced by Feyel, De La Pradelle and Mokobodzki [22] (see also [14, 6, 5] for related works). We point out, however, that Theorem 3.4 below is, in essence, independent of the latter, since the assumptions and conclusions are different. In [22], the authors have to assume that the function $|\cdot| : \mathcal{M} \rightarrow \mathbf{R}_+$ which controls the ‘size’ of $\mu_{t,s}$ in (1.8), is Lipschitz continuous with respect to the distance d . This is mostly not going to be the case for the examples treated below, since in infinite dimensions $|\cdot|$ is often going to be a norm which is stronger than d . For a further discussion, see Remark 3.5.

We will show how to use this version of the Sewing lemma to give a precise meaning to rough equations of the form (1.2). Though this particular step was already achieved in

the previous works [31, 17, 27], the merit of our approach is that our integrands belong to a class of paths which are independent of the propagator $(S_{t,s})_{(s,t) \in \Delta_2}$ generated by $(L_t)_{t \in [0, T]}$. This paves the way to a possible treatment of quasilinear equations where the propagator will depend on the solution itself (we will address this problem in future work). It also has a potential attractive application to unify Lyons' theory of multiplicative functionals with that of Gubinelli and Tindel, for the construction of the rough convolutions

$$z_t := \int_0^t S_{t,r} F(y_r) \cdot d\mathbf{X}_r. \quad (1.9)$$

A basic role in this construction is played by the *affine group* \mathcal{M} , defined as the semi-direct product

$$\mathcal{M} := \mathcal{G} \ltimes \mathcal{B} \quad (1.10)$$

where $\mathcal{G} \subset \mathcal{L}(\mathcal{B})$ is a group of linear bijections operating on \mathcal{B} via the natural action $(T, b) \mapsto Tb$. More precisely, \mathcal{M} is the set of pairs $(T, b) \in \mathcal{G} \times \mathcal{B}$ endowed with the group multiplication

$$(T_1, b_1) \circ (T_2, b_2) := (T_1 T_2, b_1 + T_1 b_2), \quad (T_j, b_j) \in \mathcal{M}, \quad j = 1, 2. \quad (1.11)$$

In practice, because of the parabolic nature of (1.1), it will be necessary to replace \mathcal{G} by an appropriate set of (non-necessarily invertible) linear operators containing $S_{t,s}$ for $0 \leq s \leq t \leq T$. In this case \mathcal{M} is only a monoid, but this is sufficient for our purposes.

Surprisingly enough, our version of the multiplicative Sewing Lemma (i.e. Theorem 3.4), is seen to be useful for purposes that are orthogonal to rough paths theory and the construction of z_t . In particular, it will allow us to construct the propagators $S_{t,s}$ ‘by hand’ (hence reproving classical results from Kato, Tanabe, and Sobolevskii), or to show a version of the Lie-Trotter product formula, for propagators which are generated by the sum of two dissipative operators. Though these results are well-known in principle, the observation that they could be deduced from a Sewing Lemma perspective is new and could be seen as one of our main contributions. Note that, apart from the new construction of the rough convolution, this first part of the paper is essentially independent of the second.

The paper is organized as follows. In Section 2 we explain our functional analytic setting and present our main results. In particular, we will give an existence and uniqueness statement for (1.1), and provide a stochastic example with random coefficients. In Section 3, we prove a general multiplicative sewing lemma and give some applications in the context of (not necessarily rough) evolution equations. Though we believe that these applications are interesting by themselves, they merely consist of establishing new proofs of already known results in functional analysis, and therefore their reading could be avoided at first. In Section 4, we introduce the space of controlled paths $\mathcal{D}_{X, \alpha}^{2\gamma}$ associated to a monotone family $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$ of interpolation spaces, and then apply the multiplicative Sewing Lemma in order to construct the ‘rough convolution’ $\int_0^t S_{t,s} y_s \cdot d\mathbf{X}_s$, where S is the propagator associated to the family $(L_t)_{t \in [0, T]}$ (its existence will be guaranteed by Assumption 2.6). Similar to [29, 31], it will be seen that $\mathcal{D}_{X, \alpha}^{2\gamma}$ is a natural space of integrands for which the rough convolution is well-defined. We point out that, though the construction of rough convolutions was already carried out in [31] (see also [27]), it does require a proof because our definition of the controlled path space is different from the above-mentioned works. As a natural continuation, Section 5 will be devoted to the proof of Theorem 2.15 where

existence, uniqueness, and stability of the solution map are stated for (1.1), in the subcritical case. This will be done via a Picard fixed point argument in the controlled path space. Finally, Section 6 will be devoted to examples. For completeness, we will also address in that section a supercritical case corresponding to the transport noise. In particular, the equivalence between the mild representation (1.2) and the weak solution theory provided in [36] will be discussed.

1.2 General notation

Throughout the paper, $T \in (0, \infty)$ is considered as a fixed, positive time horizon. We denote by $\mathbf{N} := \{1, 2, \dots\}$ and by $\mathbf{N}_0 := \mathbf{N} \cup \{0\}$. Integers will be denoted by \mathbf{Z} , while real numbers (resp. complex numbers) will be denoted by \mathbf{R} (resp. \mathbf{C}). The set $[0, \infty)$ of non-negative real numbers will be denoted by \mathbf{R}_+ . We also denote by $\mathbf{T}^n = \mathbf{R}^n / \mathbf{Z}^n$ the n -dimensional torus. We write $a \lesssim_\gamma b$ provided there exists $C = C(\gamma)$ such that $a \leq Cb$. When C can be chosen independently of γ we write simply $a \lesssim b$.

If X, Y are Banach spaces, we denote by $\mathcal{L}(X, Y)$ the space of continuous linear operators from X to Y , endowed with the operator-norm topology. Similarly, we denote by $\mathcal{L}_s(X, Y)$ the same space as above but equipped with the strong topology. Recall that the strong topology is the coarsest topology that makes the maps $X \ni x \mapsto Sx \in Y$ continuous when S varies in $\mathcal{L}(X, Y)$. For simplicity, we write $\mathcal{L}(X) = \mathcal{L}(X, X)$ and likewise $\mathcal{L}_s(X) = \mathcal{L}_s(X, X)$.

We shall frequently work with the usual Sobolev spaces $W^{\alpha,p}(\mathcal{O}, \mathbf{R}^n)$, $p \geq 1$, $\alpha \in \mathbf{R}$ and a domain $\mathcal{O} \subset \mathbf{R}^n$, equipped with the standard norm $|\cdot|_{W^{\alpha,p}}$. Moreover, we will denote by $W_0^{\alpha,p}(\mathcal{O}, \mathbf{R}^n)$ the closure of $\mathcal{C}_c^\infty(\mathcal{O}, \mathbf{R}^n)$ (the smooth, compactly supported functions) in the topology of $W^{\alpha,p}(\mathcal{O}, \mathbf{R}^n)$. For notational simplicity we write $W^{k,p}(\mathcal{O}) := W^{k,p}(\mathcal{O}, \mathbf{R})$ and similarly for Bessel potential spaces $H^{k,p}(\mathcal{O}) = H^{k,p}(\mathcal{O}, \mathbf{R})$ (see Example 2.4 where these spaces are introduced).

We use the symbol π to denote a generic partition of $[0, T]$ and we shall sometimes blur the difference between thinking of partitions as a set of points and as a set of intervals.

In the sequel, we will need to introduce various norms and seminorms. The ‘rule of thumb’ is that quantities, which are only semi-norms, will be denoted by the brackets $[\cdot]$ while norms will usually be denoted by the simple bars $|\cdot|$. The double bars $\|\cdot\|$ will be used only for the controlled paths spaces $\mathcal{D}_{X,\alpha}^{2\gamma}$ defined in Section 4 and for the norms on functions from the Definition 2.14.

Acknowledgements

The authors would like to thank M. Hairer for many useful discussions and comments. The anonymous referees are warmly thanked for their careful reading of the paper, notably for having pointed out several gaps in the first version. AG gratefully acknowledges the financial support by the Leverhulme Trust through Hendrik Weber’s Philip Leverhulme Prize. AH and TN gratefully acknowledge the financial support by the DFG via Research Unit FOR 2402.

2 Settings and main results

2.1 Functional analytic framework

We will first define a concept of monotone family of interpolation spaces $(\mathcal{B}_\alpha, |\cdot|_\alpha)_{\alpha \in \mathbf{R}}$, which encodes a notion of ‘spatial regularity’ for (1.1).

Definition 2.1. A family of separable Banach spaces $(\mathcal{B}_\alpha, |\cdot|_\alpha)_{\alpha \in \mathbf{R}}$ is called a *monotone family of interpolation spaces* if for every $\alpha \leq \beta$, \mathcal{B}_β is a continuously embedded, dense subspace of \mathcal{B}_α , and if the following interpolation inequality holds: for $\alpha \leq \beta \leq \gamma$ and $x \in \mathcal{B}_\alpha \cap \mathcal{B}_\gamma$:

$$|x|_\beta^{\gamma-\alpha} \lesssim |x|_\alpha^{\gamma-\beta} |x|_\gamma^{\beta-\alpha}. \quad (2.1)$$

The main interest in considering a family as above is the following property, whose proof is evident by interpolation and therefore left to the reader. We write $\Delta_2 = \{(t, s) : T \geq t \geq s \geq 0\}$. If $S : \Delta_2 \rightarrow \mathcal{L}(\mathcal{B}_\alpha) \cap \mathcal{L}(\mathcal{B}_{\alpha+1})$ is such that for each $x \in \mathcal{B}_{\alpha+1}$, and any $(t, s) \in \Delta_2$, $|(S_{t,s} - \text{id})x|_\alpha \lesssim |t-s||x|_{\alpha+1}$ while $|S_{t,s}x|_{\alpha+1} \lesssim |t-s|^{-1}|x|_\alpha$, then for every $\sigma \in [0, 1]$, $S_{t,s}$ belongs to $\mathcal{L}(\mathcal{B}_{\alpha+\sigma})$ and the following estimate holds true:

$$|(S_{t,s} - \text{id})x|_\alpha \lesssim |t-s|^\sigma |x|_{\alpha+\sigma}, \quad |S_{t,s}x|_{\alpha+\sigma} \lesssim |t-s|^{-\sigma} |x|_\alpha. \quad (2.2)$$

Example 2.2 (Hilbert space setting). Let $(\mathcal{H}, |\cdot|)$ be a separable Hilbert space, on which we are given a closed densely defined unbounded operator $L : D(L) \subset \mathcal{H} \rightarrow \mathcal{H}$ whose resolvent set contains a sector $\Sigma_{\vartheta, \lambda} := \{\zeta \in \mathbf{C}, |\arg(\zeta - \lambda)| < \pi/2 + \vartheta\}$ for some $\vartheta > 0$, $\lambda \in \mathbf{R}$, and such that for every $\zeta \in \Sigma_{\vartheta, \lambda}$, $|(\zeta - L)^{-1}| \leq \Lambda(1 + |\zeta|)^{-1}$, with $\Lambda > 0$ independent on ζ . An operator L satisfying this property is called sectorial.

For $\alpha > 0$ we can define the fractional powers $(-L)^{-\alpha}$ through the formula [52, Eq. (6.3)]. Next, introduce the space

$$\mathcal{H}_\alpha := \text{Im}(-L)^{-\alpha} \subseteq \mathcal{H}, \quad (2.3)$$

endowed with the norm $|x|_\alpha = |(-L)^\alpha x|$. Additionally, we define $\mathcal{H}_{-\alpha}$ as the completion of \mathcal{H} with respect to the norm $|((-L)^\alpha)^{-1} \cdot |$ (the fractional powers of L are one-to-one thanks to the sector condition). The interpolation inequality (2.1) can be proved using spectral decomposition and Hölder inequality (see [33, Sec 6]).

Example 2.3. Let $(\mathcal{B}_k, |\cdot|_k)_{k \in \mathbf{Z}}$ be a monotone family of reflexive Banach spaces, in the sense that for each $k \in \mathbf{Z}$, $\mathcal{B}_{k+1} \hookrightarrow \mathcal{B}_k$ (densely) and (2.1) is satisfied for every $\alpha, \beta, \gamma \in \mathbf{Z}$. Then for $\theta \in [0, 1]$ and $k \in \mathbf{Z}$ we can define a space $\mathcal{B}_{k+\theta}$ by complex interpolation:

$$\mathcal{B}_{k+\theta} := [\mathcal{B}_k, \mathcal{B}_{k+1}]_\theta. \quad (2.4)$$

For the precise definition and properties of complex interpolation spaces see [11, 49]. With this definition, it can be shown that $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$ is a monotone family of interpolation spaces. The reflexivity of \mathcal{B}_k is necessary in order to guarantee the consistency relation $[\mathcal{B}_k, \mathcal{B}_{k+1}]_0 = \mathcal{B}_k$ and $[\mathcal{B}_k, \mathcal{B}_{k+1}]_1 = \mathcal{B}_{k+1}$ implying that for all $\alpha \leq \beta < \gamma$ we have $\mathcal{B}_\beta = [\mathcal{B}_\alpha, \mathcal{B}_\gamma]_{\frac{\beta-\alpha}{\gamma-\alpha}}$ which in turns implies (2.1) by [49, Thm 2.6].

Note that given a Hilbert space \mathcal{H} and L as in Example 2.2 which is self-adjoint, one can construct $(\mathcal{H}_\alpha)_{\alpha \in \mathbf{R}}$ using the so called *Sobolev tower*. First construct $(\mathcal{H}_k)_{k \in \mathbf{Z}}$ (which do not require fractional powers of L) and then define $(\mathcal{H}_\alpha)_{\alpha \in \mathbf{R}}$ via the complex interpolation

as above. If L is self-adjoint then $(\mathcal{H}_k)_{k \in \mathbf{Z}}$ will be reflexive and [49, Thm 4.17] guarantees this construction of $(\mathcal{H}_\alpha)_{\alpha \in \mathbf{R}}$ is equivalent to the one in Example 2.2.

Another example, is to take $\mathcal{B}_k = W^{k,p}(\mathcal{O})$ for $k \in \mathbf{Z}$ and $p \in (1, \infty)$ and where \mathcal{O} is equal to \mathbf{R}^n or \mathbf{T}^n , or is a smooth domain inside \mathbf{R}^n .

Example 2.4. Another useful family of Banach spaces consists of the Bessel potential spaces $H^{s,p}(\mathbf{R}^n)$, for $1 < p < \infty$ and $\alpha \in \mathbf{R}$, that can be defined in terms of Fourier transform \mathcal{F} of tempered distributions $\mathcal{S}'(\mathbf{R}^n)$ as

$$H^{\alpha,p}(\mathbf{R}^n) := \left\{ f \in \mathcal{S}'(\mathbf{R}^n), \|f\|_{H^{\alpha,p}} := \|\mathcal{F}^{-1}(1 + |\xi|^2)^{\alpha/2} \mathcal{F}f\|_{L^p} < \infty \right\}.$$

We recall the well known characterization using the complex interpolation $[\cdot, \cdot]$ (see [49])

$$H^{k\theta,p}(\mathbf{R}^n) = [L^p(\mathbf{R}^n), W^{k,p}(\mathbf{R}^n)]_\theta,$$

for each $\theta \in [0, 1]$ and $k \in \mathbf{Z}$. Moreover, it holds $W^{\alpha,p}(\mathbf{R}^n) = H^{\alpha,p}(\mathbf{R}^n)$ when α is an integer, or when $p = 2$ for any $\alpha \in \mathbf{R}$, and for any $\epsilon > 0$ we have the continuous embedding $W^{\alpha+\epsilon,p}(\mathbf{R}^n) \hookrightarrow H^{\alpha,p}(\mathbf{R}^n) \hookrightarrow W^{\alpha-\epsilon,p}(\mathbf{R}^n)$ for every $\alpha \in \mathbf{R}$ and $p > 1$. One can define $H^{\alpha,p}(\mathbf{T}^n)$ similarly through the Fourier transform on the torus and establish analogous properties.

For a smooth and bounded domain $\mathcal{O} \subset \mathbf{R}^n$, the space $H^{\alpha,p}(\mathcal{O})$ can be defined algebraically as the set of restrictions $u = f|_{\mathcal{O}}$ of elements $f \in H^{\alpha,p}(\mathbf{R}^n)$. One then defines the norm of $u \in H^{\alpha,p}(\mathcal{O})$ to be the infimum of $\|f\|_{H^{\alpha,p}(\mathbf{R}^n)}$ over all such functions (see [58, Chap. 3]). We also define $H_0^{\alpha,p}(\mathcal{O})$ to be the closure of $\mathcal{C}_c^\infty(\mathcal{O})$ in $H^{\alpha,p}(\mathcal{O})$. An important consequence of the above definition is that Bessel spaces respect complex interpolation

$$H^{\alpha+\theta,p}(\mathcal{O}) = [H^{\alpha,p}(\mathcal{O}), H^{\alpha+1,p}(\mathcal{O})]_\theta, \quad (2.5)$$

for every $\alpha \in \mathbf{R}$, $\theta \in [0, 1]$ and same holds for $H_0^{\alpha,p}(\mathcal{O})$. This together with reflexivity of Bessel potential spaces guarantees that both $H^{\alpha,p}(\mathcal{O})$ and $H_0^{\alpha,p}(\mathcal{O})$ generate a monotone family of interpolations spaces.

2.2 Hölder spaces and controls

For $n \geq 2$ and a Banach space V , we define $\mathcal{C}_n(0, T; V)$ to be the space of continuous functions from the simplex $\Delta_n = \{T \geq t_n \geq t_{n-1} \geq \dots \geq t_1 \geq 0\}$ to V . For $n = 1$ we adopt the convention that $\Delta_1 := [0, T]$ while $\mathcal{C}(0, T; V) \equiv \mathcal{C}_1(0, T; V)$ is just the usual space of continuous functions taking values in V . In the sequel, we will be only interested in the cases $n = 1, 2, 3$. If V is a Banach space and

$$S : \Delta_2 \rightarrow \mathcal{L}(V)$$

is a two-parameter family of bounded linear maps, we define the increment operator δ^S for $f : \Delta_1 \rightarrow V$ and $g : \Delta_2 \rightarrow V$ as

$$\delta^S f_{t,s} := f_t - S_{t,s} f_s, \quad \text{for } (t, s) \in \Delta_2; \quad (2.6)$$

while

$$\delta^S g_{t,u,s} := g_{t,s} - g_{t,u} - S_{t,u} g_{u,s}, \quad \text{for } (t, u, s) \in \Delta_3, \quad (2.7)$$

and we recall (see [31]) that $\text{Im } \delta^S = \text{Ker } \delta^S$.¹ When $S = \text{id}$, δ^S corresponds to the usual increment operator from controlled paths theory, and we shall use the notation $\delta := \delta^{\text{id}}$.

For $f \in \mathcal{C}_1(0, T; \mathcal{B}_\alpha)$ we let

$$|f|_{0,\alpha} = \sup_{0 \leq t \leq T} |f_t|_\alpha.$$

If $\gamma > 0$, the norm in $\mathcal{C}_1^\gamma(0, T; \mathcal{B}_\alpha)$ is defined as the usual Hölder norm, namely

$$|f|_{\gamma,\alpha} := |f|_{0,\alpha} + [\delta f]_{\gamma,\alpha}. \quad (2.8)$$

where for $g = (g_{t,s}) : \Delta_2 \rightarrow \mathcal{B}_\alpha$, we let $[g]_{\gamma,\alpha}$ be the quantity

$$[g]_{\gamma,\alpha} := \sup_{0 \leq s < t \leq T} \frac{|g_{t,s}|_\alpha}{|t - s|^\gamma}. \quad (2.9)$$

Equipped with $[\cdot]_{\gamma,\alpha}$, the space $\mathcal{C}_2^\gamma(0, T; \mathcal{B}_\alpha)$ of all families such that the above quantity is finite forms a Banach space. If $h = (h_{t,u,s}) : \Delta_3 \rightarrow \mathcal{B}_\alpha$, we let

$$[h]_{\gamma_1,\gamma_2,\alpha} := \sup_{(t,u,s) \in \Delta_3} \frac{|h_{t,u,s}|_\alpha}{|t - u|^{\gamma_1} |u - s|^{\gamma_2}}, \quad (2.10)$$

and we denote by $\mathcal{C}_3^{\gamma_1,\gamma_2}(0, T; \mathcal{B}_\alpha)$ the Banach space formed by all 3-indices elements as above such that $[h]_{\gamma_1,\gamma_2,\alpha} < \infty$. All the previous norms can be taken with \mathcal{B}_α replaced by \mathcal{B}_α^m for $m \in \mathbf{N}$, though we will still use the notations $|\cdot|_{\gamma,\alpha}$, $[\cdot]_{\gamma,\alpha}$ etc. throughout the paper. For paths living in a more general Banach space V , the notations $|\cdot|_{\gamma,V}$, $[\cdot]_{\gamma,V}$, $[\cdot]_{\gamma_1,\gamma_2,V}$ will be used instead (with obvious changes in the definitions).

We will also work with functions that exhibit a uniform continuity ‘similar to Hölder’ but in a weaker sense.

Definition 2.5. We say that a function $\omega : \Delta_2 \rightarrow \mathbf{R}_+$ is a control if it is a continuous map with $\omega(s, s) = 0$ for all $s \in [0, T]$, superadditive in the sense that

$$\omega(t, r) + \omega(r, s) \leq \omega(t, s),$$

for all $0 \leq s \leq r \leq t \leq T$.

For $p \geq 1$ and a Banach space $(V, |\cdot|_V)$ define a space $\mathcal{C}_2^{p\text{-var}}(0, T; V)$ of all $g : \Delta_2 \rightarrow V$ such that there exists a control ω with $|g_{t,s}|_V \leq \omega^{1/p}(t, s)$.

Note that functions in $\mathcal{C}_2^{p\text{-var}}$ are continuous by the definition of a control. Note also that $\mathcal{C}_2^\gamma \subset \mathcal{C}_2^{1/\gamma\text{-var}}$ because $\omega(t, s) = C|t - s|$ is a control for all $C > 0$. Another example of a control can be obtained using p -variation. Let $g : \Delta_2 \rightarrow V$ be a continuous function such that:

$$\omega_g(t, s) = \sup_{\pi(t,s)} \sum_{t_i \in \pi(t,s)} |g_{t_{i+1}, t_i}|_V^p < \infty,$$

where the above supremum ranges over all partitions $\pi(t, s)$ of $[s, t]$. Then $\omega_g(t, s)$ defines a control and $|g_{t,s}|_V \leq \omega_g^{1/p}(t, s)$, thus continuous functions of finite p -variation lie in the space $\mathcal{C}_2^{p\text{-var}}$.

¹Here δ^S on the left is from 2.6 and δ^S on the right is from 2.7.

2.3 Assumptions and main results

We first state our assumptions on the family $(L_t)_{t \in [0, T]}$. In what follows, we shall fix a number $\vartheta > 0$, $\lambda \in \mathbf{R}$ and define a sector $\Sigma_{\vartheta, \lambda}$ of the complex plane as follows:

$$\Sigma_{\vartheta, \lambda} := \{\zeta \in \mathbf{C}, |\arg(\zeta - \lambda)| < \pi/2 + \vartheta\} \quad (2.11)$$

with the convention adopted throughout the paper that $\arg 0 = 0$, thus in particular $[\lambda, \infty) \subset \Sigma_{\vartheta, \lambda}$.

In the next statements, we assume that we are given two Banach spaces

$$(\mathcal{X}_1, |\cdot|_1) \subset (\mathcal{X}_0, |\cdot|_0),$$

where the embedding is continuous and dense.

Assumption 2.6. Let $(L_t)_{t \in [0, T]}$ be a family of closed, densely defined linear operators on \mathcal{X}_0 with domains containing \mathcal{X}_1 , and such that there exist constants $\Lambda, M > 0$ (depending only on T) such that:

- (L1) For each $t \in [0, T]$, the resolvent set $\rho(L_t)$ contains $\Sigma_{\vartheta, \lambda}$ and there exists a constant $\Lambda > 0$ such that for $i = 0, 1$,

$$|(\zeta - L_t)^{-1}|_{\mathcal{L}(\mathcal{X}_i)} \leq \Lambda(1 + |\zeta|)^{-1}, \quad \forall \zeta \in \Sigma_{\vartheta, \lambda}. \quad (2.12)$$

- (L2) For any $t \in [0, T]$, we have

$$|(\zeta - L_t)^{-1}|_{\mathcal{L}(\mathcal{X}_0, \mathcal{X}_1)} \leq M, \quad \forall \zeta \in \Sigma_{\vartheta, \lambda}.$$

- (L3) There exists a control ω and $\varrho \in (0, 1]$ such that for all $(t, s) \in \Delta_2$:

$$|L_t - L_s|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} \leq \omega^\varrho(t, s).$$

- (L4) The above control satisfies the following integrability property: for all $(t, s) \in \Delta_2$

$$\int_s^t \frac{\omega^\varrho(r, s) dr}{r - s} < \infty.$$

Let $(L_t)_{t \in [0, T]}$ be such that Assumption 2.6 is satisfied. From classical results obtained independently by Tanabe and Sobolevskii in the 60's [57, 55] (see also the seminal work of Kato [40]), it is known that for $x \in \mathcal{X}_0$, the equation

$$\partial_t u = L_t u_t, \quad u_s := x \in \mathcal{X}_0, \quad (2.13)$$

admits a unique solution (in the usual weak, PDE sense) denoted by $S_{t,s}x$ and which depends linearly on $x \in \mathcal{X}_0$. The two parameter mapping $S: \Delta_2 \rightarrow \mathcal{L}(\mathcal{X}_0) \cap \mathcal{L}(\mathcal{X}_1)$ is usually referred to as the *propagator* associated to the family $(L_t)_{t \in [0, T]}$ (it is sometimes called the ‘parabolic fundamental solution’ associated with L ., but for conciseness we adopt the terminology used in [53]). The family $(S_{t,s})_{(t,s) \in \Delta_2}$ should be heuristically understood as ‘ $\exp(\int_s^t L_r dr)$ ’ and its main interest lies in the existence of the Duhamel-type formula (1.2).

The precise definition of a propagator is as follows.

Definition 2.7. We say that $S: \Delta_2 \rightarrow \mathcal{L}(\mathcal{X}_0)$ is a *propagator* associated to the family $(L_t)_{t \in [0, T]}$ of unbounded, closed operators with dense domains containing \mathcal{X}_1 if and only if the following holds:

(P1) $S \in \mathcal{C}_2(0, T; \mathcal{L}_s(\mathcal{X}_0))$ and there exist constants $\lambda, \Lambda > 0$ such that for every $(t, s) \in \Delta_2$

$$|S_{t,s}|_{\mathcal{L}(\mathcal{X}_0)}, |S_{t,s}|_{\mathcal{L}(\mathcal{X}_1)} \leq \Lambda e^{\lambda(t-s)}. \quad (2.14)$$

(P2) $S_{t,t} = \text{id}$ and $S_{t,s} = S_{t,u}S_{u,s}$ for $(t, u, s) \in \Delta_3$.

(P3) There exists $C_T > 0$ such that for every $(t, s) \in \Delta_2, s \neq t$:

$$|S_{t,s} - \text{id}|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} \leq C_T |t - s|.$$

(P4) For all $s, t \in [0, T]$ and $x \in \mathcal{X}_1$ we have:

$$\frac{d}{dt} S_{t,s} x = L_t S_{t,s} x \quad \text{and} \quad \frac{d}{ds} S_{t,s} x = -S_{t,s} L_s x,$$

where the differentiation is taking place in the Banach space \mathcal{X}_0 .

(P5) For every $(t, s) \in \Delta_2, s \neq t$ we have that $S_{t,s} \mathcal{X}_0 \subset \mathcal{X}_1$ and for some constant $N_T > 0$

$$|L_t S_{t,s}|_{\mathcal{L}(\mathcal{X}_0)} \lesssim |S_{t,s}|_{\mathcal{L}(\mathcal{X}_0, \mathcal{X}_1)} \leq N_T |t - s|^{-1}. \quad (2.15)$$

Remark 2.8. If (L_t) satisfies Assumption 2.6 then the first inequality in (2.15) can be justified as follows. We claim that for each $t \in [0, T]$ and $y \in \mathcal{X}_1$

$$|L_t y|_0 \leq C_t |y|_1. \quad (2.16)$$

Using this bound at $t = 0$ together with (L3) implies uniformly for $t \in [0, T]$

$$|L_t y|_0 \leq (C_0 + \omega^q(0, T)) |y|_1.$$

It remains to show (2.16). Fix $t \in [0, T]$, the assumption on the closure of L_t shows that $\mathcal{Y} = D(L_t)$ is a complete Banach space with respect to the graph norm $x \mapsto |x|_0 + |L_t x|_0$. It is therefore enough to show that the inclusion $\mathcal{X}_1 \subset \mathcal{Y}$ is continuous i.e. that $|x|_0 + |L_t x|_0 \lesssim |x|_1$ for every $x \in \mathcal{X}_1$. By the Closed Graph Theorem it is equivalent to show that if $x_n \rightarrow x$ in \mathcal{X}_1 and $x_n \rightarrow x'$ in \mathcal{Y} then $x = x'$. Then, since $(\mathcal{X}_1, |\cdot|_1)$ is continuously embedded in $(\mathcal{X}_0, |\cdot|_0)$, we have for such sequences

$$|x_n - x|_0 \rightarrow 0 \quad \text{and} \quad |x_n - x'|_0 + |L_t(x_n - x')|_0 \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

This trivially implies $x = x'$ by the uniqueness of the limit in \mathcal{X}_0 , thus showing (2.16).

As is well-known when $\omega(t, s) = C|t - s|$ for some $C > 0$, Assumption 2.6 guarantees that the family $(L_t)_{t \in [0, T]}$ generates a propagator (see [57, 55]). Though the general case should be known in principle, we are not aware of any reference in the p -variation setting (see nevertheless [40] for $p = 1$).

Remark 2.9. Note that in general if L_t is the generator of an analytic semigroup then it satisfies a resolvent bound of the form (2.12) for some $\Lambda \geq 1$. As will be seen later, in order to be able to recover the existence and uniqueness of the propagators from the Sewing Lemma (Theorem 3.4), we need to restrict ourselves to the case when $\Lambda = 1$ in (L1). This is, however, not restrictive as far as Theorems 2.15 and 2.18 are concerned, since for the case $\Lambda > 1$ we can simply refer to the classical results of Tanabe and Sobolevskii [57, 55]. Note that these results also guarantee that Λ in (2.12) and Λ in (2.14) are the same. We shall see that this is also true in our case $\Lambda = 1$.

A criterion that guarantees that $\Lambda = 1$ in (L1), is the dissipativity of the operators L_t for each $t \in [0, T]$. Recall that an unbounded operator A on a Banach space V is *dissipative* if

$$\langle Au, u^* \rangle \leq 0 \quad \forall u \in D(A) \subset V, \quad u^* \in V^* \quad \text{s.t.} \quad \langle u, u^* \rangle = |u|^2 = |u^*|^2, \quad (2.17)$$

where $\langle u, u^* \rangle$ denotes the value of u^* at u . Then, it is well-known (see [52, 28]) that if for a dissipative operator A and $\zeta \in \rho(A)$ one has the image of $\zeta - A$ being equal to V , then $|(\zeta - A)^{-1}|_{\mathcal{L}(V)} \leq (1 + |\zeta|)^{-1}$. For an overview on dissipative operators on Banach spaces, we refer for instance to [48].

Theorem 2.10 (Tanabe/Sobolevskii). *Assume that the Banach spaces \mathcal{X}_0 and \mathcal{X}_1 are reflexive. Let $(L_t)_{t \in [0, T]}$ be a family of operators satisfying the properties (L1), (L3) of Assumption 2.6, with the additional hypothesis that $\Lambda = 1$ in (2.12). Then, there exists a unique map*

$$S: \Delta_2 \rightarrow \mathcal{L}(\mathcal{X}_0),$$

satisfying properties (P1), (P2), (P3), (P4), and in addition for all $(t, s) \in \Delta_2$ we have the property²

$$|S_{t,s} - e^{(t-s)L_s}|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} \lesssim_T |t - s| \omega^\varrho(t, s). \quad (2.18)$$

Moreover, one has in (2.14) that $\Lambda = 1$ and λ is the same as in the sector $\Sigma_{\vartheta, \lambda}$. Finally, if (L4) is also satisfied, then S is the propagator associated to the family $(L_t)_{t \in [0, T]}$ (in the sense of Definition 2.7).

In Section 3, we will provide a whole new proof of the above result which is based on the ‘multiplicative sewing lemma’ Theorem 3.4. We would like to point out that in order to meet the hypotheses of Theorem 3.4, in the above theorem we have to assume that both \mathcal{X}_0 and \mathcal{X}_1 are reflexive (see Lemma 3.7). Note that [57] does not need to assume reflexivity, while [40] does assume reflexivity as well.

For our purposes, we will need the propagators to have stronger properties and act not only on two spaces \mathcal{X}_0 and \mathcal{X}_1 but on a continuous range of interpolation spaces $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$ for $\alpha \in I$ where $I \subset \mathbf{R}$ is an interval. For this purpose, stronger assumptions on the family L_t will be required. We have the following definition.

Definition 2.11. Let $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$ be a monotone family of interpolation spaces, and fix an interval $I \subset \mathbf{R}$. We say that S is a propagator on the full range $(\mathcal{B}_\alpha)_{\alpha \in I}$ if for every $\alpha \in I$, S restricted to \mathcal{B}_α is itself a propagator, in the sense of Definition 2.7 for the pair $(\mathcal{X}_0, \mathcal{X}_1) = (\mathcal{B}_\alpha, \mathcal{B}_{\alpha+1})$.

We now give a concrete example of a family $(L_t)_{t \in [0, T]}$ where Assumption 2.6 is fulfilled with $\Lambda = 1$.

²The existence of the exponential in (2.18) is implied by the property (L1), as will be seen in Section 3.

Example 2.12. Let $1 < p < \infty$ and define

$$\mathcal{X}_0 := L^p(\mathbf{T}^n) \quad \text{and} \quad \mathcal{X}_1 := W^{2,p}(\mathbf{T}^n).$$

Then \mathcal{X}_1 is a constant domain associated to the family $(L_t)_{t \in [0, T]}$ defined for each $t \in [0, T]$ as:

$$L_t u := \nabla \cdot (a_t(x) \nabla u), \quad u \in \mathcal{X}_1,$$

where $a_t(x) \in \mathbf{R}^{n \times n}$ is a matrix satisfying the following uniform ellipticity condition: there exists a constant $\varkappa > 0$ such that

$$\sum_{j,k} a_t^{jk}(x) \xi^j \xi^k \geq \varkappa |\xi|^2, \quad (2.19)$$

for every $t \in [0, T]$, $x \in \mathbf{T}^n$ and $\xi \in \mathbf{R}^n$. Moreover, we assume that for every $t \in [0, T]$,

$$a_t(\cdot) \in \mathcal{C}^2(\mathbf{T}^n; \mathcal{L}(\mathbf{R}^n)). \quad (2.20)$$

The results of [52, Sec. 7.3] imply that for every $t \in [0, T]$ the operator L_t is sectorial, and since above \varkappa is independent of t one can choose the same ϑ and λ in $\Sigma_{\vartheta, \lambda}$ for every $t \in [0, T]$. Moreover, it is classical (see for instance [15]) that for each $y \in [0, T]$ the operator L_t is dissipative and satisfies property (L1) with $\Lambda = 1$, but since the proof is relatively simple, we now give a short argument why this is true in the case $p \geq 2$ (the case when $p \in (1, 2)$ can be done similarly). Fix $t \in [0, T]$. The fact that $\text{Im}(\zeta - L_t) = \mathcal{X}_0$ for all $\zeta \in \rho(L_t)$ is well-known for spatially \mathcal{C}^2 coefficients (see again [52, Sec. 7.3]). We now want to show that L_t is dissipative in the sense of (2.17).

Let $f \in \mathcal{C}^\infty(\mathbf{T}^n)$ be non zero and define

$$g(x) = \frac{|f(x)|^{p-2}}{|f|_{L^p}^{p-2}} f(x).$$

Let q be such that $p^{-1} + q^{-1} = 1$ then we have

$$|f|_{L^p}^2 = |g|_{L^q}^2 = \int_{\mathbf{T}^n} f(x) g(x) dx.$$

We can now use this g and the uniform ellipticity (2.19) to deduce

$$\begin{aligned} \langle L_t f, g \rangle &= - \int_{\mathbf{T}^n} a_t(x) \nabla f(x) \cdot \nabla g(x) dx \\ &= - |f|_{L^p}^{2-p} \int_{\mathbf{T}^n} a_t(x) \nabla f(x) \cdot \nabla (|f|^{p-2}(x) f(x)) \\ &= - |f|_{L^p}^{2-p} (p-1) \int_{\mathbf{T}^n} |f(x)|^{p-2} a_t(x) \nabla f(x) \cdot \nabla f(x) dx \\ &\leq -\varkappa |f|_{L^p}^{2-p} (p-1) \int_{\mathbf{T}^n} |f(x)|^{p-2} |\nabla f(x)|^2 \leq 0. \end{aligned}$$

By density, this inequality can be extended to an inequality for all $f \in W^{2,p}(\mathbf{T}^n)$. This shows that L_t is dissipative, hence $\Lambda = 1$.

Moreover, the operators $(L_t)_{t \in [0, T]}$ satisfy (L2) but potentially not uniformly in t . If in addition, there exists $p \geq 1$ and a control ω such that for all $(t, s) \in \Delta_2$

$$\sup_{x \in \mathbf{T}^n} |a_t(x) - a_s(x)| \leq \omega^{1/p}(t, s),$$

then (L2) is satisfied uniformly and moreover (L3) holds true with $\varrho = 1/p$. (For justification in a case of Hölder continuous $t \mapsto a_t(x)$ see [1, Section 14], a general control case would follow by using a time change.) The property (L4) is satisfied for example in the case of Hölder continuity i.e when $\omega(t, s) = C|t - s|$.

Finally, if the condition (2.20) is replaced by the stronger assumption that $a_t(\cdot) \in \mathcal{C}^\infty(\mathbf{T}^n; \mathcal{L}(\mathbf{R}^n))$, then S extends uniquely to a propagator on the full range $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$ (in the sense of Definition 2.11), where the $\mathcal{B}_\alpha = H^{2\alpha, p}(\mathbf{T}^n)$ are the Bessel potential spaces from Example 2.4. See Example 3.10 for more details.

Next, in order to present our main results on the evolution equation (1.1), we first need to recall the definition of a two-step rough path.

Definition 2.13 (Rough Path). Let $\gamma > 1/3$. We define the space of rough paths $\mathcal{C}^\gamma(0, T; \mathbf{R}^d)$ to consist of the pairs $(X, \mathbb{X}) =: \mathbf{X}$ such that $X \in \mathcal{C}^\gamma(0, T; \mathbf{R}^d)$, $\mathbb{X} \in \mathcal{C}_2^{2\gamma}(0, T; \mathbf{R}^{d \times d})$ are satisfying the Chen's relation:

$$\delta \mathbb{X}_{t,u,s}^{i,j} := \mathbb{X}_{t,s}^{i,j} - \mathbb{X}_{t,u}^{i,j} - \mathbb{X}_{u,s}^{i,j} = \delta X_{t,u}^i \delta X_{u,s}^j, \quad (2.21)$$

for every $(t, u, s) \in \Delta_3$ and $1 \leq i, j \leq d$. The rough paths space is equipped with the pseudometric

$$\varrho_\gamma(\mathbf{X}, \tilde{\mathbf{X}}) = [X - \tilde{X}]_\gamma + [\mathbb{X} - \tilde{\mathbb{X}}]_{2\gamma},$$

where the quantities $[X]_\gamma$ resp. $[\mathbb{X}]_{2\gamma}$ are the Hölder seminorms defined in (2.9). For simplicity, we shall write in the sequel $\varrho_\gamma(\mathbf{X}) := \varrho_\gamma(0, \mathbf{X})$.

Now, we need to restrict our study to a suitable class of non-linearities.

Definition 2.14. Let $\mathcal{B} := (\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$ be a monotone family of interpolation spaces. For some fixed $\alpha, \beta \in \mathbf{R}$ and $k \in \mathbf{N}_0$ we define the space $\mathcal{C}_{\alpha, \beta}^k(\mathcal{B}^m, \mathcal{B}^n)$ as the space of k -differentiable functions $G: \mathcal{B}_\theta^m \rightarrow \mathcal{B}_{\theta+\beta}^n$ for every $\theta \geq \alpha$, and $n, m \in \mathbf{N}_0$ and such that $D^i G$ sends bounded subsets of \mathcal{B}_θ^m to bounded sets of $\mathcal{B}_{\theta+\beta}^n$, for all $i = 0, \dots, k$.

Similarly, define $\text{Lip}_{\alpha, \beta}(\mathcal{B}^m, \mathcal{B}^n)$ to be a space of Lipschitz functions $\mathcal{B}_\theta^m \rightarrow \mathcal{B}_{\theta+\beta}^n$ for all $\theta \geq \alpha$ that send bounded sets to bounded sets and such that the Lipschitz constant is uniformly bounded on bounded sets. When $m = n$ we will simply write $\mathcal{C}_{\alpha, \beta}^k(\mathcal{B}^m)$ or $\text{Lip}_{\alpha, \beta}(\mathcal{B}^m)$.

We now have all at hand to introduce our main result on the existence and uniqueness of solutions.

Theorem 2.15 (Solvability of (1.1)). *Fix a two-step rough path $\mathbf{X} = (X, \mathbb{X})$ in $\mathcal{C}^\gamma(0, T; \mathbf{R}^d)$ with $\gamma > 1/3$, and consider a monotone family of interpolation spaces $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$. Assume that $(L_t)_{t \in [0, T]}$ is a given family of linear operators generating the propagator $(S_{t,s})_{(t,s) \in \Delta_2}$ on $(\mathcal{B}_\alpha)_{\alpha \in (-3\gamma, 0]}$. For some $\sigma < \gamma$, assume that we are given a non-linearity $F \in \mathcal{C}_{-2\gamma, -\sigma}^3(\mathcal{B}, \mathcal{B}^d)$ and $N \in \text{Lip}_{\alpha, -\delta}(\mathcal{B})$ for some $0 \leq \delta < 1$.*

Then, for every $x \in \mathcal{B}_0$, there exists a maximal time $\tau \in (0, T]$ and a unique function $u \in \mathcal{C}([0, \tau], \mathcal{B}_0)$ such that $(u, F(u))$ is a controlled rough path in the sense of Definition 4.3 and u is a mild solution to the Rough PDE (1.1), namely

$$u_t = S_{t,0}x + \int_0^t S_{t,r}N(u_r)dr + \int_0^t S_{t,r}F(u_r) \cdot d\mathbf{X}_r, \quad t < \tau, \quad (2.22)$$

where the latter integral is understood in the rough integral sense of Theorem 4.5.

Remark 2.16. When the rough path X is β -Hölder with $\beta > 1/2$, then u is a solution to the mild equation (2.22) where all the integrals are well-defined using Young integration.

Remark 2.17. It was recently shown by Lê [43] that the additive sewing Lemma of [29] has a natural extension in a stochastic setting, exploiting a certain martingale decomposition. As already emphasized in (1.7), the value $\sigma = \gamma$ is critical, even though Itô solutions exist and are unique for $\sigma = \frac{1}{2}$ and X is Brownian so that $\gamma = \frac{1}{2} - \varepsilon$ for any $\varepsilon > 0$ (to avoid issues related to stochastic parabolicity, note however that (1.7) has to be understood first as a Stratonovich equation, then corrected to an equivalent Itô form). It should be possible to deal simultaneously with the semigroup $\exp(t\Delta)$ and the semi-martingale structure of the right-hand side, in order to recover such solvability results. We chose nevertheless to leave this question for future investigations.

When talking about the mild solutions it is natural to ask whether these coincide with the weak solutions. The following statement can be considered as an answer to this question.

Theorem 2.18. *Let (\mathcal{B}_α) , (L_t) , N , F , x be as in Theorem 2.15 and let $\nu = \max\{1, \sigma + 2\gamma\}$. Then for all $\varphi \in \mathcal{B}_{-\nu}^*$ the following integral formula holds:*

$$\langle u_t, \varphi \rangle = \langle x, \varphi \rangle + \int_0^t \langle L_s u_s, \varphi \rangle ds + \int_0^t \langle N(u_s), \varphi \rangle ds + \int_0^t \langle F(u_s), \varphi \rangle \cdot d\mathbf{X}_s, \quad (2.23)$$

where the last integral is the usual rough integral in the sense of [25, Theorem 4.10] and $L_s u_s$ is viewed as an element of \mathcal{B}_{-1} . Conversely, if $(u, F(u))$ is a controlled rough path in the sense of Definition 4.3 and (2.23) holds for all $\varphi \in \mathcal{B}_{-\nu}^*$, then it is also a mild solution, namely (2.22) holds.

For the precise statement, we refer the reader to the Theorem 5.9, where we also show that weak solutions are mild solutions.

2.4 An illustrating example: non-autonomous stochastic reaction-diffusion equations

Consider the non-autonomous evolution problem

$$du_t(x) = \nabla \cdot (a_t(x) \nabla u_t(x)) dt + f(u_t(x)) dt + \sum_{i=1}^d p_i(u_t(x)) d\mathbf{X}_t^i, \quad (2.24)$$

$$u_0 \in H^{k,p}(\mathbf{T}^n),$$

with $(a^{ij}) \in \mathcal{C}^{\varrho,\infty}([0, T] \times \mathbf{T}^n; \mathcal{L}(\mathbf{R}^n))$, $1 \leq i, j \leq n$, for some $\varrho > 0$ as in Example 2.12. We assume that f and p_i are polynomials with coefficients in $H^{k,p}(\mathbf{T}^n)$ i.e.

$$f(x, u(x)) = \sum_{j=0}^m h_j(x) u^j(x) \quad x \in \mathbf{T}^n,$$

for some $m \in \mathbf{N}_0$ and $h_j \in H^{k,p}(\mathbf{T}^n)$ for every $j \in \{1, \dots, m\}$, and p_i has a similar form for all $i \in \{1, \dots, d\}$. Furthermore, $X_t = (X_t^1, X_t^2, \dots, X_t^d)$ is assumed to be endowed with a rough path enhancement $\mathbf{X} = (X, \mathbb{X}) \in \mathcal{C}^\gamma(\mathbf{R}_+, \mathbf{R}^d)$ with $\gamma > 1/3$.

Fix $p \in (1, \infty)$ and for $\alpha \in \mathbf{R}$, define $\mathcal{B}_\alpha := H^{k+2\alpha,p}(\mathbf{T}^n)$ (this is indeed a monotone family of interpolation spaces, by definition of the Bessel potential spaces and Example 2.3).

By Theorem 3.9, the assumptions on (a^{ij}) guarantee that $L_t = \nabla \cdot (a_t \nabla)$ is the generator of a propagator $S_{t,s}$ on the full range $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$, in the sense of Definition 2.11.

We need to check that the required assumptions on the non-linearities hold. If k is such that $p(k - 4\gamma) > n$, then it is easily observed from the Sobolev Embedding Theorem that for every $\alpha \geq -2\gamma$, multiplication is a smooth operation from $H^{k+2\alpha,p}(\mathbf{T}^n) \times H^{k+2\alpha,p}(\mathbf{T}^n) \rightarrow H^{k+2\alpha,p}(\mathbf{T}^n)$. Therefore, the operator induced by f is smooth from $\mathcal{B}_\alpha = H^{k+2\alpha,p}(\mathbf{T}^n)$ to itself for every $\alpha \geq 0$. Moreover, since f is a polynomial it is easy to see that it sends bounded sets of $H^{k+2\alpha,p}(\mathbf{T}^n)$ to bounded sets for all $\alpha \geq 0$ therefore showing that $f \in \mathcal{C}_{0,0}^\infty(\mathcal{B})$. Similarly, since $H^{k,p}(\mathbf{T}^n) \subset H^{k-4\gamma,p}(\mathbf{T}^n)$ continuously, we see that p_i induce smooth operators from $\mathcal{B}_\alpha = H^{k+2\alpha,p}(\mathbf{T}^n)$ to itself for every $\alpha \geq -2\gamma$ and that $p_i \in \mathcal{C}_{-2\gamma,0}^\infty(\mathcal{B})$ for each $i = 1, \dots, d$.

We now want to specialize further our results, by introducing a stochastic context for (1.1) (which constitutes an important motivation for introducing a rough paths formulation, see the discussion in the introduction). We have the following.

Theorem 2.19. *Let $(\Omega, \mathcal{F}, \mathbb{P}, (\mathcal{F}_t)_{t \in [0,T]})$ be a filtered probability space satisfying the usual assumptions, let $(B^i)_{1 \leq i \leq d}$ be a multidimensional Brownian motion on Ω , for some $d \geq 1$. Let $(k, p) \in \mathbf{N} \times (1, \infty)$ be such that $k > n/p + 4/3$. Let $f, p_i : \mathbf{R} \rightarrow \mathbf{R}$ be polynomials with coefficients in $H^{k,p}(\mathbf{T}^n)$ for every $i \in \{1, \dots, d\}$. Consider an adapted process $a : \Omega \times [0, T] \rightarrow \mathcal{C}^\infty(\mathbf{T}^n; \mathcal{L}(\mathbf{R}^n))$ such that for \mathbb{P} -a.e. ω , $a(\omega) \in \mathcal{C}^{\varrho,\infty}([0, T] \times \mathbf{T}^n; \mathcal{L}(\mathbf{R}^n))$ satisfies the assumptions of Example 2.12 with the coercivity constant $\varkappa(\omega) > 0$ in (2.19).*

Then there exists a stopping time τ such that the equation

$$\begin{aligned} du_t(x) &= \nabla \cdot (a_t(\omega, x) \nabla u_t(x)) dt + f(u_t(x)) dt + \sum_{i=1}^d p_i(u_t(x)) dB_t^i, \\ u_0 &\in H_0^{k,p}(\mathbf{T}^n), \end{aligned} \quad (2.25)$$

has a weak Itô solution in the following sense: a stochastic process $u : \Omega \times [0, T] \rightarrow H^{k,p}(\mathbf{T}^n)$ is adapted and satisfies:

- \mathbb{P} -almost surely either $\tau = T$, or $\limsup_{t \nearrow \tau} |u_t|_{H^{k,p}} = \infty$;
- for every $t \in [0, T]$ and any $\varphi \in \mathcal{C}^\infty(\mathbf{T}^n)$, we have on $\{t < \tau\}$:

$$\begin{aligned} &\int_{\mathbf{T}^n} (u_t(x) - u_0(x)) \varphi(x) dx + \int_0^t \int_{\mathbf{T}^n} a_s(x) \nabla u_s(x) \cdot \nabla \varphi(x) dx ds \\ &= \int_0^t \int_{\mathbf{T}^n} f(u_s(x)) \varphi(x) dx ds + \sum_{i=1}^d \int_0^t \int_{\mathbf{T}^n} p_i(u_s(x)) \varphi(x) dx dB_s^i. \end{aligned} \quad (2.26)$$

If in addition, the following moment estimate holds true: for every $m \geq 1$ there exists $K_m > 0$

$$\mathbb{E} \left[\sup_{t \in [0, T]} |a_t(\cdot)|_{\mathcal{C}^{k-1}(\mathbf{T}^n)}^m \right] \leq K_m, \quad (2.27)$$

*then the solution $u \in \mathcal{C}([0, \tau], H^{k,p}(\mathbf{T}^n))$ is unique in the class previously defined.*³

³Note that this does not rule out the possibility of existence of discontinuous in time weak solutions.

The existence part is a simple consequence of Theorem 2.18 and the fact that in this context $\varphi \in \mathcal{B}_{-1}^*$ for $\varphi \in \mathcal{C}^\infty(\mathbf{T}^n)$. The uniqueness statement is a bit more involved and requires some probabilistic tools like Kolmogorov extension theorem. A full proof will be presented in Section 5.3. We want to point out once again that the importance of the above theorem comes from the fact that the equation (2.25) can not be solved as a mild Itô equation because in this case the propagator $S_{t,s}$ is itself random, thus making the stochastic integrand $S_{t,s} p_i(u_s)$ non adapted.

Note that the results of Example 2.12 and Section 2.4 can be generalized to general smooth bounded domains $\mathcal{O} \subset \mathbf{R}^n$ instead of the torus \mathbf{T}^n , considering for instance (homogeneous) Dirichlet boundary conditions. In order to replicate the above results one can construct a scale $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$ similarly as in Example 2.2 for the Dirichlet Laplacian $(L, D(L)) = (\Delta_D, H^{2,p}(\mathcal{O}) \cap H_0^{1,p}(\mathcal{O}))$. In order to simplify the presentation, we have decided to postpone talking about general domains until Section 6.2, where we look at the Cahn-Hilliard equation with Dirichlet boundary conditions. The case of homogeneous Neumann boundary conditions could be handled in a similar fashion.

3 A multiplicative sewing Lemma and some by-products

Needless to say, the Sewing Lemma is a central result in rough paths theory. It is the core argument that permits to deal with the integration of controlled paths [29] (not even mentioning its implicit use in the original ‘multiplicative functional’-formalism by Lyons [50]). There is a vast literature on extensions of this result, e.g. in the context of reduced increments associated to a semigroup [31, 19], or flows [5], see also the recent works [43, 6, 7, 8]. We remark that the original statement in [22] is fairly general and applies in particular to non-commutative settings. In this section, we introduce a new version of this result which fits well in multiplicative, infinite-dimensional contexts.

3.1 The main result

We are going to present a version of a multiplicative sewing lemma which can be considered as a generalization of the non-commutative sewing lemma by Feyel, De La Pradelle, and Mokobodzki [22]. In the sequel, we denote by (\mathcal{M}, \circ) a monoid (for convenience we will mostly omit to write \circ). For a function $\mu: \Delta_2 \rightarrow \mathcal{M}$ and an arbitrary partition $\pi = \{s = t_0 < t_1 < \dots < t_k = t\}$ of $[s, t]$ set:

$$\mu_{t,s}^\pi = \prod_{i=0}^{k-1} \mu_{t_{i+1}, t_i}.$$

Definition 3.1. Let (\mathcal{M}, \circ) be a monoid. We call a triple $(\mathcal{M}, |\cdot|, d)$ a submultiplicative monoid if d is a metric on \mathcal{M} and a function $|\cdot|: \mathcal{M} \rightarrow \mathbf{R}_+$ is such that for all $a, b, c \in \mathcal{M}$:

$$d(ac, bc) \leq d(a, b)|c|, \quad \text{and} \quad d(ca, cb) \leq |c|d(a, b). \quad (3.1)$$

Moreover, we assume that for all $C \in \mathbf{R}_+$ the sets $B_C = \{a \in \mathcal{M} : |a| \leq C\}$ are complete with respect to the metric d .

Remark 3.2. Let \mathcal{M} be a submultiplicative monoid. As a consequence of (3.1), the multiplication is continuous with respect to d on the sets B_C . Indeed if $|b_n| \leq C$ and

$d(a_n, a) \rightarrow 0$ and $d(b_n, b) \rightarrow 0$ as $n \rightarrow \infty$ then:

$$d(a_n b_n, ab) \leq d(a_n, a)|b_n| + |a|d(b_n, b) \leq Cd(a_n, a) + |a|d(b_n, b) \rightarrow 0,$$

as $n \rightarrow \infty$. The same is true if instead of $|b_n| \leq C$ we have $|a_n| \leq C$.

Definition 3.3. Let \mathcal{M} be a submultiplicative monoid. We say that a function $\mu : \Delta_2 \rightarrow \mathcal{M}$ is

(i) *multiplicative* if for every $(t, u, s) \in \Delta_3$:

$$\mu_{t,s} = \mu_{t,u}\mu_{u,s}.$$

(ii) *almost-multiplicative* if there exists a control $\omega : \Delta_2 \rightarrow \mathbf{R}_+$ and $z > 1$ such that for each $(t, u, s) \in \Delta_3$:

$$d(\mu_{t,s}, \mu_{t,u}\mu_{u,s}) \leq \omega^z(t, s). \quad (3.2)$$

Next, let $\epsilon : \Delta_2 \rightarrow \mathbf{R}_+$ be such that $\epsilon(t, s)$ is continuous, increasing in t and decreasing in s . We say that

(iii) μ has *moderate growth* with growth rate ϵ if

$$|\mu_{t,s}^\pi| \leq \epsilon(t, s), \quad \text{for each } (t, s) \in \Delta_2, \quad (3.3)$$

for every $(t, s) \in \Delta_2$, independently of the choice of partition π of $[s, t]$.

We denote by $\text{BG}_2(0, T; \mathcal{M})$ the set of all functions $\mu : \Delta_2 \rightarrow \mathcal{M}$ with moderate growth for some ϵ as above.

Theorem 3.4 (Multiplicative sewing Lemma). *Let $(\mathcal{M}, |\cdot|, d)$ be a submultiplicative monoid with unit $\mathbf{1}$. Let $\epsilon : \Delta_2 \rightarrow \mathbf{R}_+$ be increasing in the first argument and decreasing in the second, and let $\mu \in \text{BG}_2(0, T; \mathcal{M})$ with a growth rate ϵ . Assume that there exists a control ω and a constant $z > 1$ so that (3.2) holds.*

Then, there exists a unique multiplicative $\varphi \in \text{BG}_2(0, T; \mathcal{M})$ such that for every $(t, s) \in \Delta_2$:

$$d(\varphi_{t,s}, \mu_{t,s}) \lesssim_{z,T} \omega^z(t, s). \quad (3.4)$$

The function φ has the same growth rate ϵ and for all $(t, s) \in \Delta_2$, for every sequence of partitions $\pi_n = \{s = t_0 < t_1 < \dots < t_n = t\}$ with mesh-size $|\pi_n| = \max_i |t_{i+1}^n - t_i^n| \rightarrow 0$ as $n \rightarrow \infty$ we have:

$$\varphi_{t,s} = d\text{-}\lim_{n \rightarrow \infty} \prod_{i=0}^{k_n-1} \mu_{t_{i+1}^n, t_i^n}. \quad (3.5)$$

Moreover, if $\mu_{t,t} = \mathbf{1}$ for all $t \in [0, T]$ then $\varphi_{t,t} = \mathbf{1}$ and if in addition $\mu : \Delta_2 \rightarrow (\mathcal{M}, d)$ is continuous then so is $\varphi : \Delta_2 \rightarrow (\mathcal{M}, d)$.

Proof. Our proof is reminiscent to that of the additive Sewing Lemma in [25].

Existence: Let $(t, s) \in \Delta_2$ and $\pi = \{s = t_0 < t_1 < \dots < t_k = t\}$ be a partition of $[s, t]$. Since ω is a control there exists $0 < l < k$ such that:

$$\omega(t_{l+1}, t_{l-1}) \leq \frac{2\omega(t, s)}{k}.$$

This is true since assuming the opposite would contradict the superadditivity of ω . Denote by $\hat{\pi}$ the partition of $[s, t]$ obtained by removing the point t_l from π . Then using (3.2) and submultiplicativity:

$$\begin{aligned} d(\mu_{t,s}^{\hat{\pi}}, \mu_{t,s}^{\pi}) &\leq \left| \prod_{i=l+1}^k \mu_{t_{i+1}, t_i} \right| d(\mu_{t_{l+1}, t_{l-1}}, \mu_{t_{l+1}, t_l} \circ \mu_{t_l, t_{l-1}}) \left| \prod_{i=0}^{l-2} \mu_{t_{i+1}, t_i} \right| \\ &\leq \epsilon(t, t_{l+1}) \omega^z(t_{l+1}, t_{l-1}) \epsilon(t_{l-1}, s) \leq \epsilon^2(t, s) 2^z \omega^z(t, s) k^{-z}. \end{aligned}$$

Repeating this procedure recursively until we arrive at trivial partition $\pi_0 = \{s, t\}$ we obtain the so called maximal inequality

$$\sup_{\pi} d(\mu_{t,s}, \mu_{t,s}^{\pi}) \leq 2^z \zeta(z) \epsilon^2(t, s) \omega^z(t, s), \quad (3.6)$$

where the supremum is taken over all partitions of $[s, t]$ and $\zeta(z) = \sum_{k=1}^{\infty} k^{-z}$ is the Riemann zeta function. We claim that to show the existence of the limit (3.5) and its independence of the sequence of the partitions it suffices to show that

$$\sup_{|\pi| \vee |\pi'| < \varepsilon} d(\mu_{t,s}^{\pi}, \mu_{t,s}^{\pi'}) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \quad (3.7)$$

Indeed, this would imply that $\mu_{t,s}^{\pi_n}$ is Cauchy for every sequence of partitions π_n with $|\pi_n| \rightarrow 0$. Then since $|\mu_{t,s}^{\pi_n}| \leq \epsilon(t, s)$ the completeness of the sets $\{a \in \mathcal{M} : |a| \leq C\}$ with respect to the metric d would imply that μ^{π_n} converges to some element $\varphi_{t,s}$ such that $|\varphi_{t,s}| \leq \epsilon(t, s)$. The independence of the limit from the sequence of partitions is obvious once (3.7) holds.

To show (3.7) we assume without loss of generality that π' is a refinement of π (since otherwise we can simply use the triangle inequality with the term $\mu_{t,s}^{\pi \cup \pi'}$). If $\pi = \{s = t_0 < t_1 < \dots < t_k = t\}$ we define

$$M_{t,s}^l = \prod_{i=l}^{k-1} \mu_{t_{i+1}, t_i}^{\pi' \cap [t_i, t_{i+1}]} \prod_{i=0}^{l-1} \mu_{t_{i+1}, t_i}.$$

Note that $M_{t,s}^k = \mu_{t,s}^{\pi}$ and $M_{t,s}^0 = \mu_{t,s}^{\pi'}$. Then, by the triangle inequality, submultiplicativity and the maximal inequality (3.6):

$$\begin{aligned} d(\mu_{t,s}^{\pi}, \mu_{t,s}^{\pi'}) &\leq \sum_{i=0}^{k-1} d(M_{t,s}^i, M_{t,s}^{i+1}) \leq \epsilon^2(t, s) \sum_{i=0}^{k-1} d(\mu_{t_{i+1}, t_i}, \mu_{t_{i+1}, t_i}^{\pi' \cap [t_i, t_{i+1}]}) \\ &\leq 2^z \epsilon^4(t, s) \zeta(z) \sum_{i=0}^{k-1} \omega^z(t_{i+1}, t_i) \\ &\lesssim_{z,T} \omega(t, s) \max_i \{\omega^{z-1}(t_{i+1}, t_i)\} \rightarrow 0 \quad \text{as } |\pi| \rightarrow 0, \end{aligned}$$

thus showing (3.7).

To show (3.4) it is enough to take the limit in (3.6) as $|\pi| \rightarrow 0$. Note that by Remark 3.2, the multiplication of $\mu_{t,r}^{\pi_1}$ with $\mu_{r,s}^{\pi_2}$ is continuous with respect to d . This, together with the independence of the limit in 3.5 with respect to the sequence of partitions, immediately implies the multiplicativity of φ and thus $\varphi \in \text{BG}_2(0, T; \mathcal{M})$.

If $\mu_{t,t} = \mathbb{1}$ then it is clear from (3.4) that $\varphi_{t,t} = \mathbb{1}$. One can also use (3.4) to show that this together with continuity of $\mu : \Delta_2 \rightarrow (\mathcal{M}, d)$ will imply continuity of $\varphi : \Delta_2 \rightarrow (\mathcal{M}, d)$. Details are left to the reader.

Uniqueness: Let $\psi \in \text{BG}_2(0, T; \mathcal{M})$ be another multiplicative map satisfying

$$d(\psi_{t,s}, \mu_{t,s}) \lesssim_{z,T} \omega^z(t, s) ,$$

then by the triangle inequality:

$$d(\varphi_{t,s}, \psi_{t,s}) \lesssim_{z,T} \omega^z(t, s) .$$

Let $\tilde{\epsilon}$ be a growth rate of ψ and denote $\epsilon_0 = 1 + \tilde{\epsilon} + \epsilon$. Let $\pi = \{s = t_0 < t_1 < \dots < t_k = t\}$ be an arbitrary partition and define now for $1 \leq l \leq k-1$, $D_{t,s}^l = \phi_{t,t_l} \psi_{t_l s}$. We also denote $D_{t,s}^0 = \varphi_{t,s}$ and $D_{t,s}^k = \psi_{t,s}$. Then

$$\begin{aligned} d(\varphi_{t,s}, \psi_{t,s}) &\leq \sum_{l=0}^{k-1} d(D_{t,s}^l, D_{t,s}^{l+1}) \leq \sum_{l=0}^{k-1} |\varphi_{t,t_{l+1}}| d(\varphi_{t_{l+1},t_l}, \psi_{t_{l+1},t_l}) |\psi_{t_l,s}| \\ &\lesssim \epsilon_0^2(t, s) \sum_{l=0}^{k-1} \omega^z(t_{l+1}, t_l) , \end{aligned}$$

which converges to zero as $|\pi| \rightarrow 0$ similarly as before. Thus, we must have $\varphi = \psi$ which concludes the proof. \blacksquare

Remark 3.5. Though our result is new, it is similar in spirit to [22, Theorem 10], which itself is a generalization of the construction of the whole signature from the lower order ‘iterated integrals’ by Lyons in [50, Thm 2.2.1]⁴.

In [22], the authors make the assumption that the function $|\cdot|$ is Lipschitz continuous with respect to the distance d , while in our setting this assumption is replaced by the growth condition (3.3) and the assumption that the closed balls of $|\cdot|$ are complete with respect to d . This becomes a necessary modification if one wants to apply this sewing lemma on infinite dimensional spaces since as we will see later, in most examples $|\cdot|$ will induce a stronger topology than the one generated by d . It implies that the assumption of Lipschitz continuity as in [22] can no longer be satisfied in general.

An important example of submultiplicative monoids is provided by some class of (unital) *Banach algebras* $(\mathcal{A}, |\cdot|)$ (recall that by definition $|ab| \leq |a||b|$ for any $a, b \in \mathcal{A}$). In this case, the property (3.3) can be replaced by a suitable exponential growth assumption, which is easier to verify in practice.

Corollary 3.6 (Multiplicative Sewing Lemma in a Banach Algebra). *Let $(\mathcal{A}, |\cdot|)$ be a Banach algebra with a unit $\mathbb{1}$ and let p be a norm on \mathcal{A} (possibly different from $|\cdot|$). Let d_p be the metric induced by p i.e. $d_p(a, b) = p(a - b)$, and assume that the triple $(\mathcal{A}, |\cdot|, d_p)$ is a submultiplicative monoid.*

Let $\mu : \Delta_2 \rightarrow (\mathcal{A}, p)$ be continuous such that $\mu_{t,t} = \mathbb{1}$ for all $0 \leq t \leq T$. Assume that there exists a control $\bar{\omega}$ so that

$$|\mu_{t,s}| \leq \exp \bar{\omega}(t, s), \quad \text{for every } (t, s) \in \Delta_2. \quad (3.8)$$

⁴One could also show that [50, Thm 2.2.1] follows from our version of the multiplicative sewing lemma, Theorem 3.4, by using the so-called ‘neo-classical inequality’ to prove the growth condition (3.3).

Assume also that there exists another control ω and $z > 1$ such that

$$p(\mu_{t,s} - \mu_{t,u}\mu_{u,s}) \lesssim_{z,T} \omega^z(t, s), \quad (3.9)$$

for every $(t, u, s) \in \Delta_2$. Then, there exists a unique multiplicative and continuous $\varphi: \Delta_2 \rightarrow (\mathcal{A}, p)$ such that for every $(t, s) \in \Delta_2$:

$$|\varphi_{t,s}| \leq \exp \bar{\omega}(t, s), \quad (3.10)$$

$$p(\varphi_{t,s} - \mu_{t,s}) \lesssim_{z,T} \omega^z(t, s). \quad (3.11)$$

Proof. We can conclude from Theorem 3.4 once we show $\mu \in \text{BG}_2(0, T; \mathcal{A})$ and that one can take a growth rate for μ to be $\epsilon(t, s) := \exp \bar{\omega}(t, s)$. For this we use submultiplicativity of the norm $|\cdot|$ to deduce for every partition $\pi = \{s = t_0 < t_1 < \dots < t_k = t\}$

$$|\mu_{t,s}^\pi| \leq \prod_{i=0}^{k-1} |\mu_{t_{i+1}, t_i}| \leq \prod_{i=0}^{k-1} \exp \bar{\omega}(t_{i+1}, t_i) = \exp \left\{ \sum_{i=0}^{k-1} \bar{\omega}(t_{i+1}, t_i) \right\} \leq \exp \{ \bar{\omega}(t, s) \},$$

which concludes the proof. \blacksquare

3.2 A new proof of Tanabe/Sobolevskii's Theorem (proof of Theorem 2.10)

We start with a lemma.

Lemma 3.7. *Let $(X, |\cdot|_X)$, $(Y, |\cdot|_Y)$ be two reflexive Banach spaces such that $X \subseteq Y$ and $|x|_Y \leq |x|_X$ for all $x \in X$. Define $\mathcal{A} = \mathcal{L}(X) \cap \mathcal{L}(Y)$ with the norm*

$$|\cdot|_{\mathcal{A}} = \max\{|\cdot|_{\mathcal{L}(X)}, |\cdot|_{\mathcal{L}(Y)}\},$$

and seminorm $p(\cdot) = |\cdot|_{\mathcal{L}(X, Y)}$, then $(\mathcal{A}, |\cdot|_{\mathcal{A}}, p)$ is a submultiplicative monoid.

Proof. The fact that $(\mathcal{A}, |\cdot|_{\mathcal{A}})$ is an algebra is trivial. For submultiplicativity of p :

$$\begin{aligned} p(ab) &= |ab|_{\mathcal{L}(X, Y)} \leq |a|_{\mathcal{L}(Y, Y)} |b|_{\mathcal{L}(X, Y)} \leq |a|_{\mathcal{A}} p(b), \\ p(ab) &= |ab|_{\mathcal{L}(X, Y)} \leq |a|_{\mathcal{L}(X, Y)} |b|_{\mathcal{L}(X, X)} \leq p(a) |b|_{\mathcal{A}}. \end{aligned}$$

Now, without loss of generality let us show that the unit ball $B = \{a \in \mathcal{A} : |a|_{\mathcal{A}} \leq 1\}$ is complete with respect to p . Let (a_n) be a Cauchy sequence for p with $|a_n|_{\mathcal{A}} \leq 1$ for every $n \geq 0$. By completeness of $\mathcal{L}(X, Y)$ we infer the existence of a limit $a \in \mathcal{L}(X, Y)$ such that $p(a_n - a) \rightarrow 0$. It remains to show that a belongs to B . Since the operator-norm topology is stronger than the weak operator topology, we have that for all $x \in X, y^* \in Y^*$:

$$\langle y^*, a_n x \rangle \rightarrow \langle y^*, a x \rangle.$$

The fact that Y is reflexive implies that the unit ball in $\mathcal{L}(Y)$ is compact with respect to the weak operator topology (see [13, Thm 2.19]), therefore since $|a_n|_{\mathcal{L}(Y)} \leq 1$ there exists $b \in \mathcal{L}(Y)$ such that $|b|_{\mathcal{L}(Y)} \leq 1$ and for each $y \in Y, y^* \in Y^*$, it holds

$$\langle y^*, a_n y \rangle \rightarrow \langle y^*, b y \rangle.$$

Now, $X \subset Y$, so that convergence in the weak operator topology in $\mathcal{L}(Y)$ implies convergence in the weak operator topology in $\mathcal{L}(X, Y)$ and hence $a = b$, thus $|a|_{\mathcal{L}(Y)} \leq 1$. Similarly, using the fact that $Y^* \subset X^*$ we have that the weak operator topology in $\mathcal{L}(X, Y)$ is stronger than the weak operator topology in $\mathcal{L}(X)$, thus $|a|_{\mathcal{L}(X)} \leq 1$ and therefore $a \in B$. \blacksquare

With this lemma, we can now proceed to the proof of one of our main results.

Proof of Theorem 2.10. We introduce the Banach algebra $\mathcal{A} = \mathcal{L}(\mathcal{X}_0) \cap \mathcal{L}(\mathcal{X}_1)$ whose norm is defined as $|\cdot|_{\mathcal{A}} = \max\{|\cdot|_{\mathcal{L}(\mathcal{X}_0)}, |\cdot|_{\mathcal{L}(\mathcal{X}_1)}\}$. By Lemma 3.7 if we further define $p = |\cdot|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)}$ then the triple $(\mathcal{A}, |\cdot|_{\mathcal{A}}, p)$ is a submultiplicative monoid.

Next, from Assumption (L1), it is classical (see [52, 28]) that for each $t \in [0, T]$, one can define an analytic semigroup $e^{sL_t} \in \mathcal{A}$. It is given by the so-called Dunford-Taylor integral formula (see [52, 28]):

$$e^{tL_s} = \frac{1}{2\pi i} \int_{\mathcal{C}} e^{\zeta t} (\zeta - L_s)^{-1} d\zeta, \quad (3.12)$$

where \mathcal{C} is any contour running from $\infty e^{-i\theta}$ to $\infty e^{i\theta}$ in the sector $\Sigma_{\vartheta, \lambda}$ for $\theta \in (0, \pi/2 + \vartheta)$. Moreover, we have

$$|e^{tL_s}|_{\mathcal{L}(\mathcal{X}_0)}, |e^{tL_s}|_{\mathcal{L}(\mathcal{X}_1)} \leq e^{\lambda t}, \quad \forall t \geq 0, \quad (3.13)$$

for some constant λ being the same as in $\Sigma_{\vartheta, \lambda}$. In addition, the following estimates hold

$$|e^{(t-s)L_s} - \text{id}|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} \lesssim_T |t-s|, \quad \text{and} \quad |e^{(t-s)L_s}|_{\mathcal{L}(\mathcal{X}_0, \mathcal{X}_1)} \lesssim_T |t-s|^{-1}. \quad (3.14)$$

We now show an auxiliary estimate that is going to be useful: for every $u, s \in [0, T]$ and $\tau > 0$ the following estimate holds in $|\cdot|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)}$:

$$p(e^{\tau L_s} - e^{\tau L_u}) \lesssim \tau \omega^\varrho(u, s). \quad (3.15)$$

Indeed, using the formula (3.12), we have

$$e^{\tau L_s} - e^{\tau L_u} = \frac{1}{2\pi i} \int_{\mathcal{C}} e^{\zeta \tau} (\zeta - L_u)^{-1} (L_s - L_u) (\zeta - L_s)^{-1} d\zeta.$$

By (L1), $|(\zeta - L_u)^{-1}|_{\mathcal{A}} \lesssim 1$, and using Hölder continuity (L3) and submultiplicativity of p we indeed get

$$\begin{aligned} p(e^{\tau L_s} - e^{\tau L_u}) &\leq \frac{1}{2\pi} \int_{\mathcal{C}} |e^{\zeta \tau}| |(\zeta - L_u)^{-1}|_{\mathcal{A}} p(L_s - L_u) |(\zeta - L_s)^{-1}|_{\mathcal{A}} d|\zeta| \\ &\lesssim \omega^\varrho(u, s) \int_{\mathcal{C}} \frac{|e^{\zeta \tau}| d|\zeta|}{(1 + |\zeta|)^2} \lesssim \tau \omega^\varrho(u, s), \end{aligned}$$

which shows (3.15).

Next, in order to apply the multiplicative sewing lemma we define

$$\mu_{t,s} := e^{(t-s)L_s}, \quad (t, s) \in \Delta_2.$$

Clearly $\mu_{t,t} = \text{id}$ for every $t \in [0, T]$. By (3.13) we have $|\mu_{t,s}|_{\mathcal{A}} \leq e^{\lambda(t-s)}$ for some $\lambda \in \mathbf{R}$, and since $\bar{\omega}(t, s) = \lambda(t-s)$ is a control then μ satisfies (3.8). We now show that $\mu : \Delta_2 \rightarrow (\mathcal{A}, p)$ is continuous. For pairs $(t, s), (v, u) \in \Delta_2$, assuming without loss of generality that $t-s-v+u > 0$, we have

$$\begin{aligned} p(\mu_{t,s} - \mu_{v,u}) &\leq p(e^{(v-u)L_s} (e^{(t-s-v+u)L_s} - \text{id})) + p(e^{(v-u)L_s} - e^{(v-u)L_u}) \\ &\lesssim |t-s-v+u| + |v-u| \omega^\varrho(u, s), \end{aligned}$$

where for the first term we used (3.14) and for the second term we use (3.15). This clearly implies continuity after taking $|t-v| + |s-u| \rightarrow 0$. Now, note that

$$\mu_{t,s} - \mu_{t,u} \mu_{u,s} = (e^{(t-u)L_s} - e^{(t-u)L_u}) e^{(u-s)L_s}.$$

Thus,

$$\begin{aligned} p(\mu_{t,s} - \mu_{t,u}\mu_{u,s}) &\leq p\left(e^{(t-u)L_s} - e^{(t-u)L_u}\right) |e^{(u-s)L_s}|_{\mathcal{A}} \\ &\lesssim (t-u)\omega^\varrho(u,s) \leq (t-s)\omega^\varrho(t,s), \end{aligned}$$

where going from the second to the third line we used (3.15). Since both $|t-s|$ and $\omega(t,s)$ are controls then so is $|t-s|^a\omega^b(t,s)$ for all $a, b > 0$ such that $a+b \geq 1$ (see [26, p.22]). In particular this is true for $a = 1/(1+\varrho)$ and $b = \varrho/(1+\varrho)$. Therefore,

$$p(\mu_{t,u,s} - \mu_{t,u}\mu_{t,u}) \lesssim \left(|t-s|^{1/(1+\varrho)}\omega^{\varrho/(1+\varrho)}(t,s)\right)^{1+\varrho},$$

thus (3.9) is satisfied with $z = 1 + \varrho$. We can thus apply Corollary 3.6, and obtain the existence of the unique continuous multiplicative function $S: \Delta_2 \rightarrow (\mathcal{A}, p)$ such that $S_{t,t} = \text{id}$ for every $t \in [0, T]$ and there exists $C > 0$ such that for all $(t, s) \in \Delta_2$, $|S_{t,s}|_{\mathcal{A}} \leq e^{\lambda(t-s)}$ and

$$|S_{t,s} - \mu_{t,s}|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} \leq C|t-s|\omega^\varrho(t,s). \quad (3.16)$$

One can use density of \mathcal{X}_1 in \mathcal{X}_0 and continuity of S with respect to the topology induced by the $\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)$ -norm to prove that $S \in \mathcal{C}_2(0, T; \mathcal{L}_s(\mathcal{X}_0))$, proving that S satisfies (P1) and (P2). To show (P3) we simply use (3.16):

$$\begin{aligned} |S_{t,s} - \text{id}|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} &\leq |S_{t,s} - \mu_{t,s}|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_1)} + |\mu_{t,s} - \text{id}|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} \\ &\lesssim |t-s|\omega^\varrho(t,s) + |t-s| \\ &\lesssim_T |t-s|. \end{aligned}$$

We now show (P4). Let $x \in \mathcal{X}_1$ and take any $\epsilon > 0$, then by multiplicativity:

$$\begin{aligned} \epsilon^{-1}(S_{t+\epsilon,s} - S_{t,s})x &= \epsilon^{-1}(S_{t+\epsilon,t} - \text{id})S_{t,s}x \\ &= \epsilon^{-1}(\mu_{t+\epsilon,t} - \text{id})S_{t,s}x + \epsilon^{-1}(S_{t+\epsilon,t} - \mu_{t+\epsilon,t})S_{t,s}x. \end{aligned} \quad (3.17)$$

We have shown that $S_{t,s} \in \mathcal{L}(\mathcal{X}_0) \cap \mathcal{L}(\mathcal{X}_1)$ and therefore $S_{t,s}x \in \mathcal{X}_1$ for $x \in \mathcal{X}_1$. Using $\mu_{t+\epsilon,t} = e^{\epsilon L_t}$ we conclude that the first term in (3.17) converges to $L_t S_{t,s}x$ as $\epsilon \rightarrow 0$. For the second term, using (3.16):

$$\begin{aligned} \epsilon^{-1}|(S_{t+\epsilon,t} - \mu_{t+\epsilon,t})S_{t,s}x|_0 &\leq \epsilon^{-1}|S_{t+\epsilon,t} - \mu_{t+\epsilon,t}|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} |S_{t,s}x|_1 \\ &\lesssim \omega^\varrho(t+\epsilon, t)|S_{t,s}x|_1, \end{aligned}$$

which vanishes as ϵ goes to 0. Putting it all together we conclude that for every $x \in \mathcal{X}_1$ and $(t, s) \in \Delta_2$ with $s \neq t$:

$$\frac{d}{dt}S_{t,s}x = L_t S_{t,s}x.$$

The proof that $\frac{d}{ds}(S_{t,s}x) = -S_{t,s}L_s x$ is similar, hence (P4) follows.

The first inequality of (2.15) follows by Remark 2.8. Now, assuming in addition that (L4) holds, we will show that S satisfies (P5). Let $x \in \mathcal{X}_1$, since $\mu_{t,s} \in \mathcal{L}(\mathcal{X}_1)$ we can use (P4) to differentiate in \mathcal{X}_0 :

$$\frac{d}{dr}(S_{t,r}\mu_{r,s}x) = S_{t,r}(L_s - L_r)\mu_{r,s}x.$$

Thus, integrating with respect to r we obtain an equation for all $\tau < t$:

$$S_{t,s}x = S_{t,\tau}\mu_{\tau,s}x + \int_s^\tau S_{t,r}(L_r - L_s)\mu_{r,s}x dr. \quad (3.18)$$

By density of \mathcal{X}_1 in \mathcal{X}_0 we can extend this integral equation for all $x \in \mathcal{X}_0$. For simplicity denote $|\cdot| = |\cdot|_{\mathcal{L}(\mathcal{X}_0, \mathcal{X}_1)}$. Since 3.18 holds for all $x \in \mathcal{X}_0$ we can use this equation together with $|S_{t,s}|_{\mathcal{L}(\mathcal{X}_1)} \lesssim_T 1$ to obtain:

$$\begin{aligned} |S_{t,s}| &\lesssim |\mu_{\tau,s}| + \int_\tau^s |S_{t,r}| |L_r - L_s|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} |\mu_{r,s}| dr \\ &\lesssim (\tau - s)^{-1} + \int_\tau^s |S_{t,r}| \omega^\varrho(r, s) |r - s|^{-1} dr, \end{aligned} \quad (3.19)$$

where we swapped limits of integration, used Hölder regularity (L3) and a semigroup bound $|\mu_{t,s}| \lesssim |t - s|^{-1}$. Now multiply both sides of (3.19) by $|t - s|$, set $f(s) = |t - s| |S_{t,s}|$ and choose $\tau = \frac{t+s}{2}$ to obtain:

$$f(s) \lesssim 2 + \int_{\frac{t+s}{2}}^s f(r) |t - s| |t - r|^{-1} \omega^\varrho(r, s) |r - s|^{-1} dr.$$

We now use Gronwall's inequality to derive:

$$\begin{aligned} f(s) &\lesssim \exp \left(\int_{\frac{t+s}{2}}^s |t - s| |t - r|^{-1} \omega^\varrho(r, s) |r - s|^{-1} dr \right) \\ &= \exp \left(- \int_s^{\frac{t+s}{2}} \frac{\omega^\varrho(r, s) dr}{r - s} - \int_s^{\frac{t+s}{2}} \frac{\omega^\varrho(r, s) dr}{t - r} \right). \end{aligned}$$

The first integral in the exponential is uniformly bounded on $[0, T]$ by assumption (L4), and for the second integral we simply observe that there is no blow up of the integrand over the integrated area. Therefore, $f(s) \lesssim_T 1$ and recalling that $f(s) = |t - s| |S_{t,s}|$ we conclude $|S_{t,s}| \lesssim_T |t - s|^{-1}$, thus finishing the proof. \blacksquare

Remark 3.8. A similar construction of S satisfying (P1), (P2), (P3), (P4) through the limiting infinite product of $e^{(t-s)L_s}$ for a family of operators L_t with $\Lambda = 1$ is present in the paper of T.Kato [40], but only in the case where the family L_t is of bounded variation (as a function of t) in $\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)$. Tanabe himself proved the existence of a propagator S for all operators satisfying Assumption 2.6 but the more general $1/\varrho$ -variation assumption is replaced by ϱ -Hölder regularity. His approach is different and relies on the construction of the approximate solutions to equation (2.13). These approximations do not use a limiting infinite product which allows to relax the dissipativity assumption on L_t . The above proof using the multiplicative sewing lemma allows to unify constructions of S when restricted to operators with $\Lambda = 1$.

The following straightforward generalization of Theorem 2.10 will be extensively used in the sequel.

Theorem 3.9. *Let $(\mathcal{B}_\alpha, |\cdot|_\alpha)_{\alpha \in \mathbf{R}}$ be a monotone family of interpolation spaces such that \mathcal{B}_α is reflexive for each $\alpha \in \mathbf{R}$. Assume that there exists a set of indices $K := \{k_-, k_- + 1, \dots, k_+\} \subset \mathbf{Z}$ such that for every $k \in K$, $(L_t)_{t \in [0, T]}$ satisfies Assumption 2.6 where $(\mathcal{X}_0, \mathcal{X}_1)$ is replaced by $(\mathcal{B}_k, \mathcal{B}_{k+1})$ (with control ω possibly depending on $k \in \mathbf{Z}$).*

Then the family S , as constructed in Theorem 2.10, extends uniquely to a propagator on the full range $(\mathcal{B}_\alpha)_{\alpha \in [k_-, k_+]}$ (in the sense of Definition 2.11). More explicitly, we have

(P1*) $S \in \mathcal{C}(\Delta_2, \mathcal{L}_s(\mathcal{B}_\alpha))$ and there exists λ_α such that $\|S_{t,s}\|_{\mathcal{L}(\mathcal{B}_\alpha)} \leq e^{\lambda_\alpha(t-s)}$ for every $\alpha \in [k_-, k_+ + 1]$,

(P2*) $S_{t,t} = \text{id}$ and $S_{t,s} = S_{t,u}S_{u,s}$ for all $(s, u, t) \in \Delta_3$.

(P3*) For $(s, t) \in \Delta_2$, $\alpha \in [k_-, k_+]$, and $x \in \mathcal{B}_{\alpha+1}$ the following differential equations hold true in \mathcal{B}_α :

$$\frac{d}{dt}S_{t,s}x = L_t S_{t,s}x, \quad \frac{d}{ds}S_{t,s}x = -S_{t,s}L_sx.$$

(P4*) For all $(s, t) \in \Delta_2$, $\alpha, \beta \in [k_-, k_+ + 1]$ and $\beta \geq \alpha$ the propagator $S_{t,s}$ maps \mathcal{B}_α to \mathcal{B}_β , and in addition for $\sigma \in [0, 1]$ the following smoothing inequalities are true:

$$|S_{t,s}x|_\beta \lesssim |t-s|^{-(\beta-\alpha)}|x|_\alpha, \quad |(S_{t,s} - \text{id})x|_\alpha \lesssim |t-s|^\sigma|x|_{\alpha+\sigma}. \quad (3.20)$$

Proof. If α is an integer, the proof follows by Theorem 2.10 by taking $(\mathcal{X}_0, \mathcal{X}_1) = (\mathcal{B}_\alpha, \mathcal{B}_{\alpha+1})$. The general case follows by interpolation (2.1), since $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$ is a monotone family of interpolation spaces. \blacksquare

Example 3.10. Consider the family of operators given in Example 2.12 and for each $k \in \mathbf{Z}$, let $\mathcal{B}_k = H^{2k,p}(\mathbf{T}^n)$ where $p \in (1, \infty)$, and assume that for all $t \in [0, T]$, $a(t, \cdot) \in \mathcal{C}^\infty(\mathbf{T}^n; \mathbf{R}^{n \times n})$. Then, one can check similarly as in Example 2.12 that the assumptions of Theorem 3.9 are fulfilled for $K = \mathbf{Z}$, which means that the associated propagator satisfies the properties (P1*)–(P4*) for the full scale $(\mathcal{B}_\alpha)_{\alpha \in \mathbf{R}}$.

3.3 Lie-Trotter product formula

In this subsection we present another application of the multiplicative sewing lemma - the proof of a Lie-Trotter-type formula for families $(S_{t,s})$ such that (P1)–(P4) hold. We shall call such a family a quasipropagator. ⁵

Recall that a semigroup S_t is called contractive if $|S_t| \leq 1$ for all times t . The classical Lie-Trotter product formula for the exponential of two (not necessary commuting) operators states that, if $A, B : \mathcal{X}_0 \rightarrow \mathcal{X}_0$ are two closed (unbounded) operators with common domain \mathcal{X}_1 such that they form (respectively) contractive semigroups S_t^A and S_t^B , and such that their sum $A + B$ is a closable operator generating a contractive semigroup S_t^{A+B} , then for every t and every $x \in \mathcal{X}_0$:

$$\lim_{n \rightarrow \infty} \prod_{i=0}^{n-1} S_{\frac{t}{n}}^A S_{\frac{t}{n}}^B x = S_t^{A+B} x. \quad (3.21)$$

A proof of this result which is based on so called Chernoff \sqrt{n} -Lemma can be found in [28, p. 53]. Here we present an alternative proof based on the multiplicative sewing lemma under the form of Corollary 3.6 together with simple estimates on semigroups and commutators. In addition, our proof extends to the quasipropagators and does not assume contractivity of the underlying semigroups but only the exponential bounds like in Definition 3.1, which is weaker. Strictly speaking, every semigroup that satisfies such an exponential bound can be shifted to become contractive, but our proof does not require that and can be applied instantly.

⁵Though this result is a non-trivial consequence of Theorem 3.4, it will not be needed in the rest of the paper.

Assumption 3.11. We are given continuously embedded reflexive Banach spaces $\mathcal{X}_2 \subseteq \mathcal{X}_1 \subseteq \mathcal{X}_0$ and $(A_t), (B_t)$ for $t \in [0, T]$ are two families of unbounded closed operators satisfying the properties (L1) with $\Lambda = 1$ for the range of indices $K = \{0, 1, 2\}$ and (L3) with $K = \{0, 1\}$. Moreover, we assume that for every $t \in [0, T]$, the domains $D(A_t^i)$ and $D(B_t^i)$ both contain \mathcal{X}_i for $i = 1, 2$.

Prior to proving a generalized Lie-Trotter formula, we will need a commutator estimate of two semigroups.

Lemma 3.12 (Commutator estimate). *Let the families $(A_t)_{t \in [0, T]}, (B_t)_{t \in [0, T]}$ satisfy Assumption 3.11. For every $v, u \in [0, T]$ define a commutator*

$$C(t, s) := \exp\{tA_v\} \exp\{sB_u\} - \exp\{sB_u\} \exp\{tA_v\}.$$

Then the following estimate holds true uniformly in $v, u \in [0, T]$:

$$|C(t, s)|_{\mathcal{L}(\mathcal{X}_2, \mathcal{X}_0)} \lesssim (t \vee s)^2. \quad (3.22)$$

Proof. The fact that these estimates are uniform in $v, u \in [0, T]$ will follow from the uniform bounds in assumption 2.6, so without loss of generality we will show this for the constant in time families A and B . Define the second order Taylor remainder

$$R^A(t) := e^{tA} - \text{id} - tA,$$

and similarly define R^B . Then such a remainder satisfies

$$|R^A(t)|_{\mathcal{L}(\mathcal{X}_k, \mathcal{X}_j)} \lesssim t^{k-j}, \quad \text{for } (k, j) \in \{(2, 0), (2, 1), (1, 0)\}. \quad (3.23)$$

Indeed, for $(k, j) = (1, 0)$ we can see that by the triangle inequality and (3.14):

$$|R^A(t)|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} \leq |\exp\{tA\} - \text{id}|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} + t|A|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} \lesssim t + t|A|_{\mathcal{L}(\mathcal{X}_1, \mathcal{X}_0)} \lesssim t.$$

Likewise, Assumption (L1) and Theorem 3.9 with $k = 1$ implies that $|\exp\{tA\} - \text{id}|_{\mathcal{L}(\mathcal{X}_2, \mathcal{X}_1)} \lesssim |t - s|$ and we can use it to show (3.23) for $(k, j) = (2, 1)$. For $(k, j) = (2, 0)$ we simply use the formula $\frac{d}{dt}e^{tA} = e^{tA}A$ twice and then the uniform boundedness of $|e^{tA}|_{\mathcal{L}(\mathcal{X}_0)}$ in t . For $x \in \mathcal{X}_2$

$$|R^A(t)x|_0 = \left| \int_0^t \int_0^s e^{rA} A^2 x \, dr ds \right|_0 \lesssim \int_0^t \int_0^s dr ds |A^2 x|_0 \lesssim t^2 |x|_2. \quad (3.24)$$

We use this now to rewrite our commutator $C(t, s)$ as:

$$\begin{aligned} C(t, s) &= (\text{id} + tA + R^A(t)) (\text{id} + sB + R^B(s)) \\ &\quad - (\text{id} + sB + R^B(s)) (\text{id} + tA + R^A(t)) \\ &= ts(AB - BA) + R^A(t) + sR^A(t)B + R^A(t)R^B(s) \\ &\quad - R^B(s) - tR^B(s)A - R^B(s)R^A(t). \end{aligned}$$

Using $|AB - BA|_{\mathcal{L}(\mathcal{X}_2, \mathcal{X}_0)} < \infty$ and the bounds (3.23) the result follows. ■

Theorem 3.13 (Lie-Trotter product formula). *Let families $(A_t)_{t \in [0, T]}$, $(B_t)_{t \in [0, T]}$ satisfy Assumption 3.11. Assume that $(A_t + B_t)_{t \in [0, T]}$ is a family of closable operators and that this closure also satisfies Assumption 3.11. Then (A_t) , (B_t) and the closure of $(A_t + B_t)$ all generate quasipropagators which we respectively call $S_{t,s}^A$, $S_{t,s}^B$ and $S_{t,s}^{A+B}$ for $(t, s) \in \Delta_2$. Moreover, for any sequence of partitions π^n of $[s, t]$ with $|\pi^n| \rightarrow 0$ as $n \rightarrow \infty$ the following Lie-Trotter formula holds for every $(t, s) \in \Delta_2$ and every $x \in \mathcal{X}_0$:*

$$S_{t,s}^{A+B}x = \lim_{n \rightarrow \infty} \prod_{[u,v] \in \pi^n} S_{v,u}^A S_{v,u}^B x, \quad (3.25)$$

where the limit is taken in \mathcal{X}_0 and the product over $[u, v] \in \pi^n$ means that it runs over all two neighbouring points in the partition.

Proof. Take $\mathcal{A} = \mathcal{L}(\mathcal{X}_2) \cap \mathcal{L}(\mathcal{X}_0)$, $|\cdot|_{\mathcal{A}} = \max\{|\cdot|_{\mathcal{L}(\mathcal{X}_0)}, |\cdot|_{\mathcal{L}(\mathcal{X}_2)}\}$ and $p(\cdot) = |\cdot|_{\mathcal{L}(\mathcal{X}_2, \mathcal{X}_0)}$, then using again Lemma 3.7 we have that $(\mathcal{A}, |\cdot|_{\mathcal{A}}, p)$ is a submultiplicative monoid. Without loss of generality we assume that the resolvent sets of A_t and B_t both contain $\Sigma_{\vartheta, \lambda}$ for the same $\vartheta > 0$ and $\lambda \in \mathbf{R}$. Moreover, without loss of generality we also assume that the control ω and exponent ϱ in (L3) is the same for A and B . Define $\mu: \Delta_2 \rightarrow \mathcal{A}$ by

$$\mu_{t,s} := S_{t,s}^A S_{t,s}^B, \quad (t, s) \in \Delta_2.$$

It follows from Remark 3.2 and by Tanabe's Theorem that $|\mu_{t,s}|_{\mathcal{A}} \leq e^{2\lambda(t-s)}$. We now study the quantity $p(\mu_{t,u,s} - \mu_{t,u}\mu_{u,s})$. Denote $\mu_{t,s}^A = e^{(t-s)A_s}$ and $\mu_{t,s}^B = e^{(t-s)B_s}$. First, by the multiplicativity of S^A and S^B we have for any $(t, u, s) \in \Delta_3$,

$$\mu_{t,s} - \mu_{t,u}\mu_{u,s} = S_{t,u}^A (S_{u,s}^A S_{t,u}^B - S_{t,u}^B S_{u,s}^A) S_{u,s}^B.$$

Second, by the triangle inequality and submultiplicativity of p :

$$\begin{aligned} p(\mu_{t,s} - \mu_{t,u}\mu_{u,s}) &\leq |S_{t,u}^A|_{\mathcal{A}} p(S_{u,s}^A S_{t,u}^B - S_{t,u}^B S_{u,s}^A) |S_{u,s}^B|_{\mathcal{A}} \\ &\leq e^{\lambda(t-s)} p(S_{u,s}^A S_{t,u}^B - S_{t,u}^B S_{u,s}^A) \\ &\lesssim_T p(S_{u,s}^A - \mu_{u,s}^A) |S_{t,u}^B|_{\mathcal{A}} + |\mu_{u,s}^A|_{\mathcal{A}} p(S_{t,u}^B - \mu_{t,u}^B) \\ &\quad + p(\mu_{u,s}^A \mu_{t,u}^B - \mu_{t,u}^B \mu_{u,s}^A) \\ &\quad + p(\mu_{t,u}^B - S_{t,u}^B) |\mu_{u,s}^A|_{\mathcal{A}} + |S_{t,u}^B|_{\mathcal{A}} p(\mu_{u,s}^A - S_{u,s}^A) \\ &\lesssim_T p(S_{u,s}^A - \mu_{u,s}^A) + p(\mu_{t,u}^B - S_{t,u}^B) + p(C(u-s, t-u)). \end{aligned} \quad (3.26)$$

Here the commutator

$$C(u-s, t-u) = \exp\{(u-s)A_s\} \exp\{(t-u)B_u\} - \exp\{(t-u)B_u\} \exp\{(u-s)A_s\},$$

is exactly of the form considered in Lemma 3.12, so we have

$$p(C(u-s, t-u)) \lesssim |t-s|^2.$$

The construction of S^A and S^B using Theorem 2.10 guarantees that

$$p(S_{u,s}^A - \mu_{u,s}^A) \lesssim_T |u-s| \omega^\varrho(u, s), \quad p(S_{t,u}^B - \mu_{t,u}^B) \lesssim_T |t-u| \omega^\varrho(t, u),$$

for some $\varrho \in (0, 1)$. Applying these to (3.26) we get:

$$p(\mu_{t,s} - \mu_{t,u}\mu_{u,s}) \lesssim_{\lambda, T} |t-s| \omega^\varrho(t, s) + |t-s|^2 \lesssim |t-s| \bar{\omega}^\varrho(t, s),$$

where $\bar{\omega}(t, s) = \omega(t, s) + |t - s|^{\frac{1}{e}}$ is a control. We now have all the necessary ingredients to apply the multiplicative sewing lemma (Corollary 3.6) which implies that the limit $\lim_{|\pi| \rightarrow 0} \prod_{[u, v] \in \pi} S_{v, u}^A S_{v, u}^B$ exists with respect to the semi-norm p , is multiplicative and independent of the partitions. Call this limit φ . First, we show that $\varphi_{t, s} x = S_{t, s}^{A+B} x$ for $x \in \mathcal{X}_2$ and for that it is enough to show $\frac{d}{dt} \varphi_{t, s} x = \frac{d}{dt} S_{t, s}^{A+B} x$ since $\varphi_{t, t} = S_{t, t}^{A+B} = \text{id}$. Like in the proof of Tanabe's theorem note that there is $R(\epsilon)$ such that $|R(\epsilon)|_0 \lesssim 1$ uniformly in (t, s) and ϵ such that

$$\begin{aligned} \epsilon^{-1}(\varphi_{t+\epsilon, s} x - \varphi_{t, s} x) &= \epsilon^{-1}(\mu_{t+\epsilon, t} - \text{id})\varphi_{t, s} x + \omega^\epsilon(t + \epsilon, t)R(\epsilon) \\ &= \epsilon^{-1}((S_{t+\epsilon, t}^A - \text{id})S_{t+\epsilon, t}^B(S_{t+\epsilon, t}^B - \text{id}))\varphi_{t, s} x \\ &\quad + \bar{\omega}^\epsilon(t + \epsilon, t)R(\epsilon) \\ &\rightarrow (A_t + B_t)\varphi_{t, s} x \quad \text{as } \epsilon \rightarrow 0. \end{aligned}$$

By the assumption that S^{A+B} is a quasipropagator we have $\frac{d}{dt} S_{t, s}^{A+B} = (A_t + B_t)S_{t, s}^{A+B}$ on \mathcal{X}_1 . Using the inclusion $\mathcal{X}_2 \subset \mathcal{X}_1$ this implies that $\varphi_{t, s} x = S_{t, s}^{A+B} x$ for $x \in \mathcal{X}_2$, since φ satisfies the same equation on \mathcal{X}_2 . It remains to show that the limit in (3.25) can be taken for all $x \in \mathcal{X}_0$. Let $x \in \mathcal{X}_0$ and denote $S_{t, s}^{A+B, n} = \prod_{[u, v] \in \pi^n} S_{v, u}^A S_{v, u}^B$. Since \mathcal{X}_2 is dense in \mathcal{X}_0 , for every $\epsilon > 0$ we can choose $y \in \mathcal{X}_2$ such that $|x - y|_0 \leq \epsilon$. But then

$$\begin{aligned} |(S_{t, s}^{A+B} - S_{t, s}^{A+B, n})x| &\leq |S_{t, s}^{A+B} - S_{t, s}^{A+B, n}|_{\mathcal{L}(\mathcal{X}_2, \mathcal{X}_0)}|y|_2 + |S_{t, s}^{A+B, n}|_{\mathcal{L}(\mathcal{X}_0)}|x - y|_0 \\ &\lesssim p(S_{t, s}^{A+B} - S_{t, s}^{A+B, n})|y|_2 + e^{\lambda(t-s)}\epsilon. \end{aligned}$$

Letting first $n \rightarrow \infty$ and then $\epsilon \rightarrow 0$ gives the desired result. \blacksquare

Note that in case when the families (A_t) and (B_t) are just constant operators A and B , we simply recover the usual Lie-Trotter formula (3.21) for unbounded operators. Moreover, the estimates are a bit better in this case since we have $S_{t, s}^A = \mu_{t, s}^A$, and therefore we can obtain the bound

$$|e^{(t-s)(A+B)} - e^{(t-s)A}e^{(t-s)B}|_{\mathcal{L}(\mathcal{X}_2, \mathcal{X}_0)} \lesssim e^{2\lambda(t-s)}|t - s|^2.$$

Remark 3.14 (Strang Splitting). If A, B are two infinitesimal generators which are independent of the time-like variable, then one can find an even better approximation of $e^{t(A+B)}$. It was noticed by Strang in [56] that for $h \geq 0$ small enough, the operator $\exp\{\frac{h}{2}A\}\exp\{hB\}\exp\{\frac{h}{2}A\}$ yields an approximation of the semigroup $\exp h(A + B)$ which is of higher order than the 'naïve' choice $\exp hA \exp hB$. Towards using the Sewing Lemma, for $(t, s) \in \Delta_2$ we let

$$\mu_{t, s} := \exp\left\{\frac{t-s}{2}A\right\}\exp\{(t-s)B\}\exp\left\{\frac{t-s}{2}A\right\},$$

so that

$$\begin{aligned} \mu_{t, s} - \mu_{t, u}\mu_{u, s} &= e^{\frac{t-u}{2}A} \left[e^{\frac{u-s}{2}A} e^{(t-s)B} e^{\frac{t-u}{2}A} - e^{(t-u)B} e^{\frac{t-s}{2}A} e^{(u-s)B} \right] e^{\frac{u-s}{2}A}. \\ &=: e^{\frac{t-u}{2}A} C^\sharp(u-s, t-u) e^{\frac{u-s}{2}A}. \end{aligned}$$

Expanding the exponentials like in Lemma 3.12 one can deduce that formally: $C^\sharp(u-s, t-u) = O(|t-s|^3)$, for an appropriate norm (the choice of $|\cdot|_{\mathcal{L}(\mathcal{X}_3, \mathcal{X}_0)}$ would do if one generalizes assumption (3.11) to \mathcal{X}_3 , and generalizes (3.24) accordingly).

The Sewing Lemma implies in turn that the iterated products of μ converge faster than the former first order approximation.

4 Controlled Path according to a monotone family of interpolation spaces

In this section, we build a framework for studying rough evolution equations of the form (1.1). Towards this end, we will define the space of paths that locally ‘look like’ a rough path X , which will be the natural space where the solution of (1.1) lives. In order to set up a mild formulation of (1.1) we also need to show that there is a notion of integration with respect to \mathbf{X} on such spaces. This will be done using a version of the classical sewing lemma for semigroups and propagators [31]. The connection between this result (which we will refer to as ‘affine sewing lemma’ in the sequel) and Theorem 3.4 will be established, having observed the role played by the affine group as introduced in (1.10)-(1.11).

Starting from this section and till the end of the article, we fix a monotone family of interpolations spaces $(\mathcal{B}_\beta)_{\beta \in \mathbf{R}}$. From now on, the index β in \mathcal{B}_β will refer to the varying parameter of the scale and the index α in \mathcal{B}_α will denote some fixed level thereof. Moreover, we will fix a family of unbounded operators $(L_t)_{t \in [0, T]}$ that acts on the monotone family $(\mathcal{B}_\beta)_{\beta \in \mathbf{R}}$, such that they generate the propagator $(S_{t,s})_{(t,s) \in \Delta_2}$ on the full range $(\mathcal{B}_\beta)_{\beta \in \mathbf{R}}$ in the sense of Definition 2.11. We do not necessarily assume that $S_{t,s}$ was constructed using the Theorem 3.9. This potentially can allow us to apply the forthcoming results to linear operators L_t that do not necessarily have $\Lambda = 1$ in (L1) or to Banach spaces \mathcal{B}_β that are not necessarily reflexive.

4.1 Affine Sewing Lemma

The affine sewing lemma is going to allow us to define integrals of the form $z_t := \int_0^t S_{t,r} y_r \cdot d\mathbf{X}_r$. Before we state the affine sewing lemma let us describe the algebraic properties of such integrals. Note that the linearity of the integral does not patch together nicely with the usual increment operator δ from (2.6). This is due to the fact that in this case: $\delta z_{t,s} \equiv z_t - z_s = \int_s^t S_{t,r} y_r \cdot d\mathbf{X}_r + S_{t,s} z_s - z_s \neq \int_s^t S_{t,r} y_r \cdot d\mathbf{X}_r$. Instead, we have the relation

$$\delta^S z_{t,s} \equiv z_t - S_{t,s} z_s = \int_s^t S_{t,r} y_r \cdot d\mathbf{X}_r .$$

As a matter of fact, the integral $\int_s^t S_{t,r} y_r \cdot d\mathbf{X}_r$ has a multiplicative structure. Indeed, letting $\beta \in \mathbf{R}$ (to be chosen later) and defining

$$\mathcal{M}(\beta) := \mathcal{L}(\mathcal{B}_\beta) \times \mathcal{B}_\beta, \tag{4.1}$$

we see that the multiplication of two elements $\mu_j \equiv (S_j, x_j) \in \mathcal{M}(\beta)$, $j = 1, 2$, can be defined as $\mu_1 \circ \mu_2 := (S_1 S_2, x_1 + S_1 x_2)$. Defining addition componentwise, we note that $\mathcal{M}(\beta)$ is a near-ring, namely $(\mathcal{M}(\beta), \circ)$ forms a monoid and the multiplication is right-distributive: $(\mu_1 + \mu_2) \circ \nu = \mu_1 \circ \nu + \mu_2 \circ \nu$ (though it is not left distributive in general).

With this at hand, and assuming that the rough convolution $\int_s^t S_{t,r} y_r \cdot d\mathbf{X}_r$ is meaningful,

we should have

$$\begin{aligned}
& (S_{t,u}, \int_u^t S_{t,r} y_r \cdot d\mathbf{X}_r) \circ (S_{u,s}, \int_s^u S_{u,r} y_r \cdot d\mathbf{X}_r) \\
&= (S_{t,u} S_{u,s}, \int_u^t S_{t,r} y_r dX_r + S_{t,u} \int_s^u S_{u,r} y_r \cdot d\mathbf{X}_r) \\
&= (S_{t,s}, \int_s^t S_{t,r} y_r \cdot d\mathbf{X}_r),
\end{aligned}$$

meaning that $\varphi_{t,s} := (S_{t,s}, \int_s^t S_{t,r} y_r \cdot d\mathbf{X}_r)$ is multiplicative in $\mathcal{M}(\beta)$. This suggests that using an appropriate approximation of the second component, we might be able to use Theorem 3.4 in order to construct the rough convolution.

Now, prior to define the integration map we need to specify which type of integrand shall be considered in the sequel. Let $\gamma \in [0, 1]$ and $\alpha \in \mathbf{R}$, we introduce the space $\mathcal{Z}_\alpha^\gamma$ as follows: $\mathcal{Z}_\alpha^\gamma$ consists of each 2-index element $\xi = (\xi_{t,s}) \in \mathcal{C}_2^\gamma(\mathcal{B}_\alpha) + \mathcal{C}_2^{2\gamma}(\mathcal{B}_{\alpha-\gamma})$ with the property that $\delta\xi \in \mathcal{C}_3^{2\gamma,\gamma}(\mathcal{B}_{\alpha-2\gamma}) + \mathcal{C}_3^{\gamma,2\gamma}(\mathcal{B}_{\alpha-2\gamma})$. Namely, there exist ξ^1, ξ^2 and h^1, h^2 with

$$\begin{aligned}
\xi_{t,s} &= \xi_{t,s}^1 + \xi_{t,s}^2, & (t, s) \in \Delta_2, \\
\delta\xi_{t,u,s} &= h_{t,u,s}^1 + h_{t,u,s}^2, & (t, u, s) \in \Delta_3,
\end{aligned} \tag{4.2}$$

and such that $[\xi^1]_{\gamma,\alpha} + [\xi^2]_{2\gamma,\alpha-\gamma} + [h^1]_{2\gamma,\gamma,\alpha-2\gamma} + [h^2]_{\gamma,2\gamma,\alpha-2\gamma} < \infty$, see (2.10). The space $\mathcal{Z}_\alpha^\gamma$ is then equipped with the natural norm

$$\|\xi\|_{\mathcal{Z}_\alpha^\gamma} := \inf_{\xi^1, \xi^2, h^1, h^2} \left([\xi^1]_{\gamma,\alpha} + [\xi^2]_{2\gamma,\alpha-\gamma} + [h^1]_{2\gamma,\gamma,\alpha-2\gamma} + [h^2]_{\gamma,2\gamma,\alpha-2\gamma} \right),$$

where infimum is taken over every decomposition of the form (4.2). (Note that analogous spaces were introduced in [31].)

We also identify the space which is going to be the image of the integration map by $\mathcal{E}_\alpha^{0,\gamma} = \mathcal{C}(\mathcal{B}_\alpha) \cap \mathcal{C}^\gamma(\mathcal{B}_{\alpha-\gamma})$ equipped with the norm being the maximum of norms on $\mathcal{C}(\mathcal{B}_\alpha)$ and $\mathcal{C}^\gamma(\mathcal{B}_{\alpha-\gamma})$. With this at hand we get:

Theorem 4.1 (Affine Sewing Lemma). *Consider the propagator $(S_{t,s})_{(s,t) \in \Delta_2}$ as in Theorem 3.9, let $\alpha \in \mathbf{R}$ and $\gamma \in (1/3, 1/2]$.*

There exists a unique continuous linear map $\mathcal{I} : \mathcal{Z}_\alpha^\gamma \rightarrow \mathcal{E}_\alpha^{0,\gamma}$ such that $\mathcal{I}_0(\xi) = 0$ for any $\xi \in \mathcal{Z}_\alpha^\gamma$ and moreover, for every $0 \leq \beta < 3\gamma$, it holds

$$|\delta^S \mathcal{I}_{t,s}(\xi) - S_{t,s} \xi_{t,s}|_{\alpha-2\gamma+\beta} \lesssim \|\xi\|_{\mathcal{Z}_\alpha^\gamma} |t-s|^{3\gamma-\beta}. \tag{4.3}$$

Finally, one has

$$\mathcal{I}_t(\xi) = \lim_{|\pi| \rightarrow 0} \sum_{[u,v] \in \pi} S_{t,u} \xi_{v,u}, \tag{4.4}$$

where the limit is taken in the sense of topology of $\mathcal{B}_{\alpha-2\gamma}$, over arbitrary partitions π of $[0, t]$ whose mesh-size $|\pi| \equiv \max\{v-u, : [u, v] \in \pi\}$ goes to 0.

The proof of Theorem 4.1 in the context of semigroups can be found in [27] or [31]. The proof in the case of propagators is carried out *mutatis mutandis*, using for instance the smoothing property (2.15). For the sake of completeness, we provide an alternative proof based on the multiplicative sewing lemma.

Proof. Define the monoid $\mathcal{M} = \mathcal{M}(\alpha - 2\gamma)$. We first show that \mathcal{M} can be endowed with a submultiplicative monoid structure, in the sense of Definition 3.1. Given $\mu_j \equiv (S_j, x_j) \in \mathcal{M}$, $j = 1, 2$, one defines a distance (which turns out to be also a norm):

$$d(\mu_1, \mu_2) := |S_1 - S_2|_{\mathcal{L}(\mathcal{B}_{\alpha-2\gamma})} + |x_1 - x_2|_{\alpha-2\gamma},$$

while we let

$$|\mu| := \max(1, d(\mu, 0)).$$

Using the right-distributivity of \circ one can show $d(\mu_1 \circ \nu, \mu_2 \circ \nu) \leq d(\mu_1, \mu_2)|\nu|$ and the inequality $d(\nu \circ \mu_1, \nu \circ \mu_2) \leq |\nu|d(\mu_1, \mu_2)$ can also be shown easily. Then, since $|\cdot|$ is continuous with respect to d , the completeness assumption from Definition 3.1 is satisfied and one concludes that $(\mathcal{M}, |\cdot|, d)$ is a submultiplicative monoid.

Next, define $\mu : \Delta_2 \rightarrow \mathcal{M}$ as

$$\mu_{t,s} := (S_{t,s}, S_{t,s}\xi_{t,s}).$$

For every $(t, u, s) \in \Delta_3$, observe that

$$\mu_{t,s} - \mu_{t,u} \circ \mu_{u,s} \equiv (0, S_{t,s}\delta\xi_{t,u,s} + S_{t,u}(S_{u,s} - \text{id})\xi_{t,u}).$$

Hence, using (3.20):

$$\begin{aligned} d(\mu_{t,s}, \mu_{t,u} \circ \mu_{u,s}) &\lesssim_S |\delta\xi_{t,u,s}|_{\alpha-2\gamma} + |(S_{u,s} - \text{id})\xi_{t,u}|_{\alpha-2\gamma} \\ &\leq \|\xi\|_{\mathcal{Z}_\alpha^\gamma} [|t - u|^{2\gamma}|u - s|^\gamma + |t - u|^\gamma|u - s|^{2\gamma}], \end{aligned}$$

showing in particular that μ is almost-multiplicative.

We now note that for every partition π of $[s, t]$, $\mu_{t,s}^\pi = (S_{t,s}, \mathcal{S}_{t,s}^\pi(\xi))$, where $\mathcal{S}_{t,s}^\pi(\xi) := \sum_{(u,v) \in \pi} S_{t,u}\xi_{v,u}$ is the partial sum associated with π . Along the same lines as the proof of (3.6), one can show that $|\mu_{t,s}^\pi| \lesssim 1 + |t - s|^\gamma + |t - s|^{3\gamma}$ uniformly over every partition π of $[s, t]$, thus showing $\mu \in \text{BG}_2(0, T; \mathcal{M})$. We can then either use $\mu \in \text{BG}_2(0, T; \mathcal{M})$ and apply Theorem 3.4 or use the fact that $|\cdot|$ is Lipschitz with respect to d and apply [22, Theorem 10] to obtain existence of the unique multiplicative $\varphi_{t,s} = (S_{t,s}, I_{t,s})$ such that $|I_{t,s} - S_{t,s}\xi_{t,s}|_{\alpha-2\gamma} = d(\varphi_{t,s}, \mu_{t,s}) \lesssim |t - s|^{3\gamma}$. Letting $\mathcal{S}_t(\xi) := I_{0,t}$, it is seen, thanks to multiplicativity of φ that $\delta^S \mathcal{S}_{t,s}(\xi) = I_{t,s}$, so that (4.3) holds with $\beta = 0$.

We now go over the proof of (4.3) for general $\beta \in (0, 3\gamma)$ which also implies the continuity of \mathcal{S} as a map $\mathcal{Z}_\alpha^\gamma \rightarrow \mathcal{E}_\alpha^{0,\gamma}$. To show this we take the dyadic partitions, namely $\pi_k := \{s = t_0 < t_1 < \dots < t_{2^k} = t\}$ where $t_i = s + 2^{-k}i(t - s)$. Denoting by $m = (u + v)/2$, and decomposing ξ as in (4.2), we have

$$\begin{aligned} \mathcal{S}_{t,s}^{\pi_k} - \mathcal{S}_{t,s}^{\pi_{k+1}} &= \sum_{[u,v] \in \pi_k} S_{t,u}\delta\xi_{v,m,u} + S_{t,m}(S_{m,u} - \text{id})\xi_{v,m} \\ &= \sum_{[u,v] \in \pi_k} S_{t,u}h_{v,m,u}^1 + \sum_{[u,v] \in \pi_k} S_{t,u}h_{v,m,u}^2 \\ &\quad + \sum_{[u,v] \in \pi_k} S_{t,m}(S_{m,u} - \text{id})\xi_{v,m}^1 + \sum_{[u,v] \in \pi_k} S_{t,m}(S_{m,u} - \text{id})\xi_{v,m}^2 \\ &=: \text{I} + \text{II} + \text{III} + \text{IV}. \end{aligned}$$

Denote $\beta' = \alpha - 2\gamma + \beta$ for shorthand. From the definition of the space $\mathcal{Z}_\alpha^\gamma$, and from (3.20) we can bound the first two terms as follows:

$$|\text{I} + \text{II}|_{\beta'} \leq \|\xi\|_{\mathcal{Z}_\alpha^\gamma} \sum_{[u,v] \in \pi_k} |t - m|^{-\beta} [|v - m|^{2\gamma} |m - u|^\gamma + |v - m|^\gamma |m - u|^{2\gamma}].$$

For the third term, we have

$$\begin{aligned} |\text{III}|_{\beta'} &\lesssim_S \sum_{[u,v] \in \pi_k} |t - m|^{-\beta} |(S_{m,u} - \text{id})\xi_{m,v}^1|_{\alpha-2\gamma} \\ &\leq [\xi^1]_{\gamma,\alpha} \sum_{[u,v] \in \pi_k} |t - m|^{-\beta} |v - m|^\gamma |m - u|^{2\gamma}, \end{aligned}$$

and similarly for the fourth term:

$$|\text{IV}|_{\beta'} \leq [\xi^2]_{2\gamma,\alpha-\gamma} \sum_{[u,v] \in \pi_k} |t - m|^{-\beta} |v - m|^{2\gamma} |m - u|^\gamma.$$

Now choose $\delta \geq 0$ such that $3\gamma - 1 > \delta > \beta - 1$. Summing all contributions, and observing that $v - m = m - u = \frac{|t-s|}{2^{k+1}} \leq t - m$ we have

$$\begin{aligned} |\mathcal{I}_{t,s}^{\pi_k} - \mathcal{I}_{t,s}^{\pi_{k+1}}|_{\beta'} &\lesssim \|\xi\|_{\mathcal{Z}_\alpha^\gamma} \sum_{[u,v] \in \pi_k} |t - m|^{-\beta} |v - m|^{3\gamma-1} |m - u| \\ &\lesssim \|\xi\|_{\mathcal{Z}_\alpha^\gamma} \sum_{[u,v] \in \pi_k} |t - m|^{\delta-\beta} |v - m|^{3\gamma-1-\delta} |m - u| \\ &\lesssim \|\xi\|_{\mathcal{Z}_\alpha^\gamma} 2^{-k(3\gamma-1-\delta)} |t - s|^{3\gamma-1-\delta} \sum_{[u,v] \in \pi_k} |t - m|^{\delta-\beta} |m - u| \\ &\lesssim \|\xi\|_{\mathcal{Z}_\alpha^\gamma} 2^{-k(3\gamma-1-\delta)} |t - s|^{3\gamma-1-\delta} \int_s^t |t - r|^{\delta-\beta} dr \\ &\lesssim \|\xi\|_{\mathcal{Z}_\alpha^\gamma} 2^{-k(3\gamma-1-\delta)} |t - s|^{3\gamma-\beta}, \end{aligned}$$

where we used the convexity of the integrand in the above Riemann sum since $\delta - \beta > -1$. Since δ above is chosen so that $3\gamma - 1 - \delta > 0$ we can sum over $k \in \mathbf{N}_0$ to finally obtain (4.3) and finish the proof. \blacksquare

Remark 4.2. The monoid defined above can be endowed with a more complex submultiplicative structure which involves two indices of spatial regularity $\beta' \leq \beta$. Let

$$\mathcal{M} := \mathcal{M}(\beta', \beta) := (\mathcal{L}(\mathcal{B}_{\beta'}) \cap \mathcal{L}(\mathcal{B}_\beta)) \times \mathcal{B}_{\beta'},$$

and given $\mu_j \equiv (S_j, x_j) \in \mathcal{M}, j = 1, 2$, introduce the distance $d(\mu_1, \mu_2) := |S_1 - S_2|_{\mathcal{L}(\mathcal{B}_\beta, \mathcal{B}_{\beta'})} + |x_1 - x_2|_{\beta'}$. Defining further $|\mu| := \max(1, |S|_{\mathcal{L}(\mathcal{B}_\beta) \cap \mathcal{L}(\mathcal{B}_{\beta'})} + |x|_\beta)$, it is easily checked, using the same compactness argument as that of the proof of Lemma 3.7, that $(\mathcal{M}, |\cdot|, d)$ is a submultiplicative monoid. Making the choice $(\beta', \beta) := (\alpha - 2\gamma, \alpha + \gamma - \kappa)$, $\kappa > 0$ being arbitrary, one can obtain an alternative proof of Theorem 4.1 (the main difference is that the condition (3.3) is shown to hold on dyadic partitions only, in which case a version of Theorem 3.4 still holds).

The advantage of introducing $\mathcal{M}(\beta', \beta)$ in comparison with the above (simpler) proof is that it allows obtaining approximation results for rough convolutions. More precisely, given

an almost multiplicative approximation $\tilde{S}_{t,s}$ of $S_{s,t}$ (think for instance of a semigroup $e^{(t-s)L_s}$ or of the resolvent approximation of such semi-group), the multiplicative sewing Lemma then tells us that $\sum_{[u,v] \in \pi} \tilde{S}_{t,u} \xi_{v,u}$ converges towards the same limit as (4.4) as $|\pi| \rightarrow 0$ by considering $\tilde{\mu}_{t,s} = (\tilde{S}_{t,s}, \tilde{S}_{t,s} \xi_{t,s})$. This fact could certainly be useful in the quasilinear case, or for numerical analysis purposes. On the other hand, potential applications of this observation go beyond the scope of this paper, and hence we chose to avoid such a level of generality.

4.2 Controlled rough paths

In this paragraph, we introduce a notion of controlled paths that live in some \mathcal{B}_α . Our definition differs from the ones given in [31, 27] since it is independent of the propagator S and its characterization does not involve the reduced increments δ^S . Prior to introduce the notion of controlled paths in addition to the $\mathcal{E}_\alpha^{0,\gamma}$ we define a space $\mathcal{E}_\alpha^{\gamma,2\gamma} = \mathcal{C}_2^\gamma(\mathcal{B}_{\alpha-\gamma}) \cap \mathcal{C}_2^{2\gamma}(\mathcal{B}_{\alpha-2\gamma})$. Both $\mathcal{E}_\alpha^{0,\gamma}$ and $\mathcal{E}_\alpha^{\gamma,2\gamma}$ reflect the parabolic nature of (1.1) and show an interplay between the time and spatial regularity.

Definition 4.3 (Controlled path according to a monotone family). Let $\mathcal{B} := (\mathcal{B}_\beta)_{\beta \in \mathbf{R}}$ be a monotone family of interpolation spaces. Assume that $\mathbf{X} \equiv (X, \mathbb{X}) \in \mathcal{C}^\gamma(0, T; \mathbf{R}^d)$ for some $\gamma > 1/3$, and let $\alpha \in \mathbf{R}$. We say that a pair (y, y') is *controlled by \mathbf{X}* according to \mathcal{B}_α if the following holds:

- (i) We have $(y, y') \in \mathcal{C}(\mathcal{B}_\alpha) \times (\mathcal{E}_{\alpha-\gamma}^{0,\gamma})^d$;
- (ii) The remainder R^y defined as:

$$R_{t,s}^y := \delta y_{s,t} - y'_s \cdot \delta X_{s,t} \equiv \delta y_{t,s} - \sum_{i=1}^d y'_s{}^i \delta X_{t,s}^i,$$

belongs to $\mathcal{E}_\alpha^{\gamma,2\gamma}$.

We will denote the space of all such controlled rough paths by $\mathcal{D}_X^{2\gamma}([0, T], \mathcal{B}_\alpha)$ or simply $\mathcal{D}_{X,\alpha}^{2\gamma}([0, T])$. When $T > 0$ is fixed we will simplify further and make an abuse of notation by writing simply $\mathcal{D}_{X,\alpha}^{2\gamma}$. Sometimes we will refer to y' as the Gubinelli derivative.

We endow the space $\mathcal{D}_{X,\alpha}^{2\gamma}$ with the norm:

$$\|y, y'\|_{\mathcal{D}_{X,\alpha}^{2\gamma}} = |y|_{0,\alpha} + |y'|_{\mathcal{E}_{\alpha-\gamma}^{0,\gamma}} + |R^y|_{\mathcal{E}_\alpha^{\gamma,2\gamma}}.$$

With this definition, it is easy to check that $\mathcal{D}_{X,\alpha}^{2\gamma}$ is a Banach space (we leave the details to the reader).

One can actually see from the above definition that $y \in \mathcal{E}_\alpha^{0,\gamma}$ and

$$[\delta y]_{\gamma,\alpha-\gamma} \leq |y'|_{0,\alpha-\gamma} [\delta X]_\gamma + [R^y]_{\gamma,\alpha-\gamma}. \quad (4.5)$$

For that reason, we do not make Hölder regularity of y as a part of the definition of a controlled rough path.

Note that one can recover the usual definition of the controlled rough path (see [29, 25]) from the above definition if one takes $\mathcal{B}_\beta = \mathcal{B}_0$ for all $\beta \in \mathbf{R}$. In the notation of [25, Definition 4.6] we then have $\mathcal{D}_X^{2\gamma}(0, T; \mathcal{B}_0) = \mathcal{D}_X^{2\gamma}([0, T], \mathcal{B}_0)$. Therefore, all our later analysis applies to the finite-dimensional case by choosing such constant family \mathcal{B}_0 and the propagator to be the identity: $S_{t,s} = \text{id}$.

Remark 4.4. The definition of controlled rough paths can be reformulated using the ‘reduced increment’ δ^S instead of $\delta = \delta^{\text{id}}$, which might look more natural when talking about the integrals $\int_0^t S_{t,s} y_s \cdot d\mathbf{X}_s$. Following [31, 19, 27], it is indeed possible to introduce the space $\mathcal{D}_{X,S,\alpha}^{2\gamma}$ consisting of pairs $(y, y') \in \mathcal{C}(0, T; \mathcal{B}_\alpha) \times \mathcal{C}(0, T; \mathcal{B}_{\alpha-\gamma})$, such that $\tilde{R}_{t,s}^y := \delta^S y_{t,s} - S_{t,s} y'_s \delta X_{t,s}$ belongs to $\mathcal{C}_2^{2\gamma}(0, T; \mathcal{B}_{\alpha-2\gamma}) \cap \mathcal{C}_2^\gamma(0, T; \mathcal{B}_{\alpha-\gamma})$ while $\delta^S y'$ belongs to $\mathcal{C}_2^\gamma(0, T; \mathcal{B}_{\alpha-2\gamma})$.

However, this has the inconvenience that the spaces considered depend on the propagator S while, as seen in the present paper, it is possible to get rid of this dependency (the spaces $\mathcal{D}_{X,\alpha}^{2\gamma}$ do however, depend on the scale $(\mathcal{B}_\beta)_{\beta \in \mathbf{R}}$). This would in addition make some proofs (like the proof of Lemma 4.7) more tedious. Finally, the space $\mathcal{D}_{X,\alpha}^{2\gamma}$ is much closer to the usual notion of controlled rough path; for instance its elements are controlled rough paths in the sense of [25, Definition 4.6] at the level of $\mathcal{B}_{\alpha-2\gamma}$.

In the sequel we will extensively use the following interpolation inequality, which is an immediate consequence of (2.1); for every $\beta \in [\gamma, 2\gamma]$, we have:

$$|R_{t,s}^y|_{\alpha-\beta} \leq |R^y|_{\mathcal{E}^{\gamma,2\gamma}} |t-s|^\beta. \quad (4.6)$$

We now state the fundamental result for the above controlled rough paths, namely that for such paths the ‘rough convolution’ with respect to \mathbf{X} is well-defined.

Theorem 4.5 (Integration). *Let $\mathbf{X} = (X, \mathbb{X}) \in \mathcal{C}^\gamma(0, T; \mathbf{R}^d)$ for some $\gamma > 1/3$ and let $\alpha \in \mathbf{R}$. Let $(y^i, y^{i'}) \in \mathcal{D}_{X,\alpha}^{2\gamma}$ for $i = 1, \dots, d$. Then the integral*

$$\int_0^t S_{t,s} y_s \cdot d\mathbf{X}_s := \lim_{|\pi| \rightarrow 0} \sum_{[u,v] \in \pi} S_{t,u} (y_u \cdot \delta X_{v,u} + y'_u : \mathbb{X}_{v,u}), \quad (4.7)$$

exists as an element of \mathcal{B}_α , where we denote $y'_u : \mathbb{X}_{v,u} := \sum_{1 \leq i,j \leq d} y^{i,j} \mathbb{X}_{v,u}^{ij}$.

Moreover, for every $0 \leq \beta < 3\gamma$ the above integral satisfies the estimate

$$\left| \int_s^t S_{t,u} y_u \cdot d\mathbf{X}_u - S_{t,s} (y_s \cdot \delta X_{t,s} + y'_s : \mathbb{X}_{t,s}) \right|_{\alpha-2\gamma+\beta} \lesssim \varrho_\gamma(\mathbf{X}) \|y, y'\|_{\mathcal{D}_{X,\alpha}^{2\gamma}} |t-s|^{3\gamma-\beta}, \quad (4.8)$$

for all $(s, t) \in \Delta_2$.

Proof. It suffices to apply the affine sewing lemma (Theorem 4.1) to

$$\xi_{t,s} = y_s \cdot \delta X_{t,s} + y'_s : \mathbb{X}_{t,s}, \quad (t, s) \in \Delta_2,$$

for which we need to show that $\xi \in \mathcal{Z}_\alpha^\gamma$. Indeed, the existence of the integral (4.7) will follow immediately by (4.4) while (4.8) is a consequence of (4.3).

First, note that ξ is indeed an element of $\mathcal{C}_2^{\gamma,\alpha} + \mathcal{C}_2^{2\gamma,\alpha-\gamma}$ and that moreover

$$\|\xi\|_{\mathcal{C}_2^{\gamma,\alpha} + \mathcal{C}_2^{2\gamma,\alpha-\gamma}} \leq |y \cdot \delta X|_{\gamma,\alpha} + |y' : \mathbb{X}|_{2\gamma,\alpha-\gamma} \leq \varrho_\gamma(\mathbf{X}) \|y, y'\|_{\mathcal{D}_{X,\alpha}^{2\gamma}},$$

by definition of $\mathcal{D}_{X,\alpha}^{2\gamma}$. Next, thanks to Chen’s relation we have the algebraic identity

$$\delta \xi_{t,u,s} = \delta X_{t,u} \cdot R_{u,s}^y + \mathbb{X}_{t,u} : \delta y'_{u,s},$$

from which we infer

$$|\delta\xi_{t,u,s}|_{\alpha-2\gamma} \leq [X]_\gamma |t-u|^\gamma |u-s|^{2\gamma} [R^y]_{2\gamma, \alpha-2\gamma} + [\mathbb{X}]_{2\gamma} |t-u|^{2\gamma} |u-s|^\gamma [\delta y']_{\gamma, \alpha-2\gamma}.$$

Hence, it follows that $\delta\xi \in \mathcal{C}_2^{2\gamma, \gamma}(\mathcal{B}_{\alpha-2\gamma}) + \mathcal{C}_2^{\gamma, 2\gamma}(\mathcal{B}_{\alpha-2\gamma})$ with

$$\|\delta\xi\|_{\mathcal{C}_2^{2\gamma, \gamma}(\mathcal{B}_{\alpha-2\gamma}) + \mathcal{C}_2^{\gamma, 2\gamma}(\mathcal{B}_{\alpha-2\gamma})} \leq \varrho_\gamma(\mathbf{X}) \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}}.$$

Summing the above contributions, we find that $\xi \in \mathcal{Z}_\alpha^\gamma$, and the conclusion of Theorem 4.1 yields the claimed estimate. \blacksquare

The following result not only describes the stability of integration but also tells us that the ‘rough convolution’ improves the spatial regularity of the controlled rough path.

Corollary 4.6. *The integration map defined in Theorem 4.5 is continuous from $\mathcal{D}_{X, \alpha}^{2\gamma}$ into itself. In addition, for $T \leq 1$ and for every σ, γ' such that $0 < \sigma < \gamma' \leq \gamma$, the linear map*

$$\mathcal{D}_{X, \alpha}^{2\gamma'}([0, T]) \rightarrow \mathcal{D}_{X, \alpha+\sigma}^{2\gamma'}([0, T]), \quad (y, y') \mapsto (z, z') := \left(\int_0^\cdot S_{\cdot, u} y_u \cdot d\mathbf{X}_u, y \right),$$

is well-defined, bounded, and it satisfies the following estimate:

$$\|z, z'\|_{\mathcal{D}_{X, \alpha+\sigma}^{2\gamma'}} \leq |y_0|_\alpha + C_{\gamma, \sigma} T^\varepsilon (1 + \varrho_\gamma(\mathbf{X})) \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma'}},$$

where $\varepsilon := \min\{\gamma - \gamma', \gamma' - \sigma\}$.

Proof. The first step is to show that z is indeed controlled by X if one lets $z' = y$. For this we need to evaluate the remainder $R_{t,s}^z := \delta z_{t,s} - y_s \cdot \delta X_{t,s}$ and show that it has the correct regularity. Denote by

$$\mathcal{R}_{t,s} = \int_s^t S_{t,u} y_u \cdot d\mathbf{X}_u - S_{t,s} (y_s \cdot \delta X_{t,s} + y'_s : \mathbb{X}_{t,s}), \quad (t, s) \in \Delta_2,$$

where the first integral is understood in the sense of Theorem 4.5. Using the fact that $\varrho_{\gamma'}(\mathbf{X}) \leq \varrho_\gamma(\mathbf{X})$ for $T \leq 1$ and using the estimate (4.8) with $\beta = \sigma + (2-i)\gamma'$ for $i \in \{1, 2\}$, we get

$$\begin{aligned} |\mathcal{R}_{t,s}|_{\alpha+\sigma-i\gamma'} &= |\mathcal{R}_{t,s}|_{\alpha-2\gamma'+\sigma+(2-i)\gamma'} \lesssim \varrho_{\gamma'}(\mathbf{X}) \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma'}} |t-s|^{\gamma'-\sigma+i\gamma'} \\ &\leq \varrho_\gamma(\mathbf{X}) \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma'}} |t-s|^{i\gamma'} T^{\gamma'-\sigma}, \end{aligned}$$

uniformly over $(t, s) \in \Delta_2$. Next, observe that

$$\begin{aligned} R_{t,s}^z &\equiv \int_s^t S_{t,u} y_u \cdot d\mathbf{X}_u - y_s \cdot \delta X_{s,t} + (S_{t,s} - \text{id}) \int_0^s S_{s,u} y_u \cdot d\mathbf{X}_u \\ &= \left(\int_s^t S_{t,u} y_u \cdot d\mathbf{X}_u - S_{t,s} (y_s \cdot \delta X_{t,s} + y'_s : \mathbb{X}_{t,s}) \right) \\ &\quad + (S_{t,s} - \text{id}) y_s \cdot \delta X_{t,s} + (S_{t,s} - \text{id}) \int_0^s S_{s,u} y_u \cdot d\mathbf{X}_u + S_{t,s} y'_s : \mathbb{X}_{t,s}, \\ &=: \mathcal{R}_{t,s} + \text{I}_{t,s} + \text{II}_{t,s} + \text{III}_{t,s}. \end{aligned}$$

Using the smoothing property (2.15) for S , we see that for $i = 1, 2$:

$$|\mathbf{I}_{t,s}|_{\alpha+\sigma-i\gamma'} \leq [X]_\gamma |t-s|^\gamma |S_{t,s} - \text{id}|_{\mathcal{L}(\mathcal{B}_\alpha, \mathcal{B}_{\alpha-(i\gamma'-\sigma)})} |y_s|_\alpha \lesssim |t-s|^{i\gamma'} T^{\gamma-\sigma} |y|_{0,\alpha}.$$

For the second term, we have, thanks to Theorem 4.5:

$$\begin{aligned} |\mathbf{II}_{t,s}|_{\alpha+\sigma-i\gamma'} &\lesssim |S_{t,s} - \text{id}|_{\mathcal{L}(\mathcal{B}_{\alpha+\sigma}, \mathcal{B}_{\alpha+\sigma-i\gamma'})} |S_{s,0} y_0|_{\alpha+\sigma} [X]_\gamma s^\gamma \\ &\quad + |S_{t,s} - \text{id}|_{\mathcal{L}(\mathcal{B}_{\alpha+\sigma}, \mathcal{B}_{\alpha+\sigma-i\gamma'})} |S_{s,0} y'_0|_{\alpha+\sigma} [\mathbb{X}]_{2\gamma} s^{2\gamma} \\ &\quad + |S_{t,s} - \text{id}|_{\mathcal{L}(\mathcal{B}_{\alpha+\sigma}, \mathcal{B}_{\alpha+\sigma-i\gamma'})} |\mathcal{R}_{s,0}|_{\alpha+\sigma} \\ &\lesssim \varrho_\gamma(\mathbf{X}) |t-s|^{i\gamma'} \left\{ s^{\gamma-\sigma} |y|_{0,\alpha} + |y'|_{0,\alpha-\gamma'} s^{2\gamma-\gamma'-\sigma} + s^{\gamma'-\sigma} \|y, y'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} \right\} \\ &\lesssim \varrho_\gamma(\mathbf{X}) \|y, y'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} |t-s|^{i\gamma'} T^{\gamma'-\sigma}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} |\mathbf{III}_{t,s}|_{\alpha+\sigma-\gamma'} &\leq [\mathbb{X}]_{2\gamma} |t-s|^{2\gamma} |S_{t,s}|_{\mathcal{L}(\mathcal{B}_{\alpha-\gamma'}, \mathcal{B}_{\alpha-\gamma'+\sigma})} |y'_s|_{\alpha-\gamma'} \\ &\lesssim \varrho_\gamma(\mathbf{X}) |t-s|^{2\gamma} |y'|_{0,\alpha-\gamma'} T^{2\gamma-\gamma'-\sigma} \\ |\mathbf{III}_{t,s}|_{\alpha+\sigma-2\gamma'} &\leq [\mathbb{X}]_{2\gamma} |t-s|^{2\gamma} |S_{t,s} y'_s|_{\alpha-\gamma'} \\ &\lesssim \varrho_\gamma(\mathbf{X}) |t-s|^{2\gamma} |y'|_{0,\alpha-\gamma'} T^{2\gamma-2\gamma'}. \end{aligned}$$

Combining the above estimates, we obtain:

$$\max_{i=1,2} [R^z]_{i\gamma', \alpha+\sigma-i\gamma'} \lesssim \varrho_\gamma(\mathbf{X}) T^\varepsilon \|y, y'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}}.$$

In particular, we see that z is controlled by X according to \mathcal{B}_α and that $z' = y$.

Next, to estimate the Hölder norm of the Gubinelli derivative, we first observe that

$$[R^y]_{\gamma', \alpha+\sigma-2\gamma'} \leq C |R^y|_{\mathcal{E}_\alpha^{\gamma', 2\gamma'}} T^{\gamma'-\sigma},$$

as can be easily seen by the interpolation inequality (4.6). Then, using that $\gamma' > \sigma$ we have

$$\begin{aligned} |\delta z'_{t,s}|_{\alpha+\sigma-2\gamma'} &= |\delta y_{t,s}|_{\alpha+\sigma-2\gamma'} \leq |y'_s|_{\alpha+\sigma-2\gamma'} |\delta X_{t,s}| + |R^y_{t,s}|_{\alpha+\sigma-2\gamma'}. \\ &\lesssim [X]_\gamma |y'|_{0,\alpha-\gamma'} |t-s|^{\gamma'} T^{\gamma-\gamma'} + |t-s|^{\gamma'} |R^y|_{\mathcal{E}_\alpha^{\gamma', 2\gamma'}} T^{\gamma'-\sigma}, \end{aligned}$$

which, by writing $z'_t = z'_0 + \delta z'_{t,0}$, yields the estimate

$$|z'|_{\mathcal{E}_{\alpha+\sigma-\gamma'}^{0,\gamma'}} \equiv |z'|_{0,\alpha+\sigma-\gamma'} + [\delta z']_{\gamma', \alpha+\sigma-2\gamma'} \lesssim |y_0|_\alpha + \varrho_\gamma(\mathbf{X}) T^\varepsilon \|y, y'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}}.$$

To conclude that (z, z') is controlled by X according to $\mathcal{B}_{\alpha+\sigma}$ it remains to estimate $|z|_{0,\alpha+\sigma}$, for which we use:

$$z_t = \mathcal{R}_{t,0} + S_{t,0} y_0 \cdot \delta X_{t,0} + S_{t,0} y'_0 : \mathbb{X}_{t,0}.$$

Therefore, using (4.8), smoothing properties of the propagator and $\varrho_{\gamma'}(\mathbf{X}) \leq \varrho_\gamma(\mathbf{X})$ we get

$$|z_t|_{\alpha+\sigma} \lesssim \varrho_\gamma(\mathbf{X}) \|y, y'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} t^{3\gamma'-2\gamma'-\sigma} + |y|_{0,\alpha} [X]_\gamma t^{\gamma-\sigma} + |y'|_{0,\alpha-\gamma'} [\mathbb{X}]_{2\gamma} t^{2\gamma-\gamma'-\sigma},$$

which implies that $|z|_{0,\alpha+\sigma} \lesssim \varrho_\gamma(\mathbf{X}) \|y, y'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} T^\varepsilon$, thus finishing the proof. \blacksquare

We are now going to see that a controlled path composed with some sufficiently regular function is again a controlled path.

Lemma 4.7. *Let $\gamma \in (1/3, 1/2]$, $\alpha \in \mathbf{R}$ and fix $\sigma \geq 0$. Let $F \in \mathcal{C}_{\alpha-2\gamma, -\sigma}^2(\mathcal{B})$ be some non-linearity with bounded derivatives up to second order. For $(y, y') \in \mathcal{D}_{X, \alpha}^{2\gamma}$, define*

$$(z_t, z'_t) := (F(y_t), DF(y_t) \circ y'_t), \quad \text{for every } t \in [0, T].$$

Then the following assertions are true.

(i) *One has $(z, z') \in \mathcal{D}_{X, \alpha-\sigma}^{2\gamma}$ and moreover:*

$$\|z, z'\|_{\mathcal{D}_{X, \alpha-\sigma}^{2\gamma}} \lesssim \|F\|_{\mathcal{C}^2} (1 + \varrho_\gamma(\mathbf{X}))^2 \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}} (1 + \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}}). \quad (4.9)$$

(ii) *If we assume further that $F \in \mathcal{C}_{\alpha-2\gamma, -\sigma}^3(\mathcal{B})$ with bounded third derivative, and if $(\tilde{z}, \tilde{z}') := (F(\tilde{y}), DF(\tilde{y}) \circ \tilde{y}')$ for another such pair $(\tilde{y}, \tilde{y}') \in \mathcal{D}_{X, \alpha}^{2\gamma}$, then the following estimate holds*

$$\begin{aligned} \|z - \tilde{z}, z' - \tilde{z}'\|_{\mathcal{D}_{X, \alpha-\sigma}^{2\gamma}} &\lesssim \|F\|_{\mathcal{C}^3} (1 + \varrho_\gamma(\mathbf{X}))^2 \|y - \tilde{y}, y' - \tilde{y}'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}} \\ &\quad \times (1 + \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}} + \|\tilde{y}, \tilde{y}'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}})^2. \end{aligned}$$

Where in both (i) and (ii) we set $\|F\|_{\mathcal{C}^k} = \max\{\|F\|_{\mathcal{C}^k(\mathcal{B}_{\alpha-\gamma}, \mathcal{B}_{\alpha-\gamma-\sigma})}, \|F\|_{\mathcal{C}^k(\mathcal{B}_{\alpha-2\gamma}, \mathcal{B}_{\alpha-2\gamma-\sigma})}\}$.

Proof. First, observe that because of the continuity of F and the inclusion $\mathcal{B}_\alpha \subset \mathcal{B}_{\alpha-\gamma}$, we have $(z, z') \in \mathcal{C}(0, T; \mathcal{B}_{\alpha-\sigma}) \times \mathcal{C}(0, T; \mathcal{B}_{\alpha-\sigma-\gamma}^d)$. We can view $D^k F(y_t)$ as an element of $\mathcal{L}(\mathcal{B}_{\alpha-i\gamma}^{\otimes k}, \mathcal{B}_{\alpha-i\gamma-\sigma})$ for $k = 1, 2, 3$ and $i = 1, 2$. With this at hand, we write $\delta z'_{t,s} = DF(y_s) \circ \delta y'_{t,s} + (DF(y_t) - DF(y_s)) \circ y'_t$, and since $|\cdot|_{\alpha-2\gamma} \leq |\cdot|_{\alpha-\gamma}$ we obtain:

$$\begin{aligned} |\delta z'|_{\gamma, \alpha-\sigma-2\gamma} &\lesssim |DF|_{\mathcal{L}(\mathcal{B}_{\alpha-2\gamma}, \mathcal{B}_{\alpha-2\gamma-\sigma})} [\delta y']_{\gamma, \alpha-2\gamma} \\ &\quad + |D^2 F|_{\mathcal{L}(\mathcal{B}_{\alpha-2\gamma}^{\otimes 2}, \mathcal{B}_{\alpha-2\gamma-\sigma})} [\delta y]_{\gamma, \alpha-2\gamma} |y'|_{0, \alpha-\gamma} \\ &\lesssim \|F\|_{\mathcal{C}^2} (1 + \varrho_\gamma(\mathbf{X})) \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}} (1 + \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}}), \end{aligned}$$

where we used (4.5) to estimate $[\delta y]_{\gamma, \alpha-2\gamma}$. Next, we estimate the remainder term

$$R_{t,s}^z := F(y_t) - F(y_s) - DF(y_s) \circ y'_s \cdot \delta X_{t,s},$$

and show that it belongs to $\mathcal{C}_2^{2\gamma}(0, T; \mathcal{B}_{\alpha-\sigma-2\gamma}) \cap \mathcal{C}_2^\gamma(0, T; \mathcal{B}_{\alpha-\sigma-\gamma})$. We rewrite R^z as

$$\begin{aligned} R_{t,s}^z &= F(y_t) - F(y_s) - DF(y_s) \circ \delta y_{t,s} + DF(y_s) \circ R_{t,s}^y \\ &= T_{t,s} + DF(y_s) \circ R_{t,s}^y, \end{aligned}$$

where, by Taylor's formula

$$T_{t,s} := \left(\int_0^1 \int_0^1 D^2 F(y_s + \theta\theta' \delta y_{t,s}) d\theta' d\theta \right) \circ (\delta y_{t,s} \otimes \delta y_{t,s}).$$

By definition of the spaces $\mathcal{C}_{\alpha-2\gamma, -\sigma}^2$, we have for $i = 1, 2$:

$$\begin{aligned} |R^z|_{i\gamma, \alpha-\sigma-i\gamma} &\leq |D^2F|_{\mathcal{L}(\mathcal{B}_{\alpha-i\gamma}^{\otimes 2}, \mathcal{B}_{\alpha-i\gamma-\sigma})} [\delta y]_{\gamma, \alpha-i\gamma}^2 \\ &\quad + |DF|_{\mathcal{L}(\mathcal{B}_{\alpha-i\gamma}, \mathcal{B}_{\alpha-i\gamma-\sigma})} [R^y]_{i\gamma, \alpha-i\gamma} \\ &\lesssim \|F\|_{\mathcal{C}^2} (1 + \varrho_\gamma(\mathbf{X}))^2 \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}} (1 + \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}}), \end{aligned}$$

which implies (i).

For (ii), we write (with obvious notations)

$$R_{t,s}^z - R_{t,s}^{\tilde{z}} = T_{t,s} - \tilde{T}_{t,s} + (DF(y_s) - DF(\tilde{y}_s)) \circ R_{t,s}^y + DF(\tilde{y}_s) \circ (R_{t,s}^y - R_{t,s}^{\tilde{y}}).$$

For $i = 1, 2$ we have

$$\begin{aligned} |T_{t,s} - \tilde{T}_{t,s}|_{\alpha-\sigma-i\gamma} &\leq |D^3F|_{\mathcal{L}(\mathcal{B}_{\alpha-i\gamma}^{\otimes 3}, \mathcal{B}_{\alpha-i\gamma-\sigma})} |y - \tilde{y}|_{0, \alpha-i\gamma} |\delta y_{t,s}|_{\alpha-i\gamma}^2 \\ &\quad + |D^2F|_{\mathcal{L}(\mathcal{B}_{\alpha-i\gamma}^{\otimes 2}, \mathcal{B}_{\alpha-i\gamma-\sigma})} |\delta y_{t,s} - \delta \tilde{y}_{t,s}|_{\alpha-i\gamma} |\delta \tilde{y}_{t,s}|_{\alpha-i\gamma} \\ &\lesssim \|F\|_{\mathcal{C}^3} |y - \tilde{y}|_{\mathcal{E}_\alpha^{0, \gamma}} (1 + [y]_{\gamma, \alpha-\gamma} + [\tilde{y}]_{\gamma, \alpha-\gamma})^2 |t - s|^{2\gamma}. \end{aligned}$$

Similarly,

$$|(DF(y_s) - DF(\tilde{y}_s)) \circ R_{t,s}^y|_{\alpha-i\gamma} \leq |D^2F|_{\mathcal{L}(\mathcal{B}_{\alpha-i\gamma}^{\otimes 2}, \mathcal{B}_{\alpha-i\gamma-\sigma})} |y - \tilde{y}|_{0, \alpha} |R^y|_{\mathcal{E}_\alpha^{\gamma, 2\gamma}} |t - s|^{i\gamma},$$

while

$$|DF(\tilde{y}_s) \circ (R_{t,s}^y - R_{t,s}^{\tilde{y}})|_{\alpha-i\gamma-\sigma} \leq |DF|_{\mathcal{L}(\mathcal{B}_{\alpha-i\gamma}, \mathcal{B}_{\alpha-i\gamma-\sigma})} |R^y - R^{\tilde{y}}|_{\mathcal{E}_\alpha^{\gamma, 2\gamma}}.$$

Summing the above three estimates yields the correct bound for $|R^z - R^{\tilde{z}}|_{\mathcal{E}_{\alpha-\sigma}^{\gamma, 2\gamma}}$.

For the Gubinelli derivatives, we have

$$\begin{aligned} &\delta(DF(y)y' - DF(\tilde{y})\tilde{y}')_{t,s} \\ &= \int_0^1 (D^2F(y_s + \theta\delta y_{t,s}) - D^2F(\tilde{y}_s + \theta\delta \tilde{y}_{t,s})) d\theta \circ (\delta y_{t,s} \otimes y'_t) \\ &\quad + \int_0^1 D^2F(\tilde{y}_s + \theta\delta \tilde{y}_{t,s}) d\theta \circ ((\delta y_{t,s} - \delta \tilde{y}_{t,s}) \otimes y'_t) \\ &\quad + \int_0^1 D^2F(\tilde{y}_s + \theta\delta \tilde{y}_{t,s}) d\theta \circ (\delta \tilde{y}_{t,s} \otimes (y'_t - \tilde{y}'_t)) \\ &\quad + (DF(y_s) - DF(\tilde{y}_s)) \circ \delta y'_{t,s} + DF(\tilde{y}_s) \circ (\delta y_{t,s} - \delta \tilde{y}'_{t,s}). \end{aligned}$$

This gives

$$|\delta z'_{t,s} - \delta \tilde{z}'_{t,s}|_{\alpha-2\gamma} \lesssim \|F\|_{\mathcal{C}^3} \|y - \tilde{y}, y' - \tilde{y}'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}} \left(1 + \|y, y'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}} + \|\tilde{y}, \tilde{y}'\|_{\mathcal{D}_{X, \alpha}^{2\gamma}}\right)^2.$$

This finishes the proof of Lemma 4.7. ■

5 Equations with subcritical multiplicative noise: proof of Theorem 2.15

In this section we fix $\gamma \in (1/3, 1/2]$, $\alpha \in \mathbf{R}$, a rough path $\mathbf{X} = (X, \mathbb{X}) \in \mathcal{C}^\gamma(0, T; \mathbf{R}^d)$, and we let $\sigma \in [0, \gamma)$. We will address the proof of local existence and uniqueness for the rough PDE

$$du_t = L_t u_t dt + N(u_t) dt + \sum_{i=1}^d F_i(u_t) d\mathbf{X}_t^i \quad \text{and} \quad u_0 = x \in \mathcal{B}_\alpha, \quad (5.1)$$

under suitable conditions on the non-linearities. The proof of Theorem 2.15 is a simple consequence of Theorem 5.1 below. Further properties of the solution map will be also given in Theorems 5.4 and 5.5.

In the sequel, an equation of the form (5.1) will be referred to as ‘subcritical’ provided that the function F sends \mathcal{B}_α to $\mathcal{B}_{\alpha-\sigma}$ with some $\sigma \in [0, \gamma)$. Concrete examples of subcritical equations will be given in Section 6.

5.1 Solutions to subcritical RPDEs

For $i = 1, \dots, d$, let $F_i \in \mathcal{C}_{\alpha-2\gamma, -\sigma}^2$, and for each $(y, y') \in \mathcal{D}_{X, \alpha}^{2\gamma}$ let

$$(z_t, z'_t) := \left(\int_0^t S_{t,s} F(y_s) \cdot d\mathbf{X}_s, F(y_t) \right), \quad \text{for every } t \in [0, T].$$

Then, Lemma 4.7 together with Corollary 4.6 gives us that (z, z') is again an element of the controlled paths space $\mathcal{D}_{X, \alpha}^{2\gamma}$. This is due to the fact that, though F reduces the spatial regularity by σ , the lost regularity is recovered from the smoothing properties of the integration map associated with S . This observation suggests that we might be successful in applying a Banach fixed point argument in order to solve (5.1) locally.

Theorem 5.1 (Local solution of subcritical RPDEs). *Fix $\alpha \in \mathbf{R}$, $\gamma \in (1/3, 1/2]$ and $\sigma \in [0, \gamma)$. Assume that we are given a non-linearity $F = (F_1, \dots, F_d)$ such that $F_i \in \mathcal{C}_{\alpha-2\gamma, -\sigma}^3(\mathcal{B})$ for $i = 1, \dots, d$ and $N \in \text{Lip}_{\alpha, -\delta}(\mathcal{B})$ for some $n \geq 1$ and $1 > \delta \geq 0$.*

For every $x \in \mathcal{B}_\alpha$, there exists $0 < \tau \leq T$ and a unique $(u, u') \in \mathcal{D}_{X, \alpha}^{2\gamma}([0, \tau])$ such that $u' = F(u)$ and

$$u_t = S_{t,0}x + \int_0^t S_{t,r} N(u_r) dr + \int_0^t S_{t,r} F(u_r) \cdot d\mathbf{X}_r, \quad t < \tau. \quad (5.2)$$

Proof. Assume first that $T \leq 1$ and let us define the map

$$\mathcal{M}_T(y, y')_t := \left(S_{t,0}x + \int_0^t S_{t,s} N(y_s) ds + \int_0^t S_{t,s} F(y_s) \cdot d\mathbf{X}_s, F(y_t) \right). \quad (5.3)$$

Instead of solving the equation directly in the space $\mathcal{D}_{X, \alpha}^{2\gamma}$ we will show that the map \mathcal{M}_T is invariant and contractive inside a ball of a larger space. We now fix a parameter $\gamma' \in (\sigma, \gamma)$ and let $\varepsilon = \min\{\gamma - \gamma', \gamma' - \sigma\}$. We further define two continuous paths $\xi : [0, T] \rightarrow \mathcal{B}_\alpha$ and $\xi' : [0, T] \rightarrow \mathcal{B}_{\alpha-\gamma'}$ as

$$\xi_t := S_{t,0}x + \int_0^t S_{t,r} F(x) \cdot d\mathbf{X}_r, \quad \xi'_t := F(x), \quad t \in [0, T],$$

and observe that $(\xi, \xi') \in \mathcal{D}_{X,\alpha}^{2\gamma'}$. This is indeed a consequence of Corollary 4.6 applied to the constant path $(F(x), 0) \in \mathcal{D}_{X,\alpha-\sigma}^{2\gamma'}$, and of the fact that $(S_{t,0}x, 0)$ belongs to $\mathcal{D}_{X,\alpha}^{2\gamma'}$ (using the smoothing properties of the propagator).

The first step is to show the existence of a positive $T_*(\varepsilon, \gamma, |x|_\alpha, \varrho_\gamma(\mathbf{X})) \leq 1$ such that for every $T \in [0, T_*]$ the map \mathcal{M}_T leaves the ball $B_T(x)$ invariant, where

$$B_T(x) = \left\{ (y, y') \in \mathcal{D}_{X,\alpha}^{2\gamma'}([0, T]) : (y_0, y'_0) = (x, F(x)) \text{ and } \|y - \xi, y' - \xi'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} \leq 1 \right\}.$$

Recall that from the definition of $\mathcal{C}_{\alpha-2\gamma, -\sigma}^3$ it follows that F_i together with its derivatives sends bounded sets of \mathcal{B}_β to $\mathcal{B}_{\beta-\sigma}$ for $\beta \geq \alpha - 2\gamma$. Therefore, without loss of generality one can view F_i and its derivatives to be bounded since we restrict ourselves to the ball $B_T(x)$. Similarly, one can assume that N is globally Lipschitz.

Step 1: Stability of $B_T(x)$. Let $(y, y') \in B_T(x)$ and let $(z, z') = \mathcal{M}_T(y, y')$. For simplicity denote the drift term $\mathcal{N}_t := \int_0^t S_{t,r} N(y_r) dr$ and define:

$$(\zeta, \zeta') := (z - \xi - \mathcal{N}, z' - \xi').$$

Note that $\zeta_0 = \zeta'_0 = 0$ and $\zeta_t = \int_0^t S_{t,r} (F(y_r) - F(x)) \cdot d\mathbf{X}_r$. Furthermore, it is readily checked by definition of $\|\cdot\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}}$ and the triangle inequality, that

$$\begin{aligned} \|z - \xi, z' - \xi'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} &\leq \|\mathcal{N}, 0\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} + \|\zeta, \zeta'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} \\ &= |\mathcal{N}|_{0,\alpha} + |\mathcal{N}|_{\mathcal{E}_\alpha^{\gamma', 2\gamma'}} + \|\zeta, \zeta'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}}. \end{aligned} \quad (5.4)$$

In order to estimate the drift term, we note that

$$\delta \mathcal{N}_{t,s} = (S_{t,s} - \text{id}) \int_0^s S_{s,r} N(y_r) dr + \int_s^t S_{t,r} N(y_r) dr.$$

The first term is easily estimated thanks to (2.15). We have indeed for $i = 0, 1, 2$ using $N \in \text{Lip}_{\alpha, -\delta}(\mathcal{B})$ and denoting by $\|N\|_1$ a Lipschitz constant for N as a function $\mathcal{B}_\alpha \rightarrow \mathcal{B}_{\alpha-\delta}$

$$\begin{aligned} \left| (S_{t,s} - \text{id}) \int_0^s S_{s,r} N(y_r) dr \right|_{\alpha-i\gamma'} &\lesssim \|N\|_1 |t-s|^{i\gamma'} \int_0^s (s-r)^{-\delta} (N(0) + |y_r|_\alpha) dr \\ &\lesssim_N |t-s|^{i\gamma'} T^{1-\delta} (1 + |y|_{0,\alpha}). \end{aligned}$$

For the second term, we have similarly

$$\begin{aligned} \left| \int_s^t S_{t,r} N(y_r) dr \right|_{\alpha-i\gamma'} &\lesssim \int_s^t |t-r|^{-\max\{0, \delta-i\gamma'\}} |N(y_r)|_{\alpha-\delta} dr \\ &\lesssim_N |t-s|^{\min\{1, 1+i\gamma'-\delta\}} (1 + |y|_{0,\alpha}). \end{aligned}$$

Note that thanks to our hypothesis that $1 - \delta > 0$ and $\gamma' < 1/2$ it follows that $\kappa := \min\{1 - 2\gamma', 1 - \delta\}$ is positive and

$$\max_{i=1,2,3} |\mathcal{N}|_{i\gamma', \alpha-i\gamma'} \lesssim_N T^\kappa (1 + |y|_{0,\alpha}) \leq T^\kappa (2 + \|\xi, \xi'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}}). \quad (5.5)$$

Next, note that Corollary 4.6 together with Lemma 4.7-(i) imply

$$\begin{aligned}
\|\zeta, \zeta'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} &\lesssim \varrho_\gamma(\mathbf{X}) T^\varepsilon \|F(y_\cdot) - F(x), DF(y_\cdot) \circ y'\|_{\mathcal{D}_{X,\alpha-\sigma}^{2\gamma'}} \\
&\lesssim \|F\|_{\mathcal{C}^2} (1 + \varrho_\gamma(\mathbf{X}))^3 T^\varepsilon \|y_\cdot - x, y'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} \\
&\leq \|F\|_{\mathcal{C}^2} (1 + \varrho_\gamma(\mathbf{X}))^3 T^\varepsilon \left(1 + \|\xi - x, \xi'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}}\right). \tag{5.6}
\end{aligned}$$

Putting together (5.6), (5.4) and (5.5), and noting that the bound on $\|\xi, \xi'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}}$ only depends on $F, x, \varrho_\gamma(\mathbf{X})$ we see that there is some constant $C > 0$ which only depends on $N, F, x, \varrho_\gamma(\mathbf{X})$ and the indices $\gamma, \gamma', \sigma, \delta$ such that:

$$\|z - \xi, z' - \xi'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} \leq CT^{\varepsilon \wedge \kappa}.$$

Taking T small enough, we see that $\mathcal{M}_T(B_T(x)) \subset B_T(x)$, which shows the claimed stability.

Step 2: contraction property. Consider now $(y^j, y^{j'}) \in B_T(x)$ and let $(z^j, z^{j'}) := \mathcal{M}_T(y^j, y^{j'})$, $j = 1, 2$. For every $(t, s) \in \Delta_2$, we have

$$\begin{aligned}
z_t^1 - z_t^2 &= \int_0^t S_{t,r} [N(y_r^1) - N(y_r^2)] dr + \int_0^t S_{t,r} [F(y_r^1) - F(y_r^2)] \cdot d\mathbf{X}_r \\
&= \bar{\mathcal{N}}_t + \bar{\zeta}_t.
\end{aligned}$$

Similarly, as above, using the Lipschitz property of N , we have for the drift term:

$$\begin{aligned}
|\delta \bar{\mathcal{N}}_{t,s}|_{\alpha-i\gamma'} &\lesssim \left| (S_{t,s} - \text{id}) \int_0^s S_{s,r} [N(y_r^1) - N(y_r^2)] dr + \int_s^t S_{t,r} [N(y_r^1) - N(y_r^2)] dr \right|_{\alpha-i\gamma'} \\
&\lesssim \|N\|_1 \left(|t-s|^{i\gamma'} \int_0^s (s-r)^{-\delta} dr + \int_s^t |t-r|^{-\max\{0, \delta-i\gamma'\}} dr \right) |y^1 - y^2|_{0,\alpha} \\
&\lesssim_{N,\alpha,|x|_\alpha, \mathbf{X}} T^\kappa \|y^1 - y^2, y^{1'} - y^{2'}\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}}.
\end{aligned}$$

Next, applying again Corollary 4.6 and then using Lemma 4.7-(ii), we get

$$\begin{aligned}
\|\bar{\zeta}, \bar{\zeta}'\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}} &\lesssim \varrho_\gamma(\mathbf{X}) T^\varepsilon \|y^1 - y^2, DF(y^1) \circ y^{1'} - DF(y^2) \circ y^{2'}\|_{\mathcal{D}_{X,\alpha-\sigma}^{2\gamma'}} \\
&\lesssim \|F\|_{\mathcal{C}^3} (1 + \varrho_\gamma(\mathbf{X}))^3 T^\varepsilon \|y^1 - y^1, y^{1'} - y^{2'}\|_{\mathcal{D}_{X,\alpha}^{2\gamma'}}.
\end{aligned}$$

This shows the claimed contraction property for $T \leq T_*$ small enough.

Applying Banach Fixed point Theorem, we see that there exists a unique fixed point $(u, u') \in \mathcal{D}_{X,\alpha}^{2\gamma'}$ for \mathcal{M}_T , and it is clearly a solution of (5.2) with $u' = F(u)$. Repeating the argument with $(T^1, x^1) := (T_*, u_{T_*})$ in place of $(0, x)$, we construct a sequence (T^n, x^n) and it is easily seen that $\tau := \sup_{n \in \mathbf{N}} T^n < T$ implies $\limsup_{n \rightarrow \infty} |x^n|_\alpha = \infty$. We then construct a solution on $[0, \tau)$ by ‘gluing together’ each solution on $[T^n, T^{n+1}]$. To show that this solution actually lives in the space $\mathcal{D}_{X,\alpha}^{2\gamma'}$ it suffices to use the equation, together with the bounds (4.8) and (4.9). Details are left to the reader. \blacksquare

Remark 5.2. Note that the techniques of the proof do not restrict us to the range of the Hölder regularity of the rough path to be $\gamma \in (1/3, 1/2]$ and one could generalize the statement and the whole theory above to the range $\gamma \in (1/n, 1/2]$ for $n \geq 3$. However, this would require a change in the definition of a controlled rough path in order to add a control of the higher order (up to $n - 2$) iterated integrals of X as explored in [30]. In turn, one would also need better smoothness properties on F (\mathcal{C}^n would do). We chose to avoid these considerations for computational convenience.

5.2 Continuity of the solution map and smoothing away from zero

Here we will briefly mention properties of the mild solution to the equation (5.1) that we constructed above. We will not present any proofs here since they are simply technical modifications of the analogous statements from [27] in the case of a constant differential operator L and a nonlinearity F which sends \mathcal{B}_α to itself (i.e there is no loss of space regularity from F). Our first result is going to be on the stability of the solution map (5.3) with respect to noise and the initial condition. In the case of a Brownian rough path, the continuity of the solution map is one of the main advantages of our pathwise formulation compared with Itô calculus (where only measurability holds in general). We now define a notion of distance between two controlled rough paths:

Definition 5.3. Let $\gamma \in (1/3, 1/2]$, $0 < \sigma < \gamma' \leq \gamma$ and let $\mathbf{X}, \tilde{\mathbf{X}} \in \mathcal{C}^\gamma(0, T; \mathbf{R}^d)$. For $(y, y') \in \mathcal{D}_{X, \alpha}^{2\gamma}([0, T])$ and $(z, z') \in \mathcal{D}_{\tilde{X}, \alpha}^{2\gamma}([0, T])$ define a distance $d_{2\gamma', 2\gamma, \alpha}$ as follows

$$d_{2\gamma', 2\gamma, \alpha}(y, z) = |y - z|_{0, \alpha} + |y' - z'|_{0, \alpha - \gamma} \\ + |y' - z'|_{\gamma', \alpha - 2\gamma} + [R^y - R^z]_{\gamma', \alpha - \gamma} + |R^y - R^z|_{2\gamma', \alpha - 2\gamma},$$

where we make an abuse of notation by not writing the dependence of $d_{2\gamma', 2\gamma, \alpha}(y, z)$ on $\mathbf{X}, \tilde{\mathbf{X}}$ and y', z' .

Theorem 5.4 (Stability of the solution to subcritical RPDE). *Let $\gamma \in (1/3, 1/2]$ and $\mathbf{X}, \tilde{\mathbf{X}} \in \mathcal{C}^\gamma(0, T; \mathbf{R}^d)$. Let $\alpha \in \mathbf{R}$ and $x, \tilde{x} \in \mathcal{B}_\alpha$. Assume that $F \in \mathcal{C}_{\alpha - 2\gamma, -\sigma}^3(\mathcal{B}, \mathcal{B}^d)$ for $0 \leq \sigma < \gamma$, and N be as in Theorem 5.1. Define $(u, F(u)) \in \mathcal{D}_{X, \alpha}^{2\gamma}([0, \tau_1])$ to be the solution to the RPDE:*

$$du_t = L_t u_t dt + N(u_t) dt + F(u_t) \cdot d\mathbf{X}_t, \quad u_0 = x \in \mathcal{B};$$

and $(z, F(z)) \in \mathcal{D}_{\tilde{X}, \alpha}^{2\gamma}([0, \tau_2])$ to be a solution of the same RPDE but driven by the rough path $\tilde{\mathbf{X}}$ and initial condition \tilde{x} . Assume that $\varrho_\gamma(\mathbf{X}), \varrho_\gamma(\tilde{\mathbf{X}}), |x|_\alpha, |\tilde{x}|_\alpha \leq M$. Then for every γ' such that $0 < \sigma < \gamma' \leq \gamma$ there exists time $\tau < 1 \wedge \tau_1 \wedge \tau_2$ such that for the following seminorm taken with respect to this time τ we have:

$$d_{2\gamma', 2\gamma, \alpha}(u, z) \lesssim_M \varrho_\gamma(\mathbf{X}, \tilde{\mathbf{X}}) + |x - \tilde{x}|_\alpha. \quad (5.7)$$

Moreover, if both solutions do not blow up before time T i.e. $(u, F(u)) \in \mathcal{D}_{X, \alpha}^{2\gamma}([0, T])$ and $(z, F(z)) \in \mathcal{D}_{\tilde{X}, \alpha}^{2\gamma}([0, T])$, then (5.7) holds on $[0, T]$.

The proof relies on the continuity properties of the integration map and composition with the smooth functions similar to Lemma 4.7 and Corollary 4.6 modified for the above metrics. The small sacrifice of time-regularity, which is represented by taking $\gamma' < \gamma$, and the fact

that u is a fixed point of the solution map (5.3) then allows us to use continuity properties of integration and composition to deduce for $\varepsilon = \min(\gamma - \gamma', \gamma' - \sigma, 1 - \delta, 1 - 2\gamma')$:

$$d_{2\gamma', 2\gamma, \alpha}(u, v) \lesssim d_{2\gamma', 2\gamma, \alpha}(u, v)\tau^\varepsilon + \varrho_\gamma(\mathbf{X}, \tilde{\mathbf{X}}) + |x - \tilde{x}|_\alpha.$$

Therefore, taking τ small enough, we obtain the desired result. The global Lipschitz estimate on the whole $[0, T]$ in (5.7) can be obtained simply by iteration of the result.

From classical PDE theory, it is expected that away from zero the solution is going to be infinitely smooth in space because of the smoothing property of the propagator. The following proposition tells us that it is indeed the case for our subcritical equations.

Proposition 5.5. *Let $\gamma \in (1/3, 1/2]$, $0 \leq \sigma < \gamma$, and $\mathbf{X} \in \mathcal{C}^\gamma(0, T; \mathbf{R}^d)$. Let $\alpha \in \mathbf{R}$, $x \in \mathcal{B}_\alpha$ and F, N be as in Theorem 5.4 and $(u, F(u)) \in \mathcal{D}_{X, \alpha}^{2\gamma}([0, \tau])$ be the solution to the equation (5.1). Denote $M_t := |u|_{0, \alpha, [0, t]}$ then for every $0 < s < t < \tau$ and $\beta > 0$ we have that $(u, F(u)) \in \mathcal{D}_{X, \alpha + \beta}^{2\gamma}([s, t])$ and there exist $\nu = \nu(\delta, \gamma, \sigma, \alpha)$, and a finite constant $C(M_t) = C(M_t, \tau, F, N, X)$ such that:*

$$|u|_{0, \alpha + \beta, [s, t]} \lesssim s^{-\beta} |u|_{0, \alpha, [0, t]} + C(M_t) t^\nu.$$

Remark 5.6. In [27] in the case of $F_i \in \mathcal{C}_{\alpha - 2\gamma, 0}^3$ and the operators L independent of time, the authors also show that solution to RPDE (2.22) when driven by the Brownian rough path coincides almost surely with the mild solution to the corresponding SPDE where the integration is given now in Itô sense. The authors also use Lipschitz continuity of the solution map Theorem 5.4 in order to show Malliavin differentiability of the solution to rough partial differential equations driven by a (regular enough) Gaussian rough path. We believe that the analogous results should hold true in the case of time dependent families L_t and $F_i \in \mathcal{C}_{\alpha - 2\gamma, -\sigma}^3$ with $\sigma < \gamma$.

5.3 Weak solutions. Proofs of Theorems 2.18 and 2.19

In this subsection, we are going to show the equivalence between mild solutions and the weak solutions that were already mentioned in Section 2.3. For notational convenience we will now restrict ourselves to the case when the initial condition belongs to the space \mathcal{B}_0 . With this at hand, we give a notion of the weak solution:

Definition 5.7. Let $(\mathcal{B}_\beta)_{\beta \in \mathbf{R}}$ be a monotone family of interpolation spaces and let $x \in \mathcal{B}_0$. Let \mathbf{X}, L, N, F be as in Theorem 5.1 and let $\nu = \max\{1, \sigma + 2\gamma\}$. We say that $(u, F(u)) \in \mathcal{D}_X^{2\gamma}([0, T], \mathcal{B}_0)$ is a weak solution of the rough PDE (1.1) if for all $\varphi \in \mathcal{B}_{-\nu}^*$ the following integral formula holds for all $0 \leq t \leq T$:

$$\langle u_t, \varphi \rangle = \langle u_0, \varphi \rangle + \int_0^t \langle L_s u_s, \varphi \rangle ds + \int_0^t \langle N(u_s), \varphi \rangle ds + \int_0^t \langle F(u_s), \varphi \rangle \cdot d\mathbf{X}_s, \quad (5.8)$$

where as in Theorem 4.5, $\langle F(u_s), \varphi \rangle$ denotes the vector $(\langle F_1(u_s), \varphi \rangle, \dots, \langle F_d(u_s), \varphi \rangle)$.

Note that all the above integrals make sense. Since $-\nu \leq -1$ and because of the dense inclusions $\mathcal{B}_0 \subset \mathcal{B}_{-\delta} \subset \mathcal{B}_{-1}$ we also get the reverse inclusions $\mathcal{B}_{-1}^* \subset \mathcal{B}_{-\delta}^* \subset \mathcal{B}_0^*$ and thus all the terms $\langle u_0, \varphi \rangle, \langle L_s u_s, \varphi \rangle, \langle N(u_s), \varphi \rangle$ are well-defined. Moreover, denoting by $\mathcal{D}_X^{2\gamma}(0, T; V)$ the usual controlled rough path spaces in the Banach space V (in the sense of [25, Definition 4.6]) we note that $(F(u), DF(u)F(u)) \in \mathcal{D}_X^{2\gamma}([0, T], \mathcal{B}_{-\sigma})$ implies that

$(F(u), DF(u)F(u)) \in \mathcal{D}_X^{2\gamma}(0, T; \mathcal{B}_{-\sigma-2\gamma})$. Using the fact that $\varphi \in \mathcal{B}_{-\nu}^* \subseteq \mathcal{B}_{-\sigma-2\gamma}^*$ we can easily show that

$$\left(\langle F(u), \varphi \rangle, \langle DF(u)F(u), \varphi \rangle \right) \in \mathcal{D}_X^{2\gamma}(0, T; \mathbf{R}^d),$$

which implies that the rough integral in (5.8) is well-defined.

Before showing the equivalence of mild and weak solutions, we need the following technical lemma:

Lemma 5.8. *Let $\mathbf{X} \in \mathcal{C}^\gamma(0, T; \mathbf{R}^d)$ for $\gamma \in (1/3, 1/2]$. Then for every $\varphi \in \mathcal{C}(0, T; \mathcal{B}_0^*)$ and $(y, y') \in \mathcal{D}_X^{2\gamma}([0, T], \mathcal{B}_0)$ we have for each $0 \leq t \leq T$:*

$$\int_0^t \left\langle \int_0^s S_{r,s} y_r \cdot d\mathbf{X}_r, \varphi_s \right\rangle ds = \int_0^t \int_r^t \langle S_{r,s} y_r, \varphi_s \rangle ds \cdot d\mathbf{X}_r.$$

The proof can be carried out *mutatis mutandis* as in [27, Lemma 6.2]. The only difference is that instead of the inner product of Hilbert space \mathcal{H} we use testing with the functions from \mathcal{B}_0^* , thus one should simply replace the Cauchy-Schwarz inequality with the boundedness of the functional φ . With this at hand:

Theorem 5.9. *Let $(\mathcal{B}_\beta)_{\beta \in \mathbf{R}}$ be a monotone family of interpolation spaces and let $u_0 \in \mathcal{B}_0$. Let \mathbf{X}, L, N, F be as in Theorem 5.1. Assume that for all $\beta \in \mathbf{R}$ one has $L \in \mathcal{C}(0, T; \mathcal{L}(\mathcal{B}_\beta, \mathcal{B}_{\beta-1}))$, Then $(u, F(u)) \in \mathcal{D}_X^{2\gamma}([0, T], \mathcal{B}_0)$ is a mild solution of the rough PDE (1.1) namely it satisfies*

$$u_t = S_{t,0} x + \int_0^t S_{t,r} N(u_r) dr + \int_0^t S_{t,r} F(u_r) \cdot d\mathbf{X}_r,$$

if and only if it is a weak solution in the sense of Definition 5.7.

Proof. Without loss of generality we can assume in both cases that $x = 0$ by replacing (u_t, u_t') by $(u_t + S_{t,0}x, u_t')$ (using $\delta^S S_{\cdot,0}x = 0$).

Mild \Rightarrow Weak. Assume also for simplicity that $N = 0$ since dealing with the drift term is easier than with the diffusion term F . Now let $(u, F(u)) \in \mathcal{D}_X^{2\gamma}([0, T], \mathcal{B}_0)$ satisfy for $0 \leq t \leq T$

$$u_t = \int_0^t S_{t,s} F(u_s) \cdot d\mathbf{X}_s.$$

Let $\varphi \in \mathcal{B}_{-\nu}^*$ be arbitrary for $\nu = \max\{1, \sigma + 2\gamma\}$. Note that since $\mathcal{B}_{-\nu}$ is continuously embedded in \mathcal{B}_{-1} then $L \in \mathcal{C}(0, T; \mathcal{L}(\mathcal{B}_0, \mathcal{B}_{-\nu}))$ implying $L^* \varphi \in \mathcal{C}(0, T; \mathcal{B}_0^*)$. Thus, testing with $L_s^* \varphi$ and integrating from 0 to t gives

$$\begin{aligned} \int_0^t \langle L_s u_s, \varphi \rangle ds &= \int_0^t \left\langle L_s \int_0^s S_{s,r} F(u_r) \cdot d\mathbf{X}_r, \varphi \right\rangle ds \\ &= \int_0^t \int_r^t \langle L_s S_{s,r} F(u_r), \varphi \rangle ds \cdot d\mathbf{X}_r \\ &= \int_0^t \left\langle \int_r^t L_s S_{s,r} F(u_r) ds, \varphi \right\rangle \cdot d\mathbf{X}_r \\ &= \int_0^t \left\langle \int_r^t \frac{d}{ds} S_{s,r} F(u_r) ds, \varphi \right\rangle \cdot d\mathbf{X}_r \\ &= \int_0^t \langle S_{t,r} F(u_r), \varphi \rangle d\mathbf{X}_r - \int_0^t \langle F(u_r), \varphi \rangle \cdot d\mathbf{X}_r \\ &= \langle u_t, \varphi \rangle - \int_0^t \langle F(u_r), \varphi \rangle \cdot d\mathbf{X}_r, \end{aligned}$$

where we used Lemma 5.8 in the second equality together with the fact that $L_s^* \varphi$ is a continuous function in time with values in \mathcal{B}_0^* , and the fourth equality is justified by (P3*). **Weak \Rightarrow Mild.** The proof is similar to the standard proof for SPDEs found either in [16] or [33]. All the additional difficulties are similar to the ones in the proof of Lemma 6.1 in the case of supercritical noise. \blacksquare

We now come back to the applications of this Theorem to the stochastic setting and address the proof of Theorem 2.19.

Proof of existence in Theorem 2.19. First, note that the multidimensional Brownian motion can be lifted almost surely to an ω -dependent rough path $(B(\omega), \mathbb{B}(\omega))$ by setting for $(t, s) \in \Delta_2$:

$$\mathbb{B}_{t,s}^{i,j} := \int_s^t (B_r^i - B_s^i) dB_r^j,$$

in the sense of Itô integration. The corresponding random variable is supported in the space $\mathcal{C}^\gamma(0, T; \mathbf{R}^d)$ for any $\gamma < \frac{1}{2}$. For details, we refer to [25, Section 10].

Fix $\gamma \in (1/3, 1/2)$ such that $1/3 < \gamma < \frac{1}{4}(k - n/p)$, which is possible by assumption $k > n/p + 4/3$. With all the above requirements satisfied, we see that pathwise, the equation (2.24) falls into the framework of Theorem 5.1. Indeed, apply Theorem 5.1 \mathbb{P} -almost surely, with the family $(L_t(\omega))_{t \in [0, T]}$ defined as $L_t(\omega) := \nabla \cdot (a_t(\omega, \cdot) \nabla(\cdot))$ to solve equation (2.24) on $\mathcal{D}_{B,0}^{2\gamma}$ for $\mathcal{B}_0 = H^{k,p}(\mathbf{T}^n)$. Note that we have chosen γ so that $k > n/p + 4\gamma$. Therefore, the spaces $\mathcal{B}_\alpha = H^{k+\alpha,p}(\mathbf{T}^n)$ are Banach algebras for every $\alpha > -2\gamma$. It is now easy to see that $f \in \mathcal{C}_{0,0}^\infty(\mathcal{B})$ and $p_i \in \mathcal{C}_{-2\gamma,0}^\infty(\mathcal{B})$ (see Section 2.4). We conclude that there exists a unique local solution $u(\omega)$ to (2.25) with a potential blow up time $\tau(\omega)$, such that $u \in \mathcal{C}([0, \tau(\omega)), H^{k,p}(\mathbf{T}^n))$.

Now one can easily see that for $\varphi \in \mathcal{C}^\infty(\mathbf{T}^n)$, the functional $\langle \cdot, \varphi \rangle_{L^2(\mathbf{T}^n)}$ belongs to \mathcal{B}_{-1}^* (here we need \mathcal{B}_{-1}^* since in the case of reaction diffusion equations $\sigma = 0$ and therefore $\nu = \max\{1, \sigma + 2\gamma\} = 1$). This implies that u satisfies almost surely on $\{t < \tau\}$ the integral formula:

$$\begin{aligned} \langle u_t, \varphi \rangle_{L^2(\mathbf{T}^n)} &= \langle u_0, \varphi \rangle_{L^2(\mathbf{T}^n)} + \int_0^t \langle \nabla \cdot (a_t(\cdot) \nabla u_s), \varphi \rangle_{L^2(\mathbf{T}^n)} ds \\ &\quad + \int_0^t \langle f(u_s), \varphi \rangle_{L^2(\mathbf{T}^n)} ds + \sum_{i=1}^d \int_0^t \langle p_i(u_s), \varphi \rangle_{L^2(\mathbf{T}^n)} \cdot d\mathbf{B}_s^i. \end{aligned}$$

From [25, Section 9] we see that the above rough integrals coincide with the usual Itô integral almost surely since the processes $\langle p_i(u_s), \varphi \rangle_{L^2(\mathbf{T}^n)}$ are adapted. This implies that the constructed solution is indeed a weak Itô solution. \blacksquare

Proof of uniqueness in Theorem 2.19. In this proof we assume that $f = 0$ since working with the diffusion term only adds notational difficulty. For notational simplicity, in the sequel we write $\mathcal{C}(\mathcal{B}_0)$ instead of $\mathcal{C}([0, T], \mathcal{B}_0)$, and similarly for other spaces. The proof requires a localization argument: fix an arbitrary $\varrho > 0$, define the stopping time $\tau_\varrho := \inf\{t \in (0, \tau) : |u_t|_0 \geq \varrho\}$ and denote by u^ϱ the stopped process $u_t^\varrho := u_{t \wedge \tau_\varrho}$, $t \in [0, T]$.

We will prove that for any⁶ $\gamma \in (1/3, 1/2)$ and every $\gamma > \varepsilon > 0$, the remainder

⁶In fact it is enough to show this for some $\gamma \in (1/3, 1/2)$, but we will prove a slightly stronger statement.

$$\begin{aligned}
R_{t,s}^{u^\varrho}(x) &= u_t^\varrho(x) - u_s^\varrho(x) - \sum_{i=1}^d p_i(u_s^\varrho(x)) \delta B_{t \wedge \tau_\varrho, s \wedge \tau_\varrho}^i \\
&= \int_{s \wedge \tau_\varrho}^{t \wedge \tau_\varrho} L_r u_r dr + \sum_{i=1}^d \int_{s \wedge \tau_\varrho}^{t \wedge \tau_\varrho} (p_i(u_r^\varrho) - p_i(u_s^\varrho)) dB_r^i,
\end{aligned}$$

belongs to $\mathcal{C}_2^{2\gamma-\varepsilon}(\mathcal{B}_{-2\gamma}) \cap \mathcal{C}_2^{\gamma-\varepsilon}(\mathcal{B}_{-\gamma})$, \mathbb{P} -almost surely. The uniqueness then follows from a slight modification of Theorem 2.18. Indeed, since we have that the Brownian rough path (B, \mathbb{B}) lives in $\mathcal{C}^{\frac{1}{2}-\delta}$ for every $\delta > 0$, \mathbb{P} -almost surely, then one can use some extra regularity of the Brownian rough path to show that the solution map (5.3) for the equation (2.25) has a unique fixed point in the space

$$\{(v, v', R^v) \in \mathcal{C}(\mathcal{B}_0) \times \mathcal{C}^{\gamma-\varepsilon}(\mathcal{B}_0) \times (\mathcal{C}_2^{2\gamma-\varepsilon}(\mathcal{B}_{-2\gamma}) \cap \mathcal{C}_2^{\gamma-\varepsilon}(\mathcal{B}_{-\gamma}))\}. \quad (5.9)$$

The rest of the proof is split in two parts. First, we show that for every $\gamma \in (1/3, 1/2)$ and $\varepsilon \in (0, \gamma)$, u^ϱ is $(\gamma - \varepsilon)$ -Hölder with values in $\mathcal{B}_{-\gamma}$ with full probability. Then, we use this information to deduce that the two-parameter quantity R^{u^ϱ} defined above lies in $\mathcal{C}_2^{2\gamma-\varepsilon}(\mathcal{B}_{-2\gamma})$. This, together with the assumption that the weak solution u belongs to $\mathcal{C}(\mathcal{B}_0)$, easily implies that $(u^\varrho, p(u^\varrho), R^{u^\varrho})$ belongs to the space defined in (5.9) and therefore must be unique solution on the interval $[0, \tau_\varrho]$. Therefore, since such uniqueness is true for every $\varrho > 0$ and since $u^\varrho = u$ on $[0, \tau_\varrho]$ one must have that u is a unique weak Itô solution to (2.25) on $[0, \tau)$.

Step 1: proof that $\mathbb{P}(u^\varrho \in \mathcal{C}^{\gamma-\varepsilon}(\mathcal{B}_{-\gamma})) = 1$. Recall the notations of Theorem 2.18. Since here $\sigma = 0$, we have that $\nu = \max(1, \sigma + 2\gamma) = 1$. Let $M_t := \sum_{i=1}^d \int_0^{t \wedge \tau_\varrho} p_i(u_s) dB_s^i$, we will first show that for every $\gamma \in [0, 1/2)$, $M \in \mathcal{C}^\gamma(\mathcal{B}_0)$, \mathbb{P} -a.s. Observe by a standard density argument that (2.26) holds for any $\varphi \in \mathcal{B}_{-1}^* \simeq H^{-k+2, q}(\mathbf{T}^n)$, where here q is the conjugate exponent i.e. $p^{-1} + q^{-1} = 1$. Fix $\varphi \in \mathcal{B}_{-1}^* \subset \mathcal{B}_0^*$, take $(t, s) \in \Delta_2$ and use Burkholder–Davis–Gundy inequality to obtain

$$\mathbb{E}|\langle \delta M_{t,s}, \varphi \rangle|^m \leq C(p, \varrho)^m |t - s|^{m/2} |\varphi|_{\mathcal{B}_0^*}^m, \quad (5.10)$$

where C is a constant depending on the polynomials p_i and ϱ . Kolmogorov's continuity criterion implies that for any $\gamma \in [0, \frac{1}{2} - \frac{1}{m})$, there is an event Ω^φ of full probability such that $\langle M, (\omega), \varphi \rangle$ belongs to $\mathcal{C}^\gamma(\mathbf{R})$, for any $\omega \in \Omega^\varphi$. Let $(\varphi_n)_{n \in \mathbf{N}}$ be a sequence of elements in \mathcal{B}_{-1}^* such that $\{\varphi_n\}_{n \in \mathbf{N}}$ is dense in \mathcal{B}_0^* , define $\Omega_\infty := \bigcap_{n \in \mathbf{N}} \Omega^{\varphi_n}$ and observe that $\mathbb{P}(\Omega_\infty) = 1$. By the previous discussion, we further remark that for each $n \in \mathbf{N}$, $(t, s) \in \Delta_2$, and $\omega \in \Omega$:

$$\mathbf{1}_{\Omega_\infty}(\omega) |t - s|^{-\gamma} |\langle \delta M_{t,s}(\omega), \varphi_n \rangle| \leq \tilde{C}_\varrho(\omega) |\varphi_n|_{\mathcal{B}_0^*}, \quad (5.11)$$

where \tilde{C}_ϱ is a random constant with $\mathbb{E} \tilde{C}_\varrho^m \lesssim_m C(p, \varrho)^m$. Since both sides of (5.11) are continuous with respect to the \mathcal{B}_0^* -norm, we conclude that the same relation holds true for each $\varphi \in \mathcal{B}_0^*$. Taking the supremum with respect to $\varphi \in \mathcal{B}_0^*$ and $(t, s) \in \Delta_2$ shows the desired conclusion. Taking m big enough, we can guarantee in particular that γ can be taken arbitrary on the interval $(1/3, 1/2)$.

Now set $g_t := L_t M_t = L_t \sum_{i=1}^d \int_0^{t \wedge \tau_\varrho} p_i(u_s) dB_s^i$. By the above computation and since $L \in \mathcal{C}(0, T; \mathcal{L}(\mathcal{B}_0, \mathcal{B}_{-1}))$ we have $g \in \mathcal{C}(\mathcal{B}_{-1})$, \mathbb{P} -almost surely. To be more precise note the following estimate for every $h \in \mathcal{B}_0 = H^{k,p}(\mathbf{T}^n)$

$$|L_t h|_{-1} = |(1 - \Delta)^{\frac{k}{2}-1} \nabla \cdot (a_t \nabla h)|_{L^p(\mathbf{T}^n)} \lesssim |a_t|_{\mathcal{C}^{k-1}(\mathbf{T}^n)} |h|_{H^{k,p}(\mathbf{T}^n)} \leq A |h|_0, \quad (5.12)$$

where we use notation $A = \sup_{t \in [0, T]} |a_t|_{\mathcal{C}^{k-1}(\mathbf{T}^n)}$. Therefore, $|g|_{0,-1} \leq A|M|_{0,0}$. Next, it follows that \mathbb{P} -almost surely, $u_t^\varrho - M_t = v_{t \wedge \tau_\varrho}$ where v is defined as a weak (hence a mild) solution of the following parabolic equation with random coefficients

$$\begin{aligned} \partial_t v_t - L_t v_t &= g_t \quad \text{on } [0, T] \\ v_0 &= u_0 \in \mathcal{B}_0. \end{aligned} \tag{5.13}$$

Now, for each $t \in [0, T]$ we have

$$\delta v_{t,s} = (S_{t,s} - \text{id})S_{s,0}u_0 + \int_s^t S_{t,r}g_r dr + (S_{t,s} - \text{id}) \int_0^s S_{s,r}g_r dr.$$

It is easy to see that the first two terms can be bounded by $(|u_0|_0 + |g|_{0,-1})|t-s|^\gamma$ in $\mathcal{B}_{-\gamma}$. For the third term, for any $0 < \varepsilon < \gamma$ we have

$$\begin{aligned} |(S_{t,s} - \text{id}) \int_0^s S_{s,r}g_r dr|_{-\gamma} &\lesssim |t-s|^{\gamma-\varepsilon} \int_0^s |S_{s,r}g_r|_{-\varepsilon} dr \\ &\lesssim |t-s|^{\gamma-\varepsilon} \int_0^s |s-r|^{-1+\varepsilon} |g_r|_{-1} dr \lesssim |t-s|^{\gamma-\varepsilon} T^\varepsilon |g|_{0,-1}, \end{aligned}$$

where we used that $s \leq T$. Therefore, $v \in \mathcal{C}^{\gamma-\varepsilon}(\mathcal{B}_{-\gamma})$ and thus $u^\varrho = v_{\cdot \wedge \tau_\varrho} + M \in \mathcal{C}^{\gamma-\varepsilon}(\mathcal{B}_{-\gamma})$ with probability one. Moreover, using (5.12) we obtain

$$[u^\varrho]_{\gamma-\varepsilon, -\gamma} \lesssim_T |u_0|_0 + |g|_{0,-1} + [M]_{\gamma, -\gamma} \lesssim |u_0|_0 + (A+1)\tilde{C}_\varrho. \tag{5.14}$$

Step 2: proof that $\mathbb{P}(R^{u^\varrho} \in \mathcal{C}_2^{2\gamma-\varepsilon}(\mathcal{B}_{-2\gamma})) = 1$. First, note two elementary bounds

$$\begin{aligned} \left| \int_{s \wedge \tau_\varrho}^{t \wedge \tau_\varrho} L_r u_r \right|_{-1} &\lesssim |t-s| A |u^\varrho|_{0,0}, \\ \left| \int_{s \wedge \tau_\varrho}^{t \wedge \tau_\varrho} L_r u_r \right|_0 &= |\delta u_{t,s}^\varrho - \delta M_{t,s}|_0 \lesssim |u^\varrho|_{0,0} + |M|_{0,0}. \end{aligned}$$

The interpolation inequality (2.1) implies then for any $\theta \in [0, 1]$ and every $(t, s) \in \Delta_2$

$$\left| \int_{s \wedge \tau_\varrho}^{t \wedge \tau_\varrho} L_r u_r \right|_{-\theta} \leq C_\varrho |t-s|^\theta A^\theta (1 + |M|_{0,0})^{1-\theta}, \tag{5.15}$$

with potentially different constant C_ϱ . Using the equation for u and applying (5.15) with $\theta = 2\gamma$ yields for any $\varphi \in \mathcal{B}_{-1}^* \subset \mathcal{B}_{-2\gamma}^*$

$$\begin{aligned} \mathbb{E} |\langle R_{t,s}^{u^\varrho}, \varphi \rangle|^m &\lesssim \mathbb{E} \left| \int_{s \wedge \tau_\varrho}^{t \wedge \tau_\varrho} \langle L_r u_r, \varphi \rangle dr \right|^m + \sum_{i=1}^d \mathbb{E} \left| \int_{s \wedge \tau_\varrho}^{t \wedge \tau_\varrho} \langle \delta p_i(u^\varrho)_{r,s}, \varphi \rangle dB_r^i \right|^m \\ &\lesssim_{\varrho, m, p} |\varphi|_{\mathcal{B}_{-2\gamma}^*}^m [|t-s|^{2\gamma m} \mathbb{E} A^{2\gamma m} (1 + |M|_{0,0})^{(1-2\gamma)m} \\ &\quad + |t-s|^{(\gamma-\varepsilon+\frac{1}{2})m} \mathbb{E} [u^\varrho]_{\gamma-\varepsilon, -\gamma}^m] \\ &\leq |\varphi|_{\mathcal{B}_{-2\gamma}^*}^m C(\varrho, K_m, p, T) |t-s|^{((\gamma-\varepsilon+\frac{1}{2}) \wedge 2\gamma)m}, \end{aligned}$$

where we used Burkholder–Davis–Gundy in the second inequality, the moments estimate (2.27) on A together with moments bounds on $|M|_{0,0}$ and the estimate (5.14) on $[u^\varrho]_{\gamma-\varepsilon, -\gamma}$ in the last inequality.

Using a two-parameter version of Kolmogorov’s criterion (see [23]), we obtain by a similar argument as in the previous step, that R^{u^ϱ} belongs to $\mathcal{C}_2^\beta(\mathcal{B}_{-2\gamma})$ with probability one, for any $\beta < (\gamma - \varepsilon + \frac{1}{2}) \wedge 2\gamma - \frac{1}{m}$. Taking m large enough guarantees $R^{u^\varrho} \in \mathcal{C}_2^{2\gamma-\varepsilon}(\mathcal{B}_{-2\gamma})$. This concludes our second step, thus finishing the proof of uniqueness. \blacksquare

6 Further examples

In this section, we present two equations that can be covered by Theorem 2.15, and we quickly explain why the required assumptions are satisfied. For simplicity only, the examples given here are autonomous, namely L_t is constant in time (non-autonomous versions of the following are easily seen to be covered as well, we refer for instance to section 2.4).

For completeness, and because it was quickly discussed in the introduction, we will also provide an example of an equation where the diffusion term is not subcritical, and thus our framework does not apply (though existence and uniqueness were shown in previous papers, based on a priori estimates). Nevertheless, we will show that the variational solution satisfies also a suitable mild formulation, which could be useful to prove regularity results.

We recall that, given a filtered probability space $(\Omega, \mathcal{F}, \mathbb{P}, (\mathcal{F}_t)_{t \in [0, T]})$, a fractional Brownian motion (fBM) with Hurst parameter $H \in (1/3, 1/2]$ is defined as a Gaussian process B^H with covariance such that

$$\mathbb{E}[B_t^H B_s^H] = \frac{1}{2}(s^{2H} + t^{2H} - |t - s|^{2H}),$$

which covers the standard Brownian motion if one lets $H = 1/2$.

6.1 Stochastic Navier-Stokes equation

The two-dimensional Navier-Stokes equation describes the time evolution of the velocity $u_t(x)$ of an incompressible fluid on a surface (here represented by the flat torus \mathbf{T}^2 for simplicity). A perturbation of the latter by a singular, finite-dimensional noise term can be written, for parameters $1/3 < H$ and $\sigma < H$, as

$$\begin{aligned} du_t(x) &= \Delta u_t(x) dt + \mathcal{B}(\mathcal{K}u_t(x), u_t(x)) dt + f(x)(-\Delta)^\sigma u_t(x) dB_t^H, \\ u_0 &\in L^p(\mathbf{T}^2, \mathbf{R}^2) \quad \text{with} \quad \int_{\mathbf{T}^2} u_0(x) dx = 0. \end{aligned} \quad (6.1)$$

where f is smooth and bounded, while B_t^H denotes a fractional Brownian motion enhanced to a rough path $\mathbf{B}^H := (B^H(\omega), \mathbb{B}^H(\omega)) \in \mathcal{C}^\gamma(0, T; \mathbf{R}^d)$ for every $\gamma < H$ (see Theorem 2.19). The bilinear operator \mathcal{B} is defined formally as $\mathcal{B}(u, w) = -(u \cdot \nabla)w$, while \mathcal{K} is the continuous linear mapping defined in the Fourier space as

$$\mathcal{F}(\mathcal{K}w)(k) = \frac{-\mathcal{F}(w)(k_2, -k_1)}{k_1^2 + k_2^2}, \quad k \equiv (k_1, k_2) \in \mathbf{Z}^2,$$

\mathcal{F} being the discrete Fourier transform. Fix $p \in (1, \infty)$ and introduce the scale of Banach spaces $\mathcal{B}_\beta = H_0^{2\beta, p}(\mathbf{T}^2, \mathbf{R}^2)$ which consists of the Bessel potential spaces on \mathbf{T}^2 intersected with the space of distributions u such that $\mathcal{F}(u)(0) = 0$. It is well known (see [52]) that the Laplacian Δ generates a semigroup $(S_t)_{t \in [0, \infty)}$ on the full range $(\mathcal{B}_\beta)_{\beta \in \mathbf{R}}$ in the sense of Definition 2.11. Moreover, the non-linearity $N(u) = \mathcal{B}(\mathcal{K}u, u)$ lies in $\text{Lip}_{0, -\delta}(\mathcal{B})$ and any $1 > \delta > 1/2$ (we refer for instance to Section 8 in [34] for details).

The operator $(-\Delta)^\sigma$ sends \mathcal{B}_β to $\mathcal{B}_{\beta-\sigma}$ for all $\beta \in \mathbf{R}$, while multiplication with the smooth function f is a smooth operation from $\mathcal{B}_{\beta-\sigma}$ to itself. Hence, we obtain that for every initial condition $u_0 \in L^p(\mathbf{T}^2, \mathbf{R}^2)$ with $\int_{\mathbf{T}^2} u_0(x) dx = 0$ (meaning $u_0 \in \mathcal{B}_0$) and for almost every realization $\mathbf{B}^H(\omega)$ of the fractional Brownian Motion, there is a unique maximal $\tau(\omega)$ and $u(\omega)$, solution of (6.1) on $[0, \tau(\omega))$.

For related investigations on rough Navier-Stokes equations, we also refer to [38, 39], where a slightly different model is investigated based on a priori estimates.

6.2 Cahn-Hilliard equation

Let again $H > 1/3$, fix a dimension $n \geq 1$, a let $\mathcal{O} \subset \mathbf{R}^n$ be a bounded domain with smooth boundary. We consider the semilinear equation

$$du_t(x) = -\Delta(\Delta u_t(x) - u_t^3(x) + u_t(x))dt + \sum_{i=1}^d f_i(x) \cdot \nabla u_t(x) dB_t^{H,i}, \quad (6.2)$$

$$u_t(x) = \Delta u_t(x) = 0 \quad \text{on} \quad [0, T] \times \partial\mathcal{O}, \quad (6.3)$$

where $f_i \in \mathcal{C}_c^\infty(\mathcal{O}; \mathbf{R}^n)$ and the pair $(u_0, \Delta u_0)$ lies in $H_0^{k,p}(\mathcal{O}) \times H_0^{k-2,p}(\mathcal{O})$ and with $(k, p) \in [0, \infty) \times (1, \infty)$ chosen so that $k > \frac{n}{p} \vee (2 + \frac{1}{p})$ (the second condition guarantees that the trace at $\partial\mathcal{O}$ of Δu_0 is meaningful). Because of the mixed boundary conditions, it is more convenient in this case to work within a scale constructed from the fractional powers of the bi-Laplacian. For that purpose, we introduce the following operator on $L^p(\mathcal{O})$

$$(L, D(L)) = (-\Delta^2, \{u \in H^{4,p}(\mathcal{O}) : u = \Delta u = 0 \text{ on } \partial\mathcal{O}\}). \quad (6.4)$$

Next, we introduce the scale of Banach spaces

$$(\mathcal{B}_\beta, |\cdot|_\beta) := (D(-L)^{\frac{k}{4}+\beta}, |(-L)^{\frac{k}{4}+\beta} \cdot |), \quad \text{if } \beta \geq 0, \quad (6.5)$$

and if $\beta < 0$ one proceeds by completion as in Example 2.2. With the definition (6.4), the bi-Laplacian coincides with the square $(i\Delta_D)^2$ where Δ_D is the Dirichlet Laplacian, namely

$$(\Delta_D, D(\Delta_D)) = (\Delta, H^{2,p}(\mathcal{O}) \cap H_0^{1,p}(\mathcal{O})).$$

This implies that, $(L, D(L))$ satisfies the hypotheses of [3, Corollary 3.7.15], and we infer that for any $\beta \in \mathbf{R}$ it satisfies Assumption 2.6 with $(\mathcal{X}_0, \mathcal{X}_1) = (\mathcal{B}_\beta, \mathcal{B}_{\beta-1})$. Hence, the operator L generates a semigroup on the full range $(\mathcal{B}_\beta)_{\beta \in \mathbf{R}}$, and moreover it is easily checked that the non-linearity $N(u) := -\Delta(-u^3 + u)$ sends \mathcal{B}_0 to $\mathcal{B}_{-1/2}$ (this can be seen as a consequence of the fact that \mathcal{B}_0 forms an algebra for $kp > n$, which is inherited from the same property for the Bessel potential space $H^{k,p}(\mathcal{O})$).

Concerning the noise term, $B^H(\omega)$ is again lifted to a rough path as in the previous paragraph, and similarly as before $F_i(u) := f_i(x) \cdot \nabla u$ sends \mathcal{B}_β to $\mathcal{B}_{\beta-1/4}$, for any $\beta \in \mathbf{R}$. This means in particular that the assumptions of Theorem 2.15 are satisfied and hence we conclude existence and uniqueness of local solutions, for each $u_0 \in \mathcal{B}_0$.

6.3 Parabolic equations with transport noise

As in Example 2.12, assume that we are given a second order operator $L_t = \nabla \cdot (a(t)\nabla)$ satisfying the uniform ellipticity condition (2.19) and assume in addition that $a(t)$ is symmetric for all t . Recall that L_t satisfies Assumption 2.6 with the scale of spaces defined as $\mathcal{B}_\beta := H^{2\beta}(\mathbf{R}^n)$. Consider the parabolic equation with transport noise

$$du_t(x) = L_t u_t(x) dt + \sum_{i=1}^d f_i(x) \cdot \nabla u_t(x) d\mathbf{X}_t^i, \quad u_0 \in L^2(\mathbf{R}^n), \quad (6.6)$$

where $X = (X^1, \dots, X^d) \in \mathcal{C}^\gamma([0, T]; \mathbf{R}^d)$ is a *geometric* rough path with Hölder exponent $\gamma \in (\frac{1}{3}, \frac{1}{2}]$, and $f = (f_1, \dots, f_d)$ is a collection of smooth vector fields on \mathbf{R}^d .

Notice that this equation is ‘supercritical’ and does not fall into the framework of Theorem 2.15. Indeed, note that $F(u) = (f_1 \cdot \nabla u, \dots, f_d \cdot \nabla u)$ sends \mathcal{B}_β into $\mathcal{B}_{\beta-\frac{1}{2}}^d$, thus violating the subcriticality requirement since $\gamma < \frac{1}{2}$. Nevertheless, this type of rough partial differential equation with transport rough input has been investigated in the literature [4, 20, 36, 37] (see also [35] for a similar quasilinear ansatz), relying on a priori estimates and techniques from the theory of transport equations as introduced by DiPerna and Lions in [21]. The approach uses the variational formulation instead of the mild formulation of the equation, i.e.

$$\langle \delta u_{t,s}, \phi \rangle = \int_s^t \langle u_r, L_r \phi \rangle dr - \sum_{i=1}^d \int_s^t \langle u_r, \operatorname{div}(f_i \phi) \rangle d\mathbf{X}_r^i,$$

for $\phi \in H^3(\mathbf{R}^n)$, meaning that the rough integral $\int_0^\cdot F(u_r) \cdot d\mathbf{X}_r$ has to be understood as a $H^{-3}(\mathbf{R}^n)$ -valued path. Above and below $\langle \cdot, \cdot \rangle$ denotes dual pairing of $H^{-k}(\mathbf{R}^n)$ and $H^k(\mathbf{R}^n)$.

We now show that, provided we have a variational solution to (6.6), this equation may be equivalently formulated using the propagators.

Lemma 6.1. *Let u be the solution to (6.6) as described in [36, Theorem 1]. Then u allows for the following representation in $H^{-3}(\mathbf{R}^n)$*

$$\delta^S u_{t,s} = \sum_{i=1}^d \int_s^t S_{t,r}(f_i \cdot \nabla u_r) d\mathbf{X}_r^i, \quad (6.7)$$

where S denotes the propagator associated to $(L_t)_{t \in [0, T]}$.

Proof. As in [4], we introduce the *unbounded rough drivers*:

$$A_{t,s}^1 \phi = \sum_{i=1}^d f_i \cdot \nabla \phi X_{t,s}^i \quad A_{t,s}^2 \phi = \sum_{i,j=1}^d f_i \cdot \nabla (f_j \cdot \nabla \phi) \mathbb{X}_{t,s}^{j,i},$$

for every $(s, t) \in \Delta_2$. Recall that the solution to (6.6) is defined as the path $u: [0, T] \rightarrow L^2(\mathbf{R}^n)$ such that the linear functional $u_{t,s}^\sharp$ defined for every $\phi \in H^3(\mathbf{R}^n)$ and $(s, t) \in \Delta_2$ by

$$\langle u_{t,s}^\sharp, \phi \rangle := \langle \delta u_{t,s}, \phi \rangle - \int_s^t \langle u_r, L_r \phi \rangle dr - \langle u_s, [A_{t,s}^{1,*} + A_{t,s}^{2,*}] \phi \rangle,$$

satisfies the estimate $|u_{t,s}^\sharp|_{-3/2} \lesssim |t-s|^\lambda$ for some $\lambda > 1$ (we recall that $|\cdot|_\alpha := |\cdot|_{H^{2\alpha}(\mathbf{R}^n)}$).

Notice that since $a(t)$ is symmetric, then L_t is self-adjoint and so is the propagator $S_{t,s}$. Fix $t > 0$, $\phi \in H^3(\mathbf{R}^n)$ and define $f_s = S_{t,s} \phi$ so that $\delta f_{t,s} = - \int_s^t L_r f_r dr$. Straightforward computations give

$$\begin{aligned} \delta \langle u, f \rangle_{t,s} &= \langle \delta u_{t,s}, f_s \rangle + \langle u_t, \delta f_{t,s} \rangle \\ &= \int_s^t \langle u_r, L_r f_s \rangle dr + \langle u_s, A_{t,s}^{1,*} f_s + A_{t,s}^{2,*} f_s \rangle + \langle u_{t,s}^\sharp, f_s \rangle - \int_s^t \langle u_t, L_r f_r \rangle dr \\ &= \langle u_s, A_{t,s}^{1,*} f_s + A_{t,s}^{2,*} f_s \rangle + u(f)_{t,s}^\sharp, \end{aligned} \quad (6.8)$$

where we have defined

$$u(f)_{t,s}^\sharp = \int_s^t (\langle u_r, L_r f_s \rangle - \langle u_t, L_r f_r \rangle) dr + \langle u_{t,s}^\sharp, f_s \rangle.$$

For the second term above we use the definition of u^\natural and the continuity of $S_{t,s}$ on $H^3(\mathbf{R}^n)$ to get

$$|\langle u_{t,s}^\natural, f_s \rangle| \lesssim |t-s|^\lambda |S_{t,s}\phi|_{3/2} \lesssim |t-s|^\lambda |\phi|_{3/2}.$$

For the first term, we write

$$\begin{aligned} \left| \int_s^t (\langle u_r, L_r f_s \rangle - \langle u_t, L_r f_r \rangle) dr \right| &\leq \int_s^t |\langle \delta u_{t,r}, L_r f_r \rangle| dr + \int_s^t |\langle u_r, L_r \delta f_{r,s} \rangle| dr \\ &\leq |t-s|^{\lambda-1} |u|_{\lambda-1, -1/2} \int_s^t |L_r f_r|_{1/2} dr + |u|_{0,0} \int_s^t |L_r \delta f_{r,s}|_0 dr \\ &\lesssim |t-s|^\lambda |u|_{\lambda-1, -1/2} |\phi|_{3/2} + |u|_{0,0} |t-s|^{3/2} |\phi|_{3/2}, \end{aligned}$$

where we have used

$$|L_r f_r|_{1/2} \lesssim |\phi|_{3/2} \quad \text{and} \quad |\delta f_{r,s}|_1 = |S_{t,r}(\text{id} - S_{r,s})\phi|_1 \lesssim |r-s|^{1/2} |\phi|_{3/2}.$$

Now note that $f_t = \phi$, thus $\delta \langle u, f \rangle_{t,s} = \langle u_t, \phi \rangle - \langle u_s, S_{t,s}\phi \rangle = \langle \delta^S u_{t,s}, \phi \rangle$. Using the fact that S is self-adjoint and rewriting (6.8) we get the following equality on $H^{-3}(\mathbf{R}^n)$

$$\delta^S u_{t,s} = S_{t,s} A_{t,s}^1 u_s + S_{t,s} A_{t,s}^2 u_s + u_{t,s}^{S,\natural},$$

where $u_{t,s}^{S,\natural}$ is defined on $H^3(\mathbf{R}^n)$ as

$$u_{t,s}^{S,\natural}(\phi) = \int_s^t (\langle u_r, L_r S_{t,s}\phi \rangle - \langle u_t, L_r S_{t,r}\phi \rangle) dr + \langle u_{t,s}^\natural, S_{t,s}\phi \rangle.$$

By the uniqueness part of Theorem 4.1 this is exactly (6.7) as constructed in Theorem 4.5. \blacksquare

References

- [1] H. Amann. Existence and regularity for semilinear parabolic evolution equations. *Annali della Scuola Normale Superiore di Pisa-Classe di Scienze*, **11**(4): 593–676, 1984.
- [2] H. Amann. Quasilinear evolution equations and parabolic systems. *Transactions of the American Mathematical Society*, **293**(1): 191–227, 1986.
- [3] W. Arendt, C.J. Batty, M. Hieber and F. Neubrander. *Vector-valued Laplace Transforms and Cauchy Problems* (2nd edition). Monographs in Mathematics Vol. **96**, Birkhäuser, 2011.
- [4] I. Bailleul and M. Gubinelli. Unbounded rough drivers. *Annales Mathématiques de la Faculté des Sciences de Toulouse*, **26**(4): 795–830, 2017.
- [5] I. Bailleul and S. Riedel. Rough flows. *Journal of the Mathematical Society of Japan*, 2019.
- [6] A. Brault and A. Lejay. The non-linear sewing lemma I: weak formulation. *Electronic Journal of Probability*, **24**, 2019.
- [7] A. Brault and A. Lejay. The non-linear sewing lemma II: Lipschitz continuous formulation. Preprint arXiv:1810.11988, 2018.

- [8] A. Brault and A. Lejay. The non-linear sewing lemma III: Stability and generic properties. *Forum Mathematicum*, **1**. No. ahead-of-print. De Gruyter, 2020.
- [9] C.E. Bréhier, and A. Debussche. Kolmogorov equations and weak order analysis for SPDEs with nonlinear diffusion coefficient. *Journal de Mathématiques Pures et Appliquées*, **119**: 193–254, 2018.
- [10] Z. Brzezniak and M. Veraar. Is the stochastic parabolicity condition dependent on p and q ?. *Electronic Journal of Probability*, **17**, 2012.
- [11] A. Calderón. Intermediate spaces and interpolation, the complex method. *Studia Mathematica*, **24**(2): 113–190, 1964.
- [12] M. Caruana, P.K. Friz, and H. Oberhauser. A (rough) pathwise approach to a class of non-linear stochastic partial differential equations. *Annales de l’Institut Henri Poincaré (C) Non Linear Analysis*, **28**: 27–46, 2011.
- [13] C. Choi and J.M. Kim. Locally convex vector topologies on $b(x,y)$. *Journal of the Korean Mathematical Society*, **45**(6): 1677–1703, 2008.
- [14] L. Coutin and A. Lejay. Perturbed linear rough differential equations [Équations différentielles linéaires rugueuses perturbées]. In *Annales mathématiques Blaise Pascal*, **21**:103–150, 2014.
- [15] A. Cialdea and V. Mazya. Criteria for the L^p -dissipativity of systems of second order differential equations. *Ricerche di matematica*, **55**(2): 73–105, 2006.
- [16] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*. Cambridge University Press, 2008.
- [17] A. Deya. A discrete approach to Rough Parabolic Equations. *Electronic Journal of Probability*, **16**: 1489–1518, 2011.
- [18] A. Deya. Numerical schemes for rough parabolic equations. *Applied Mathematics & Optimization*, **65**(2): 253–292, 2012.
- [19] A. Deya, M. Gubinelli and S. Tindel. Non-linear rough heat equations. *Probability Theory and Related Fields*, **153**(1-2): 97–147, 2012.
- [20] A. Deya, M. Gubinelli, M. Hofmanová, and S. Tindel. A priori estimates for rough PDEs with application to rough conservation laws. To appear in *Journal of Functional Analysis* (Arxiv preprint available at arXiv:1604.00437).
- [21] R.J. DiPerna and P.L. Lions. Ordinary differential equations, transport theory and Sobolev spaces. *Inventiones mathematicae*, **98**(3): 511–547, 1989.
- [22] D. Feyel, A. de La Pradelle and G. Mokobodzki. A non-commutative sewing lemma. *Electronic Communications in Probability*, **13**: 24–34, 2008.
- [23] P.K. Friz, A. Hocquet and K. Lê. Rough Markov diffusions and stochastic differential equations. In progress.
- [24] P.K. Friz and H. Oberhauser. Rough path stability of (semi-) linear SPDEs. *Probability Theory and Related Fields*, **158**(1-2):401–434, 2014.

- [25] P.K. Friz and M. Hairer. *A course on rough paths: with an introduction to regularity structures*. Springer, 2014.
- [26] P.K. Friz and N.B. Victoir. *Multidimensional stochastic processes as rough paths: theory and applications*, volume **120**. Cambridge University Press, 2010.
- [27] A. Gerasimovičs and M. Hairer. Hörmander’s theorem for semilinear SPDEs. *Electronic Journal of Probability*, volume **24**, 2019.
- [28] J.A. Goldstein. *Semigroups of linear operators and applications*. Oxford University Press, USA, 1985.
- [29] M. Gubinelli. Controlling rough paths. *Journal of Functional Analysis*, **216**: 86–140, 2004.
- [30] M. Gubinelli. Ramification of rough paths. *Journal of Functional Analysis*, **248**, 4: 693–721, 2010.
- [31] M. Gubinelli and S. Tindel. Rough evolution equations. *The Annals of Probability*, **38**(1): 1–75, 2010.
- [32] M. Gubinelli, A. Lejay and S. Tindel. Young Integrals and SPDEs. *Potential Analysis*, **25**, 307–326 (2006).
- [33] M. Hairer. An introduction to stochastic PDEs. *arXiv preprint*, 2009.
- [34] M. Hairer and J.C. Mattingly. A theory of hypoellipticity and unique ergodicity for semilinear stochastic PDEs. *Electronic Journal of Probability*, **16**(23): 658–738, 2011.
- [35] A. Hocquet. Quasilinear rough partial differential equations with transport noise. Preprint arXiv:1808.09867 (2018).
- [36] A. Hocquet, M. Hofmanová. An energy method for rough partial differential equations. *Journal of Differential Equations*, **265**(4): 1407–1466, 2018.
- [37] A. Hocquet and T. Nilssen. An Itô Formula for rough partial differential equations and some applications. *Potential Analysis*: 1–56, 2020.
- [38] M. Hofmanová, J-M. Leahy and T. Nilssen. On the Navier–Stokes equation perturbed by rough transport noise. *Journal of Evolution Equations* **19**(1): 203–247, 2019.
- [39] M. Hofmanová, J-M. Leahy and T. Nilssen. On a rough perturbation of the Navier–Stokes system and its vorticity formulation. *Annals of Applied Probability*. To appear.
- [40] T. Kato. Integration of the equation of evolution in a Banach space. *Journal of the Mathematical Society of Japan*, **5**(2): 208–234, 1953.
- [41] N.V. Krylov, B.L. Rozovskii. Stochastic evolution equations. *Journal of Soviet Mathematics* **16**(4): 1233–1277, 1981.
- [42] C. Kuehn and A. Neamțu. Pathwise mild solutions for quasilinear stochastic partial differential equations. *Journal of Differential Equations*, 2020.
- [43] K. Lê. A stochastic sewing lemma and applications. *Electronic Journal of Probability* **25**, 2020.

- [44] J.A. León. On equivalence of solution to stochastic differential equation with anticipating evolution system. *Stochastic analysis and applications*, **8**(3): 363–387, 1990.
- [45] J.A. León and D. Nualart. Stochastic evolution equations with random generators. *The Annals of Probability*, **26**(1): 149–186, 1998.
- [46] P.L. Lions and P.E. Souganidis. Fully nonlinear stochastic partial differential equations. *Comptes Rendus de l'Académie des Sciences-Series I-Mathematics*, **326**(9): 1085–1092, 1998.
- [47] P.L. Lions and P.E. Souganidis. Uniqueness of weak solutions of fully nonlinear stochastic partial differential equations. *Comptes Rendus de l'Académie des Sciences-Series I-Mathematics*, **331**(10): 783–790, 2000.
- [48] G. Lumer and R.S. Phillips. Dissipative operators in a Banach space. *Pacific Journal of Mathematics*, **11**(2): 679–698, 1961.
- [49] A. Lunardi. *Interpolation theory*, volume 16. Springer, 2018.
- [50] T.J. Lyons. Differential equations driven by rough signals. *Revista Matemática Iberoamericana*, **14**(2): 215–310, 1998.
- [51] E. Pardoux. Stochastic partial differential equations and filtering of diffusion processes. *Stochastics*. **3**(1-4):127–167, 1980.
- [52] A. Pazy. *Semigroups of linear operators and applications to partial differential equations*. Springer-Verlag New York, 1983.
- [53] M. Reed and B. Simon. *Methods of modern mathematical physics, Vol. III: Scattering theory*. New York, San Francisco, London, 1979.
- [54] F. Russo and P. Vallois. Forward, backward and symmetric stochastic integration. *Probability theory and related fields*, **97**(3):403–421, 1993.
- [55] P.E. Sobolevskii. Equations of parabolic type in a Banach space. *American Mathematical Society Translations*, **49**:1–62, 1966.
- [56] G. Strang. On the construction and comparison of difference schemes. *SIAM Journal on Numerical Analysis*, **5**(3):506–517, 1968.
- [57] H. Tanabe. On the equations of evolution in a Banach space. *Osaka Mathematical Journal*, **12**(2):363–376, 1960.
- [58] H. Triebel. *Theory of Function Spaces*. Springer Science & Business Media, 1983.
- [59] R. Hesse, and A. Neamțu. Local mild solutions for rough stochastic partial differential equations *Journal of Differential Equations*, 2019