Citation for published version:

Weiss, MC 2011, 'Diagnostic decision making: the last refuge for general practitioners?', *Social Science and Medicine*, vol. 73, no. 3, pp. 375-382. <https://doi.org/10.1016/j.socscimed.2011.05.038>

DOI:

[10.1016/j.socscimed.2011.05.038](https://doi.org/10.1016/j.socscimed.2011.05.038)

Publication date:

2011

Document Version

Peer reviewed version

[Link to publication](https://doi.org/10.1016/j.socscimed.2011.05.038)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Title: Diagnostic Decision Making: The Last Refuge for General Practitioners?

Authors:

Marjorie Cecilia Weiss, University of Bath, United Kingdom*

*Corresponding Author's Contact Details: prsmw@bath.ac.uk

Keywords: UK, Clinical decision-making; pharmacist prescribing; nurse prescribing; general practice; uncertainty; professional judgement

Acknowledgements:

The author is grateful for the help and support from Miss Hannah Dawson in developing the argument in this paper. The author also gratefully acknowledges the assistance of Dr Jane Sutton and the four anonymous reviewers for their useful comments in refining and clarifying the themes presented here.

Abstract

Nurses and pharmacists gained the right to prescribe as independent prescribers in the UK in 2007. Independent prescribers are responsible for the initial assessment of patients with diagnosed and undiagnosed conditions. Public policy discourse and the views expressed by health care professionals have conceptualised diagnostic decision making as being at a 'higher level' and more difficult than prescribing decision making. This paper presents five themes related to this premise. Firstly diagnostic decision making is put into the broader context of clinical reasoning which underpins all types of clinical decisions including both diagnostic and prescribing decisions. Secondly, the nature of diagnostic decision making is discussed as to whether it is indeed separable from the prescribing decision making process. Thirdly, the conception that all diagnostic decisions are inherently more difficult is contested when difficulty in decision making is more appropriately applied to all types of clinical decisions which involve greater complexity and uncertainty. The fourth topic concerns whether this perception of diagnoses as being more difficult is a response by the medical profession to the threat of independent prescribing, reflecting their wish to maintain professional power and dominance over other professions. The final section considers how expertise in diagnoses could be developed in nurse and pharmacist independent prescribers. To develop their expertise in making accurate diagnoses, medicine uses the model of learning basic science mechanisms followed by engagement with patient clinical problems followed by years of clinical experience. However this may be just one way of achieving

1 diagnostic expertise. Other approaches, such as the use of deliberate
2 practice and feedback, may be more suitable to the diverse range of
3
4 knowledge and experiences of nurse and pharmacist prescribers.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

In the UK, appropriately trained nurses and pharmacists have been able to legally sign prescriptions for prescription-only medicines since 2004. This followed from the Crown Report in 1999 which recommended that the 'legal authority to prescribe should extend beyond currently authorised prescribers' (Department of Health, 1999, pp. 36). Legislative changes in 2003 led to the extension of prescribing authority to nurses and pharmacists with the introduction of supplementary prescribing in 2004 (Department of Health, 2005) and with further legal changes in 2006, to allow nurse and pharmacist independent prescribing in 2007 (Department of Health, 2006). Previously both nurses and pharmacists had been able to advise medical practitioners on medicine use and, particularly in community pharmacies, to provide advice on over-the-counter medicines. However these changes were a major shift in policy which enabled health care professionals, other than a doctor or dentist, to have legal authority to write prescriptions for the full range of prescription-only medicines including, for nurses, controlled drugs. The aims of these developments were to give patients quicker access to medicines, to decrease general practitioners' workload and to better use the skills of nurses and pharmacists (Department of Health, 2005).

Supplementary prescribing is described as 'a voluntary partnership between the responsible independent prescriber (a doctor or a dentist) and a supplementary prescriber (nurse or pharmacist) to implement an agreed patient specific clinical management plan with the patient's

1 agreement' (Department of Health, 2005, pp.11). The clinical
2 management plan details, amongst other issues, the types of medicines to
3 be prescribed, limitations or restrictions on prescribing, relevant warnings
4 and the circumstances under which the supplementary prescriber should
5 refer or seek advice from the independent prescriber. In 2006,
6 independent prescribing (IP) by nurses and pharmacists was introduced
7 with the independent prescriber defined as the 'practitioner responsible
8 and accountable for the assessment of patients with undiagnosed or
9 diagnosed conditions and for decisions about the clinical management
10 required, including prescribing' (Department of Health, 2006, pp.2). As
11 independent prescribers, nurses and pharmacists are no longer required
12 to use clinical management plans, although many in practice will use
13 condition-specific protocols or guidelines to inform, and limit, the range of
14 medicines they feel competent to prescribe.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 Anxieties raised by doctors to these new roles have centred on concerns
36 about patient safety and that nurses and pharmacists lack training in
37 diagnosing (British Medical Association, 2005; Day 2005). More moderate
38 medical voices have expressed a need for appropriate training, support
39 and governance for new prescribers but that it could benefit patients
40 (Avery & Pringle, 2005). Other authors expressed fears over new
41 prescribers taking on a diagnostic role and that patients could be
42 endangered by 'the reckless expansion of nurse and pharmacist
43 prescribing' (Anonymous, 2006). Making diagnoses has been called the
44 most important responsibility carried out by medical doctors (Gutkin,
45 2009) and that diagnosis is almost the only skill that still defines them
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

(Godlee, 2008). On the other hand, the changes in the types of activities traditionally carried out by doctors and nurses have been noted by Ghislaine Young, a nurse practitioner and salaried partner in a GP practice (Young, 2005). She has argued that the distinction between medicine (diagnosis and cure) and nursing (care) has become increasingly blurred and, further, that professional groups should work together to enhance the care of patients.

Research exploring the views of medical doctors about nurse and pharmacist prescribing have found that, although they were generally positive about the role of new prescribers, some expressed concern about nurses' and pharmacists' ability to diagnose (Child & Cantrill, 1999, Bissell et al., 2008, Stewart et al., 2009, Weiss et al., 2006). Doctors acting as mentors for pharmacist prescribers expressed concern about boundary encroachment and medical deskilling. They preferred the model of supplementary prescribing where they could retain ultimate control over the process by setting barriers within which pharmacists could prescribe (Lloyd & Hughes, 2007). Similar views were expressed by doctors working in secondary care who wanted to set limits to nurse and pharmacist prescribing by only allowing them to work within protocols (Buckley et al. 2006). In Bissell's (2008) study, diagnosing by doctors was considered to be a 'more skilled, uncertain and difficult task' with prescribing seen as a lesser task than 'de novo' diagnosis (Bissell et al., 2008, pp. 57). Evidence from the evaluation of nurse and pharmacist supplementary prescribing suggested that prescribing was considered by doctors to have diminished as a skilled practice and, with the widespread use of protocols, much of it

1 was considered routine. These doctors saw their role, with its superior
2 knowledge, training and experience, to be reserved for diagnosis and *plus*
3 *ultra* prescribing (Cooper et al., 2011).
4
5
6
7
8

9 Views of nurse and pharmacist prescribers in previous research have
10 echoed similar reservations about diagnosis. In an evaluation of
11 pharmacist supplementary prescribing, while many new prescribers were
12 keen to become independent prescribers, the reason for this was not so
13 they could diagnose, but rather so they could avoid the need for individual
14 patient clinical management plans. Some of these new prescribers had no
15 desire to take on the diagnosis of new conditions and felt unqualified to do
16 so (Lloyd et al., 2010, Weiss et al., 2006). In Luker's early evaluation of
17 nurse prescribing, nurses expressed fear about making an incorrect
18 diagnosis, although this research took place before the introduction of
19 supplementary prescribing and was during the time when nurses were
20 able to prescribe from a limited formulary (Luker et al., 1997).
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Supplementary prescribing may have been more acceptable to doctors
given the framework in which it was conducted with patient-specific
clinical management plans. This arrangement allows doctors to retain
control over the process (hence the earlier word used for supplementary
prescriber was a 'dependent' prescriber) and circumscribed the area of
clinical practice for the new prescriber (Lloyd & Hughes, 2007). The word
supplementary implies something both sub-ordinate and non-essential. In
this sense both context and terminology are important. In policy
documents the distinction between supplementary prescribing and

1 independent prescribing created an explicit hierarchy between diagnostic
2 decision making and prescribing decision making, privileging the
3
4 superiority of diagnosis as a cognitive decision making process. The
5
6 responses of health care professionals noted above reinforce this
7
8 distinction and suggest that such an assumption may be pervasive. There
9
10 are several issues, related to this concern about diagnostic decision
11
12 making, that this paper seeks to address: (1) to provide some background
13
14 into what is meant by diagnostic decision making and how this relates to
15
16 broader issues of clinical reasoning, (2) to consider whether diagnostic
17
18 decision making is truly able to be regarded as a separate process distinct
19
20 from prescribing, and other, patient management decision making, (3) to
21
22 suggest that it may be more appropriate to focus on 'difficult' clinical
23
24 decisions, which includes diagnostic as well as other patient management
25
26 decisions, rather than diagnostic decisions per sé, (4) to consider the
27
28 threat to the medical profession posed by nurse and pharmacist
29
30 independent prescribers who may wish to diagnose and, finally (5) to
31
32 explore educational and cognitive psychology perspectives on the
33
34 development of diagnostic expertise amongst nurse and pharmacist
35
36 prescribers.
37
38
39
40
41
42
43
44
45
46

47 While prescribing by allied health care professionals has also been
48
49 implemented, this paper will explicitly take the perspective of nurse and
50
51 pharmacist prescribing in a UK primary care setting.
52
53
54
55

56 Diagnostic Decision Making
57
58
59
60
61
62
63
64
65

1 Elstein and Schwartz describe diagnosis as a process of generating one or
2 more hypotheses, using these to predict what additional findings should
3
4 be present and undertaking further data collection to test out these
5
6 hypotheses (Elstein & Schwartz, 2002). This hypothetico-deductive
7
8 approach generates hypotheses early during the initial presentation of a
9
10 problem and draws upon existing knowledge, associations and experience
11
12 of the medical doctor (Round, 2000). This process of generating
13
14 hypotheses and drawing upon existing knowledge, associations and
15
16 experience is part of the process of clinical reasoning (Norman et al.,
17
18 2009). Clinical reasoning also informs other kinds of clinical decisions
19
20 including decisions about management such as treatment or referral
21
22 decisions as well as the interpretation, and weight given to, clinical
23
24 research evidence. Experience and knowledge both inform clinical
25
26 reasoning and are influenced by previous applications of it. For the
27
28 purposes of this paper, clinical reasoning will be used as the broader term
29
30 and the underlying process informing all types of clinical decisions
31
32 including diagnostic decisions, as well as prescribing decisions, referral
33
34 decisions or other decisions about management. Clinical reasoning is
35
36 usually seen as encompassing both analytic and non-analytic processes,
37
38 both of which will be reviewed here.
39
40
41
42
43
44
45
46
47
48

49 In the analytic model, medical work is seen as a logical, step by step
50
51 cognitive process such that when a doctor is confronted with a particular
52
53 patient situation, details of the patient problem are compared with
54
55 professional knowledge. In this model the key elements are the
56
57 professional's knowledge, cognitive capacities and limitations and the data
58
59
60
61
62
63
64
65

required for diagnosis and intervention (Berg, 1997). This analytic approach characterises the decision as a two-stage process. The first stage consists of collecting and analysing the evidence with regard to benefits, harms and costs of each potential outcome option. The second stage consists of comparing the desirability of the different outcomes of each option (Eddy, 1990). In the analytic model, these processes are performed explicitly or implicitly, may be subject to imperfect clinical information and (if so) are likely to require subjective professional judgement. Analytic models include those which use Bayes' Theorem, where a priori probabilities associated with the known prevalence of a diagnosis and the conditional probabilities associated with each sign or symptom with each diagnosis are used to calculate the probability of each diagnosis under consideration (Elstein & Schwartz, 2002; Eva, 2004). This approach assumes that causal rules linking clinical features to diagnoses can be extracted from clinical practice and that, with experience, these rules can become more refined and attuned to reality (Eva, 2004).

The non-analytic model, which comprises pattern recognition or direct automatic retrieval models, compares a current patient situation to past patient cases (or abstractions of such) to make a judgement as to the probability that a particular case belongs to a specific diagnostic category. A new case is categorised by its resemblance to memories of instances previously seen (Brooks et al., 1991) or to a more abstract prototype (Elstein & Schwartz, 2002). While usually termed a non-analytic model, a more accurate description may be that such reasoning is rapid and unconscious/subconscious, but nonetheless still analytic. Some authors

1 suggest that the ability to use pattern recognition increases with
2 experience and expertise, as novice decision makers have no experience
3 to draw upon (Luker et al., 1998) but rather base their decisions on
4 taught procedures and guidelines (Benner, 1982). Novices think
5 analytically through guiding principles whereas experts can make
6 decisions 'intuitively' through rapid, unconscious or subconscious analytic
7 processes. However other research has shown no qualitative difference
8 between the reasoning strategies used by junior medical students
9 compared with experienced doctors (Neufeld et al., 1981).

10
11
12
13
14
15
16
17
18
19
20
21
22
23 The interplay between age (or clinical experience) and clinical reasoning is
24 complex. A systematic review of the relationship between clinical
25 experience and quality of care found that increasing experience resulted in
26 a decline in performance as measured by physician clinical knowledge,
27 adherence to guidelines and in some patient outcomes (Choudhry, et al.,
28 2005). Other evidence suggests that increasing age may result in
29 increased non-analytic thinking without a loss of diagnostic accuracy (Eva
30 et al., 2010; Groves et al., 2003). More experienced doctors may be more
31 likely to consider less common or less stereotypical conditions and more
32 able to integrate complex, social and behavioural information (Feltovich
33 1981; Elstad et al., 2010). However age can also result in reliance on data
34 gathered early in the consultation and less willingness to re-assess a
35 diagnostic decision when presented with new information (Eva &
36 Cunningham, 2006).

1 Recently Norman et al. (2009) have developed the idea of iterative
2 diagnosis which draws upon elements from both the analytic and non-
3
4 analytic models of reasoning. Drawing upon the original hypo-thetico-
5
6 deductive model by Elstein and Schwartz (2002), the process involves the
7
8 clinician generating hypotheses (often impressionistically using non-
9
10 analytic reasoning). These are then iteratively tested through additional
11
12 patient questioning for their alignment with the clinician's clinical
13
14 knowledge and experience using analytical reasoning. According to
15
16 Norman et al. (2009) analytic and non-analytic reasoning usually
17
18 integrate smoothly and unconsciously throughout the decision making
19
20 process and subsequent consultation interaction.
21
22
23
24
25
26
27

28 The literature on clinical reasoning recognises that human beings are
29
30 subject to cognitive biases or errors through the use of heuristics or
31
32 cognitive shortcuts which try to make the complex more simple (Hall,
33
34 2002). Heuristics can be seen as a highly adaptive and appropriate
35
36 response to complex decision making in the real world. However heuristics
37
38 are more often viewed negatively and seen as the process through which
39
40 clinical evidence becomes biased or weighted inappropriately. Authors
41
42 acknowledge that heuristics are most commonly employed when making
43
44 judgements under uncertainty (Tversky & Kahneman, 1974, Hall 2002).
45
46 These include the availability bias where the probability of an event (or
47
48 diagnosis) occurring is rated more likely if it can be related to a case
49
50 easily recalled. There is the confirmation bias which directs the line of
51
52 questioning towards confirming the diagnosis under consideration instead
53
54 of investigating evidence which may refute it (Norman et al., 2009). A
55
56
57
58
59
60
61
62
63
64
65

1 third example is the framing bias where patients are swayed to give
2 answers to questions which support a diagnosis because of the way in
3 which a particular question has been asked. The consequence of these
4 biases can be an inaccurate diagnosis due to premature closure of
5 questioning without the critical data having been collected (Norman et al.,
6 2009).

7
8
9
10
11
12
13
14
15
16 Hamm (1988) views analytic and non-analytic clinical reasoning as being
17 on a continuum while more recent experts suggest that the optimal form
18 of reasoning is an additive model in which both analytic and non-analytic
19 processes play a role (Eva, 2004). However, as noted by Eva, accuracy of
20 diagnoses relies not only on being able to draw upon a range of analytic
21 and non-analytic strategies, but also the context within which a problem is
22 encountered. Context in this sense includes the clinical setting, recent
23 cases and personal factors such as clinician experience and the current
24 state of medical opinion (Eva, 2004). The influence of social and
25 psychological factors (e.g. patient recently bereaved, consultation before
26 a holiday weekend) on doctors' decision making has been widely
27 recognised (Howie, 1976; Katz, 1985), as has the importance of
28 considering how social, psychological and biological systems influence
29 diagnostic certainty (Lutfey & McKinlay, 2009).

30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51 Having considered some of the key features of the clinical reasoning
52 process, I will now consider the intertwined nature of how diagnostic and
53 prescribing decisions are made.

Diagnostic and Prescribing Decision Making

Howie (1972) was one of the first investigators to hypothesise that a treatment decision might be derived directly from signs, symptoms and investigations without the intervening stage of a diagnostic decision. Conventional wisdom suggested a sequential process involving a diagnostic decision followed by a treatment decision but Howie's radical proposition was that a diagnostic label was less a reason for treatment but rather a justification for it (Howie, 1972). Howie found that diagnostic labels had a poorer predictive value for antibiotic treatment than the original symptoms-sign complex. In a follow-up study using simulated patients, Howie found that doctors required less information to make a decision on management than to make a diagnostic decision (Howie, 1974). He notes that a decision to prescribe an antibiotic relates less to a decision on diagnosis but rather to a decision not to prescribe an antibiotic. Indeed, non-analytic reasoning may play a part in such processes where diagnoses are generated impressionistically (and potentially unconsciously) and are used to inform clinical management decisions. However these findings also suggest that there is not a separate process of clinical reasoning used to inform diagnostic decisions which is distinct from those processes informing the decision to prescribe. Privileging the aspect of clinical reasoning which informs diagnostic decisions, as noted in the comments by the medical profession described earlier, may not be appropriate given that these same processes inform decisions to prescribe.

1 Further evidence of the complex relationship between diagnostic and
2 treatment decision making comes from Bloor's study of ENT specialists
3
4 (Bloor, 1976; Bloor, 1978). He found that ENT specialists', when assessing
5
6 children's suitability for possible adeno-tonsillectomy surgery, did not
7
8 make decisions by considering and weighing up all the available
9
10 information. Rather the specialists used a series of specifically situated
11
12 routines which were used 'to construct images of the clinical signs,
13
14 symptoms and circumstances of each patient' (Bloor, 1976, pp.45). These
15
16 routines, which varied widely between each specialist, were the means by
17
18 which they rendered 'unproblematic their decisions on patient disposal'
19
20 (Bloor, 1976, pp.45). For example, the specialists differed in what signs
21
22 were considered important, how important these signs were weighted,
23
24 how specific and extensive their search procedures were and the decision
25
26 rules used to denote whether or not surgery was indicated. Other general
27
28 properties of decision making noted by Bloor were its repetitive, routinised
29
30 and idiosyncratic nature, and its orientation towards action (Bloor, 1978).
31
32 Evidence-based medicine can be seen as a development which sought to
33
34 address this variability in specialists' use of routines. In addition these
35
36 findings suggest there may be considerable overlap in the processes of
37
38 clinical reasoning used to make diagnostic and management decisions.
39
40
41
42
43
44
45
46
47
48

49 The intertwined nature of diagnosis and treatment decisions was also
50
51 found by Berg (1992) in his ethnographic study of medical action in
52
53 practice. He noted that the phased two-step motion of searching for a
54
55 diagnosis and then deciding upon treatment did not hold. The process of
56
57
58
59
60
61
62
63
64
65

transforming the presenting patient problem into a solvable patient
problem is:

'uni-directionally geared towards the construction of disposal.

'Diagnosis' and 'therapy' are terms which can be applied to this
process in retrospect, but in an 'in situ' study of medical practice the
usage of these terms creates an artefactual distinction' (pp.169).

Berg describes how medical disposals are socially constructed such that
the type of questions asked, the way they are asked and the doctor's
interpretation of the patient's answers shape the symptoms, historical and
examination patient data towards a solvable problem. Patient data is
recast and reconstructed into a pattern which aligns with the considered
transformation. Some patient data may be ignored, devalued or
emphasised in the course of medical interaction in the quest towards
constructing a medical disposal (Berg, 1992). This may be because
doctors have generated diagnostic hypotheses through non-analytic
reasoning and these are (unconsciously) guiding their selective use, and
emphasis on, patient data. Atkinson (1995) similarly noted the social and
interactional nature of medical decision making where decisions may be
subject to debate, negotiation and revision. He considers the sequential
ordering of the decision making process from the gathering of patient
information to diagnosis to treatment to be troublesome. Even 'objective'
information such as laboratory test findings are the outcome of a process
of decision making and are themselves judgements mediated by the
interpretations and values attached to them in collegial discourse

(Atkinson, 1995). This body of evidence underscores the dynamic nature of medical decision making, the constant revision of diagnostic and management decisions in the light of new information, and the falsity in trying to ring fence the reasoning process involved in making diagnostic decisions as separate from decisions about prescribing or management. This suggests that doctors when making decisions which may appear overtly management oriented may have made a diagnostic decision using rapid non-analytic reasoning to underlie that management decision. Equally, it may be that nurse or pharmacist prescribers when acting in their role, even as a supplementary prescriber, to make a decision about clinical management, may have done the same.

Not all Diagnoses are Created Equal

The previous section explored some of the evidence around the dynamics of the clinical reasoning process and how diagnostic decisions are deeply intertwined with decisions about management. This section will explore a range of diagnostic decisions and will argue that it is not, as noted in the introduction, that diagnostic decisions are intrinsically more difficult than prescribing decisions. Rather there are features, involving complexity and uncertainty, which are associated with some clinical decisions (e.g. both diagnostic and management) that makes them more difficult or challenging for the practitioner to make.

Clinical decisions occur in many contexts and are undertaken by both health care professionals and lay people. Self care, in terms of managing

1 minor ailments and long term conditions, is high on the NHS agenda
2 (Nazareth & Murray, 2010). Whether clinical reasoning is viewed within an
3
4 analytic or non-analytic framework, the process where an individual
5
6 compares their signs and symptoms with their medical (or lay) knowledge
7
8 of the condition, can be viewed as engaging in a process of diagnostic
9
10 decision making. Bloor and Horrobin (1975) note that patients prior to
11
12 coming to see a doctor are expected to assess their own symptoms and
13
14 seek appropriate self-care, and yet become passive and deferential once
15
16 they enter the surgery. This, the authors suggest, places patients here in
17
18 unique 'double bind' situation. Patients need to self diagnose their
19
20 symptoms and decide, using their personal knowledge, experience and lay
21
22 referral networks, upon an appropriate course of action. This might
23
24 include self care or a visit to the GP. If the patient decides to seek help
25
26 from the GP, they are then expected to defer to the clinical reasoning and
27
28 course of action decided by the GP.
29
30
31
32
33
34
35
36

37
38 Community pharmacists undertake a similar cognitive diagnostic process
39
40 to patients when they provide advice and recommend over-the-counter
41
42 medicines in response to patient requests for advice about particular
43
44 symptoms (Tully et al., 1997). Community pharmacists have a well
45
46 described role in managing minor ailments and responding to patient
47
48 symptoms (Hassell et al., 2000). Evidence from cognitive psychology
49
50 suggests that where decisions are routine and repeated often, as in the
51
52 case with customers consulting pharmacists about common situations
53
54 needing minor ailment advice, they are easier as they require less effort
55
56 and little deliberation (Lehto & Nah, 2006). The cognitive load or burden
57
58
59
60
61
62
63
64
65

1 associated with a decision is also less when the presenting information
2 has already been organised into schemata and stored in long term
3 memory (van Merriënboër & Sweller, 2005). Such schemata reduce the
4 burden on working memory which is both capacity and time limited. In
5 clinical practice the storage of complex schemata is likely to be aided by
6 previous exposure to protocols, guidelines or flow diagrams which exist for
7 many clinical conditions and help organise incoming information. When a
8 customer requests advice on a common minor ailment, familiarity and
9 existing protocols and procedures relevant to the encounter are likely to
10 reduce the cognitive load on the pharmacist's decision making process.
11 Also potentially influencing the effort required is that decisions regarding
12 minor ailments involve an acute self-limiting condition, and thus involve
13 minimal uncertainty about the outcome (e.g. the patient is likely to get
14 better regardless).

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35 These 'easier' diagnostic decisions can also occur with some chronic
36 conditions such as hypertension. Current guidance from the British
37 Hypertension Society and the National Institute for Health and Clinical
38 Excellence states that a diagnosis of uncomplicated hypertension is made
39 when three readings above 140/90mm Hg are obtained on three separate
40 visits (National Institute for Health and Clinical Excellence, 2006). Lifestyle
41 or pharmacological interventions may be initiated, depending upon the
42 patient's level of cardiovascular risk. This, like the example above, relies
43 on an existing schemata (a hypertension guideline) which helps organise
44 presenting information and exerts less cognitive load on working memory
45 (van Merriënboër & Sweller, 2005). An increasing number of novel
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

elements to process increase the burden on working memory and makes decision making more difficult. Uncertainty is also one of the primary reasons why decisions can be difficult (Lehto & Nah, 2006). As described by Light (1979), there are different kinds of uncertainty: that arising from incomplete mastery of knowledge, that arising from indeterminacy of the clinical area itself and, as noted by Fox (1957), not being able to determine the difference between imperfect mastery of available knowledge and imperfections in the knowledge in the clinical area itself. Uncertainty can be exacerbated when there is complexity, for example when there is a lack of clarity in terms of patient presentation (e.g. atypical patient presentation or lack of symptom specificity to a particular illness) and / or increased complexity of the patient situation (e.g. the presence of co-morbidities or a complicated patient social situation). Increasing uncertainty and complexity require greater deliberation and impose a greater cognitive load on a practitioner's decision making. This may lead to the practitioner seeking additional information from the patient or to use the cognitive biases or heuristics described earlier (Round, 2000; Tversky & Kahneman, 1974). Intuitive decisions made under conditions of uncertainty are prone to cognitive biases and may lead to error (Norman, et al., 2009; Hall, 2002).

A Threat to Medical Dominance?

The arguments put forward so far have suggested that the process of making diagnostic and treatment decisions are intertwined within the process of clinical reasoning and are not separable as distinct reasoning

1 entities. Further that it is not diagnosis itself that is difficult but rather
2 clinical decisions, both diagnostic and management, that involve greater
3 complexity and uncertainty which are the more cognitively challenging. In
4 considering the reasons why doctors may consider diagnostic decision
5 making to be superior to other forms of clinical reasoning, one obvious
6 interpretation is that they feel threatened by the expansion of nurses and
7 pharmacists into prescribing roles. They wish to maintain control over
8 diagnostic decision making so they can retain control over the prescribing
9 process. Analogous to Pope's argument on medicine's response to the
10 evidence based medicine movement (Pope, 2003), medicine's response to
11 new prescribers' expansion into diagnostic decision making could be
12 viewed as a 'last stand' and wish to maintain power in the face of external
13 threats.

14 Abbott (1988) discusses how professionals, such as nurse and pharmacist
15 prescribers, make jurisdictional claims to an area of practice previously
16 under full jurisdictional control of the medical profession. As independent
17 prescribers, nurses and pharmacists can be seen as making a
18 jurisdictional claim to engage in diagnostic decision making. The outcome
19 of such jurisdictional claims is mediated through interactions between the
20 various stakeholders at the public, legal or workplace level (Abbott,
21 1988). The public outcry by doctors regarding diagnostic decision making
22 may reflect their desire to maintain their jurisdictional boundaries, done in
23 a manner designed to enlist the support of the wider public. Equally,
24 nurses and pharmacists, under the dominance of medicine, may feel
25 uncomfortable encroaching on territory identified with doctors and may

1 self-limit their areas of practice by indicating their unwillingness to engage
2 in diagnoses (Bissell et al., 2008, Weiss et al., 2006, Lloyd & Hughes,
3 2007).
4
5
6
7
8

9 Previous researchers have investigated the threat posed by nurse and
10 pharmacist prescribers (Fisher, 2010, Weiss & Sutton, 2009, Cooper et.
11 al., 2011). These researchers concluded that power relationships still
12 played a part and that the dominance of medicine has not been
13 threatened. Allsop (2006) has suggested that medicine has
14 accommodated to change resulting from the introduction of new
15 prescribers and has maintained overall control of patient management.
16 Cooper et al. (2011), in their evaluation of nurse and pharmacist
17 supplementary prescribing, identified five ways of working which
18 supported the continued dominance of medicine. These included:
19 patients' and supplementary prescribers' views of doctors as being
20 hierarchically superior; doctors' initial legitimization of nurses' and
21 pharmacists' prescribing; doctors' belief that they could control,
22 particularly nurse, prescribers' access to training; nurse and pharmacist
23 prescribers' frequent recourse to seeking advice from doctors (and
24 doctors' encouragement of this); and doctors' denigration of most routine
25 prescribing, making it therefore more suitable for nurses and pharmacists
26 to undertake (Cooper et al., 2011). However, this study and the previous
27 studies have only explored the dominance of medicine under
28 supplementary prescribing arrangements (or earlier nurse prescribing
29 arrangements). With the introduction of independent prescribing, and the
30 more explicit possibility for nurses and pharmacists to make diagnoses,
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

doctors may indeed feel under threat especially when, under pressure from supplementary prescribing, doctors made much of the skill and defining nature of diagnosing as key to their role (Cooper et al., 2011).

Developing Expertise

Nonetheless, it is important to consider the evidence regarding pharmacists' and nurses' ability to diagnose and prescribe, and whether they are safe prescribers. In Latter et al.'s (2007) study of the clinical appropriateness of nurse prescribing, nurses were found to be generally making clinically appropriate decisions. However some negative comments were made by the medical doctors on the expert panel which questioned nurses' assessment and diagnosing skills. However, the method used annotated transcripts of nurse consultations and some examples of incomplete assessment may have been because nurses did not verbalise their assessments. Concerns have been expressed about nurses' knowledge of pharmacology (Morrison-Griffiths et al., 2002, Buckley et al., 2006, Offredy et al., 2008, Cooper et al., 2008), pharmacists' lack of training in clinical examination skills and diagnoses (Weiss et al., 2006, Hoti et al., 2010, Cooper et al., 2008), variations in the preparation and academic qualifications of nurses going on prescribing courses (Latter & Courtenay, 2004) and pharmacists' lack of contact with patients (Buckley et al., 2006, Child et al., 1998). However, there is also some evidence, although not extensive, supporting the decision making skills of nurses and pharmacists. In addition to the study above (Latter et al., 2007), other research has found that nurses wrote prescriptions to a high

1 standard with most prescriptions determined to be appropriately and
2 safely prescribed (Drennan et al., 2009). In their assessment of the safety
3 and appropriateness of nurse and pharmacist prescribing, no errors were
4 identified across the 71 medicines prescribed and prescriptions were
5 judged overall to be appropriate (Bissell et al., 2008). The clinical
6 effectiveness and costs of mental health nurse supplementary prescribers
7 compared with psychiatrists' prescribing, similarly found no difference
8 between the two groups (Norman et al., 2010). Evidence also suggests
9 that nurse and pharmacist prescribers, in comparison with doctors, are
10 more likely to adhere to guidelines (Shulman & Jani, 2005, MacDonald et
11 al., 2005).

27 Underlying nurses' and pharmacists' performance as prescribers is their
28 training which has a strong emphasis on competence. The first required
29 learning outcome for pharmacist prescribing courses states independent
30 pharmacist prescribers should be able to:

40 'Understand the responsibility that the role of independent prescriber
41 entails, be aware of their own limitations, and work within the limits
42 of their professional competence – knowing when and how to refer /
43 consult / seek guidance from another member of the health care
44 team' (General Pharmaceutical Council, 2010, pp.1).

54 In nursing, section 2, practice standard 2.2 of the Standards for
55 Prescribing Practice states that 'you must only ever prescribe within your
56 level of experience and competence' (Nursing & Midwifery Council, 2010,

1 pp.29). The emphasis put upon competence, coupled with the dominance
2 of a patient safety agenda within the culture of the NHS, has meant that
3 self-limitation and restraint in terms of breadth of prescribing, has become
4 a professional ideal for some new prescribers (Weiss & Sutton, 2009).
5
6 Restrictions to the range of clinical areas in which prescribing occurs and
7 the use of protocols to guide prescribing decisions facilitate new
8 prescribers' competence in their areas of practice. This also makes
9 possible the development of doctors' overseer role in patient
10 management, helps maintain doctors' overall control over the prescribing
11 process and preserves their higher status (Allsop, 2006). However with
12 the maturation of nurse and pharmacist prescribing and the potential for
13 their expertise to develop, as well as anecdotal examples of nurses and
14 pharmacists who have expanded their areas of practice (Young, 2005,
15 Anonymous, 2004, Anonymous, 2009, Coombes, 2008, Tomlin, 2009,
16 Sibbald, 2008), it is uncertain how long doctors' control over patient
17 management will be maintained.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The traditional approach to medical clinical teaching, and the development
of diagnostic expertise, consists of two stages: mastery of basic
biomedical sciences followed by the assessment of patient clinical
problems (Eva, 2004). Experience, following on from medical training, is
recognised as important to the development of expertise. As noted by
Freidson (1970), medical knowledge is justified in terms of the doctor's
personal knowledge and their professional experience. This practical
reasoning is resistant to change on the basis of abstractions or statistical
considerations (Freidson, 1970). Clinical experience is seen as an

important source of control over uncertainty, with the ultimate respect given to judgement based on experience (Light, 1979). Increasing age, correlating with increasing expertise, has been shown to increase reliance on non-analytic clinical reasoning (Elstad et al., 2010), with GPs identified as diagnostic experts relying on relatively few elements of clinical data (Groves et al., 2003).

The importance and salience of experience has been contested in more recent years with the work of Archie Cochrane, David Sackett and the rise of evidence-based medicine (EBM). Cochrane supported the use of randomised controlled trials, where appropriate, to test the effectiveness of medical interventions and for these findings to guide clinical decision making (Cochrane, 1972). The emphasis on identifying, appraising and implementing relevant clinical research into individual patient management situations was further refined by David Sackett and central to the evidence-based medicine movement (Sackett et al., 1996). What the evidence based movement proposed was that clinical experience may not be the best method of developing expertise and that EBM with its 'conscientious, explicit and judicious use of current best evidence' in making decisions about patients (Sackett et al., 1996, pp.71), is a more robust method of developing clinical expertise.

Ericsson and Towne (2010) take a cognitive psychology approach to the study of expertise. They describe two distinct perspectives, one of which is the traditional approach, or accumulated experience model, where expertise is developed through increased training and experience over a

1 period of years. This model has been criticised because the relationship
2 between increasing experience does not correlate in a linear fashion with
3 increasing expertise. At a certain point, performance reaches a level of
4 automaticity and is effortless, and increasing experience will not improve
5 accuracy further. The second approach, called expert performance,
6 focuses on individuals who have superior performance at reproducible
7 tasks that capture the expertise representative of a particular domain
8 such as medicine. In this model, experts, such as doctors, have superior
9 anticipation skills and more refined cognitive representations which enable
10 them to access and execute appropriate actions quickly. These
11 representations are stored in long-term memory but, because they can be
12 rapidly accessed like short term working memory yet are not likewise
13 capacity-limited, this storage / retrieval aspect of memory has been called
14 long term working memory. Most intriguingly, this expert performance
15 approach can be enhanced through 'deliberate practice' where an
16 individual intensely concentrates on a specific aspect of performance
17 which leads to modification of the mechanisms responsible for
18 improvement. In this model, informative feedback and coaching are key
19 to the development of expertise. In medicine this could include expert
20 performers working through descriptions of previous diagnostic situations
21 where the correct outcome is known so that they receive immediate
22 feedback on the accuracy of their decisions (Ericsson & Towne, 2010).

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54 In contrast to the medical training model of the learning of basic sciences
55 followed by engagement with patient clinical problems, the nurse and
56 pharmacist prescribing training model emphasises competence. This
57
58
59
60
61
62
63
64
65

1 emphasis on competence is at least in part necessary due to the diverse
2 knowledge, qualifications and experiences of the nurses and pharmacists
3 who may decide to become prescribers. For the development of further
4 expertise, evidence suggests that for nurse and pharmacist prescribing to
5 succeed, they need support from peers (through professional networks,
6 multidisciplinary working and collaborative practice), they need good
7 mentorship arrangements with doctors and they need to have access to
8 continuous professional development (Cooper et al., 2008, Weiss et al.,
9 2006, Stewart et al., 2009, Carey, et al. 2009, Otway, 2001, Bradley et
10 al., 2007). Alongside these, the methods for developing expertise
11 previously mentioned including increased clinical experience, use of tools
12 derived from evidence-based medicine, and enhancement through
13 'deliberate practice' are also likely to play a role.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 Medical educators have similarly recognised that there might be a number
34 of routes for developing expertise (Eva, 2004). Eva (2004) suggests that
35 there might be several strategies for developing diagnostic expertise such
36 as accumulating a mental database of patient cases, obtaining experience
37 with a diverse range of diagnostic categories and identifying similarities in
38 the underlying concepts of superficially distinct problems. These strategies
39 bear similarity to Ericsson and Towne's (2010) notion of 'deliberate
40 practice' where diagnostic skill is gained through concentration on specific
41 aspects of diagnostic performance and the development of refined
42 cognitive representations stored in long term working memory. As
43 discussed by Eva (2004), while the learning of basic science mechanisms
44 contributes towards making accurate diagnoses, it may not be the only
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 way of doing so. For nurse and pharmacist prescribers to develop
2 diagnostic expertise, it is likely that a range of approaches, which may
3 include newer ideas such as the use of deliberate practice, will be shown
4 to be effective in enhancing diagnostic performance. Independent
5 prescribing by nurses and pharmacists is still a relatively new
6 phenomenon and new prescribers need time to mature and develop in
7 their roles. Given the starting point of new prescribers, further research is
8 needed on what methods or combination of methods, derived from
9 medical training, medical practice or cognitive psychology, are most
10 effective in enhancing their diagnostic skill.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

26 Conclusions

27
28
29
30 Clinical reasoning underlies all types of clinical decisions including
31 diagnostic decisions, as well as prescribing decisions, referral decisions or
32 other decisions about patient management. Previous research suggests
33 that diagnostic decisions are deeply intertwined with decisions about
34 management, using the same clinical reasoning processes. In addition,
35 everyone, when they are confronted with having to manage new
36 symptoms, make diagnostic decisions. This suggests that privileging
37 diagnostic decision making as being superior to, or more difficult than,
38 prescribing decision making is conceptually flawed. This is not to say there
39 are not difficult diagnostic and patient management decisions. Rather
40 there are features, involving complexity and uncertainty, which are
41 associated with some clinical decisions (e.g. both diagnostic and
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

management) that makes them more difficult or challenging for a practitioner to make.

The medical profession's response to nurse and pharmacist independent prescribing could be viewed as a 'last stand' and wish to maintain their professional power in the face of external threats. Independent prescribing explicitly allows for the possibility for nurses and pharmacists to make diagnoses. Indeed, doctors' perception of this professional threat may have been exacerbated by their earlier response to supplementary prescribing, when they denigrated the routine nature of the prescribing undertaken by pharmacists and nurses, and made much of the skill and defining nature of diagnosing. Although ostensibly concerned about the risks to public safety of nurse and pharmacist independent prescribing, the evidence to date, although not extensive, suggests that nurses and pharmacists can prescribe safely and appropriately.

The model for training nurse and pharmacist prescribers relies on their recognition of their areas of competence and to prescribe only within these clinical areas. Protocols and guidelines may be helpful in supporting new prescribers' decision making, particularly when they are new to their role, less confident and are able to use guidelines to help organise the incoming patient clinical information. However, as new independent prescribers mature in their role and as independent prescribing becomes more commonplace, consideration needs to be given as to how to develop the diagnostic expertise of nurse and pharmacist prescribers. The traditional medical model of learning basic science mechanisms followed

1 by engagement with patient clinical problems is one approach to learning
2 how to make accurate diagnoses. But it may not be the only way. Indeed,
3
4 research suggests that diagnostic performance can be enhanced through
5
6 'deliberate practice' whereby an individual intensely concentrates on a
7
8 specific aspect of performance and, through feedback and coaching, is
9
10 able to improve their diagnostic accuracy. Further research is needed to
11
12 determine if such an approach will prove effective in developing the
13
14 expertise of nurses and pharmacists in diagnosing. In conclusion, the
15
16 question becomes, not can pharmacist and nurse prescribers make
17
18 diagnoses, because clearly they already do. Rather the question is, is it
19
20 possible for nurse and pharmacist prescribers to develop expertise and
21
22 become expert diagnosticians? The answer to this surely is a cautious
23
24 'yes'.
25
26
27
28
29
30
31
32

33 References

34
35
36
37 Abbott, A. (1988). *The system of professions: an essay on the division of*
38
39 *expert labour*. Chicago and London: the University of Chicago Press.
40
41
42

43
44 Allsop, J. (2006). Medical dominance in a changing world: the UK case.
45
46 *Health Sociology Review*, 15, 444-57.
47
48
49

50
51 Anonymous. (2004). Pharmacist is made a partner in Devon GP
52
53 dispensing practice. *The Pharmaceutical Journal*, 272, 463.
54
55
56
57
58
59
60
61
62
63
64
65

1 Anonymous. (2006). Nurse prescribing 'a threat to safety'. *Pulse* 5
2 October. Available from:
3
4 <http://www.pulsetoday.co.uk/story.asp?sectioncode=23&storycode=4010>
5
6 [567](#) (Accessed 10 May 2011).
7
8
9

10
11 Anonymous. (2009). Pharmaceutical Care Awards: Making a difference in
12 care homes. *The Pharmaceutical Journal*, 283, 13.
13
14
15
16

17
18 Atkinson P. (1995). *Medical Talk and Medical Work – The Liturgy of the*
19
20
21 *Clinic*. Sage: London.
22
23
24

25
26 Avery, A.J. & Pringle, M. (2005). Extended prescribing by UK nurses and
27 pharmacists. *British Medical Journal*, 331, 1154-5.
28
29
30
31

32
33 Benner, P. (1982). From novice to expert. *American Journal of Nursing*,
34
35 82, 402-7.
36
37
38

39
40 Berg, M. (1992). The construction of medical disposals. Medical sociology
41 and medical problem solving in clinical practice. *Sociology of Health &*
42
43 *Illness*, 14, 151-79.
44
45
46
47

48
49 Berg, M. (1997). Problems and Promises of the Protocol. *Social Science &*
50
51 *Medicine*, 44, 1081-8.
52
53
54
55

56
57 Bissell, P., Cooper, R., Guillaume, L., Anderson, C., Avery, A., Hutchinson,
58
59 A., James, V., Lynn, J., Marsden, E., Murphy, E., Ratcliffe, J., Ward, P., &
60
61
62
63
64
65

1 Woolsey, I. (2008) *An Evaluation of Supplementary Prescribing in Nursing*
2 *and Pharmacy (Final Report for the Department of Health)*, University of
3
4 Sheffield.
5
6
7
8

9 Bloor, M. (1976). Bishop Berkeley and the Adeno-tonsillectomy enigma:
10 an exploration of variation in the social construction of medical disposals.
11
12 *Sociology*, 10, 43-61.
13
14
15
16
17

18 Bloor, M. (1978). On the routinised nature of work in people-processing
19 agencies: the case of adeno tonsillectomy assessments in ENT outpatient
20 clinics. In A. Davis (Ed). *Relationships between Doctors and Patients*
21 (pp.29-47). Farnborough: Teakfield Ltd.
22
23
24
25
26
27
28
29

30 Bloor, M. & Horrobin, G. (1975). Conflict and conflict resolution in doctor-
31 patient interactions. In C. Cox & M.E. Mead (Eds), *A Sociology of Medical*
32 *Practice* (pp.271-84). London: Collier MacMillan.
33
34
35
36
37
38
39

40 Bradley, E., Hynam, B. & Nolan, P. (2007). Nurse prescribing: Reflections
41 on safety in practice. *Social Science & Medicine*, 65, 599-609.
42
43
44
45
46

47 British Medical Association. (2005). *BMA calls for urgent meeting with*
48 *Patricia Hewitt on plans to extend prescribing powers*. British Medical
49 Association, press release, 10 Nov 2005. Available from:
50
51 [http://web2.bma.org.uk/pressrel.nsf/wlu/STRE-](http://web2.bma.org.uk/pressrel.nsf/wlu/STRE-6HZJ4X?OpenDocument&vw=wfmms)
52
53 [6HZJ4X?OpenDocument&vw=wfmms](http://web2.bma.org.uk/pressrel.nsf/wlu/STRE-6HZJ4X?OpenDocument&vw=wfmms)
54
55
56
57
58
59
60
61
62
63
64
65

1 Brooks, L.R., Norman, G.R. & Allen, S.W. (1991). Role of specific similarity
2 in a medical diagnostic task. *Journal of Experimental Psychology: General*,
3
4 120, 278-87.
5
6

7
8
9 Buckley, P., Grime, J. & Blenkinsopp, A. (2006). Inter- and intra-
10 professional perspectives on non-medical prescribing in an NHS trust. *The*
11
12 *Pharmaceutical Journal*, 277, 394-8.
13
14
15

16
17
18 Carey, N., Stenner, K. & Courtenay, M. (2009). Stakeholder views on the
19 impact of nurse prescribing on dermatology services. *Journal of Clinical*
20
21 *Nursing*, 19, 498-506.
22
23
24

25
26
27 Child D., Hirsch, C. & Berry, M. (1998). Health care professionals' views
28 on hospital pharmacist prescribing in the United Kingdom. *International*
29
30 *Journal of Pharmacy Practice*, 6, 159-69.
31
32
33

34
35
36 Child, D., & Cantrill, J.A. (1999). Hospital doctors' perceived barriers to
37 pharmacist prescribing. *International Journal of Pharmacy Practice*, 7,
38
39 230-7.
40
41
42

43
44
45 Choudhry, N.K., Fletcher, R.H. & Soumerai, S.B. (2005). Systematic
46
47 Review: The relationship between clinical experience and quality of health
48
49 care. *Annals of Internal Medicine*, 142, 260-73.
50
51
52

53
54
55 Cochrane, A. (1972). *Effectiveness and efficiency: random reflections on*
56
57 *health services*. London: Nuffield Provincial Hospitals Trust.
58
59
60

1
2 Coombes, R. (2008). Dr Nurse will see you now. *British Medical Journal*,
3
4 337, a1522.
5
6

7
8
9 Cooper, R.J., Anderson, C., Avery, T., Bissell, P., Guillaume, L.,
10
11 Hutchinson, A., James, V., Lymn, J., McIntosh, A., Murphy, E., Ratcliffe,
12
13 J., Read, S. & Ward, P. (2008). Nurse and pharmacist supplementary
14
15 prescribing in the UK – a thematic review of the literature. *Health Policy*,
16
17 85, 277-92.
18
19
20

21
22
23 Cooper, R. J., Bissell, P., Ward, P., et.al. (2011). Further challenges to
24
25 medical dominance? The case of nurse and pharmacist supplementary
26
27 prescribing *Health: first published on January 13, 2011 as*
28
29 *doi:10.1177/1363459310364159*.
30
31
32

33
34
35 Day, M. (2005). UK doctors protest at extension to nurse prescribing
36
37 powers. *British Medical Journal*, 331, 1159.
38
39
40

41
42 Department of Health (1999). *Review of prescribing, supply and*
43
44 *administration of medicines, Final Report (The Crown Report)*. London:
45
46 Department of Health.
47
48
49

50
51 Department of Health (2005). *Supplementary prescribing by nurses,*
52
53 *pharmacists, chiropodists/podiatrists, physiotherapists and radiographers*
54
55 *with the NHS in England – a Guide for Implementation*. Gateway reference
56
57 4941. Updated May 2005. London: Department of Health.
58
59
60
61
62
63
64
65

1
2 Department of Health (2006). *Improving patients' access to medicines: a*
3
4 *Guide to Implementing Nurse and Pharmacist Independent Prescribing*
5
6 *within the NHS in England*. London: Department of Health.
7
8
9

10
11 Drennan, J., Naughton, C., Allen, D., Hyde, A., Felle, P., O'Boyle, K.,
12
13 Treacy, P. & Butler, M. (2009). *National Independent Evaluation of the*
14
15 *Nurse and Midwife Prescribing Initiative*. Dublin: University College Dublin.
16
17
18
19

20
21 Eddy, D.M. (1990). Anatomy of a Decision. *Journal of the American*
22
23 *Medical Association*, 263, 441-443.
24
25
26

27
28 Elstad, E.A., Lutfey, K.E., Marceau, L.D., Campbell, S.M., Kneserbeck, O.
29
30 & McKinlay, J.B. (2010). What do physicians gain (and lose) with
31
32 experience: Qualitative results from across-national study of diabetes.
33
34 *Social Science & Medicine*, 70, 1728-36.
35
36
37

38
39 Elstein, A.S. & Schwarz, A. (2002). Clinical problem solving and diagnostic
40
41 decision making: selective review of the cognitive literature. *British*
42
43 *Medical Journal*, 324, 729-82.
44
45
46

47
48 Ericsson, K.A. & Towne, T.J. (2010). Expertise. *Cognitive Science*, 1, 404-
49
50 416.
51
52
53

54
55
56 Eva, K.W. (2004). What every teacher needs to know about clinical
57
58 reasoning. *Medical Education*, 39, 98-106.
59
60
61

Eva, K.W. & Cunningham, J.P.W. (2006). The difficulty with experience:
Does practice increase susceptibility to premature closure? *Journal of
Continuing Education in the Health Professions*, 26, 192-8.

Eva, K.W., Link, C.L., Lutfey, K.E. & McKinlay, J.B. (2010). Swapping
Horses midstream: factors related to physicians' changing their minds
about a diagnosis. *Academic Medicine*, 85, 1113-7.

Feltovich, P.J. (1981). *Knowledge based components of expertise in
medical diagnosis*. Doctoral dissertation: University of Minnesota.

Fisher, R. (2010) Nurse prescribing: a vehicle for improved collaboration,
or a stumbling block to inter-professional working? *International Journal
of Nursing Practice* 16, 579-85.

Fox, R. (1957). Training for uncertainty. In: R.K. Merton, G.G. Reader & P.
Kendall (eds), *The Student Physician* (pp. 207-41). Cambridge, MA:
Harvard University Press.

Freidson, E. (1970). *Profession of Medicine*. New York: Dodd-Mead.

General Pharmaceutical Council (2010). Pharmacist Independent
Prescribing – Programme Learning Outcomes and Indicative Content.
Available from:

<http://www.pharmacyregulation.org/regulatingpharmacy/educationandprofessionaldevelopment/educationandtraining/pharmacistindependentprescriber/index.aspx>

Accessed 7 October 2010.

Godlee, F. (2008). What skills do doctors and nurses need? *British Medical Journal*, 337: a1722.

Groves, M., O'Rourke, P. & Alexander, H. (2003). The clinical reasoning characteristics of diagnostic experts. *Medical Teacher*, 25, 308-15.

Gutkin, C. (2009). To diagnose, or not to diagnose: that is the question. *Canadian Family Physician*, 55, 320.

Hall, K.H. (2002). Reviewing intuitive decision-making and uncertainty: the implications for medical education. *Medical Education*, 36, 216-24.

Hamm, R.M. (1988). Clinical intuition and clinical analysis: expertise and the cognitive continuum. In J. Dowie & A. Elstein (Eds). *Professional Judgement – A Reader in Clinical Decision Making*. Cambridge: Cambridge University Press.

Hassell, K., Rogers, A.E. & Noyce, P.R. (2000). Community pharmacy as a primary health and self-care resource: a framework for understanding pharmacy utilization. *Health and Social Care in the Community*, 8, 40-49.

Hoti, K., Sunderland, B., Hughes, J. & Parsons, R. (2010). An evaluation of Australian pharmacists' attitudes on expanding their role. *Pharmacy World & Science*, 32, 610-21.

Howie, J.G.R. (1972). Diagnosis – the Achilles heel? *Journal of the Royal College of General Practitioners*, 22, 310-15.

Howie, J.G.R. (1974). Further observations on diagnosis and management of general practice respiratory illness using simulated patient consultations. *British Medical Journal*, 2, 540-3.

Howie, J.G.R. (1976). Clinical judgement and antibiotic use in general practice. *British Medical Journal*, 2, 1061-4.

Katz, P. (1985). How surgeons make decisions. In R.A. Hahn & A.D. Gaines (Eds), *Physicians of Western Medicine: Anthropological Approaches to Theory and Practice* (pp. 55-75). Dordrecht: Reidel.

Latter, S. & Courtenay, M. (2004). Effectiveness of nurse prescribing: a review of the literature. *Journal of Clinical Nursing*, 13, 26-32.

Latter, S., Maben, I., Myall, M. & Young, A. (2007). Evaluating the clinical appropriateness of nurses prescribing practice: method development and findings from an expert panel analysis. *Quality and Safety in Health Care*, 16, 415-21.

1 Lehto, M.R. & Nah, F. (2006). Chapter 8: Decision making models and
2 decision support. In: G. Salvendy (ed), *Handbook of Human Factors and*
3
4 *Ergonomics* (pp 191-242). London: John Wiley & Sons.
5
6
7

8
9 Light, D.L. (1979). Uncertainty and Control in Professional Training.
10
11 *Journal of Health and Social Behaviour*, 20, 310-22.
12
13
14

15
16 Lloyd, F. & Hughes, C. (2007). Pharmacists' and Mentors' Views on the
17
18 introduction of pharmacist supplementary prescribing: a qualitative
19
20 evaluation of views and context. *International Journal of Pharmacy*
21
22 *Practice*, 15, 31-37.
23
24
25

26
27
28 Lloyd, F., Parsons, C. & Hughes, C. (2010). 'It showed me the skills that
29
30 he has': pharmacists and mentors views on pharmacist supplementary
31
32 prescribing. *International Journal of Pharmacy Practice*, 18, 29-36.
33
34
35

36
37
38 Luker K., Austin, L., Ferguson, B. & Smith, K. (1997). Nurse prescribing:
39
40 the views of nurses and other health care professionals. *British Journal of*
41
42 *Community Health Nursing*, 2, 69-74.
43
44
45

46
47 Luker, K.A., Hogg, C., Austin, L., Ferguson, B. & Smith, K. (1998).
48
49 Decision making: the context of nurse prescribing. *Journal of Advanced*
50
51 *Nursing*, 27, 657-665.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Lutfey, K.E. & McKinlay, J.B. (2009). What happens along the diagnostic
2 pathway to CHD treatment? Qualitative results concerning cognitive
3 processes. *Sociology of Health & Illness*, 31, 1077-92.
4
5
6
7

8
9 McDonald, R., Waring, J., Harrison, S., Walshe, K. & Boaden, R. (2005).
10 Rules and guidelines in clinical practice: a qualitative study in operating
11 theatres of doctors' and nurses' views. *Quality and Safety in Health Care*,
12 14, 290-4.
13
14
15
16
17

18
19 Morrison-Griffiths, S., Snowden, M.A. & Pirmohamed, M. (2002). Pre-
20 registration nurse education in pharmacology: is it adequate for the roles
21 that nurses are expected to fulfil? *Nurse Education Today*, 22, 447-56.
22
23
24
25
26
27

28
29 National Institute for Health and Clinical Excellence (2006). Hypertension.
30 Management of Hypertension in Adults in Primary Care (CG34). National
31 Institute for Health and Clinical Excellence: London. Available from:
32 <http://guidance.nice.org.uk/CG34/NICEGuidance/pdf/English>
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66 Nazareth, I. & Murray, E. (2010). Promoting Self Care for Minor Illness.
67 *British Medical Journal*, 340, c2913.

68 Neufeld, V.R., Norman, G.R., Feightner, J.W. & Barrows, H.S. (1981).
69 Clinical problem solving by medical students: a cross-sectional and
70 longitudinal analysis. *Medical Education*, 15, 315-22.

1 Norman, G., Barraclough, K., Dolovich, L. & Price, D. (2009). Iterative
2 Diagnosis. *British Medical Journal*, 339, b3490. doi: 10.1136/bmj.b3490.
3
4

5
6 Norman, I.J., Coster, S., McCrone, P., Sibley, A. & Whittelsea, C. (2010).
7 A comparison of the clinical effectiveness and costs of mental health nurse
8 supplementary prescribing and independent medical prescribing: a post-
9 test control group study. *BMC Health Services Research*, 10, 4.
10
11
12
13
14
15
16

17
18 Nursing and Midwifery Council. Standards of Proficiency for Nurse and
19 Midwife Prescribers. Standards 06-06. Available from:
20
21 <http://www.nmc-uk.org/Nurses-and-midwives/Prescribing/Education/>
22
23
24
25
26 Accessed 7 October 2010.
27
28
29

30
31 Offredy, M., Kendall, S. & Goodman, C. (2008) The use of cognitive
32 continuum theory and patient scenarios to explore nurse prescribers'
33 pharmacological knowledge and decision-making. *International Journal of*
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1 Round, A. (2000). Introduction to clinical reasoning. *Student British*
2 *Medical Journal*, 8, 15-16.
3
4

5
6
7 Sackett, D., Rosenberg, W., Muir Gray, J.A., Haynes, R.B. & Richardson,
8 W.S. (1996) Evidence-based medicine: what it is and what it isn't. *British*
9 *Medical Journal*, 312: 71.
10
11
12

13
14
15
16 Shulman, R. & Jani, U. (2005) Comparison of supplementary prescribers'
17 and doctors' compliance with guidelines for drug dosing in haemofiltration
18 on an intensive care unit. *Pharmaceutical Journal*, 274, 492-3.
19
20
21

22
23
24
25
26 Sibbald B. (2008). Should primary care be nurse led? Yes. *British Medical*
27 *Journal*, 337, a1157.
28
29
30

31
32
33 Stewart, D.C., George, J., Bond, C.M., Diack, H.L., McCaig, D.J. &
34 Cunningham, S. (2009). Views of pharmacist prescribers, doctors and
35 patients on pharmacist prescribing implementation. *International Journal*
36 *of Pharmacy Practice*, 17, 89-94.
37
38
39
40

41
42
43
44 Tomlin, M. (2009). 'Independent prescribing is neither for the faint-
45 hearted nor for those lacking experience as a pharmacist'. *Clinical*
46 *Pharmacist*, 1, 458-9.
47
48
49

50
51
52
53
54 Tully, M.P., Hassell, K. & Noyce, P. (1997). Advice-giving in community
55 pharmacies in the UK. *Journal of Health Services Research & Policy*, 2, 38-
56
57
58
59 50.
60
61
62
63
64
65

1
2 van Merriënboër, J.J.G. & Sweller, J. (2005). Cognitive load theory and
3
4 complex learning: recent developments and future directions. *Educational*
5
6 *Psychology Review* 17, 147-77.
7
8

9
10
11 Tversky, A. & Kahneman, D. (1974). Judgement and uncertainty:
12
13 heuristics and biases. *Science*, 185, 1124-31.
14
15

16
17
18 Weiss, M.C., Adams, C. & Sutton, J. (2006). *Exploring Innovation in*
19
20 *Pharmacy Practice: A qualitative evaluation of supplementary prescribing*
21
22 *by pharmacists*. London: Pharmacy Practice Research Trust. Available
23
24 from:
25
26 <http://www.pprt.org.uk/Publications/2006Publications.aspx>
27
28
29 Accessed 7 October 2010.
30
31

32
33
34
35 Weiss, M.C. & Sutton, J. (2009). The changing nature of prescribing:
36
37 pharmacists as prescribers and challenges to medical dominance.
38
39 *Sociology of Health & Illness*, 31, 406-421.
40
41

42
43
44 Young, G. (2005). The nursing profession's coming of age. *British Medical*
45
46 *Journal*, 331, 1415.
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65