Citation for published version:

Cazzola, D, Alberti, G, Ongaro, L & Minetti, AE 2014, 'The vertical excursion of the body visceral mass during vertical jumps is affected by specific respiratory maneuver', *Human Movement Science*, vol. 33, no. 1, pp. 369-380. <https://doi.org/10.1016/j.humov.2013.10.008>

DOI:

[10.1016/j.humov.2013.10.008](https://doi.org/10.1016/j.humov.2013.10.008)

Publication date:

2014

Document Version

Peer reviewed version

[Link to publication](#)

NOTICE: this is the author's version of a work that was accepted for publication in *Human Movement Science*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Human Movement Science*, 2013, DOI 10.1016/j.humov.2013.10.008

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 THE VERTICAL EXCURSION OF THE BODY VISCERAL MASS
2 DURING VERTICAL JUMPS IS AFFECTED BY SPECIFIC
3 RESPIRATORY MANOEUVRE
4

5 Dario Cazzola^{1,4}, Giampietro Alberti², Lucio Ongaro³, Alberto E. Minetti¹
6

7 ¹Laboratory of Physiomechanics, Department of Physiopathology and Transplantation,
8 Faculty of Medicine, University of Milan, Italy, ²Department of Sport, Nutrition and
9 Health Sciences, Faculty of Exercise and Sports Sciences, University of Milan,
10 ³Faculty of Exercise and Sports Sciences, University of Milan, ⁴Sport, Health and
11 Exercise Science, Department of Health, University of Bath, UK.
12

13 **Corresponding author:** Dr Dario Cazzola, Research Officer - Sport, Health &
14 Exercise Science - Department for Health | University of Bath - Applied
15 Biomechanics Suite, 1.308. BA2 7AY, BATH (UK). Mobile (UK) ++44 (0)
16 7450820004. E-mail (University of Bath): dc547@bath.ac.uk. E-mail
17 (home): dario.cazzola@me.com
18

19 ABSTRACT

20 Most of the modelling of body dynamics in sports assumes that every segment is
21 'rigid' and moves 'as a whole', although we know that uncontrolled wobbling masses
22 exist and their motion should be minimized, both in engineering and biology. The
23 visceral mass movement within the trunk segment potentially interferes with
24 respiration and motion acts as locomotion or jumping. The aim of this paper is to
25 refine and expand a previously published methodology to estimate that relative
26 motion by testing its ability to detect the reduced vertical viscera excursion within the
27 trunk. In fact, a respiratory-assisted jumping strategy is expected to limit viscera
28 motion stiffening the abdominal content of the bouncing body. Six subjects were
29 analysed, by using both inverse and direct dynamics, during repeated vertical jumps
30 performed before and after a specific respiratory training period. The viscera
31 excursion, which showed consistent intra-individual time courses, decreased by about
32 30% when the subjects had familiarized with the trunk-stiffening manoeuvre. We
33 conclude that: 1) the present methodology proved to detect subtle visceral mass
34 movement within the trunk during repetitive motor acts and, particularly, 2) a newly
35 proposed respiratory manoeuvre/training devoted to stiff the trunk segment can
36 reduce its vertical displacement.

37
38
39
40

41 1. INTRODUCTION

42 In biomechanical studies of human and animal motion and locomotion, the body is
43 often simplified as composed by a number of rigid segments. From the location of
44 those segments in 3D space, many important variables such as the body centre of
45 mass (BCoM), the related internal and external mechanical work (Willems, Cavagna,
46 & Heglund, 1995) are calculated to infer the characteristic dynamics of movement (A.
47 E. Minetti, Cisotti, & Mian, 2011; Saibene & Minetti, 2003). Also, rotational
48 parameters as joint net moments and segments inertial characteristics are based on the
49 same “rigid body model”. Unfortunately, such assumption can lead to experimental
50 inaccuracies (Gao & Zheng, 2008; Leardini, Chiari, Della Croce, & Cappozzo, 2005).
51 For this reason specific wobbling mass models have been proposed (Gruber, Ruder,
52 Denoth, & Schneider, 1998; Yue & Mester, 2002) to improve and to refine
53 experimental results especially during impacts (Gunther, Sholukha, Kessler, Wank, &
54 Blickhan, 2003; M. T. G. Pain & Challis, 2004), in the attempt to enhance the
55 description of the complex mechanical behaviour of the human body by including the
56 contribution of soft parts. This approach allows quantification of the soft tissue
57 deformation and displacement as a consequence of the impact forces transmission
58 along the body (Challis & Pain, 2008; Wakeling & Nigg, 2001) during walking
59 (Chen, Mukul, & Chou, 2011), running (Boyer & Nigg, 2007) and jumping (Gittoes,
60 Brewin, & Kerwin, 2006; Mills, Scurr, & Wood, 2011). Soft tissue and viscera
61 motion can also affect the external work of level and gradient walking (DeVita,
62 Helseth, & Hortobagyi, 2007; Zelik & Kuo, 2010) and of running economy and
63 stability (Daley & Usherwood, 2010). It as even be proposed that a suitable muscle-
64 tuned control of that collateral effect could minimize the overall energy dissipation
65 (Friesenbichler, Stirling, Federolf, & Nigg, 2011).

66 Thus, soft tissue and viscera movement has to be considered as a non-negligible
67 factor in modelling optimization strategies and in experimental methodology, also in
68 relation to the potential mechanical interaction with the rest of the body. For example,
69 several authors have just pointed out the role of the visceral mass movement (within
70 the trunk) in the locomotor-respiratory coupling during trotting and galloping in
71 quadrupeds (Alexander, 1993; Bramble & Carrier, 1983; Simons, 1999). A similar
72 condition occurs in humans, where some locomotor-respiratory coupling in running
73 (McDermott, Van Emmerik, & Hamill, 2003) and walking (Rassler & Kohl, 1996)
74 reflects the influence on the diaphragm function of the transient axial acceleration of
75 abdominal viscera (Brown, Lee, & Loring, 2004; Loring, Lee, & Butler, 2001; Wilson
76 & Liu, 1994). A very simple experiment illustrates this point: whoever tries to breath
77 out-of-phase with respect to the spontaneous pattern during repeatedly jumping in
78 place feels a great discomfort in achieving such a goal, mainly because respiratory
79 muscles have to fight against the volume changes imposed by the jump-induced
80 vertical accelerations of the visceral piston within its container.

81 In addition to the coupling between a cyclic activity as locomotion and respiration,
82 there are other movements where the visceral mass displacement can play a role. In
83 sport activities as volleyball, basketball or athletics, where jumping efficacy or
84 horizontal-to-vertical velocity conversion are crucial (Yu & Hay, 1996), it is
85 conceivable that controlling the wobbling mass could potentially avoid discomfort
86 and energy dissipation associated to adverse oscillations, by also lowering workload
87 perception (Bonsignore, Morici, Abate, Romano, & Bonsignore, 1998) or enhancing
88 the jump performance. In this respect training techniques have been suggested to
89 reduce the amplitude of that movement (Caufriez, 2005; Kapandji, 1977; Lumb,

2005) or even to obtain a beneficial influence on BCoM trajectory during the motion cycle.

A few years ago, a methodology using both 3D motion capture and platform dynamometry was proposed to infer the movement of the visceral mass during cyclic motor acts (A. Minetti & Belli, 1994). In short, by comparing the movement of the container (i.e. the rigid, multi-segment body) assessed by motion analysis, to the displacement of the 'true' BCoM, evaluated by double integration of the net vertical ground reaction force, it was possible to quantify the relative motion of the visceral mass within the trunk.

The aim of this paper was to apply that method to test whether a novel jumping technique, based on stiffening both chest and abdominal walls by means of a particular respiratory manoeuvre, was associated to the expected reduction in the visceral mass vertical displacement within the trunk. That would represent the first experimental evidence that the effects of a voluntary pattern of respiratory muscles activation during jumping can be accurately measured with a non-invasive approach.

2. MATERIALS AND METHODS

2.1 EXPERIMENTAL PROTOCOL

Six subjects (age 23.3 ± 2.5 , trunk length 0.570 ± 0.110 m, weight 659.4 ± 53.0 N) were selected to jump in two different sessions on a force platform (model 9281C, Kistler, CH) measuring the vertical GRF synchronized with a six-camera motion capture system (Vicon MX, Oxford Metrics, UK). All the subjects were students from the Sport Science Faculty (University of Milan), chosen for their motor/jumping skill. The institutional ethics committee had approved all the methods and procedures, and subjects gave their informed consent prior to the experiments.

115 The platform signal was sampled at 1200 Hz, while the optoelectronic system
116 captured frames at 400 Hz. The human body was modelled as a series of 14 linked,
117 rigid body segments: 18 reflective markers (radius = 14 mm) were placed bilaterally
118 on anatomical landmarks (Figure 1), nine on each side of the body (Mian, Thom,
119 Ardigò, Narici, & Minetti, 2006), while 4 'technical-markers' were placed on the
120 estimated centre of mass position of pectoral muscles, and right and left abdomen
121 surface. Segment mass fraction and proximal distance of the centre of mass were
122 taken from Dempster (Dempster, Gabel, & Felts, 1959).

123 The experiment consisted of two sessions, which were made up of 5 trials containing
124 15 consecutive jumps each, and spaced out by an adequate recovery period between
125 trials. During the first session, the subjects jumped barefoot, with the hand on their
126 hips, without any advice, to facilitate a natural jump execution. The second
127 experimental session took place according to the same protocol after a training period
128 of one month in which the subjects followed a specific learning progression devoted
129 to jump in the “controlled” way (see below). Before the second session, the specific
130 respiration technique and muscle contraction skills were tested on every subject:
131 airflow was measured with a heated Fleisch pneumotachograph (HS Electronics,
132 March-Hugstetten, Germany) connected to a facial mask and a differential pressure
133 transducer (Validyne MP45, Northridge, CA). The activity of rectus and obliquus
134 abdominis muscles was recorded via surface EMG (model ICP511, Grass
135 Technologies, US), and the rectified EMG signal was filtered by 2th order low-pass
136 Butterworth filter with cut-off frequency of 6 Hz (Clancy, Morin, & Merletti, 2002).
137 Both the signals were sampled at 1200 Hz by a 16-bit analog to-digital converter, and
138 stored on a desk computer. Volume changes (V) were obtained by numerical

integration of the digitized airflow signal, after calibration of the measuring apparatus by means of a graded cylinder and a metronome.

2.2 'CONTROLLED' JUMPING TECHNIQUE

The training technique suggested in this study was designed according to the idea that by predominantly using 'low' diaphragmatic respiration, the visceral mass could be increasingly compacted towards the pelvis (Calais-Germain, 2005). With the spine in the physiological upright posture, a proper contraction activity of the abdominal wall/pelvic floor muscles avoids the forward displacement of the compressed viscera, improves the stiffness of the abdominal belt and, consequently, of the whole body structure (Le Boulch, 1973). This is achievable through a limited pelvis anteversion position, the preparatory low diaphragmatic inspiration (Figure 2a), and the simultaneous dorsum-lumbar filling caused by an intra-abdominal pressure increase, which is amplified by the forced expiration during the impact phases (Caufriez, 2005; Kapandji, 1977). Further details about the jumping/breathing technique and training can be obtained from co-authors LO and GA. In Figure 2b the EMG activity of rectus and obliquus abdominis muscles, together with the expired volume, are shown during normal and 'controlled' jumps.

2.3 MECHANICAL MODEL

The method presented by Minetti and Belli is based on a model made up of a container with mass M , incorporating a hidden mass m (the visceral content), which oscillates periodically in the vertical or horizontal direction. In line with the original paper, we considered just vertical motion but included an 'external' wobbling mass

163 (m_e), representing mainly pectoral muscles and abdominal wall, as part of the
 164 container (see Figure 3). The new equation of motion is:

$$165 \quad (M + m + m_e)\ddot{y}_{CoM}(t) = F_v(t) - (M + m + m_e)g \quad (1)$$

166 which results from the system of equations:

$$167 \quad \begin{cases} M\ddot{y}_1(t) = F_v(t) - Mg - f_v(t) - f_e(t) \\ m\ddot{y}_2(t) = f_v(t) - mg \\ m_e\ddot{y}_3(t) = f_e(t) - m_e g \end{cases} \quad (2)$$

168

169 where F_v is the vertical component of GRF, f_v and f_e are vertical forces (unknown)
 170 exerted by the internal and 'external' masses, and y_1 , y_2 and y_3 are distances from
 171 ground level of the container, visceral mass and external mass.

172 In literature, the magnitude of the internal visceral mass ' m ' is estimated to be 16% of
 173 body mass (Martin, Janssens, Caboor, Clarys, & Marfell-Jones, 2003), while the
 174 external wobbling mass ' m_e ' is evaluated to be 4% of body mass (Burkhart, Arthurs,
 175 & Andrews, 2008).

176

177 2.4 DATA PROCESSING

178 A bespoke written software (LABVIEW 8.6, National Instrument, US) was developed
 179 to calculate the visceral mass vertical displacement, as shown in the equation (3),

180

$$\begin{aligned}
s(t) - s_0 = & \frac{(M + m + m_e)}{m} \left\{ \left[\int_0^t \left(\int_0^t \left(\frac{F_v(t)}{M + m + m_e} - g \right) dt \right) dt \right. \right. \\
& - \frac{t}{T} \int_0^T \left(\int_0^t \left(\frac{F_v(t)}{M + m + m_e} - g \right) dt \right) dt \Big] \\
& - \left(\frac{M + m}{M + m + m_e} \right) [y_1(t) - y_1(0)] \\
& \left. - \left(\frac{m_e}{M + m + m_e} \right) [y_3(t) - y_3(0)] \right\}
\end{aligned}$$

181

182 (3)

183 where “T” is the movement period and “t” the progressive time.

184 This method and its algorithm were validated by loading in our program the kinetic
 185 data obtained from a simulation software (Visual Nastran 4D, MSC Software) of a
 186 known mechanical model (oscillating cylinder containing a sphere linked to the
 187 ceiling by a spring).

188 The developed software automatically recognized and isolate every jump (jump cycle
 189 = time between two subsequent BCoM peaks), double integrated (trapezoidal rule) the
 190 net GRF, and downsampled displacement data from 1200 Hz to 400 Hz to match the
 191 sampling rate of the motion capture system. GRF signal was shifted backward to
 192 cover a time gap ($=2\Delta t/2=\Delta t$) due to double integration, to synchronize these data
 193 with kinematic acquisition. Force signal and kinematic data were filtered forward and
 194 backward by a 3rd order zero-lag low-pass Butterworth filter with cut-off frequency of
 195 30 Hz (Bisseling & Hof, 2006). The frequency of the input signal (GRF), f_{GRF} , was
 196 used to compare the dynamics of subjects' jumps (Boyer & Nigg, 2007) and its value
 197 was estimated by using the input peak value of the F_v , and the average loading rate
 198 between the 20% and 80% of the impact phase ($G_{v,ave}$), as:

$$f_{GRF} = \frac{1}{2(F_v/G_{v,ave})}$$

3. RESULTS

The biomechanical model chosen in this work allows an accurate BCoM estimation in locomotion (Halvorsen, Eriksson, Gullstrand, Tinmark, & Nilsson, 2009), and its adoption in jumping shows an error comparable to the literature. Indeed, two validation indices were estimated during the flight phase of the jumps: AV_1 (m/s^2) index represents an estimation of the gravity constant acceleration (g), expected to be $9.81 m/s^2$, while AV_2 (m) index is defined as the root mean square error among the model estimated and matched ballistic centre of mass trajectory (Rabuffetti & Baroni, 1999). Their overall mean values and s.d. are respectively AV_1 (m/s^2) = -9.836 ± 0.027 , AV_2 (m) = 0.003 ± 0.002 .

In Table 1 the results of all the experiments are shown. The visceral mass (VMD), pectoral and abdomen external mass displacements (EMD) are represented as relative to the BCoM. The VMD, for all the subjects, measured during normal jumps (0.069 ± 0.020 m), is significantly higher ($p < 0.05$, paired t-test), than in controlled jumps (0.053 ± 0.018 m). The average time courses of normal and controlled VMD are shown in Figure 4, while the mean individual curves of participants are displayed in Figure 5.

For all the subjects, VMD shows a different pattern with respect to the container displacement both in normal and in controlled jumps (Figure 4), with a detectable phase shift between the curves. A paired t-test shows no significant difference of time shift, both during the aerial (normal 50.6 ± 10.4 ms – controlled 49.3 ± 9.4 ms) and landing (normal 51.2 ± 14.4 ms - controlled 49.8 ± 8.8 ms) phases, confirming a

constant phase shift in both jumping techniques. A local maximum in visceral mass displacement ($\dot{s}(t) = 0$) is detectable at about 40-45% of jump period (time between two subsequent BCoM peaks) (Figure 4) and could be classified as a typical artefact of the foot impact on the force platform (Bisseling & Hof, 2006). The pectoral and abdominal EMD values show no significant difference in the two jumping techniques (paired t-test), but the pectoral EMD is significantly larger ($p < 0.05$, paired t-test) than the abdomen EMD in both techniques (Figure 6).

Pectoral and abdomen EMD show a different pattern with respect to BCoM oscillation and VMD. Finally, a non-significant difference of f_{GRF} , jumping frequency (f_{jump}), BCoM vertical excursion and contact time (t_c) between the techniques (Table 1), for all the subjects, reveals a comparable dynamic and kinematic of normal and controlled jumps.

4. DISCUSSION

The aim of this investigation was to test the effect of a combined respiratory/jumping strategy, properly designed for compacting viscera in the abdominal cavity, in limiting the vertical viscera motion during vertical jumps. Applying a previously developed method (A. Minetti & Belli, 1994), by concurrently using inverse and direct dynamics, we revealed that such a strategy reduced the vertical excursion up to 30%, with potential increases of the overall stiffness of the human trunk/body.

The VMD mean value measured was comparable with the literature: few quantitative analyses were conducted mostly anatomically (Beillas, Lafon, & Smith, 2009) or in slow-dynamic condition (Hostettler, Nicolau, Remond, Marescaux, & Soler, 2010), where vertical viscera motion was found to range between 0.03 m and 0.07 m. Only Minetti & Belli reported a value related to submaximal repeated jumps (0.08 m),

247 while Boussuges and collaborators (Boussuges, Gole, & Blanc, 2009) set the limit of
248 vertical displacement on maximal diaphragm motion (0.070 ± 0.011 m).

249 Regarding to the ‘controlled’ technique execution, experimental evidences of higher
250 abdominal muscle activation and comparable expiration volume (Figure 2) proved
251 that a voluntary diaphragm activation can be inferred: the volume of expired air
252 during the controlled jump sequence was small and comparable with the normal jump,
253 despite of a higher activation of expiratory muscles (obliquus and rectus abdominis),
254 implying that the diaphragm applied an opposite force to contrast the rising viscera. In
255 terms of interaction between respiration and movement, our results show that muscles
256 not directly involved in jumping could affect body dynamics, and stress their potential
257 effect on motor acts where locomotor/respiratory coupling-ratios can occur.

258 In the literature several authors have already speculated about frequency and phase
259 coupling between respiratory and locomotory rhythms as affected by training
260 (Bernasconi & Kohl, 1993) or workload (Rassler & Kohl, 1996), but no one provided
261 evidences of voluntary control of internal body dynamics through specific respiration
262 techniques, synchronously performed with body CoM oscillations. Only McDermott
263 (McDermott, et al., 2003), by investigating the relationship between
264 locomotor/respiratory coupling and training level, found that expert runners were
265 particularly skilled in synching their coupling during speed changes. Therefore, from
266 the energetic point of view, these interactions should be controlled to avoid energy
267 losses resulting in some extra-mechanical work done by muscles, and the time delay
268 calculated between BCoM and VMD curves in this investigation, reinforces this
269 hypothesis. In fact, the ‘economy’ of bouncing locomotion, such as running or
270 skipping, could be influenced and the mechanical external work calculated from
271 kinematically measured CoM displacement could be refined by adding viscera

272 contribution (Daley & Usherwood, 2010). While this is supposed to be a small
273 adjustment in normal subjects, any deviation from a mesomorphic body such as obese
274 patients with relevant internal and external wobbling masses would involve a more
275 substantial correction of the inverse dynamics approach. In this way the proposed
276 respiratory strategy could give potential benefits in terms of movement performance
277 and the non-invasive method described could be easily adopted.

278 In terms of data processing the previous method (Minetti & Belli, 1994) has been
279 refined: kinematic sampling frequency has been quadrupled (400 Hz) and chosen as a
280 submultiple of the dynamometric signal to facilitate synchronization, the signals were
281 accurately aligned (double integration time gap), and the mathematical model was
282 validated with physics laboratory simulation software. Besides, the method still
283 suffered of inaccuracies due to: 1) the rigid body model assumption (Cappozzo, Della
284 Croce, Leardini, & Chiari, 2005; Chiari, Della Croce, Leardini, & Cappozzo, 2005)
285 originating troublesome theoretical interpretations of the results: the discrepancy
286 between the BCoM estimates from direct and inverse dynamics is considered as an
287 indirect evidence of viscera motion, but this could be partially the results of
288 experimental inaccuracies, 2) the “skin marker artefact” (Cappozzo, Catani, Leardini,
289 Benedetti, & Croce, 1996), which particularly affects movements with considerable
290 joint rotation as sit-to-stand (Kuo, et al., 2011) or locomotion (Akbarshahi, et al.,
291 2010) rather than vertical jumps with the arms blocked on the trunk, 3) the “soft tissue
292 motion artefact” (Gruber, et al., 1998; Leardini, et al., 2005), which can be assessed
293 by accelerometers (Kitazaki & Griffin, 1995) or by adding extra markers for the
294 oscillating body parts, at the cost of a more complex biomechanical model. The 4
295 'technical' markers introduced here, positioned on the estimated centre of mass of the
296 most visible and bulky 'external' wobbling masses (pectorals and abdominal muscles),

297 allowed their movement to contribute to refine VMD estimation. This simplified
298 approach does not completely compensate for the rigid body assumption inaccuracies
299 and cannot separate viscera from limbs soft tissues contribution (Gunther, et al.,
300 2003), but it constitutes an acceptable trade-off between ideal VMD estimation and
301 practical feasibility.

302 A further variable affecting VMD and EMD measure is the muscle tuning during
303 jumping: the ‘controlled jump’ is comparable with a tuned landing thanks to an higher
304 pectoral and abdominal muscles activation and could decrease the absolute and
305 relative acceleration of the soft tissue compartments (Boyer & Nigg, 2006). Even
306 though a further frequency analysis of external masses acceleration signal (not
307 measured in this work) could reveal soft tissues vibrational changes between the
308 techniques, pectoral and abdominal EMD are not significantly different (Table 1), and
309 their patterns are similar in normal and controlled jumps (Figure 6). This is probably
310 due to similar pectoral-muscle activation in both techniques, and to a peculiar muscle
311 tuning effect on abdominal soft tissue: actually its vibration could be less influenced
312 by muscle contraction than other soft tissues (upper/lower limbs) because of its
313 anatomical characteristics and local physical constrains.

314 To date, soft tissues influences has already been investigated in locomotion (DeVita,
315 et al., 2007; Zelik & Kuo, 2010) and in jump landing (Gittoes, et al., 2006; M. T. Pain
316 & Challis, 2006), though its role still needs to be ultimately assessed. In this work,
317 even if there are several limitations, we compared two refined estimations of the most
318 influent soft tissue (viscera) motion in a simple motor task, repeatedly executed in the
319 same experimental condition. Indeed, subjects executed comparable jumps
320 considering the jumping frequency (f_{jump}), contact time (t_c), frequency of input force
321 (f_{GRF}) and the performance (body CoM vertical excursion). These evidences help to

minimize systematic and random errors, showing a de-noised measure of viscera vertical excursion.

In conclusion, the combination of the inverse/direct dynamics method to measure viscera motion and a novel respiration assisted jumping technique reveals, for the first time, that the vertical displacement of the abdominal wobbling mass can be modulated also in dynamic condition. Moreover, it has been demonstrated that the accuracy of this refined method is adequate to detect, with a non-invasive approach, the effects of internal forces on the kinematic of the visceral mass and could be adopted to evaluate those their impact in sport biomechanics and locomotion energetics. The results and the proposed jumping strategy could then constitute a prerequisite for further studies assessing the potential performance enhancement in a variety of motor acts.

ACKNOWLEDGEMENTS

The authors thank Dr Matteo Pecchiari for his invaluable experience in respiration physiology, and Dr Ezio Preatoni for his precious suggestions during paper revision.

REFERENCES

- Akbarshahi, M., Schache, A. G., Fernandez, J. W., Baker, R., Banks, S., & Pandy, M. G. (2010). Non-invasive assessment of soft-tissue artifact and its effect on knee joint kinematics during functional activity. *J Biomech*, 43, 1292-1301.
- Alexander, R. M. (1993). Breathing while trotting. *Science*, 262, 196-197.
- Beillas, P., Lafon, Y., & Smith, F. W. (2009). The effects of posture and subject-to-subject variations on the position, shape and volume of abdominal and thoracic organs. *Stapp Car Crash J*, 53, 127-154.
- Bernasconi, P., & Kohl, J. (1993). Analysis of co-ordination between breathing and exercise rhythms in man. *J Physiol*, 471, 693-706.
- Bisseling, R. W., & Hof, A. L. (2006). Handling of impact forces in inverse dynamics. In *J Biomech* (Vol. 39, pp. 2438-2444). United States.
- Bonsignore, M. R., Morici, G., Abate, P., Romano, S., & Bonsignore, G. (1998). Ventilation and entrainment of breathing during cycling and running in triathletes. *Med Sci Sports Exerc*, 30, 239-245.
- Boussuges, A., Gole, Y., & Blanc, P. (2009). Diaphragmatic motion studied by m-mode ultrasonography: methods, reproducibility, and normal values. In *Chest* (Vol. 135, pp. 391-400). United States.
- Boyer, K. A., & Nigg, B. M. (2006). Muscle tuning during running: implications of an un-tuned landing. *J Biomech Eng*, 128, 815-822.
- Boyer, K. A., & Nigg, B. M. (2007). Quantification of the input signal for soft tissue vibration during running. In *J Biomech* (Vol. 40, pp. 1877-1880). United States.
- Bramble, D. M., & Carrier, D. R. (1983). Running and breathing in mammals. *Science*, 219, 251-256.
- Brown, R. E., Lee, H. T., & Loring, S. H. (2004). Airflow synchronous with oscillatory acceleration reflects involuntary respiratory muscle activity. In *Respir Physiol Neurobiol* (Vol. 140, pp. 265-282). Netherlands: 2004 Elsevier B.V.
- Burkhart, T. A., Arthurs, K. L., & Andrews, D. M. (2008). Reliability of upper and lower extremity anthropometric measurements and the effect on tissue mass predictions. In *J Biomech* (Vol. 41, pp. 1604-1610). United States.

- 372 Calais-Germain, B. (2005). *Respiration anatomie-geste respiratoire*.
- 373 Cappozzo, A., Catani, F., Leardini, A., Benedetti, M. G., & Croce, U. D. (1996).
 374 Position and orientation in space of bones during movement:
 375 experimental artefacts. *Clin Biomech (Bristol, Avon)*, 11, 90-100.
- 376 Cappozzo, A., Della Croce, U., Leardini, A., & Chiari, L. (2005). Human movement
 377 analysis using stereophotogrammetry. Part 1: theoretical background.
 378 *Gait Posture*, 21, 186-196.
- 379 Caufriez, M. (2005). *Respiration anatomie-geste respiratoire* (Editions DesIris
 380 ed.).
- 381 Challis, J. H., & Pain, M. T. (2008). Soft tissue motion influences skeletal loads
 382 during impacts. In *Exerc Sport Sci Rev* (Vol. 36, pp. 71-75). United States.
- 383 Chen, S. J., Mukul, M., & Chou, L. S. (2011). Soft-tissue movement at the foot
 384 during the stance phase of walking. In *J Am Podiatr Med Assoc* (Vol. 101,
 385 pp. 25-34). United States.
- 386 Chiari, L., Della Croce, U., Leardini, A., & Cappozzo, A. (2005). Human movement
 387 analysis using stereophotogrammetry. Part 2: instrumental errors. *Gait*
 388 *Posture*, 21, 197-211.
- 389 Clancy, E. A., Morin, E. L., & Merletti, R. (2002). Sampling, noise-reduction and
 390 amplitude estimation issues in surface. *J Electromyogr Kinesiol*, 12, 1-16.
- 391 Daley, M. A., & Usherwood, J. R. (2010). Two explanations for the compliant
 392 running paradox: reduced work of bouncing viscera and increased
 393 stability in uneven terrain. In *Biol Lett* (Vol. 6, pp. 418-421). England.
- 394 Dempster, W. T., Gabel, W. C., & Felts, W. J. (1959). The anthropometry of the
 395 manual work space for the seated subject. *Am J Phys Anthropol*, 17, 289-
 396 317.
- 397 DeVita, P., Helseth, J., & Hortobagyi, T. (2007). Muscles do more positive than
 398 negative work in human locomotion. In *J Exp Biol* (Vol. 210, pp. 3361-
 399 3373). England.
- 400 Friesenbichler, B., Stirling, L. M., Federolf, P., & Nigg, B. M. (2011). Tissue
 401 vibration in prolonged running. In *J Biomech* (Vol. 44, pp. 116-120).
 402 United States: 2010 Elsevier Ltd.

- 403 Gao, B., & Zheng, N. N. (2008). Investigation of soft tissue movement during level
404 walking: translations and rotations of skin markers. In *J Biomech* (Vol. 41,
405 pp. 3189-3195). United States.
- 406 Gittoes, M. J., Brewin, M. A., & Kerwin, D. G. (2006). Soft tissue contributions to
407 impact forces simulated using a four-segment wobbling mass model of
408 forefoot-heel landings. In *Hum Mov Sci* (Vol. 25, pp. 775-787).
409 Netherlands.
- 410 Gruber, K., Ruder, H., Denoth, J., & Schneider, K. (1998). A comparative study of
411 impact dynamics: wobbling mass model versus rigid body models. In *J*
412 *Biomech* (Vol. 31, pp. 439-444). United States.
- 413 Gunther, M., Sholukha, V. A., Kessler, D., Wank, V., & Blickhan, R. (2003). Dealing
414 with skin motion and wobbling masses in inverse dynamics. *J Mech in Med*
415 *and Bio (JMMB)*, 3, 309-335.
- 416 Halvorsen, K., Eriksson, M., Gullstrand, L., Tinmark, F., & Nilsson, J. (2009).
417 Minimal marker set for center of mass estimation in running. *Gait Posture*,
418 30, 552-555.
- 419 Hostettler, A., Nicolau, S. A., Remond, Y., Marescaux, J., & Soler, L. (2010). A real-
420 time predictive simulation of abdominal viscera positions during quiet
421 free breathing. In *Prog Biophys Mol Biol*: 2010 Elsevier Ltd.
- 422 Kapandji, I. A. (1977). *Fisiologia articolare - Tronco e Rachide* (Vol. 3). Rome.
- 423 Kitazaki, S., & Griffin, M. J. (1995). A data correction method for surface
424 measurement of vibration on the human body. *J Biomech*, 28, 885-890.
- 425 Kuo, M. Y., Tsai, T. Y., Lin, C. C., Lu, T. W., Hsu, H. C., & Shen, W. C. (2011).
426 Influence of soft tissue artifacts on the calculated kinematics and kinetics
427 of total knee replacements during sit-to-stand. *Gait Posture*, 33, 379-384.
- 428 Le Boulch, J. (1973). *L'éducation par le mouvement* (12ème édition ed.). Paris.
- 429 Leardini, A., Chiari, L., Della Croce, U., & Cappozzo, A. (2005). Human movement
430 analysis using stereophotogrammetry. Part 3. Soft tissue artifact
431 assessment and compensation. *Gait Posture*, 21, 212-225.
- 432 Loring, S. H., Lee, H. T., & Butler, J. P. (2001). Respiratory effects of transient axial
433 acceleration. *J Appl Physiol*, 90, 2141-2150.
- 434 Lumb, A. B. (2005). Nunn's Applied Respiratory Physiology. In (Sixth Editions
435 ed., pp. 76-80).

- 436 Martin, A. D., Janssens, V., Caboor, D., Clarys, J. P., & Marfell-Jones, M. J. (2003).
 437 Relationships between visceral, trunk and whole-body adipose tissue
 438 weights by cadaver dissection. In *Ann Hum Biol* (Vol. 30, pp. 668-677).
 439 England.
- 440 McDermott, W. J., Van Emmerik, R. E., & Hamill, J. (2003). Running training and
 441 adaptive strategies of locomotor-respiratory coordination. *Eur J Appl*
 442 *Physiol*, 89, 435-444.
- 443 Mian, O., Thom, J., Ardigò, L., Narici, M., & Minetti, A. (2006). Metabolic cost,
 444 mechanical work, and efficiency during walking in young and older men.
 445 *Acta Physiol (Oxf)*, 186, 127-139.
- 446 Mills, C., Scurr, J., & Wood, L. (2011). A protocol for monitoring soft tissue motion
 447 under compression garments during drop landings. In *J Biomech* (Vol. 44,
 448 pp. 1821-1823). United States: 2011 Elsevier Ltd.
- 449 Minetti, A., & Belli, G. (1994). A model for the estimation of visceral mass
 450 displacement in periodic movements. *J Biomech*, 27, 97-101.
- 451 Minetti, A. E., Cisotti, C., & Mian, O. S. (2011). The mathematical description of the
 452 body centre of mass 3D path in human and animal locomotion. In *J*
 453 *Biomech* (Vol. 44, pp. 1471-1477). United States: 2011 Elsevier Ltd.
- 454 Pain, M. T., & Challis, J. H. (2006). The influence of soft tissue movement on
 455 ground reaction forces, joint torques and joint reaction forces in drop
 456 landings. In *J Biomech* (Vol. 39, pp. 119-124). United States.
- 457 Pain, M. T. G., & Challis, J. H. (2004). Wobbling mass influence on impact ground
 458 reaction forces: A simulation model sensitivity analysis. *Journal of Applied*
 459 *Biomechanics*, 20(3), 309-316.
- 460 Rassler, B., & Kohl, J. (1996). Analysis of coordination between breathing and
 461 walking rhythms in humans. In *Respir Physiol* (Vol. 106, pp. 317-327).
 462 Netherlands.
- 463 Saibene, F., & Minetti, A. E. (2003). Biomechanical and physiological aspects of
 464 legged locomotion in humans. *Eur J Appl Physiol*, 88, 297-316.
- 465 Simons, R. S. (1999). Running, breathing and visceral motion in the domestic
 466 rabbit (*Oryctolagus cuniculus*): testing visceral displacement hypotheses.
 467 *J Exp Biol*, 202, 563-577.

- 468 Wakeling, J. M., & Nigg, B. M. (2001). Soft-tissue vibrations in the quadriceps
469 measured with skin mounted transducers. In *J Biomech* (Vol. 34, pp. 539-
470 543). United States.
- 471 Willems, P. A., Cavagna, G. A., & Heglund, N. C. (1995). External, internal and total
472 work in human locomotion. *J Exp Biol*, 198, 379-393.
- 473 Wilson, T. A., & Liu, S. (1994). Effect of acceleration on the chest wall. *J Appl*
474 *Physiol*, 76, 1242-1246.
- 475 Yu, B., & Hay, J. G. (1996). Optimum phase ratio in the triple jump. In *J Biomech*
476 (Vol. 29, pp. 1283-1289). United States.
- 477 Yue, Z., & Mester, J. (2002). A model analysis of internal loads, energetics, and
478 effects of wobbling mass during the whole-body vibration. In *J Biomech*
479 (Vol. 35, pp. 639-647). United States.
- 480 Zelik, K. E., & Kuo, A. D. (2010). Human walking isn't all hard work: evidence of
481 soft tissue contributions to energy dissipation and return. In *J Exp Biol*
482 (Vol. 213, pp. 4257-4264). England.
- 483
- 484
- 485

Figure 1: Human body modelled with 22 reflective markers and 14 segments: head (1), trunk (2), abdomen (4), right upper arm (5), left upper arms (6), right fore arm (7), left fore arm (8), right thigh (9), left thigh (10), right shank (11), left thigh (12), right foot (13), left foot (14), and pectoral muscles (3).

Figure 2: (a) Mechanism used to generate an intra-abdominal pressure that compacts the visceral mass: the subject after a combined deep diaphragmatic inspiration and contraction of the abdominal “press” increases the intra-abdominal pressure also executing progressive and short exhalations. The black arrows indicate: (1) The lowering of the diaphragm that pushes on the viscera during inspiration (downward-pointing white arrow); (2) The musculature of the abdominal “press”, which contraction contributes to the elevation of intra-abdominal pressure (upward-pointing white arrows). (b) On the left the overall mean (normalized in respect of the maximal contraction value) and s.d of all the subjects, of rectus and obliquus abdominis muscle activation, in normal (light-grey) and controlled (dark-grey) jump are shown. The rectus and obliquus muscle activation is significantly higher in controlled jumps ($* = p < 0.01$). On the right the overall mean and s.d., of the expired volume (V) during a jump are shown. The expired volume is not significantly different between the techniques.

Figure 3: Model used for the estimation of visceral mass displacement: M is the container mass, m the internal visceral mass, and m_e is the external mass, while y_1 , y_2 and y_3 are distances from ground level and $s=y_2-y_1$. The whole system oscillates vertically and exerts a vertical ground reaction force F_v , while internal and external mass exerts a force f_v and f_e respectively on the container.

Figure 4: The overall mean curve of VMD (visceral mass displacement) in normal (grey solid line) and controlled (grey dashed line) jumps, and overall mean curve (controlled and normal) of body CoM (black solid line) are shown. All the curves are time-normalized with single jump duration (0-100%).

Figure 5: The mean of all the trials curves (5 trial of at least 15 jumps for every subject), presented with black bold line, and their variability (s.d. of all the trials curves), presented with light grey lines, are shown for both techniques (normal and controlled) for each subject (S1, S2, S3, S4, S5, S6). The curves are time-normalized with single jump duration.

Figure 6: The overall mean curve of pEMD (pectoral external mass displacement) in normal (black solid line) and controlled (black dashed line) jumps, the overall mean

curve of aEMD (abdominal external mass displacement) in normal (grey solid line) and controlled (grey dashed line) jumps, and the overall mean curve (controlled and normal) of body CoM (black dotted line). All the curves are time-normalized with single jump duration (0-100%). The pEMD and aEMD, for all the subjects, are not significantly different in the two techniques, but the pEMD is significantly higher ($p < 0.05$) than aEMD both in normal and in controlled jumps.

Table 1: The mean and s.d. values of (1) visceral mass displacement (VMD), (2) body CoM displacement (CoM), (3) pectorals (overall mean of right and left) external mass displacement (pEMD), (4) abdomen (overall mean of right and left) external mass displacement (EMD), (5) estimated input frequency (fGRF), (6) jumping frequency (f_{jump}) and (7) contact time (t_c) in “normal” and “controlled” jumps are presented for every subject.

Table 1

JUMP type	Subject	N		VMD (m)	CoM (m)	pEMD (m)	aEMD (m)	f _{GRF} (Hz)	f _{jump} (Hz)	t _c (s)
Normal	S1	76	Mean	0.073	0.209	0.030	0.016	7.13	2.40	0.106
			SD	0.015	0.019	0.008	0.007	0.69	0.02	0.003
	S2	70	Mean	0.089	0.347	0.042	0.026	6.79	1.66	0.114
			SD	0.005	0.049	0.009	0.008	0.51	0.07	0.003
	S3	85	Mean	0.059	0.168	0.031	0.010	8.42	1.96	0.101
			SD	0.005	0.007	0.010	0.006	0.30	0.13	0.003
	S4	85	Mean	0.056	0.216	0.029	0.018	7.67	2.09	0.109
			SD	0.008	0.026	0.008	0.008	0.53	0.06	0.006
	S5	71	Mean	0.102	0.311	0.040	0.024	7.21	1.82	0.098
			SD	0.005	0.013	0.008	0.009	0.33	0.03	0.001
	S6	90	Mean	0.051	0.137	0.049	0.041	6.76	2.65	0.067
			SD	0.006	0.008	0.010	0.010	0.67	0.10	0.002
	All	477	Mean	0.069	0.219	0.037	0.023	7.35	2.09	0.099
			SD	0.020	0.075	0.009	0.011	0.79	0.34	0.015
Controlled	S1	80	Mean	0.051	0.161	0.028	0.012	7.95	2.37	0.100
			SD	0.008	0.011	0.009	0.005	0.46	0.03	0.003
	S2	72	Mean	0.078	0.321	0.026	0.026	6.88	1.81	0.104
			SD	0.007	0.029	0.008	0.008	0.41	0.03	0.002
	S3	92	Mean	0.049	0.171	0.021	0.012	8.28	2.28	0.101
			SD	0.006	0.010	0.009	0.009	0.53	0.29	0.002
	S4	86	Mean	0.046	0.242	0.031	0.013	7.59	2.36	0.096
			SD	0.009	0.036	0.010	0.009	0.56	0.03	0.001
	S5	69	Mean	0.076	0.306	0.040	0.019	7.02	1.80	0.103
			SD	0.010	0.013	0.011	0.008	0.25	0.02	0.004
	S6	93	Mean	0.030	0.155	0.046	0.038	6.78	2.74	0.069
			SD	0.004	0.011	0.010	0.010	0.68	0.03	0.001
	All	492	Mean	0.053	0.217	0.032	0.020	7.46	2.21	0.097
			SD	0.018	0.069	0.009	0.010	0.77	0.35	0.012

Figure 1
[Click here to download high resolution image](#)

Figure 2
[Click here to download high resolution image](#)

Figure 3
[Click here to download high resolution image](#)

Figure 4
[Click here to download high resolution image](#)

Figure 6
[Click here to download high resolution image](#)

Figure 5
[Click here to download high resolution image](#)