Citation for published version:

Power, J 2011, 'Unicity of Enrichment over Cat or Gpd', *Applied Categorical Structures*, vol. 19, no. 1, pp. 293-299. <https://doi.org/10.1007/s10485-009-9202-7>

DOI:

[10.1007/s10485-009-9202-7](https://doi.org/10.1007/s10485-009-9202-7)

Publication date:

2011

[Link to publication](#)

The original publication is available at www.springerlink.com

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Unicity of Enrichment over Cat or Gpd

John Power
Department of Computer Science
University of Bath
Bath, BA2 7AY
United Kingdom
Email: A.J.Power@bath.ac.uk

Abstract

Enrichment of ordinary monads over Cat or Gpd is fundamental to Max Kelly's unified theory of coherence for categories with structure. So here, we investigate existence and unicity of enrichments of ordinary functors, natural transformations, and hence also monads, over Cat and Gpd . We show that every ordinary natural transformation between 2-functors whose domain 2-category has either tensors or cotensors with the arrow category is 2-natural. We use that to prove that an ordinary monad, or endofunctor, on such a 2-category has at most one enrichment over Cat or Gpd . We also describe a monad on Cat that has no enrichment. So enrichment over Cat is a non-trivial property of a monad rather than a structure that is additional to it. Finally, we present an example, due to Kelly, of V other than Cat or Gpd and an ordinary monad for which more than one enrichment over V exists, showing that our main theorem is specific to Cat and Gpd .

1 Introduction

A central theme, perhaps the central theme, of Max Kelly's research was the study of coherence for categories with algebraic structure. His most prominent paper on the topic was [1], complementing his definitive book on enriched categories [4]. My own scientific career has been founded on my four years, 1984-88, as his research fellow working on the topic, a fact for which I am profoundly grateful. Max's ideas have now become embedded into theoretical computer science (see [7] for an overview and [8] for a recent application of them).

In the Australian summer of 1991/92, Max invited me to Sydney to continue our work on coherence for categories with algebraic structure. He asked the following question: given an ordinary monad T on Cat , could there be at most one enrichment of T over Cat ? A variant of the question, cohering with his research on coherence, was whether an ordinary monad on Cat can have at most one enrichment over the cartesian closed category Gpd of groupoids.

We knew that there are monads on Cat that admitted no enrichment: an example appears in Section 3. But is there at most one? The question was a natural one for Max to ask, coming after his proof, with Foltz and Lair, that Cat possesses precisely two symmetric monoidal closed structures [2], and coming before his investigation with Lack of property-like structures on Cat [6]. A positive answer would indicate that 2-monadicity is a property of a monad on Cat rather than a genuine additional structure.

In fact, the answer is Yes, the answer following from an existence result with unicity implicit: if a 2-category A admits tensors or cotensors with the arrow category (see [5]), given any pair of 2-functors of the form $F, G : A \longrightarrow B$, any ordinary natural transformation $\alpha : F_0 \Rightarrow G_0 : A_0 \longrightarrow B_0$ is necessarily 2-natural. We prove that result in Section 2.

Having proved the above result, we proceed to Max's uniqueness question. The answer to his question can be framed in a more general setting than that in which he originally asked it: given any 2-category A satisfying a very mild cocompleteness or completeness condition, namely the existence of either tensors or cotensors with the arrow category, any monad on A possesses at most one enrichment over Cat , with the same applying for enrichment over Gpd . Note that there is no assumption of rank on the monad, and the base 2-category need not be either complete or cocomplete. We deduce the result in Section 3.

Inevitably, having proved the result, the question arises whether the same is true for enrichment over a more general class of symmetric monoidal closed categories V . So, in Section 4, we present a counter-example, due to Max, to show that the argument does not extend, i.e., we give a V other than Cat or Gpd together with a monad with two possible enrichments over V .

2 Existence of Enrichment over Cat or Gpd of Ordinary Natural Transformations

In this section, we show that if a 2-category A admits tensors or cotensors with the arrow category $\rightarrow$, then for any 2-functors $F, G : A \longrightarrow B$, any natural transformation

$$\alpha : F_0 \Rightarrow G_0 : A_0 \longrightarrow B_0$$

is 2-natural.

We first observe that, in order to prove the result, we do need a condition on A , i.e., if we drop the assumption on A that it admits either tensors or cotensors with $\rightarrow$, there may exist natural transformations α that are not 2-natural.

Example 2.1 *Let A be the free 2-category on a 2-cell. So A has two objects, which we denote by 0 and 1, two 1-cells $x, y : 0 \longrightarrow 1$, and a 2-cell $\gamma : x \Rightarrow y$, but no other non-trivial 1-cells or 2-cells.*

Let $\parallel$ denote the category with two objects X and Y and two maps $f, g : X \longrightarrow Y$, i.e., $\parallel$ is the free category on a pair of parallel maps.

Define $F, G : A \longrightarrow \text{Cat}$ by putting

$$\begin{aligned} F0 &= G0 = 1 & F1 &= G1 = \parallel \\ Fx &= Gx = X & Fy &= Gy = Y \\ F\gamma &= f & G\gamma &= g \end{aligned}$$

Then $F_0 = G_0$, so the identity is a natural transformation from F_0 to G_0 . But the identity is not a 2-natural transformation as F does not equal G .

We now proceed to the main result of the section via some lemmas.

Lemma 2.2 [The main lemma] *Given any categories C and D , consider functors $f : C \longrightarrow D$ and $g : C^\rightarrow \longrightarrow D^\rightarrow$ such that the following three diagrams in Cat commute:*

$$\begin{array}{ccc} \begin{array}{ccc} C^\rightarrow & \xrightarrow{g} & D^\rightarrow \\ \text{dom} \downarrow & & \downarrow \text{dom} \\ C & \xrightarrow{f} & D \end{array} & \begin{array}{ccc} C^\rightarrow & \xrightarrow{g} & D^\rightarrow \\ \text{cod} \downarrow & & \downarrow \text{cod} \\ C & \xrightarrow{f} & D \end{array} & \begin{array}{ccc} C^\rightarrow & \xrightarrow{g} & D^\rightarrow \\ \delta \uparrow & & \uparrow \delta \\ C & \xrightarrow{f} & D \end{array} \end{array}$$

Then $g = f^\rightarrow$.

Proof Consider the square

$$\begin{array}{ccc} x & \xrightarrow{\rho} & y \\ \alpha \downarrow & & \downarrow \beta \\ u & \xrightarrow{\sigma} & v \end{array} \quad (1)$$

as an arrow $(\rho, \sigma) : \alpha \longrightarrow \beta$ in $C^\rightarrow$. The commutativities above involving dom and cod ensure that $g(\rho, \sigma) : g\alpha \longrightarrow g\beta$ has the form

$$\begin{array}{ccc} fx & \xrightarrow{f\rho} & fy \\ g\alpha \downarrow & & \downarrow g\beta \\ fu & \xrightarrow{f\sigma} & fv \end{array} \quad (2)$$

The commutativity involving δ ensures that

$$g1_x = 1_{fx} \quad (3)$$

Now, take for Diagram 1 the diagram

$$\begin{array}{ccc} x & \xrightarrow{\rho} & y \\ \rho \downarrow & & \downarrow id \\ y & \xrightarrow{id} & y \end{array}$$

In the light of Equation 3, Diagram 2 yields commutativity of the diagram:

$$\begin{array}{ccc} fx & \xrightarrow{f\rho} & fy \\ g\rho \downarrow & & \downarrow id \\ fy & \xrightarrow{id} & fy \end{array}$$

so $g\rho = f\rho$, so $g = f^{\rightarrow}$ as desired.

Proposition 2.3 *Let $F, G : A \rightarrow B$ be 2-functors, where the 2-categories A and B admit tensors with the arrow category $\rightarrow$. Then, for any natural transformation $\alpha : F_0 \Rightarrow G_0$, the diagram*

$$\begin{array}{ccc} \rightarrow \otimes FX & \xrightarrow{\rightarrow \otimes \alpha_X} & \rightarrow \otimes GX \\ (\tau_F)_X \downarrow & & \downarrow (\tau_G)_X \\ F(\rightarrow \otimes X) & \xrightarrow{\alpha_{\rightarrow \otimes X}} & G(\rightarrow \otimes X) \end{array} \quad (4)$$

commutes, where τ_F and τ_G are the canonical comparison maps induced by the 2-functoriality of F and G respectively.

Proof Given any 0-cell Y in B , consider the two composites

$$A(\rightarrow \otimes X, Y) \xrightarrow{G} B(G(\rightarrow \otimes X), GY) \xrightarrow{B(f, GY)} B(\rightarrow \otimes FX, GY)$$

generated by the two maps from $\rightarrow \otimes FX$ to $G(\rightarrow \otimes X)$ given by Diagram 4. By the enriched Yoneda lemma, the two legs of Diagram 4 are equal if and only if these two composites are equal. But pre- and post-composing the two composites by the canonical isomorphisms defining tensor yield functors of the form

$$A(X, Y)^{\rightarrow} \rightarrow B(FX, GY)^{\rightarrow}$$

which, by routine application of naturality and functoriality, satisfy the coherence conditions of Lemma 2.2 relative to the functor

$$A(X, Y) \xrightarrow{G} B(GX, GY) \xrightarrow{B(\alpha_X, GY)} B(FX, GY)$$

So by Lemma 2.2, the two composites are equal, and hence Diagram 4 commutes.

Theorem 2.4 *If A admits tensors with $\rightarrow$ and $F, G : A \rightarrow B$ are 2-functors, any ordinary natural transformation $\alpha : F_0 \Rightarrow G_0$ is 2-natural.*

Proof By the 2-categorical Yoneda lemma, we may assume without loss of generality that B has tensors with $\rightarrow$.

To give a 2-cell $\lambda : h_0 \Rightarrow h_1 : X \rightarrow Y$ in a 2-category that admits tensors with $\rightarrow$ is equivalent to giving a map h from $\rightarrow \otimes X$ to Y . Post-composition with Diagram 4 together with routine diagram manipulation yields the result.

The proof of Theorem 2.4 can also be seen as an instance of general theory about the relationship between enriched categories and categories with actions [3]. By duality, Theorem 2.4 can also be written as follows:

Corollary 2.5 *Given a 2-category A that has cotensors with $\rightarrow$, and 2-functors $F, G : A \rightarrow B$, any ordinary natural transformation $\alpha : F_0 \Rightarrow G_0$ is 2-natural.*

3 Unicity of Enrichment over Cat or Gpd

The main goal of this paper is to show that, under a very mild condition on a base 2-category, every ordinary monad on it has at most one enrichment. But, for completeness, we first show that there exist ordinary monads on Cat that do not admit any enrichment at all, even over the cartesian closed category Gpd of groupoids.

Example 3.1 *Consider the endofunctor T on Cat given by*

$$T(C) = C + (N \times ob(C))$$

where N is the discrete category on the natural numbers. The action of T on maps is evident. The functor T is finitary and is the functor part of the monad on Cat for which an algebra is a category C together with a function $E : ob(C) \rightarrow ob(C)$.

But T cannot be extended to 2-cells, even to invertible ones. For instance, putting $C = 1$, taking D to be Iso , the two object category that is equivalent to 1 , and letting f and g be the two distinct functors from C to D , it follows that f is isomorphic to g but Tf is not isomorphic to Tg . So there is no possible enrichment of the ordinary functor T over either Cat or Gpd .

We now proceed towards showing that if a 2-category A admits tensors or cotensors with the arrow category $\rightarrow$, then any ordinary monad S on A_0 admits at most one enrichment over Cat .

Definition 3.2 *Given 2-categories A and B and an ordinary functor $H : A_0 \longrightarrow B_0$, an enrichment of H is any 2-functor $K : A \longrightarrow B$ for which $K_0 = H$. And given an ordinary monad S on A_0 , an enrichment of S is any 2-monad T on A for which $T_0 = S$.*

Corollary 3.3 *If A has tensors or cotensors with $\rightarrow$, then any ordinary functor of the form $H : A_0 \longrightarrow B_0$ has at most one enrichment.*

Proof Let $F, G : A \longrightarrow B$ be 2-functors for which $F_0 = G_0 : A_0 \longrightarrow B_0$. Then the identity is an ordinary natural transformation from F_0 to G_0 . So by Theorem 2.4 and Corollary 2.5, it is 2-natural, and so $F = G$.

Corollary 3.4 *If A has tensors or cotensors with $\rightarrow$, then any ordinary monad on A_0 has at most one enrichment. Moreover, an enrichment exists if and only if its underlying functor enriches.*

Proof Suppose S and T are 2-monads on A for which $S_0 = T_0$. By Corollary 3.3, S and T have the same underlying 2-functors. The unit and multiplication of S and T agree as ordinary natural transformations, so must agree as 2-natural transformations. So $S = T$. The final claim follows immediately from Theorem 2.4 and Corollary 2.5.

The results of this section all extend trivially from *Cat* to *Gpd*, the cartesian closed category of groupoids because every *Gpd*-category is, a fortiori, a 2-category, and a 2-functor between *Gpd*-categories is precisely a *Gpd*-functor between them.

4 Lack of Unicity of Enrichment for General V

The main result of this paper asserts that an ordinary monad on a 2-category, subject to a very mild condition on the 2-category, has at most one enrichment over *Cat*. The corresponding fact trivially holds in regard to enrichment over any V for which the functor $V_0(I, -) : V_0 \longrightarrow \text{Set}$ is faithful, e.g., for $V = \text{Ab}$, *Poset* or *ωCpo* . But it fails for $V = \text{Gr-Ab}$ and $V = \text{D-Gr-Ab}$, hence for some R with $V = \text{D-Gr-R-Mod}$ [4].

Example 4.1 *Consider $V = \text{D-Gr-R-Mod}$ even for R being the ring $\mathbb{Z}$ of integers. There are representable V -functors $[-, A], [-, B] : V^{op} \longrightarrow V$ together with an ordinary natural transformation from $[-, A]$ to $[-, B]$ that is not V -natural, or, equivalently by the enriched Yoneda lemma, that does not arise uniquely from a map from A to B .*

Define A by putting $A_n = \mathbb{Z}$ if $n = 0$ and 0 otherwise. Define B by putting $B_n = \mathbb{Z}$ if $n = 1$ and 0 otherwise. This is sufficient to determine the rest of the data.

Observe that the zero map is the only map from A to B . Letting $C^{-n}(X)$ denote the set of n -cochains of X , one has

$$[X, A]_n = [X_n, Z] = C^{-n}(X)$$

and

$$[X, B]_n = [X_{-n+1}, Z] = C^{-n+1}(X)$$

Consider the family of maps

$$\alpha_X : [X, A] \longrightarrow [X, B]$$

given by

$$C^{-n}(X) \xrightarrow{\delta} C^{-n+1}(X)$$

where delta is defined by composition with $\partial : X_{n+1} \longrightarrow X_n$.

Observe that δ is a chain complex map since $\delta\delta = \delta\delta = 0$. Moreover, it is natural in X because

$$\begin{array}{ccc} [X, A] & \xrightarrow{\delta} & [X, B] \\ [g, A] \downarrow & & \downarrow [g, B] \\ [Y, A] & \xrightarrow{\delta} & [Y, B] \end{array}$$

commutes whenever $\partial g = g\partial$, i.e., whenever g is a chain map.

But it is non-zero, yielding the desired counterexample.

Example 4.1 shows that the analysis of this paper is specific to *Cat* and *Gpd*. Evidently, one could extend it in very specific ways, but the proof is remarkably special, and it certainly does not extend to those V that have been oft-studied to date.

References

- [1] R. Blackwell, G.M. Kelly, and A.J. Power, *Two-dimensional monad theory*, J. Pure Appl. Algebra **59** (1989) 1–41.
- [2] F. Foltz, G.M. Kelly, and C. Lair, *Algebraic Categories with few monoidal closed structures or none*, J. Pure Appl. Algebra **17** (1980) 171–177.
- [3] R. Gordon and A.J. Power, *Enrichment through Variation*, J. Pure Appl. Algebra **120** (1997) 167–185.
- [4] G.M. Kelly, *Basic concepts of enriched category theory*, Camb. Univ. Press (1982).
- [5] G.M. Kelly, *Elementary observations on 2-categorical limits*, Bull. Austral. Math. Soc. **39** (1980) 301–317.

- [6] G.M. Kelly and S. Lack, *On property-like structures*, Theory and Applications of Categories **3** (1997) 213–250.
- [7] A.J. Power, *Why tricategories?* Information and Computation **120** (1995) 251–262.
- [8] A.J. Power and M. Tanaka, *A Unified Category-Theoretic Semantics for Binding Structures in Substructural Logics*, J. Logic and Computation **16** (2006) 5–25.