Citation for published version:

Heffernan, SJ & Carbery, DR 2012, 'Ireland-Claisen rearrangement of ynamides: Stereocontrolled synthesis of 2-amidodienes', *Tetrahedron Letters*, vol. 53, no. 38, pp. 5180-5182. <https://doi.org/10.1016/j.tetlet.2012.07.088>

DOI:

[10.1016/j.tetlet.2012.07.088](https://doi.org/10.1016/j.tetlet.2012.07.088)

Publication date:

2012

Document Version

Peer reviewed version

[Link to publication](https://doi.org/10.1016/j.tetlet.2012.07.088)

NOTICE: this is the author's version of a work that was accepted for publication in *Tetrahedron Letters*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Tetrahedron Letters*, vol 53, issue 38, 2012, DOI 10.1016/j.tetlet.2012.07.088

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Graphical Abstract

To create your abstract, type over the instructions in the template box below.
Fonts or abstract dimensions should not be changed or altered.

Ireland-Claisen rearrangement of ynamides: stereocontrolled synthesis of 2-amidodienes

Stephen J. Heffernan, David R. Carbery*

Leave this area blank for abstract info.

Ireland-Claisen rearrangement of ynamides: stereocontrolled synthesis of 2-amidodienes

Stephen J. Heffernan, David R. Carbery^{a*}

Department of Chemistry, University of Bath, Claverton Down, Bath, United Kingdom, BA2 7AY

ARTICLE INFO

Article history:

Received
Received in revised form
Accepted
Available online

Keywords:

Sigmatropic
Ireland-Claisen
Furanomycin
Amino Acid
Rearrangement

ABSTRACT

The Ireland-Claisen rearrangement of propargyl ynamido ester substrates is reported. The expected allenamide carboxylic acid products from [3,3]-sigmatropic rearrangement are not isolated, with 2-amidodienes alternatively formed in good yield with high levels of stereocontrol after decarboxylation.

2009 Elsevier Ltd. All rights reserved.

Ynamides have developed a mature and varied area of organic synthesis in recent years.¹ These compounds have been used in a raft of transition metal catalyzed reactions, such as cycloadditions and couplings.² Whilst a strongly electron-donating nitrogen is tempered by a nitrogen electron-withdrawing protecting group, the reactivity of ynamides is still pronounced. One area of synthetic chemistry where ynamides are arguably underdeveloped is in sigmatropic rearrangement chemistry. Reactions such as the Ireland-Claisen [3,3]-sigmatropic rearrangement³ offer synthetic versatility, allowing for excellent stereocontrol and the formation of congested stereocentres.

We have recently reported the use of enamides in the Ireland-Claisen [3,3]-sigmatropic rearrangement.⁴ As part of this area of research within our group,⁵ we have utilized ynamides as synthetic intermediates to the requisite enamide substrates. The availability of ynamido propargyl alcohols has allowed us to ponder the possibility of conducting an Ireland-Claisen rearrangement of these ynamido propargyl systems.^{6,7} If successful, this [3,3]-sigmatropic rearrangement would offer a novel stereocontrolled entry to allenamide carboxylic acid fragments (Scheme 1). As allenamides are important synthetic building blocks,⁸⁻¹⁰ we felt this rearrangement was worthy of investigation.

Scheme 1. Proposed Ireland-Claisen rearrangement of ynamides to allenamides.

To examine this proposal, ester **6a** was synthesized, incorporating the phenylacetate unit which had been shown to be important for the smooth rearrangement of the analogous enamide^{4b} system (Scheme 2). Accordingly, bromopropargylsilyl ether **5**¹¹ was coupled to 2-oxazolidinone **4**, promoted by Cu-catalysis.¹² Desilylation mediated by TBAF and subsequent carbodiimide-mediated esterification formed ynamido substrate **6a** in 50% over three steps.

Scheme 2. Model Substrate Synthesis.

Initial attempts to form allenamide **3** centered upon utilizing the protocol developed for the rearrangement of enamides (Table 1).^{4b} However, we were presented with a particularly complex reaction mixture, with attempted diazomethane-mediated carboxylic acid methylation observed to be non-productive,

* Corresponding author. Tel.: +44-1225-386-144; fax: +44-1225-386-231; e-mail: d.carbery@bath.ac.uk

suggesting the absence of a carboxylic acid group. After careful chromatography, oxazolidinone substituted diene **7a**, with the major isomer characterized as the *Z,E*-isomer as displayed was obtained. As discussed by Hsung, there appears to be no general synthesis of aminodienes presently available to the synthesis community and therefore new methodologies that offer a controlled entry to such systems can be viewed as valuable.¹³ Therefore, we sought to optimize the formation of this amidodiene product (Table 1).

Table 1. Rearrangement optimization

Entry	LiHMDS (equiv)	Me ₃ SiCl (equiv)	Temp (°C)	Time (h)	Yield (%)	Z/E
1	1.3	1.3	-78→rt	24	40	>95:5
2	1.3	1.3	-78→rt	48	41	>95:5
3	1.3	0	-78→rt	24	0	-
4 ^a	1.3	1.3	-78→rt	24	13	>95:5
5	1.3 ^b	1.3	-78→rt	24	25	>95:5
6	1.3	1.3	-95→rt	1.5	21	>99:5
7	1.3	1.05	-78→rt	1.5	10	>99:5
8	1.3	1.3	-40→rt	1.5	0 ^c	-
9	1.3	1.3	-20→rt	1.5	0 ^c	-
10	1.3	1.3	0→rt	1.5	0 ^c	-
11	5.2	5.2	-95→rt	24	0	-
12	2.6	2.6	-95→rt	24	31	1:1
13	2.3	2.3	-95→rt	24	40	3:1
14	2	2	-95→rt	24	42	5:1
15	1.8	1.8	-95→rt	24	83	8:1
16	1.3	1.3	-95→rt	24	61	>95:5
17	1.05	1.05	-95→rt	24	19	6:1

^aReaction conducted in PhMe. ^bNaHMDS used as base. ^cFull mass recovery of **6a**.

The rearrangement is particularly sensitive to the initial conditions employed (Table 1). For example, low loadings of base and Me₃SiCl resulted in excellent stereocontrol (entries 1-4). The reaction requires the presence of silyl chloride and therefore supports a traditional Ireland-Claisen process occurring (entry 3). The initiating temperature is crucial to any rearrangement occurring, with -95 °C offering the best results (entries 6-10). We feel it is noteworthy that **6a** is re-isolated, with full mass balance, when this reaction is initiated at -40 °C or higher (entries 8-10). Furthermore, the addition of higher loadings of base and silyl chloride is also deleterious to the final isolated yield (entries 11-17).

Previous work in our group has demonstrated that the stereocontrolled formation of aryl-substituted silylketene acetals is a more complex problem than currently is appreciated where the *E/Z* ratios are highly dependent on the loadings of base and silyl chloride, as well temperature.¹⁴ Accordingly, we feel the presently reported rearrangement is also sensitive to the

complications of forming silylketene acetals from aryl acetate esters.

This diene is presumably the result of a post-rearrangement decarboxylation. Baldwin has reported the decarboxylation of allenyl carboxylic acids, formed from the Ireland-Claisen rearrangement of propargylic esters.^{15,16} However, the decarboxylation step required forcing thermal conditions (140-250 °C) to accomplish the loss of CO₂. Therefore, the presence of the *N*-substitution has a profound effect on this decarboxylation event. With an optimized rearrangement developed on phenyl acetate **6a**, we have sought to examine the scope of the arylacetate moiety (Table 2). The conditions chosen, and in particular the loading of base and silyl chloride, represents striking a balance between optimized yield and *Z/E* selectivity, as highlighted in Table 1. Accordingly, the substrate scope has been studied with 1.8 equivalents of LiHMDS and Me₃SiCl.

Table 2. Scope of ester functionality¹⁷

Entry	Ar	Diene	Yield (%)	Z/E
1	4-Me ₂ NC ₆ H ₄ (6b)	7b	53	>95:5
2	4-MeOC ₆ H ₄ (6c)	7c	62	9:1
3	3,4-(OCH ₂ O)C ₆ H ₃ (6d)	7d	69	2:1
4	4-FC ₆ H ₄ (6e)	7e	67	3:1
5	4-ClC ₆ H ₄ (6f)	7f	65	4:1
6	4-NO ₂ C ₆ H ₄ (6g)	7g	54	2:1
7	4-MeC ₆ H ₄ (6h)	7h	73	3:1
8	3-MeC ₆ H ₄ (6i)	7i	42	6:1
9	2-MeC ₆ H ₄ (6j)	7j	43	2:1

This decarboxylative rearrangement can accommodate electron-rich aryl groups (entries 1-3) and electron-poor aryl groups (entries 4-6), with reasonable yields in each instance. The aryl moiety can also cope with *ortho*, *meta* and *para*-substitution in a range of tolylacetates (entries 7-9). We would like to point out that the final *Z/E* ratio was highly sensitive to the initial conditions, as gauged by the extensive optimization of **6a**. Therefore, it may be reasonable to judge that each individual substrate may in turn have its *Z/E* selectivity improved through a local optimization process.

This decarboxylative rearrangement has been demonstrated on alkyl ester **6k** (Scheme 3). Whilst, the rearrangement in this instance is non-optimized, we feel the excellent levels of *Z/E* control are noteworthy and suggest good substrate scope in future studies.

Scheme 3. Incorporation of alkyl functionality.

Finally, we have briefly examined the feasibility of **7a** acting as a diene component in a Diels-Alder reaction (Scheme 4).¹⁸ To assess this point, diene **7a** was refluxed in toluene for 24 hours with the reactive dienophile, maleic anhydride (**8**). Surprisingly, even though an electron-rich diene is present with an electron-deficient dienophile, no reaction was observed, with **7a** recovered with full mass balance. To account for this interesting observation, we suggest that the requisite *s-cis* conformation of **7a** cannot be accessed, even under forcing conditions, from *s-trans*-**7a**.

Scheme 4. Attempted Diels-Alder reaction and diene conformation.

In conclusion, the first Ireland-Claisen [3,3]-sigmatropic rearrangement of an ynamido ester is reported. A range of trisubstituted amidodienes has been accessed in good to excellent levels of *E/Z* selectivity and good yields. We are currently examining the synthetic utility of these amidodiene products further and investigating the observed *E/Z* stereoselectivity. Details will be reported in due course.

Acknowledgements

We acknowledge the support of University of Bath and EPSRC for studentship funding (SJH).

References and notes

- For reviews of ynamide synthetic chemistry, see: (a) De Korver, K. A.; Li, H.; Lohse, A. G.; Hayashi, R.; Lu, Z.; Zhang, Y.; Hsung, R. P. *Chem. Rev.* **2010**, *110*, 5064. (b) Evano, G.; Coste, A.; Jouvin, K. *Angew. Chem. Int. Ed.* **2010**, *49*, 2840.
- For recent synthetic usage of ynamides, see: (a) Saito, N.; Ichimaru, T.; Sato, Y. *Org. Lett.* **2012**, *14*, 1914. (b) DeKorver, K. A.; Wang, X.-N.; Walton, M. C.; Hsung, R. P. *Org. Lett.* **2012**, *14*, 1768. (c) Garcia, P.; Evanno, Y.; George, P.; Sevrin, M.; Ricci, G.; Malacria, M.; Aubert, C.; Gandon, V. *Chem. Eur. J.* **2012**, *18*, 4337. (d) Jouvin, K.; Heimburger, J.; Evano, G. *Chem. Sci.* **2012**, *3*, 756. (e) Smith, D. L.; Goundry, W. R. F.; Lam, H. W. *Chem. Commun.* **2012**, *48*, 1505. (f) Dateer, R. B.; Shaibu, B. S.; Liu, R.-S. *Angew. Chem. Int. Ed.* **2012**, *51*, 113. (g) Davies, P. W.; Cremonesi, A.; Dumitrescu, L. *Angew. Chem. Int. Ed.* **2011**, *50*, 8931. (h) Kramer, S.; Odabachian, Y.; Overgaard, J.; Rottlaender, M.; Gagosz, F.; Skrydstrup, T. *Angew. Chem. Int. Ed.* **2011**, *50*, 5090.
- For recent reviews concerning the Ireland-Claisen rearrangement, see: (a) Castro, A. M. *Chem. Rev.* **2004**, *104*, 2939. (b) Chai, Y. H.; Hong, S. P.; Lindsay, H. A.; McFarland, C.; McIntosh, M. C. *Tetrahedron* **2002**, *58*, 2905. (c) McFarland, C. M.; McIntosh, M. C. *The Claisen Rearrangement*; Hiersemann, M. N., Nubbemeyer, U., Eds.; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2007; p117. (d) Wipf, P. *Claisen Rearrangements*; *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Paquette, L. A., Eds.; Pergamon: Oxford, 1991; Vol 5, p827. (e) Ilardi, E. A.; Stivala, C. E.; Zakarian, A. *Chem. Soc. Rev.* **2009**, *38*, 3133.
- (a) Harker, W. R. R.; Carswell, E. L.; Carbery, D. R. *Org. Biomol. Chem.* **2012**, 1406. (b) Ylioja, P. M.; Mosley, A. D.; Charlot, C. E.; Carbery, D. R. *Tetrahedron Lett.* **2008**, *49*, 1111.
- (a) Fairhurst, N. W. G.; Mahon, M. F.; Munday, R. H.; Carbery, D. R. *Org. Lett.* **2012**, *14*, 756. (b) Tellam, J. P.; Carbery, D. R. *Tetrahedron Lett.* **2011**, *52*, 6027. (c) Tellam, J. P.; Kociok-Köhn, G.; Carbery, D. R. *Org. Lett.* **2008**, *10*, 5199. (d) Tellam, J. P.; Carbery, D. R. *J. Org. Chem.*, **2010**, *75*, 7809.
- For examples of Ireland-Claisen and related ester enolate rearrangements of propargyl systems, see: (a) Rogakos, V.; Georgiadis, D.; Dive, V.; Yiotakis, A. *Org. Lett.* **2009**, *11*, 4696. (b) Baldwin, J. E.; Bennett, P. A. R.; Forrest, A. K. *J. Chem. Soc., Chem. Commun.* **1987**, 250. (c) Takai, K.; Ueda, T.; Kaihara, H.; Sunami, Y.; Moriwake, T. *J. Org. Chem.* **1996**, *61*, 8728. (d) Ishihara, J.; Koyama, N.; Nishino, Y.; Takahashi, K.; Hatakeyama, S. *Synlett*, **2009**, 2351.
- For the synthesis of α -allenyl- α -amino acids through the ester enolate Claisen rearrangement of α -amino propargyl esters, see: (a) Casara, P.; Jund, K.; Bey, P. *Tetrahedron Lett.* **1984**, *25*, 1891. (b) Castelhana, A. L.; Horne, S.; Taylor, G. J.; Billedeau, R.; Krantz, A. *Tetrahedron* **1988**, *44*, 5451. (c) Kazmaier, U.; Goerbitz, C. H. *Synthesis* **1996**, 1489.
- For a recent review of allenamide chemistry, see: Hsung, R. P.; Wei, L.-L.; Xiong, H. *Acc. Chem. Res.* **2003**, *36*, 773.
- For recent syntheses of allenamides, see: (a) Armstrong, A.; Emmerson, D. P. G. *Org. Lett.* **2009**, *11*, 1547. (b) Hyland, C. J. T.; Hegedus, L. S. *J. Org. Chem.* **2005**, *70*, 8628. (c) Shen, L.; Hsung, R. P.; Zhang, Y.; Antoline, J. E.; Zhang, X. *Org. Lett.* **2005**, *7*, 3081. (d) Trost, B. M.; Stiles, D. T. *Org. Lett.* **2005**, *7*, 2117.
- For recent synthetic usage, see: (a) Hayashi, R.; Hsung, R. P.; Feltenberger, J. B.; Lohse, A. G. *Org. Lett.* **2009**, *11*, 2125. (b) Lu, T.; Hayashi, R.; Hsung, R. P.; DeKorver, K. A.; Lohse, A. G.; Song, Z.; Tang, Y. *Org. Biomol. Chem.* **2009**, *7*, 3331. (c) Lohse, A. G.; Hsung, R. P. *Org. Lett.* **2009**, *11*, 3430. (d) Manzo, A. M.; Perboni, A. D.; Brogini, G.; Rigamonti, M. *Tetrahedron Lett.* **2009**, *50*, 4696. (e) Chen, G.; Fu, C.; Ma, S. *Org. Lett.* **2009**, *11*, 2900. (f) Brummond, K. M.; Yan, B. *Synlett* **2008**, 2303. (g) Beccalli, E. M.; Brogini, G.; Clerici, F.; Galli, S.; Kammerer, C.; Rigamonti, M.; Sottocornola, S. *Org. Lett.* **2009**, *11*, 1563. (h) Song, Z.; Hsung, R. P.; Lu, T.; Lohse, A. G. *J. Org. Chem.* **2007**, *72*, 9722.
- Barbazanges, M.; Meyer, C.; Cossy, J. *Tetrahedron Lett.* **2008**, *49*, 2902.
- Zhang, Y.; Hsung, R. P.; Tracey, M. R.; Kurtz, K. C. M.; Vera, E. L. *Org. Lett.* **2004**, *6*, 1151.
- Hayashi, R.; Hsung, R. P.; Feltenberger, J. B.; Lohse, A. G. *Org. Lett.* **2009**, *11*, 2125.
- Harker, W. R. R.; Carswell, E. L.; Carbery, D. R. *Org. Lett.* **2010**, *12*, 3712.
- Baldwin, J. E.; Bennett, P. A. R.; Forrest, A. K. *J. Chem. Soc., Chem. Commun.* **1987**, 250.
- Craig has studied extensively a decarboxylative Claisen rearrangement. For a discussion and leading references, see: Camp, J. E.; Craig, D.; Funai, K.; White, A. J. P. *Org. Biomol. Chem.* **2011**, *9*, 7904.
- Representative example; synthesis of **7b**. To a flask was added LiHMDS (1 M in THF, 0.57 mL, 0.57 mmol, 1.8 equiv.) and Me₃SiCl (73 μ L, 0.57 mmol, 1.8 equiv.). The mixture was stirred at -95 °C for 0.25 h before the dropwise addition of **6b** (100 mg, 0.32 mmol, 1 equiv.) in THF (2 mL). After 0.5 h at -95 °C, the reaction mixture was allowed to warm to room temperature after which time it was stirred for an additional 24 h. The reaction was then quenched with 1:1 M HCl/brine solution (5 mL) and extracted with EtOAc (3 x 10 mL), with the combined organic extracts being dried over Na₂SO₄, filtered and concentrated *in vacuo*. Purification via flash chromatography, eluting with 2:1 petroleum ether/EtOAc, gave **7b** (46 mg, 53%). FTIR (film/cm⁻¹) $\nu_{\max}$: 2903, 2798, 1741, 1604, 1519. ¹H NMR (500 MHz, CDCl₃) δ H: 7.29 (2H, app. d, *J* = 8.5 Hz), 6.65 (2H, app. d, *J* = 8.5 Hz), 6.48 (1H, s), 6.17–6.08 (1H, m), 5.79 (1H, dqd, *J* = 11.4, 7.2, 0.8 Hz), 4.39 (1H, d, *J* = 7.7 Hz), 4.36 (1H, d, *J* = 6.7 Hz), 3.80 (1H, d, *J* = 6.7 Hz), 3.77 (1H, d, *J* = 7.7 Hz), 2.96 (6H, s), 1.66 (3H, dd, *J* = 7.2, 1.8 Hz); ¹³C NMR (75 MHz, CDCl₃) δ C: 156.1, 149.5, 146.8, 130.3, 130.2, 128.5, 125.2, 123.8, 111.9, 61.6, 46.2, 40.4, 14.7; MS (ESI+ve) *m/z*: calcd. for C₁₆H₂₀N₂O₂Na: 295.1422, found: 295.1417, [M + Na]⁺.
- For a recent example of an amidodiene Diels-Alder reaction, see: Hayashi, R.; Ma, Z.-X.; Hsung, R. P.; *Org. Lett.* **2012**, *14*, 252.