Citation for published version:

DeLisle, S, Pittet, D, Potter, BVL, Lew, PD & Welsh, MJ 1992, 'InsP3 and Ins(1,3,4,5)P4 act in synergy to stimulate influx of extracellular Ca^{2+} in *Xenopus* oocytes', *American Journal of Physiology - Cell Physiology*, vol. 262, no. 6, pp. C1456-C1463.

Publication date:

1992

Document Version

Publisher's PDF, also known as Version of record

[Link to publication](#)

APS permit archiving of final published version at time of publication

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

InsP3 and Ins(1,3,4,5)P4 act in synergy to stimulate influx of extracellular Ca²⁺ in *Xenopus* oocytes

S. DeLisle, D. Pittet, B. V. Potter, P. D. Lew and M. J. Welsh

Am J Physiol Cell Physiol 262:C1456-C1463, 1992. ;

You might find this additional info useful...

This article has been cited by 2 other HighWire-hosted articles:

<http://ajpcell.physiology.org/content/262/6/C1456#cited-by>

Updated information and services including high resolution figures, can be found at:

<http://ajpcell.physiology.org/content/262/6/C1456.full>

Additional material and information about *American Journal of Physiology - Cell Physiology* can be found at:

<http://www.the-aps.org/publications/ajpcell>

This information is current as of September 6, 2012.

InsP₃ and Ins(1,3,4,5)P₄ act in synergy to stimulate influx of extracellular Ca²⁺ in *Xenopus* oocytes

SYLVAIN DELISLE, DIDIER PITTET, BARRY V. L. POTTER, P. DANIEL LEW, AND MICHAEL J. WELSH

Howard Hughes Medical Institute, Departments of Internal Medicine and Physiology and Biophysics, University of Iowa College of Medicine, Iowa City, Iowa 52242; Infectious Diseases Division, University Hospital, CH-1211 Geneva 4, Switzerland; and School of Pharmacy and Pharmacology, University of Bath, Bath BA2 7AY, United Kingdom

DeLisle, Sylvain, Didier Pittet, Barry V. L. Potter, P. Daniel Lew, and Michael J. Welsh. InsP₃ and Ins(1,3,4,5)P₄ act in synergy to stimulate influx of extracellular Ca²⁺ in *Xenopus* oocytes. *Am. J. Physiol.* 262 (Cell Physiol. 31): C1456–C1463, 1992.—To investigate the role of D-myo-inositol 1,3,4,5-tetrakisphosphate [Ins(1,3,4,5)P₄] in the regulation of Ca²⁺ influx, we injected inositol phosphates into *Xenopus* oocytes and measured Ca²⁺-gated Cl[−] current to assay intracellular free Ca²⁺ concentration ([Ca²⁺]_i). To assess Ca²⁺ influx, we removed extracellular Ca²⁺ or added the inorganic Ca²⁺ channel blocker Mn²⁺ to the extracellular bath and measured the resulting change in Cl[−] current. Ins(1,3,4,5)P₄ did not cause Ca²⁺ influx when injected alone or when preceded by an injection of Ca²⁺. In contrast, Ins(1,3,4,5)P₄ stimulated Ca²⁺ influx when injected after the poorly metabolized inositol trisphosphate (InsP₃) analogues D-myo-inositol 1,4,5-trisphosphorothioate [Ins(1,4,5)P₃S₃] or D-myo-inositol 2,4,5-trisphosphate [Ins(2,4,5)P₃]. These results indicate that Ins(1,3,4,5)P₄ is not sufficient to stimulate Ca²⁺ influx but acts in synergy with InsP₃s to cause Ca²⁺ influx. We also studied the effect of Ca²⁺ influx on the immediate metabolism of D-myo-inositol 1,4,5-trisphosphate [Ins(1,4,5)P₃] in single oocytes. Ca²⁺ influx shunted the metabolism of Ins(1,4,5)P₃ toward the formation of Ins(1,3,4,5)P₄ and away from D-myo-inositol 1,4-bisphosphate [Ins(1,4)P₂]. These results suggest that there is a positive feedback regulatory mechanism in which Ca²⁺ influx stimulates Ins(1,3,4,5)P₄ production and Ins(1,3,4,5)P₄ stimulates further Ca²⁺ influx.

inositol trisphosphate; D-myo-inositol 1,3,4,5-tetrakisphosphate; extracellular calcium influx; intracellular calcium concentration

STIMULATION OF MANY plasma membrane receptors increases the intracellular concentration of the second messenger D-myo-inositol 1,4,5-trisphosphate [Ins(1,4,5)P₃]. By opening Ca²⁺ channels, Ins(1,4,5)P₃ releases Ca²⁺ from intracellular stores (2, 15, 34). Ins(1,4,5)P₃ also causes a sustained influx of extracellular Ca²⁺ into the cell (2). We do not know how Ins(1,4,5)P₃ regulates Ca²⁺ influx (33).

Intracellular injection of Ins(1,4,5)P₃ can stimulate Ca²⁺ influx (7, 11, 21, 22, 27, 30, 36). However, because Ins(1,4,5)P₃ is extensively metabolized by the cell, inositol phosphates (InsP) other than Ins(1,4,5)P₃ may control Ca²⁺ influx. D-Myo-inositol 1,3,4,5-tetrakisphosphate [Ins(1,3,4,5)P₄] is a direct metabolite of Ins(1,4,5)P₃ that may also release intracellular Ca²⁺ (16, 36). Ins(1,3,4,5)P₄ has been implicated in the regulation of Ca²⁺ influx in some cellular systems (7, 21, 22, 27, 31) but not in others (30, 36). The controversy surrounding the requirement of Ins(1,3,4,5)P₄ for Ca²⁺ influx has recently been emphasized when two groups of investigators using similar techniques in mouse lacrimal acinar

cells came to opposite conclusions; the results of one study suggested that Ins(1,3,4,5)P₄ was required for Ca²⁺ influx (27), and the results of the other study suggested it was not (5). In the present studies, we investigate how inositol trisphosphates (InsP₃) and Ins(1,3,4,5)P₄ control Ca²⁺ influx in *Xenopus* oocytes. We show that Ins(1,3,4,5)P₄ stimulates Ca²⁺ influx in the presence but not in the absence of poorly metabolizable InsP₃s. In addition, we tested the possibility that Ca²⁺ influx may stimulate the phosphorylation of Ins(1,4,5)P₃ into Ins(1,3,4,5)P₄. Our observations suggest a positive feedback mechanism, whereby Ins(1,3,4,5)P₄ stimulates Ca²⁺ influx and Ca²⁺ influx promotes formation of Ins(1,3,4,5)P₄.

MATERIALS AND METHODS

Electrophysiology and intracellular injections. We obtained *Xenopus* oocytes as described previously (11, 36) and defolliculated them manually. To assay intracellular Ca²⁺ concentration ([Ca²⁺]_i), we measured Ca²⁺-activated Cl[−] currents with the two-electrode voltage-clamp technique. This assay has been validated using both Ca²⁺ electrodes and fluorescent calcium indicators (10). Oocytes were impaled with two microelectrodes (3 M KCl filled, resistance = 0.5–1.0 MΩ), and membrane voltage was maintained at −50 mV with a voltage-clamp amplifier (Axoclamp 2A, Axon Instruments, Burlingame, CA). The current necessary to maintain this membrane voltage was recorded on-line (MacLab, World Precision Instruments, New Haven, CT) and analyzed by a microcomputer (Macintosh SE/30, Apple Computer, Cupertino, CA). A third micropipette (tip diameter 2–5 μm) was attached to a pneumatic microinjector (PLI-100, Medical Systems, Greenvale, NY) and used for intracellular injections. When sequential intracellular injections involving different substances were required, the first injection pipette was withdrawn from the cell and a second pipette was reinserted at the same site. Although removal of the injection pipette from the cell did not impair subsequent cellular responses, it caused a transient inward current; after 30–60 s, current returned to values similar to those obtained just before pipette removal. For the sake of clarity, this transient current was blanked from the tracings. Injection pipettes were back-filled with Ins(1,3,4,5)P₄ (Calbiochem, San Diego, CA), Ins(1,4,5)P₃ (Amersham, Arlington Heights, IL), ³H-labeled Ins(1,4,5)P₃ (Amersham), D-myo-inositol 2,4,5-trisphosphate [Ins(2,4,5)P₃] (Boehringer Mannheim Biochemicals, Indianapolis, IN), D-myo-inositol 3,4,5,6-tetrakisphosphate [Ins(3,4,5,6)P₄] (Boehringer Mannheim), D-myo-inositol 1,4,5-trisphosphorothioate [Ins(1,4,5)P₃S₃] [prepared as previously described (9)], or CaCl₂ (Sigma Chemicals, St. Louis, MO). Each injection had a volume of 10 nl or ~1% of the oocyte volume. Before and after each experiment, we calibrated the injection pipette by expelling a drop of solution in paraffin oil and calculating volume from the measured diameter of the drop.

Ca^{2+} influx measurements. For most experiments, oocytes were initially stimulated in a bath solution containing (in mM) 116 NaCl, 2 KCl, 6 CaCl_2 , 1 MgCl_2 , and 5 *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid (HEPES), pH 7.4. We assessed Ca^{2+} influx by measuring the change in Cl^- current induced by either removing external Ca^{2+} or adding the inorganic Ca^{2+} channel blocker Mn^{2+} (4 mM) to the bath solution. For most experiments, we added Mn^{2+} rather than removing bath Ca^{2+} because membrane electrical resistance often decreased with time in the absence of external Ca^{2+} (10). Because we have previously found that Ca^{2+} influx usually occurs after the disappearance of $[\text{Ca}^{2+}]_i$ oscillations (11, 36), our standard procedure was to add Mn^{2+} during the nonoscillatory current that followed the disappearance of $[\text{Ca}^{2+}]_i$ oscillations. Occasionally, the injected InsP caused prolonged $[\text{Ca}^{2+}]_i$ oscillations. In such cases, we added Mn^{2+} when the baseline current (the average current around which the oscillations occurred) had stabilized. To determine if the addition of Mn^{2+} (or the removal of extracellular Ca^{2+}) caused a change in Cl^- current, we extracted the digitized current values for the 60-s time period immediately preceding the addition of Mn^{2+} and the current values for the 60-s time period beginning 1 min after the addition of Mn^{2+} . Because we acquired data at a rate of 4 Hz, each 60-s time period contained 240 values. We compared the two groups of values (i.e., immediately before and 1 min after Mn^{2+}) using the Student's *t* test for paired data. We interpreted a statistically significant decrease in the mean Cl^- current caused by Mn^{2+} as positive evidence of Ca^{2+} influx (11, 36). The absolute difference between the means was taken as a semiquantitative index of Ca^{2+} influx.

Ins(1,4,5) P_3 metabolism in single oocytes. To measure the metabolism of *Ins(1,4,5) P_3* , we injected $[\text{H}^3]\text{Ins(1,4,5) $\text{P}_3}$$ into single oocytes and arrested InsP metabolism at fixed time points. The injectate contained 5 μCi of lyophilized $[\text{H}^3]\text{Ins(1,4,5) $\text{P}_3}$$ (sp act 20–60 Ci/mmol) mixed with 10 μl of cold *Ins(1,4,5) P_3* (10^{-4} M). Total calculated concentration of *Ins(1,4,5) P_3* (cold and ^3H -labeled) in the injection pipette was ~ 0.11 mM. Given an injection volume of 10 nl and an oocyte volume of 1 μl , the calculated intracellular *Ins(1,4,5) P_3* concentration was ~ 1.1 μM , i.e., comparable with what we have used in the past to cause Ca^{2+} influx (36). At predetermined time points (0.25, 0.5, 1, 2, 3, 5, 10, 20, and 40 min), InsP metabolism was arrested by crushing the oocyte in 1 ml of ice-cold trichloroacetic acid solution (15%). The sample was then centrifuged (4°C , 800 *g*) for 10 min, and the supernatant was transferred into a glass tube. Three successive diethyl ether extractions (5:1, vol/vol) were performed to remove lipids and phospholipids, and the aqueous phase was frozen at -70°C . When feasible, i.e., time points of 1 min or more, we measured the cellular response to the $[\text{H}^3]\text{Ins(1,4,5) P_3 }/\text{Ins(1,4,5) $\text{P}_3}$$ injection mixture by recording Cl^- current. In these cases, recording electrodes were withdrawn from the cell 30 s before InsP metabolism was stopped.

High-performance liquid chromatography (HPLC) separation of InsPs. After addition of ATP (10 μM) and EDTA (1 mM), the thawed samples were chromatographed on two sequential anion exchange columns, SAX-10, 0.4×25 cm (Technicol, Stockport, Cheshire, UK), at a flow rate of 1.6 ml/min as previously described (31). An initial 5-min washing period in H_2O was followed by a discontinuous gradient of phosphate buffer prepared from phosphoric acid and adjusted to pH 4.2 with NaOH. Elution occurred over a 60-min period. The gradient was held at a concentration of 0.36 M from 12 to 20 min, at 0.84 M from 24 to 31 min, and at 1.8 M from 54 to 59 min; changes between these isocratic steps were linear gradients. Fractions were collected at 0.2-min intervals and radioactivity assessed by liquid scintillation counting. This HPLC procedure was optimized for the separation of the three distinct inositol tetrakisphosphates:

D-myo-inositol 1,3,4,6-tetrakisphosphate [*Ins(1,3,4,6) P_4*], *Ins(1,3,4,5) P_4* , and *Ins(3,4,5,6) P_4* (31). ^3H -labeled *D-myo*-inositol 1,4-bisphosphate [*Ins(1,4) P_2*], *Ins(1,4,5) P_3* , and *Ins(1,3,4,5) P_4* standards were purchased from Amersham.

Statistical analysis. Means were compared using the Student's *t* test for unpaired data. Differences in current within single cells were compared using the Student's *t* test for paired data. Difference between the outcome in two groups of cells (intervention vs. control) was assessed with Fisher's exact test (two-tailed). Significance was assumed when $P < 0.05$.

RESULTS AND DISCUSSION

Effect of *Ins(1,3,4,5) P_4* on Ca^{2+} influx. Injection of 1 pmol of *Ins(1,4,5) P_3* (estimated intracellular concentration of 1 μM) into *Xenopus* oocytes caused a biphasic increase in $[\text{Ca}^{2+}]_i$ (Fig. 1A). We have previously shown that the initial, transient increase in $[\text{Ca}^{2+}]_i$ represents the release of Ca^{2+} from intracellular stores and is not affected by extracellular $[\text{Ca}^{2+}]$ (36). The more sustained increase in $[\text{Ca}^{2+}]_i$ that follows can be blocked either by removing extracellular Ca^{2+} or by adding Mn^{2+} (4 mM) to the external bath (Fig. 1A). This result indicates that the sustained increase in $[\text{Ca}^{2+}]_i$ is due to the influx of extracellular Ca^{2+} . Thus, as previously reported in *Xenopus* oocytes (25, 36), *Ins(1,4,5) P_3* causes Ca^{2+} influx.

To assess the role of *Ins(1,3,4,5) P_4* in *Ins(1,4,5) P_3* -induced Ca^{2+} influx, we injected 10 pmol of *Ins(1,3,4,5) P_4* (estimated intracellular concentration of 10 μM). *Ins(1,3,4,5) P_4* caused $[\text{Ca}^{2+}]_i$ oscillations (12/12 cells) (Fig. 1, B and C). In previous experiments, we (11, 36) and others (28, 29) have found that either removing extracellular Ca^{2+} or adding Mn^{2+} to the bath solution had no effect on *Ins(1,3,4,5) P_4* -induced $[\text{Ca}^{2+}]_i$ oscillations (for an example, qualitatively compare Fig. 1B with Fig. 1C). Because Ca^{2+} influx could possibly occur following completion of the intracellular Ca^{2+} release phase [as occurs with *Ins(1,4,5) P_3* ; Fig. 1A], we also added Mn^{2+} during the steady-state current that followed the $[\text{Ca}^{2+}]_i$ oscillations (Fig. 1C). Mn^{2+} did not alter Ca^{2+} -gated Cl^- current under these conditions (no change in 11/12 cells, current decreased by 22 nA in one cell). Thus, under these conditions, *Ins(1,3,4,5) P_4* is not sufficient to cause Ca^{2+} influx. These results agree with previous reports in *Xenopus* oocytes (11, 28, 29, 36).

The synthetic *Ins(1,4,5) P_3* analogue, *Ins(1,4,5) P_3S_3* (37), is a poor substrate for the *Ins(1,4,5) P_3* 3-kinase (38), the enzyme responsible for the conversion of *Ins(1,4,5) P_3* to *Ins(1,3,4,5) P_4* . As previously observed (11, 37), injection of 10 pmol of *Ins(1,4,5) P_3S_3* (estimated intracellular concentration of 10 μM) caused prolonged $[\text{Ca}^{2+}]_i$ oscillations (30/30 cells) (Fig. 1D). Removal of Ca^{2+} from the bath or addition of Mn^{2+} (4 mM) did not change the average Cl^- current response (14/14 cells) (Fig. 1D). This result indicates that *Ins(1,4,5) P_3S_3* does not cause Ca^{2+} influx.

To mimic the physiological sequence where *Ins(1,4,5) P_3* is metabolized to *Ins(1,3,4,5) P_4* , we first injected *Ins(1,4,5) P_3S_3* (10 pmol) and then *Ins(1,3,4,5) P_4* (10 pmol) in the same cell. When injected after *Ins(1,4,5) P_3S_3* , *Ins(1,3,4,5) P_4* stimulated a Cl^- current that could be inhibited by removing extracellular Ca^{2+} or by adding Mn^{2+} (4 mM) to the bath solution (11/13 cells) (Fig. 1E). These results suggest that the combination of

Ins(1,4,5) P_3S_3 and Ins(1,3,4,5) P_4 stimulates Ca^{2+} influx. This contrasts with results obtained when either compound was injected alone. The magnitude of Ins(1,4,5) P_3S_3 /Ins(1,3,4,5) P_4 -induced Ca^{2+} influx varied

considerably from cell to cell [191 ± 46 (SE) nA, range = 57–440 nA]. There was no relationship between the amplitude of Ins(1,3,4,5) P_4 -induced Ca^{2+} influx and the time interval between Ins(1,4,5) P_3S_3 and Ins(1,3,4,5) P_4 injections (from 5 to 30 min).

To ensure that Ca^{2+} influx did not simply result from a greater total amount of injected InsP, we performed two types of experiments. First, we injected twice the amount (20 pmol) of either Ins(1,3,4,5) P_4 (7 cells) or Ins(1,4,5) P_3S_3 (3 cells). This failed to cause Ca^{2+} influx. Second, we injected Ins(1,4,5) P_3S_3 (10 pmol) twice, sequentially. This also did not cause Ca^{2+} influx (2 cells). These results suggest that Ca^{2+} influx does not result from doubling the total amount of injected InsPs.

To determine whether the potential to cause Ca^{2+} influx following Ins(1,4,5) P_3S_3 was specific to Ins(1,3,4,5) P_4 , we injected Ins(3,4,5,6) P_4 , the predominant InsP₄ in nonstimulated *Xenopus* oocytes (24). Ins(3,4,5,6) P_4 (10 pmol) did not cause $[\text{Ca}^{2+}]_i$ oscillations (12/13 cells) when injected alone and did not cause Ca^{2+} influx when injected after Ins(1,4,5) P_3S_3 (5/5 cells). These results suggest that the ability to cause Ca^{2+} influx following an injection of Ins(1,4,5) P_3S_3 is relatively specific to the Ins(1,3,4,5) P_4 isomer.

Ins(2,4,5) P_3 is another InsP₃ analogue not expected to be metabolized to Ins(1,3,4,5) P_4 (26). Ins(2,4,5) P_3 , like Ins(1,4,5) P_3S_3 , releases Ca^{2+} from intracellular stores (11, 13, 36). In contrast to Ins(1,4,5) P_3S_3 , however, Ins(2,4,5) P_3 causes Ca^{2+} influx (Fig. 2A and Ref. 36). Although we do not know why the response to Ins(2,4,5) P_3 differs from Ins(1,4,5) P_3S_3 , we hypothesized that Ins(2,4,5) P_3 could also enable Ins(1,3,4,5) P_4 to stimulate Ca^{2+} influx. To test this hypothesis, we first injected Ins(2,4,5) P_3 and waited until there was no increase in the nonoscillatory Cl^- current for a period of at least 3 min. We then injected Ins(1,3,4,5) P_4 . Compared with the average amount of current present during the 1-min time period immediately preceding Ins(1,3,4,5) P_4 injection, Ins(1,3,4,5) P_4 increased the current by 115 ± 21 nA (the increase in current was statistically significant in 24/26 cells; Fig. 2B). Injecting water instead of Ins(1,3,4,5) P_4 did not produce a significant increase in

Fig. 1. Cl^- current response to intracellular injection of inositol phosphates (InsP) as a function of time. Inward current (downward deflection) represents an increase in intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$). Bar, addition of Mn^{2+} (4 mM) to extracellular bath. A: injection of D-myo-inositol 1,4,5-trisphosphate [Ins(1,4,5) P_3] (1 pmol) caused a brief increase in $[\text{Ca}^{2+}]_i$ followed by a more sustained increase in $[\text{Ca}^{2+}]_i$. Mn^{2+} (bar) blocked sustained increase in $[\text{Ca}^{2+}]_i$. Thus Ins(1,4,5) P_3 caused Ca^{2+} influx. B and C: injection of D-myo-inositol 1,3,4,5-tetrakisphosphate [Ins(1,3,4,5) P_4] (10 pmol). To show resulting $[\text{Ca}^{2+}]_i$ oscillations more clearly, we have expanded time scale of boxed area (insert in C). Addition of Mn^{2+} during $[\text{Ca}^{2+}]_i$ oscillations (B) or following the disappearance of $[\text{Ca}^{2+}]_i$ oscillations (C) did not change the average Cl^- current. Thus Ins(1,3,4,5) P_4 alone did not cause Ca^{2+} influx. D: injection of Ins(1,4,5) P_3S_3 (10 pmol) caused prolonged $[\text{Ca}^{2+}]_i$ oscillations. Addition of Mn^{2+} (bar) did not cause any change in average current. Thus D-myo-inositol 1,4,5-trisphosphorothioate [Ins(1,4,5) P_3S_3] alone did not cause Ca^{2+} influx. E: injection of Ins(1,4,5) P_3S_3 (10 pmol) followed by Ins(1,3,4,5) P_4 (10 pmol). In this case, a nonoscillatory current developed after $[\text{Ca}^{2+}]_i$ oscillations had stopped. Addition of Mn^{2+} (bar) caused a reversible decrease in Cl^- current. Thus Ins(1,3,4,5) P_4 stimulated influx of extracellular Ca^{2+} in the presence of Ins(1,4,5) P_3S_3 .

Fig. 2. *A*: injection of D-myo-inositol 2,4,5-trisphosphate [$\text{Ins}(2,4,5)\text{P}_3$] (10 pmol) caused $[\text{Ca}^{2+}]_i$ oscillations followed by a gradual increase in nonoscillatory current. This current could be blocked by adding Mn^{2+} (bar). Thus $\text{Ins}(2,4,5)\text{P}_3$ caused Ca^{2+} influx. *B*: injection of $\text{Ins}(2,4,5)\text{P}_3$ (10 pmol) followed by $\text{Ins}(1,3,4,5)\text{P}_4$ (10 pmol). $\text{Ins}(1,3,4,5)\text{P}_4$ caused further increase in Cl^- current. Mn^{2+} (bar) blocked nonoscillatory current, and $[\text{Ca}^{2+}]_i$ oscillations resumed. Because resulting oscillatory current did not go back to baseline, this experiment did not rule out the possibility that $\text{Ins}(1,3,4,5)\text{P}_4$ released intracellular Ca^{2+} instead of causing Ca^{2+} influx. *C*: same experiment as *B* except that Mn^{2+} was present throughout experiment, preventing Ca^{2+} influx. After $\text{Ins}(2,4,5)\text{P}_3$ (10 pmol), $\text{Ins}(1,3,4,5)\text{P}_4$ (10 pmol) had no further effect on $[\text{Ca}^{2+}]_i$. This indicated that $\text{Ins}(1,3,4,5)\text{P}_4$ -induced increase in Cl^- current seen in *B* was due to Ca^{2+} influx (see text).

Cl^- current ($n = 5$). The difference between the effect of $\text{Ins}(1,3,4,5)\text{P}_4$ and control injections was statistically significant ($P < 0.001$). These results contrast with previous work in rat hepatocytes that suggested that a predominant effect of $\text{Ins}(1,3,4,5)\text{P}_4$ was to stimulate reuptake of free Ca^{2+} back into intracellular stores (18). If that had been the case, $\text{Ins}(1,3,4,5)\text{P}_4$ would have reduced the Cl^- current caused by $\text{Ins}(2,4,5)\text{P}_3$; we found the opposite.

To determine whether the $\text{Ins}(1,3,4,5)\text{P}_4$ -induced increase in Cl^- current was due to intracellular Ca^{2+} release and/or to Ca^{2+} influx, we used the same experimental protocol [i.e., injection of $\text{Ins}(2,4,5)\text{P}_3$ followed by an injection of $\text{Ins}(1,3,4,5)\text{P}_4$] but this time in the continuous presence of extracellular Mn^{2+} . Under these conditions, the $\text{Ins}(1,3,4,5)\text{P}_4$ -induced increase in Cl^- cur-

rent would be inhibited if it was due to Ca^{2+} influx. Conversely, if $\text{Ins}(1,3,4,5)\text{P}_4$ released Ca^{2+} from intracellular stores, Cl^- current would increase. As previously reported (11, 36), Mn^{2+} did not inhibit the initial $\text{Ins}(2,4,5)\text{P}_3$ -induced $[\text{Ca}^{2+}]_i$ oscillations but abolished $\text{Ins}(2,4,5)\text{P}_3$ -induced Ca^{2+} influx (compare the tracings shown in Fig. 2A and Fig. 2C). As shown in Fig. 2C, subsequent injection of $\text{Ins}(1,3,4,5)\text{P}_4$ did not increase Cl^- current (4/4 cells). This indicates that $\text{Ins}(1,3,4,5)\text{P}_4$ does not release intracellular Ca^{2+} when injected after $\text{Ins}(2,4,5)\text{P}_3$; instead, it causes Ca^{2+} influx.

In four experiments, we injected increasing doses of $\text{Ins}(1,3,4,5)\text{P}_4$ after $\text{Ins}(2,4,5)\text{P}_3$. As the dose of injected $\text{Ins}(1,3,4,5)\text{P}_4$ increased, the magnitude of Ca^{2+} influx increased (Fig. 3). These results suggest that when pre-

Fig. 3. Injection of $\text{Ins}(2,4,5)\text{P}_3$ (10 pmol) followed by five sequential injections of $\text{Ins}(1,3,4,5)\text{P}_4$ (doses of 0.3, 1.8, 3, 30, and 150 pmol at closed arrows) (top trace). First $\text{Ins}(1,3,4,5)\text{P}_4$ injection inhibited $[\text{Ca}^{2+}]_i$ oscillations. Magnitude of $\text{Ins}(1,3,4,5)\text{P}_4$ -induced extracellular Ca^{2+} influx increased with amount of $\text{Ins}(1,3,4,5)\text{P}_4$ injected. Note that Mn^{2+} caused a recurrence of $[\text{Ca}^{2+}]_i$ oscillations before blocking current. Area enclosed by box is enlarged in bottom trace.

ceded by $\text{Ins}(2,4,5)\text{P}_3$, $\text{Ins}(1,3,4,5)\text{P}_4$ stimulates Ca^{2+} influx in a dose-dependent manner.

We considered the possibility that it is the increase in $[\text{Ca}^{2+}]_i$ produced by $\text{Ins}(1,4,5)\text{P}_3$ and $\text{Ins}(2,4,5)\text{P}_3$ that provides $\text{Ins}(1,3,4,5)\text{P}_4$ with the ability to stimulate Ca^{2+} influx. We did not think that this possibility was very likely because $\text{Ins}(1,3,4,5)\text{P}_4$ itself increases $[\text{Ca}^{2+}]_i$ but does not cause Ca^{2+} influx. To test this possibility further, we injected CaCl_2 in doses expected to increase the average $[\text{Ca}^{2+}]_i$ to 25 μM [in *Xenopus* oocytes, $\text{Ins}(1,4,5)\text{P}_3$ stimulation increases $[\text{Ca}^{2+}]_i$ to 0.25–1 μM (6, 17)]. Subsequent injection of $\text{Ins}(1,3,4,5)\text{P}_4$ caused $[\text{Ca}^{2+}]_i$ oscillations but failed to cause Ca^{2+} influx (8 cells) (Fig. 4). Thus an increase in $[\text{Ca}^{2+}]_i$ is not sufficient to induce $\text{Ins}(1,3,4,5)\text{P}_4$ -dependent Ca^{2+} influx. The presence of InsP_3 appears to be required.

Fig. 4. Injection of CaCl_2 (5 μM) followed by an injection of $\text{Ins}(1,3,4,5)\text{P}_4$. Absence of Mn^{2+} -inhibitable current indicates that an increase in $[\text{Ca}^{2+}]_i$ was not sufficient to confer $\text{Ins}(1,3,4,5)\text{P}_4$ with ability to cause Ca^{2+} influx. Changes in Ca^{2+} -gated Cl^- -current induced by injections of a given volume and concentration of CaCl_2 varied widely from cell to cell [218 ± 78 (SE) nA, range from 28 to 503 nA, $n = 8$; Ref. 10]. Discrepancy between magnitude of current responses to CaCl_2 and to $\text{Ins}(1,3,4,5)\text{P}_4$ was not consistent.

Effect of Ca^{2+} influx on $\text{Ins}(1,3,4,5)\text{P}_4$ generation. Once formed following membrane-receptor activation, $\text{Ins}(1,4,5)\text{P}_3$ follows two main metabolic pathways. In one pathway, it is dephosphorylated to $\text{Ins}(1,4)\text{P}_2$ by the $\text{Ins}(1,4,5)\text{P}_3$ 5-phosphatase (12). In a second pathway, it is phosphorylated to $\text{Ins}(1,3,4,5)\text{P}_4$ by the Ca^{2+} /calmodulin-dependent $\text{Ins}(1,4,5)\text{P}_3$ 3-kinase (4, 8, 39). The second pathway might be favored by the cell, in part, because the $\text{Ins}(1,4,5)\text{P}_3$ -induced increase in $[\text{Ca}^{2+}]_i$ would activate the 3-kinase and not the 5-phosphatase (3, 32). Because Ca^{2+} influx increases $[\text{Ca}^{2+}]_i$, we hypothesized that it might shunt the metabolism of $\text{Ins}(1,4,5)\text{P}_3$ toward the formation of $\text{Ins}(1,3,4,5)\text{P}_4$. To examine the effect of Ca^{2+} influx on the metabolism of $\text{Ins}(1,4,5)\text{P}_3$, we injected $[^3\text{H}]\text{Ins}(1,4,5)\text{P}_3$ into single *Xenopus* oocytes ($2\text{--}20 \times 10^3$ counts $\cdot \text{min}^{-1} \cdot \text{cell}^{-1}$) incubated in the presence or absence of extracellular Ca^{2+} . We then stopped InsP metabolism at predetermined time points. In cells with time points ≥ 1 min ($n = 56$), we recorded the electrophysiological response and found typical changes in $[\text{Ca}^{2+}]_i$ (Fig. 1A). In cells incubated in the absence of extracellular Ca^{2+} , the sustained increase in $[\text{Ca}^{2+}]_i$ shown in Fig. 1A was not present. HPLC separation of the $[^3\text{H}]\text{InsP}$ was performed for each individual cell ($n = 72$).

By 40 min after injection, $[^3\text{H}]\text{Ins}(1,4,5)\text{P}_3$ was almost completely metabolized (Fig. 5A). The rate of $\text{Ins}(1,4,5)\text{P}_3$ metabolism was comparable to that reported by Irvine et al. (20) and was not significantly influenced by extracellular Ca^{2+} (Fig. 5A).

The predominant InsP_4 isomer recovered was $\text{Ins}(1,3,4,5)\text{P}_4$. In the presence of extracellular Ca^{2+} , the maximal accumulation of $[^3\text{H}]\text{Ins}(1,3,4,5)\text{P}_4$ counts [11.9% of total disintegrations/min (dpm) injected] occurred 3 min after $[^3\text{H}]\text{Ins}(1,4,5)\text{P}_3$ injection (Fig. 5B, open circles). In the absence of extracellular Ca^{2+} (Fig. 5B, closed circles), the maximum increase in

Fig. 5. Effect of Ca^{2+} influx on immediate metabolism of $\text{Ins}(1,4,5)\text{P}_3$ in single *Xenopus* oocytes. Cells were injected with $[\text{H}^3]\text{Ins}(1,4,5)\text{P}_3$ [2,000–20,000 disintegrations/min (dpm)] in the presence (open circles) or absence (closed circles) of extracellular Ca^{2+} . Cellular metabolism was then arrested at various time points and InsPs were separated by high-pressure liquid chromatography (see MATERIALS AND METHODS). For each time point, 3–5 individual cells were studied in presence of extracellular Ca^{2+} and 2–3 cells in absence of extracellular Ca^{2+} (total of 72 cells). Radioactivity specific to each InsP was expressed as a percentage of total injected dpm. Value recorded at each time point represented mean $\pm$ SE of radioactivity attributable to $\text{Ins}(1,4,5)\text{P}_3$ (A), $\text{Ins}(1,3,4,5)\text{P}_4$ (B), or $\text{Ins}(1,4)\text{P}_2$ (C). In A, extracellular $[\text{Ca}^{2+}]$ had no effect on gradual metabolism of $\text{Ins}(1,4,5)\text{P}_3$. In B, generation of $\text{Ins}(1,3,4,5)\text{P}_4$ was increased in presence of extracellular Ca^{2+} (open circles). In C, production of $\text{Ins}(1,4)\text{P}_2$ was decreased in presence of extracellular Ca^{2+} (open circles). * Statistically different ($P < 0.05$). These results suggest that Ca^{2+} influx shunted metabolism of $\text{Ins}(1,4,5)\text{P}_3$ toward formation of $\text{Ins}(1,3,4,5)\text{P}_4$ (see text).

$[\text{H}^3]\text{Ins}(1,3,4,5)\text{P}_4$ was smaller (5.6% of total dpm injected) and occurred later (10 min). These results suggest that more $\text{Ins}(1,3,4,5)\text{P}_4$ is formed from $\text{Ins}(1,4,5)\text{P}_3$ in the presence of extracellular Ca^{2+} .

We also recovered $[\text{H}^3]\text{Ins}(1,3,4)\text{P}_3$, the direct metabolite of $\text{Ins}(1,3,4,5)\text{P}_4$. $[\text{H}^3]\text{Ins}(1,3,4)\text{P}_3$ counts began to increase 3 min after the $[\text{H}^3]\text{Ins}(1,4,5)\text{P}_3$ injection, peaked at 20 min and decreased thereafter (not shown). $[\text{H}^3]\text{Ins}(1,3,4,6)\text{P}_4$ was recovered in two cells incubated in the presence of extracellular Ca^{2+} , one 10 min and the other 20 min after injection of $[\text{H}^3]\text{Ins}(1,4,5)\text{P}_3$. In these two cells, there was also maximal accumulation of $\text{Ins}(1,3,4)\text{P}_3$, the proposed precursor of $\text{Ins}(1,3,4,6)\text{P}_4$ (1, 35). We did not recover $[\text{H}^3]\text{Ins}(3,4,5,6)\text{P}_4$; to do so would have required long-term labeling of inositol phospholipids (20, 35). These data are nevertheless consistent with the notion that $\text{Ins}(3,4,5,6)\text{P}_4$ is not an immediate intracellular second messenger (24).

The product of the $\text{Ins}(1,4,5)\text{P}_3$ 5'-phosphatase, $\text{Ins}(1,4)\text{P}_2$, was the only InsP_2 isomer recovered. The effect of extracellular Ca^{2+} on $[\text{H}^3]\text{Ins}(1,4)\text{P}_2$ levels was opposite to its effect on $[\text{H}^3]\text{Ins}(1,3,4,5)\text{P}_4$ levels. In the presence of extracellular Ca^{2+} , $[\text{H}^3]\text{Ins}(1,4)\text{P}_2$ counts began to rise at 3 minutes and gradually increased to a maximum (3.3% of initial dpm) throughout the 40-min time course (Fig. 5C, open circles). In the absence of extracellular Ca^{2+} , $[\text{H}^3]\text{Ins}(1,4)\text{P}_2$ counts reached their maximum (5.3% of initial dpm) 5 min after injection and then gradually decreased (Fig. 5C, closed circles). These results suggest that less $\text{Ins}(1,4)\text{P}_2$ is formed from $\text{Ins}(1,4,5)\text{P}_3$ in the presence of extracellular Ca^{2+} .

The data represent the proportion of the initial radioactivity recovered as $[\text{H}^3]\text{Ins}(1,3,4,5)\text{P}_4$ and $[\text{H}^3]\text{Ins}(1,4)\text{P}_2$ at given points in time and not the overall mass of these metabolites. Within individual cells, however, the ratios of $\text{Ins}(1,3,4,5)\text{P}_4/\text{Ins}(1,4)\text{P}_2$ counts were different in the presence or absence of extracellular Ca^{2+} (Table 1). The presence of Ca^{2+} produced a 10-fold increase in the $[\text{H}^3]\text{Ins}(1,3,4,5)\text{P}_4/\text{Ins}(1,4,5)\text{P}_3$ ratio. Despite the limits of our method, these results suggest that $\text{Ins}(1,4,5)\text{P}_3$ is preferentially metabolized to $\text{Ins}(1,3,4,5)\text{P}_4$ in the presence of extracellular Ca^{2+} .

Summary and implications. Our results indicate that $\text{Ins}(1,3,4,5)\text{P}_4$ does not cause Ca^{2+} influx by itself but does stimulate Ca^{2+} influx when preceded by poorly metabolizable InsP_3 . These results suggest a synergistic action of InsP_3 and $\text{Ins}(1,3,4,5)\text{P}_4$ to cause Ca^{2+} influx (7, 21, 22, 27).

Our results show that two poorly metabolizable InsP_3 s had different effects; $\text{Ins}(2,4,5)\text{P}_3$ stimulated Ca^{2+} influx, whereas $\text{Ins}(1,4,5)\text{P}_3$ did not. A previous report has suggested that the two analogues also generate different patterns of intracellular Ca^{2+} release (14). We do not think that the difference between $\text{Ins}(2,4,5)\text{P}_3$ and $\text{Ins}(1,4,5)\text{P}_3$ is caused by a different metabolism because we have previously shown that neither compound is metabolized significantly in the oocyte (11). Because $\text{Ins}(2,4,5)\text{P}_3$ alone causes Ca^{2+} influx, we previously argued that $\text{Ins}(1,3,4,5)\text{P}_4$ is not required to cause Ca^{2+} influx (36). The finding that $\text{Ins}(1,4,5)\text{P}_3$ does not cause Ca^{2+} influx could lead to the opposite interpretation, i.e., that $\text{Ins}(1,3,4,5)\text{P}_4$ is required for Ca^{2+} influx. Until we know which poorly metabolizable compound best reproduces the action of $\text{Ins}(1,4,5)\text{P}_3$, we may have to

Table 1. Effect of Ca^{2+} influx on the metabolism of $\text{Ins}(1,4,5)\text{P}_3$ in single *Xenopus* oocytes: ratios of $\text{Ins}(1,3,4,5)\text{P}_4$ to $\text{Ins}(1,4)\text{P}_2$ within individual cells

Time, min	$[\text{Ca}^{2+}]_o$	$\text{InsP}_4/\text{InsP}_2$ ratios	n
3	nominal	$0.4 \pm 0.1^*$	3
3	6 mM	$4.8 \pm 0.1^*$	4
5	nominal	0.25 ± 0.05	2
5	6 mM	2.1 ± 0.6	3

Three and five minutes following the injection of $[\text{H}^3]\text{Ins}(1,4,5)\text{P}_3$, ratio of recovered $[\text{H}^3]\text{Ins}(1,3,4,5)\text{P}_4$ and $[\text{H}^3]\text{Ins}(1,4)\text{P}_2$ was calculated for each *Xenopus* oocyte. Oocytes were bathed with high (6 mM) or low (nominal) extracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_o$). Values are expressed as means $\pm$ SE. * Value at 6 mM $[\text{Ca}^{2+}]_o$ statistically different ($P < 0.05$) from value at nominal $[\text{Ca}^{2+}]_o$.

use alternative strategies to resolve the controversy about the absolute requirement of $\text{Ins}(1,3,4,5)\text{P}_4$ for Ca^{2+} influx (19). Nevertheless, we observed that $\text{Ins}(1,3,4,5)\text{P}_4$ could stimulate Ca^{2+} influx when preceded by either $\text{Ins}(2,4,5)\text{P}_3$ or $\text{Ins}(1,4,5)\text{P}_3\text{S}_3$. Thus our data suggest that $\text{Ins}(1,3,4,5)\text{P}_4$ could regulate Ca^{2+} influx, even if it was not absolutely required.

Because of the unavailability of nonmetabolizable InsP_4 , we cannot rule out the possibility that a metabolite of $\text{Ins}(1,3,4,5)\text{P}_4$ is also involved in stimulating Ca^{2+} influx. Ivorra et al. (23) have recently shown that $\text{Ins}(1,3,4,6)\text{P}_4$ causes both intracellular Ca^{2+} release and Ca^{2+} influx in *Xenopus* oocytes. We have recovered $\text{Ins}(1,3,4,6)\text{P}_4$ only in cells incubated in the presence of extracellular Ca^{2+} . Therefore, the effect of Ca^{2+} influx on the metabolism of $\text{Ins}(1,4,5)\text{P}_3$ may promote the formation of more than one compound active in Ca^{2+} homeostasis.

Our data also suggest that Ca^{2+} influx can shunt the metabolism of $\text{Ins}(1,4,5)\text{P}_3$ toward the formation of $\text{Ins}(1,3,4,5)\text{P}_4$ and away from dephosphorylation to $\text{Ins}(1,4)\text{P}_2$. Because we performed our experiments in intact cells, our data suggest that the known Ca^{2+} dependence of the $\text{Ins}(1,4,5)\text{P}_3$ 3-kinase enzyme is physiologically relevant. Ca^{2+} influx increased $\text{Ins}(1,3,4,5)\text{P}_4$ levels over and above those found in cells where $\text{Ins}(1,4,5)\text{P}_3$ -induced release of intracellular Ca^{2+} had occurred. Thus the $\text{Ins}(1,4,5)\text{P}_3$ 3-kinase may require sustained high levels of $[\text{Ca}^{2+}]_i$ to remain activated. There may therefore be a positive feedback regulatory mechanism by which Ca^{2+} influx increases the amounts of $\text{Ins}(1,3,4,5)\text{P}_4$ produced from $\text{Ins}(1,4,5)\text{P}_3$ and $\text{Ins}(1,3,4,5)\text{P}_4$ stimulates Ca^{2+} influx. However, the physiological contribution of this mechanism to the regulation of Ca^{2+} influx remains to be determined.

We thank Antoinette Monod for performing the HPLC separation of InsP .

These studies were supported in part by a grant from the Swiss National Foundation (32-30161.90) to P. D. Lew and from the Science and Engineering Research Council (Molecular Recognition Initiative; UK) to B. V. L. Potter. D. Pittet is supported by a grant from the Schweizerische Stiftung fuer Medizinisch-Biologische Stipendien.

B. V. L. Potter is a Lister Institute Fellow.

Address for reprint requests: M. J. Welsh, Howard Hughes Medical Institute, Departments of Internal Medicine and Physiology and Biophysics, University of Iowa College of Medicine, Iowa City, Iowa 52242.

Received 18 September 1991; accepted in final form 13 January 1992.

REFERENCES

1. Balla, T., A. J. Baukal, G. Guillemette, and K. J. Catt. Multiple pathways of inositol polyphosphate metabolism in angiotensin-stimulated adrenal glomerulosa cells. *J. Biol. Chem.* 263: 4083-4091, 1988.
2. Berridge, M. J., and R. F. Irvine. Inositol phosphates and cell signalling. *Nature Lond.* 341: 197-205, 1989.
3. Biden, T. J., L. Vallar, and C. B. Wollheim. Regulation of inositol 1,4,5-trisphosphate metabolism in insulin-secreting RINm5F cells. *Biochem. J.* 251: 435-440, 1988.
4. Biden, T. J., and C. B. Wollheim. Calcium regulates the inositol tris/tetrakisphosphate pathway in intact and broken preparations of insulin-secreting RINm5F cells. *J. Biol. Chem.* 261: 11931-11934, 1986.
5. Bird, G. St. J., M. F. Rossier, A. R. Hughes, S. B. Shears, D. L. Armstrong, and J. W. Putney. Activation of Ca^{2+} entry into acinar cells by a non-phosphorylatable inositol trisphosphate. *Nature Lond.* 352: 162-165, 1991.
6. Brooker, G., T. Seki, D. Croll, and C. Wahlestedt. Calcium waves evoked by activation of endogenous or exogenously expressed receptors in *Xenopus* oocytes. *Proc. Natl. Acad. Sci. USA* 87: 2813-2817, 1990.
7. Changya, L., D. Gallacher, R. F. Irvine, B. V. L. Potter, and O. H. Petersen. Inositol 1,3,4,5-tetrakisphosphate is essential for sustained activation of the calcium-dependent K^+ current in single internally perfused mouse lacrimal acinar cells. *J. Membr. Biol.* 109: 85-93, 1989.
8. Choi, K. Y., H. K. Kim, S. Y. Lee, K. H. Moon, S. S. Sim, J. W. Kim, H. K. Chung, and S. G. Rhee. Molecular cloning and expression of a complementary DNA for inositol 1,4,5-trisphosphate 3-kinase. *Science Wash. DC* 248: 64-66, 1990.
9. Cooke, A. M., R. Gigg, and B. V. L. Potter. *myo*-Inositol 1,4,5-trisphosphorothioate: a novel analogue of a biological second messenger. *J. Chem. Soc., Chem. Commun.* 1525-1526, 1987.
10. Dascal, N. The use of *Xenopus* oocytes for the study of ion channels. *CRC Crit. Rev. Biochem.* 22: 317-387, 1987.
11. DeLisle, S., K.-H. Krause, G. Denning, B. V. L. Potter, and M. J. Welsh. Effects of inositol trisphosphate and calcium on oscillating elevations of intracellular calcium in *Xenopus* oocytes. *J. Biol. Chem.* 265: 11726-11730, 1990.
12. Downes, C. P., M. C. Mussat, and R. H. Michell. The inositol trisphosphate phosphomonoesterase of the human erythrocyte membrane. *Biochem. J.* 203: 169-177, 1982.
13. Ferguson, J. E., J.-K. Han, J. P. Y. Kao, and R. Nuccitelli. The effects of inositol trisphosphates and inositol tetrakisphosphate on Ca^{2+} release and Cl^- current pattern in the *Xenopus laevis* oocyte. *Exp. Cell Res.* 192: 352-365, 1991.
14. Ferguson, J. E., B. V. L. Potter, and R. Nuccitelli. The effect of *myo*-inositol 1,4,5-trisphosphorothioate on Cl^- current pattern and intracellular Ca^{2+} in the *Xenopus laevis* oocyte. *Biochem. Biophys. Res. Commun.* 172: 229-236, 1990.
15. Ferris, C. D., R. L. Haganir, S. Supattapone, and S. H. Snyder. Purified inositol 1,4,5-trisphosphate receptor mediates calcium flux in reconstituted lipid vesicles. *Nature Lond.* 342: 87-89, 1989.
16. Gawler, D. J., B. V. L. Potter, and S. R. Nahorski. Inositol 1,3,4,5-tetrakisphosphate-induced release of intracellular Ca^{2+} in SH-SY5Y neuroblastoma cells. *Biochem. J.* 272: 519-524, 1990.
17. Han, J. K., and R. Nuccitelli. Inositol 1,4,5-trisphosphate-induced calcium release in the organelle layers of the stratified intact egg of *Xenopus laevis*. *J. Cell Biol.* 110: 1103-1110, 1990.
18. Hill, T. D., N. M. Dean, and A. L. Boynton. Inositol 1,3,4,5-tetrakisphosphate induces Ca^{2+} sequestration in rat liver cells. *Science Wash. DC* 242: 1176-1178, 1988.
19. Irvine, R. F. Calcium control and InsP_4 (Letter). *Nature Lond.* 352: 115, 1991.
20. Irvine, R. F., A. J. Letcher, J. P. Heslop, and M. J. Berridge. The inositol tris/tetrakisphosphate pathway-demonstration of $\text{Ins}(1,4,5)\text{P}_3$ 3-kinase activity in animal tissue. *Nature Lond.* 320: 631-634, 1986.
21. Irvine, R. F., and R. M. Moor. Micro-injection of inositol 1,3,4,5-tetrakisphosphate activates sea urchin egg by a mechanism dependent on external calcium. *Biochem. J.* 240: 917-920, 1986.
22. Irvine, R. F., and R. M. Moor. Inositol (1,3,4,5) tetrakisphosphate-induced activation of sea urchin eggs requires the presence of inositol trisphosphate. *Biochem. Biophys. Res. Commun.* 146: 284-290, 1987.
23. Ivorra, I., R. Gigg, R. F. Irvine, and I. Parker. Inositol 1,3,4,6-tetrakisphosphate mobilizes calcium in *Xenopus* oocytes with high potency. *Biochem. J.* 273: 317-321, 1991.
24. Ji, H., K. Sandberg, A. J. Baukal, and K. J. Catt. Metabolism of inositol pentakisphosphate to inositol hexakisphosphate in *Xenopus laevis* oocytes. *J. Biol. Chem.* 264: 20185-20188, 1989.
25. Lupu-Meiri, M., H. Shapira, and Y. Oron. Extracellular calcium participates in responses to acetylcholine in *Xenopus* oocytes. *FEBS Lett.* 262: 165-169, 1990.
26. Morris, A. J., K. J. Murray, P. J. England, C. P. Downes, and R. H. Michell. Partial purification and some properties of rat brain inositol 1,4,5-trisphosphate 3-kinase. *Biochem. J.* 251: 157-163, 1988.
27. Morris, A. P., D. V. Gallacher, R. F. Irvine, and O. H. Petersen. Synergism of inositol trisphosphate and tetrakisphos-

- phate in activating calcium dependent K-channels. *Nature Lond.* 330: 653-655, 1987.
28. **Parker, I., and I. Ivorra.** Inositol tetrakisphosphate liberates stored Ca^{2+} in *Xenopus* oocytes and facilitates responses to inositol trisphosphate. *J. Physiol. Lond.* 433: 207-227, 1991.
 29. **Parker, I., and F. R. S. Miledi.** Injection of inositol 1,3,4,5-tetrakisphosphate in *Xenopus* oocytes generates a chloride current dependent upon intracellular calcium. *Proc. R. Soc. Lond. B Biol. Sci.* 232: 59-70, 1987.
 30. **Penner, R., G. Matthews, and E. Neher.** Regulation of calcium influx by second messengers in rat mast cells. *Nature Lond.* 334: 499-504, 1988.
 31. **Pittet, D., D. P. Lew, G. W. Mayr, A. Monod, and W. Schlegel.** Chemoattractant receptor promotion of calcium influx across the plasma membrane of HL-60 cells; a role for cytosolic free calcium elevation and inositol 1,3,4,5-tetrakisphosphate production. *J. Biol. Chem.* 264: 7251-7261, 1989.
 32. **Pittet, D., W. Schlegel, D. P. Lew, A. Monod, and G. W. Mayr.** Mass changes in inositol tetrakis- and pentakisphosphate isomers induced by chemotherapeutic peptide stimulation in HL-60 cells. *J. Biol. Chem.* 264: 18489-18493, 1989.
 33. **Putney, J. W., H. Takemura, A. R. Hughes, D. A. Horstman, and O. Thastrup.** How do inositol phosphates regulate calcium signaling. *FASEB J.* 3: 1899-1905, 1989.
 34. **Satoh, T., C. A. Ross, A. Villa, S. Supattapone, T. Pozzan, S. H. Snyder, and J. Meldolesi.** The inositol 1,4,5-trisphosphate receptor in cerebellar Purkinje cells: quantitative immunogold labeling reveals concentration in an ER subcompartment. *J. Cell Biol.* 111: 615-624, 1990.
 35. **Shears, S. B., J. B. Parry, E. K. Tang, R. F. Irvine, R. H. Michell, and C. J. Kirk.** Metabolism of D-myo-inositol 1,3,4,5-tetrakisphosphate by rat liver, including the synthesis of a novel isomer of myo-inositol tetrakisphosphate. *Biochem. J.* 246: 139-147, 1987.
 36. **Snyder, P. M., K-H. Krause, and M. J. Welsh.** Inositol trisphosphate isomers, but not inositol 1,3,4,5-tetrakisphosphate, induce calcium influx in *Xenopus laevis* oocytes. *J. Biol. Chem.* 263: 11048-11051, 1988.
 37. **Taylor, C. W., M. J. Berridge, K. D. Brown, A. M. Cooke, and B. V. L. Potter.** DL-Myo-inositol 1,4,5-trisphosphorothioate mobilizes intracellular calcium in swiss 3T3 cells and *Xenopus* oocytes. *Biochem. Biophys. Res. Commun.* 150: 626-632, 1988.
 38. **Taylor, C. W., M. J. Berridge, A. M. Cooke, and B. V. L. Potter.** Inositol 1,4,5-trisphosphorothioate, a stable analogue of inositol trisphosphate which mobilizes intracellular calcium. *Biochem. J.* 259: 645-650, 1989.
 39. **Yamaguchi, K., M. Hirata, and H. Kuriyama.** Calmodulin activates inositol 1,4,5-trisphosphate 3-kinase activity in pig aortic smooth muscle. *Biochem. J.* 244: 787-791, 1987.