



Citation for published version:

Kostakis, A, Magdalinos, T & Stamatogiannis, MP 2015, 'Robust econometric inference for stock return predictability', *Review of Financial Studies*, vol. 28, no. 5, pp. 1506-1553. <https://doi.org/10.1093/rfs/hhu139>

DOI:

[10.1093/rfs/hhu139](https://doi.org/10.1093/rfs/hhu139)

Publication date:

2015

Document Version

Early version, also known as pre-print

[Link to publication](#)

This article has been accepted for publication in *Review of Financial Studies* Published by Oxford University Press.

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Robust Econometric Inference for Stock Return Predictability

Alexandros Kostakis
University of Manchester

Tassos Magdalinos
University of Southampton

Michalis P. Stamatogiannis
University of Bath

We are grateful to Geert Bekaert (Editor), David Hirshleifer (Executive Editor), and four anonymous referees for their extensive comments and helpful suggestions. We would also like to thank Karim Abadir, Paul Bekker, Walter Distaso, Christoph Hanck, Erik Hjalmarsson, Jan Jacobs, Ioannis Kasparis, Lynda Khalaf, Timos Papadopoulos, Peter Phillips, Jean-Yves Pitarakis, Marno Verbeek (discussant), and Michael Wolf (discussant), as well as conference participants at Netherlands Econometric Study group 2011 (Rotterdam), ESEM 2011 (Oslo), Statistical Week 2011 (Leipzig), EFA 2012 (Copenhagen), RES 2014 (Manchester), and Marie Curie ITN on Risk Management and Risk Reporting (Konstanz) and seminar participants at Aston Business School, Bath, Loughborough, Manchester Business School, Queen Mary (London) and Southampton for their insightful comments and suggestions. Magdalinos gratefully acknowledges financial support by the ESRC (grant RES-061-25-0517). Send correspondence to Tassos Magdalinos, Department of Economics, University of Southampton, SO17 1BJ, Southampton, U.K.; telephone: +44 (0) 2380593175. E-mail: a.magdalinos@soton.ac.uk.

Robust Econometric Inference for Stock Return Predictability

Abstract

This study examines stock return predictability via lagged financial variables with unknown stochastic properties. We propose a novel testing procedure that (1) robustifies inference to regressors' degree of persistence, (2) accommodates testing the joint predictive ability of financial variables in multiple regression, (3) is easy to implement as it is based on a linear estimation procedure, and (4) can be used for long-horizon predictability tests. We provide some evidence in favor of short-horizon predictability during the 1927-2012 period. Nevertheless, this evidence almost entirely disappears in the post-1952 period. Moreover, predictability becomes weaker, not stronger, as the predictive horizon increases. (*JEL* C12, C32, C58, G12, G14)

A fundamental issue in finance is whether future stock returns are predictable using publicly available information (see Fama 1970). The seminal studies of Keim and Stambaugh (1986), Fama and French (1988), and Campbell and Shiller (1988) empirically demonstrated that certain financial variables have significant predictive ability over future stock returns. Fama (1991) interpreted these findings as evidence of time-varying risk premia rather than as evidence against market efficiency. Despite the significant volume of subsequent research, the predictability debate still remains unsettled (see Ang and Bekaert 2007 for an insightful discussion). On the one hand, Lettau and Ludvigson (2001, 842) state that “it is now widely accepted that excess returns are predictable by variables such as dividend-price ratios, earning-price ratios, dividend-earnings ratios and an assortment of other financial indicators.” But many remain skeptical, claiming that the “profession has yet to find some variable that has meaningful and robust empirical equity premium forecasting power both in-sample and out-of-sample” (Welch and Goyal 2008, 1505).

Empirical support of arguments in favor of or against predictability crucially relies on inference from predictive regressions, and hence the size and power of the employed hypothesis tests assume fundamental importance. A series of recent studies, reviewed by Campbell and Yogo (2006) (hereafter CY), recognize that the most common problem undermining confidence in the reliability of predictability tests is the uncertainty about the (unobservable) time-series properties of the predictor variables and, in particular, their degree of persistence. Regardless of one’s prior beliefs on their order of integration, it is well documented that most of the variables used in predictive regressions are highly persistent with autoregressive roots extremely close to unity (see CY; Welch and Goyal 2008). This empirical fact casts doubt on the validity of standard t -tests based on least-squares regressions (see Cavanagh, Elliott, and Stock 1995; Torous, Valkanov, and Yan 2004). As Stambaugh (1999) has convincingly shown, this problem is exacerbated if additionally the innovations of the predictor are highly correlated with the innovations of the returns, that is, when the predictive regressor is endogenous. Endogeneity is a typical feature of commonly used predictors, such as price-scaled ratios. Because regression estimators and tests have fundamentally different properties in the presence of persistent and endogenous predictors, confidence in the reliability of predictability tests is undermined, as the quality of inference is conditional upon correct specification of the predictors’ time-series properties.

Acknowledging the uncertainty regarding the degree of predictive variables’ persistence, a

strand of the literature suggests modeling these variables as local-to-unity processes (see, inter alia, Lanne, 2002; Valkanov 2003; Torous et al. 2004; CY; Jansson and Moreira 2006; Hjalmars-son 2011). These processes assume the form of a first-order autoregression with root $\rho = 1 + c/n$, approaching a random walk as the sample size n increases to infinity. While providing flexibility in modeling, the use of explanatory variables that exhibit persistence without necessarily being random walks in finite samples raises serious technical complications. Because standard cointegration methods cannot accommodate the presence of local-to-unity roots in predictive regressions, Cavanagh, Elliott, and Stock (1995), Torous, Valkanov, and Yan (2004), CY, and Hjalmars-son (2011) have employed methods based on inverting the nonpivotal limit distribution of the t -statistic and constructing Bonferroni-type confidence intervals for the nuisance parameter c . This is the current state of the art methodology for testing the predictability of stock returns with highly persistent regressors.

Practical implementation of the above methodology presents two main drawbacks. First, the method is invalid if the regressor contains stationary or near-stationary components; the validity of the method requires each predictor to be at least as persistent as a local-to-unity process, a restrictive assumption that cannot be empirically tested. Second, because of the problems associated with the construction of multidimensional confidence intervals for c , the methodology is restricted to the case of a scalar regressor, that is, a single predictive variable. This imposes a severe restriction, because the joint predictability by combinations of financial variables cannot be tested. The above framework can only accommodate testing the predictive power of each financial variable in isolation, which may result in loss of information through omitted variables. These limitations also have been indicated by Ang and Bekaert (2007, footnote 3). We build upon this strand of the literature by proposing a methodology that successfully overcomes these limitations.

In recent work, Phillips and Magdalinos (2009) provide a framework of limit theory that can be used to validate inference in models with regressors exhibiting very general time-series characteristics. Endogeneity is successfully removed by means of a data-filtering procedure called IVX estimation. The key idea behind the method is the explicit control of the degree of persistence of data-filtered IVX instruments, restricted within the class of near-stationary processes. In this study, we prove that in the context of multivariate predictive regressions, the

IVX approach yields standard chi-squared asymptotic inference for testing general restrictions on predictive variables with a degree of persistence covering the entire range from stationarity of stable autoregressions to pure nonstationarity of unit root processes. The robustness of the IVX approach should alleviate practical concerns about the quality of inference under possible misspecification of the time-series properties of the predictive regressors. The dimensionality of the system of predictive regressions is of considerable practical importance too, because the IVX methodology enables the assessment of the joint predictive power of various combinations of regressors.¹ In summary, our study introduces and implements a testing procedure that resolves two important outstanding issues in the predictability literature: (1) robustness with respect to the time-series properties of the predictors and (2) joint testing in systems of predictive equations. Furthermore, we show that this testing procedure is also applicable to long-horizon predictive regressions, and we develop the relevant statistic.

We implement the proposed methodology by conducting a battery of short- and long-horizon predictability tests for U.S. stock returns during the 1927–2012 period, using a set of commonly employed variables. We focus on in-sample predictability tests because the proposed methodology aims to robustify in-sample inference with respect to regressors’ unknown time-series properties. In univariate tests, we find significant predictive ability with respect to one-period-ahead excess market returns for the earnings-price and book-to-market value ratios, as well as net equity expansion. However, this evidence almost entirely disappears in the post-1952 period. Only the consumption-wealth ratio is found to be strongly significant in this subperiod. Our multivariate tests show that the combination of the earnings-price ratio and T-bill rate is highly significant and robust to the choice of data frequency and examined period. Finally, with respect to long-horizon tests, we find that, if anything, predictability generally becomes weaker, not stronger, as the horizon increases. Only the consumption-wealth ratio remains strongly significant for all horizons examined.

¹It should be noted that the iterative procedure of Amihud, Hurvich, and Wang (2009) also accommodates multiple predictors under the restriction that these are stationary. Moreover, the recent contribution of Kelly and Pruitt (2013) also utilizes a multivariate system of predictive regressions. However, their focus is on extracting information regarding aggregate expected returns and dividend growth from the cross-section of price-dividend ratios using the present value relationship that has been employed for predictability tests *inter alia* by Lettau and Ludvigson (2005), Cochrane (2008), Lettau and van Nieuwerburgh (2008), and Binsbergen and Koijen (2010).

1. Robust Inference for Predictive Regressions

This section develops an econometric methodology for testing stock return predictability that is robust to uncertainty over the stochastic properties of the financial variables used as potential predictors. Accommodating this uncertainty requires a modeling framework that encompasses all empirically relevant classes of autoregressive data-generating mechanisms. To this end, we consider the following multivariate system of predictive regressions with regressors containing explanatory variables with arbitrary degree of persistence:

$$y_t = \mu + Ax_{t-1} + \varepsilon_t, \quad (1)$$

$$x_t = R_n x_{t-1} + u_t, \quad (2)$$

where A is an $m \times r$ coefficient matrix and

$$R_n = I_r + \frac{C}{n^\alpha} \quad \text{for some } \alpha \geq 0, \quad (3)$$

and some matrix $C = \text{diag}(c_1, \dots, c_r)$, where n denotes the sample size. The vector of predictive variables x_t in (2) exhibits a degree of persistence induced by the autoregressive matrix in (3) that belongs to one of the following persistence classes:

P(i) *Integrated regressors, if $C = 0$ or $\alpha > 1$ in (3),*

P(ii) *Local-to-unity regressors, if $C \neq 0$ and $\alpha = 1$ in (3),*

P(iii) *Near-stationary regressors, if $c_i < 0$ for all i and $\alpha \in (0, 1)$ in (3),*

P(iv) *Stationary regressors, if $c_i < 0$ for all i and $\alpha = 0$ in (3).*

The classes P(i)-P(iv) include predictors with very general time-series characteristics varying from purely stationary to purely non-stationary processes and accommodating all intermediate persistence regimes. The predictive regression system may be initialized at some x_0 that could be any fixed constant vector or a random process satisfying $\|x_0(n)\| = o_p(n^{1/2})$ when $\alpha \geq 1$ or $\alpha = 0$ and $\|x_0(n)\| = o_p(n^{\alpha/2})$ when $\alpha \in (0, 1)$.

Estimators and test statistics for conducting inference on the matrix A have very different properties according to the classification of the predictor process in (2) into one of the above

persistence classes. Standard tests are asymptotically valid only within each class P(i)–P(iv) and misspecification of the degree of predictor persistence may lead to severe size distortions, particularly in the presence of endogeneity, that is, correlation between the innovations ε_t and u_t of the predictive regression system (1)–(2) (see Elliott 1998).² CY have partly addressed the problem for univariate predictive regressions ($m = r = 1$ in (1)–(2)) by inverting the limit distribution of the t -statistic under a local-to-unity regime P(ii) and using the Bonferroni inequality to construct confidence intervals which are asymptotically valid under P(i) or P(ii). However, the CY method loses its asymptotic validity for predictors that lie closer to the stationary region than to local-to-unity time-series. Such predictors can be modeled either as local-to-unity processes, with c_i in (3) being large in absolute value (Phillips 1987), or, more formally, as belonging to the class P(iii) of near-stationary processes established by Phillips and Magdalinos (2007) and extended to multivariate systems of regression equations by Magdalinos and Phillips (2009).

We provide valid inference on A when there is no a priori knowledge of whether x_t belongs to class P(i), P(ii), P(iii), or P(iv). Our methodology for achieving robust inference is based on the IVX instrumentation procedure proposed by Phillips and Magdalinos (2009). The intuition behind this procedure is to construct an instrumental variable whose degree of persistence we explicitly control. In this way, the inference problems arising due to the uncertainty regarding the persistence of the original regressor are avoided. Using the constructed instrument, one then performs a standard instrumental variable estimation. The derived estimator asymptotically follows a mixed normal distribution, and hence the corresponding Wald statistic asymptotically follows a chi-squared distribution under the null, considerably simplifying inference.

To fix ideas, we construct near-stationary instruments belonging to the class P(iii) by differencing the regressor x_t and constructing a new process according to an artificial autoregressive

²In general, long-run endogeneity cannot be removed by standard cointegration methods, such as the fully modified least-squares estimation of Phillips and Hansen (1990) or the approaches of Saikkonen (1991) and Stock and Watson (1993), that apply when the regressor is a pure random walk ($c = 0$). As pointed out by Elliott (1998), such endogeneity corrected estimators lose their asymptotic mixed-normality property under a local-to-unity regime and the associated hypothesis tests have a nonstandard limit distribution, with the noncentrality parameter depending on the coefficient c of the local-to-unity root. Because c cannot be consistently estimated, the endogeneity cannot be removed, leading to asymptotically invalid predictability tests. Analogous problems arise when predictors exhibit a lower degree of persistence relative to local-to-unity processes, as is the case with the class of “near-stationary” processes introduced by Phillips and Magdalinos (2007) as well as stationary autoregressive processes.

matrix with specified persistence degree. Despite the fact that the difference

$$\Delta x_t = u_t + \frac{C}{n^\alpha} x_{t-1}$$

is not an innovation unless the regressor belongs to the class P(i) of integrated processes, it behaves asymptotically as an innovation after linear filtering by a matrix consisting of near-stationary roots of the type P(iii). Choosing an artificial matrix,

$$R_{nz} = I_r + \frac{C_z}{n^\beta}, \quad \beta \in (0, 1), \quad C_z < 0, \quad (4)$$

IVX instruments \tilde{z}_t are constructed as a first-order autoregressive process with autoregressive matrix R_{nz} and innovations Δx_t ,

$$\tilde{z}_t = R_{nz} \tilde{z}_{t-1} + \Delta x_t, \quad (5)$$

initialized at $\tilde{z}_0 = 0$. In particular, we use $C_z = -I_r$ and $\beta = 0.95$.

This choice of β follows from the size and power properties of the subsequently derived Wald test.³ Extensive Monte Carlo simulations presented in the Online Appendix show that the finite-sample size of the test is very close to the nominal 5% level regardless of the value of β . This holds true for all cases of regressor persistence considered. With respect to the power of the test, we find that it increases monotonically as β increases for all cases considered. This property is also suggested by the $n^{(1+\beta)/2}$ rate of convergence of the IVX estimator in Theorem A(i) provided in the Appendix. A closer inspection of the reported power plots suggests that starting from low or moderate values of β , there are considerable power gains when we further increase β towards its upper boundary, especially when the true value of A is closer to the null. Given this evidence, we confidently argue that high values of β yield the highest level of power for the Wald test and, at the same time, yield size very close to the nominal 5% level. Therefore, in the empirical implementation of our testing procedure, we use $\beta = 0.95$, which is among the highest values that β can take. Moreover, we strongly advise against using values of β less than 0.9, as they may lead to unnecessary loss of power for the test statistic.⁴

³To be precise, we follow the convention in prior literature and use the term "size" throughout this study to indicate the "probability of a type I error" for the various test statistics considered.

⁴We would like to thank an anonymous referee for suggesting this clarification.

As it is standard in the literature, we assume that the innovations ε_t of the predictive Equation (1) are uncorrelated, while allowing for correlation in the innovations of the predictor sequence u_t . The dependence structure of the innovations is formally presented below: part (i) provides assumptions under conditional homoscedasticity; part (ii) accommodates a general form of conditional heteroscedasticity under additional assumptions.

Assumption INNOV. Let $\varepsilon_t = (\varepsilon'_t, e'_t)'$, with ε_t as in (1), denote an R^{m+r} -valued martingale difference sequence with respect to the natural filtration $F_t = \sigma(\varepsilon_t, \varepsilon_{t-1}, \dots)$ satisfying

$$E_{\mathcal{F}_{t-1}}(\varepsilon_t \varepsilon'_t) = \Sigma_t \text{ a.s. and } \sup_{t \in \mathbb{Z}} E \|\varepsilon_t\|^{2s} < \infty \quad (6)$$

for some $s > 1$, where Σ_t is a positive definite matrix. Let u_t in (2) be a stationary linear process

$$u_t = \sum_{j=0}^{\infty} C_j e_{t-j} \quad (7)$$

where $(C_j)_{j \geq 0}$ a sequence of constant matrices such that $\sum_{j=0}^{\infty} C_j$ has full rank and $C_0 = I_r$.

We maintain one of the following assumptions:

(i) $\Sigma_t = \Sigma_\varepsilon$ for all t and $\sum_{j=0}^{\infty} \|C_j\| < \infty$.

(ii) The process $(\varepsilon_t)_{t \in \mathbb{Z}}$ is strictly stationary ergodic satisfying (6) with $s = 2$ and

$$\lim_{m \rightarrow \infty} \|Cov[\text{vec}(\varepsilon_m \varepsilon'_m), \text{vec}(\varepsilon_0 \varepsilon'_0)]\| = 0. \quad (8)$$

The sequence $(C_j)_{j \geq 0}$ in (7) satisfies

$$\sum_{j=0}^{\infty} j \|C_j\| < \infty. \quad (9)$$

The sequence $(\varepsilon_t)_{t \in \mathbb{Z}}$ admits the following vec-GARCH(p, q) representation:

$$\varepsilon_t = H_t^{1/2} \eta_t, \quad \text{vec}(H_t) = \bar{\varphi} + \sum_{i=1}^q A_i \text{vech}(\varepsilon_{t-i} \varepsilon'_{t-i}) + \sum_{k=1}^p B_k \text{vech}(H_{t-k}) \quad (10)$$

where $(\eta_t)_{t \in \mathbb{Z}}$ is an i.i.d. $(0, I_m)$ sequence, $\bar{\varphi}$ is a constant vector, A_i, B_k are symmetric positive semidefinite matrices for all i, k , and the spectral radius of the matrix

$\Gamma = \sum_{i=1}^q A_i + \sum_{k=1}^p B_k$ satisfies $\rho(\Gamma) < 1$.

Assumption INNOV(i) imposes conditional homoscedasticity on the martingale difference sequence ϵ_t and short-memory on the linear process (7). Assumption INNOV(ii) accounts for conditionally heteroscedastic ϵ_t with finite fourth-order moments of a very general form: the vec-GARCH process in (10) is the most general multivariate GARCH specification (see Chapter 11 of Francq and Zakoian 2010).⁵

Following standard notational convention, we define the short-run and long-run covariance matrices associated with the innovations ϵ_t and u_t in (1), (2) as follows:

$$\Sigma_{\epsilon\epsilon} = E(\epsilon_t \epsilon_t'), \quad \Sigma_{\epsilon u} = E(\epsilon_t u_t'), \quad \Sigma_{uu} = E(u_t u_t'), \quad (11)$$

$$\Omega_{uu} = \sum_{h=-\infty}^{\infty} E(u_t u_{t-h}'), \quad \Omega_{\epsilon u} = \Sigma_{\epsilon u} + \Lambda'_{u\epsilon}, \quad \Lambda_{u\epsilon} = \sum_{h=1}^{\infty} E(u_t \epsilon_{t-h}'). \quad (12)$$

Note that $\Omega_{\epsilon u}$ is only a one-sided long-run covariance matrix because ϵ_t is an uncorrelated sequence by Assumption INNOV. For the same reason, the long-run covariance of the ϵ_t sequence is equal to the short-run covariance $\Sigma_{\epsilon\epsilon}$. Denoting by $\hat{\epsilon}_t$ the OLS residuals from (1) and by \hat{u}_t the OLS residuals from (2), the covariance matrices in (11) can be estimated in a standard way:

$$\hat{\Sigma}_{\epsilon\epsilon} = \frac{1}{n} \sum_{t=1}^n \hat{\epsilon}_t \hat{\epsilon}_t', \quad \hat{\Sigma}_{\epsilon u} = \frac{1}{n} \sum_{t=1}^n \hat{\epsilon}_t \hat{u}_t', \quad \hat{\Sigma}_{uu} = \frac{1}{n} \sum_{t=1}^n \hat{u}_t \hat{u}_t'. \quad (13)$$

Accommodating autocorrelation in u_t that takes the general form (7) requires nonparametric estimation of the long-run covariance matrices in (12): letting M_n be a bandwidth parameter satisfying $M_n \rightarrow \infty$ and $M_n/\sqrt{n} \rightarrow 0$ as $n \rightarrow \infty$, we employ the usual Newey-West-type

⁵The positive semidefinite condition on the matrices A_i, B_k of (10) and the condition on the spectral radius of their sum are part of the standard Boussama (2006) stationarity conditions for the vec-GARCH process; see Theorem 11.5 of Francq and Zakoian (2010). Condition (8) is a mild weak dependence requirement on the process $\text{vec}(\epsilon_t \epsilon_t')$: it is satisfied if ϵ_t is given a vec-GARCH specification analogous to ϵ_t , but the results in this paper do not require a parametric specification of the conditional variance structure of the e_t process. A general discussion of the rate of decay of the autocovariance function in (8) in the case of univariate conditionally heteroscedastic time series admitting a stationary ARCH(∞) representation is included in Giraitis, Kokoszka, and Leipus (2000). The summability condition (9) is standard in the literature on short-memory linear processes (see Phillips and Solo, 1992).

estimators

$$\hat{\Lambda}_{uu} = \frac{1}{n} \sum_{h=1}^{M_n} \left(1 - \frac{h}{M_n + 1}\right) \sum_{t=h+1}^n \hat{u}_t \hat{u}'_{t-h}, \quad \hat{\Omega}_{uu} = \hat{\Sigma}_{uu} + \hat{\Lambda}_{uu} + \hat{\Lambda}'_{uu} \quad (14)$$

$$\hat{\Lambda}_{u\varepsilon} = \frac{1}{n} \sum_{h=1}^{M_n} \left(1 - \frac{h}{M_n + 1}\right) \sum_{t=h+1}^n \hat{u}_t \hat{\varepsilon}'_{t-h}, \quad \hat{\Omega}_{\varepsilon u} = \hat{\Sigma}_{\varepsilon u} + \hat{\Lambda}'_{u\varepsilon}. \quad (15)$$

Under the full generality of Assumption INNOV, we provide robust inference for the matrix of coefficients A that is invariant to the predictive variables belonging to classes P(i)-P(iv).

Allowing for the presence of an intercept in the predictive Equation (1) requires further development of IVX estimation and testing theory. The first step is to use a standard demeaning transformation of (1) that yields exact invariance of estimation of A to the presence of an intercept. We denote sample averages of variates in the system (1)–(2) by $\bar{y}_n = n^{-1} \sum_{t=1}^n y_t$, $\bar{x}_{n-1} = n^{-1} \sum_{t=1}^n x_{t-1}$, $\bar{\varepsilon}_n = n^{-1} \sum_{t=1}^n \varepsilon_t$, the demeaned variates by $Y_t = y_t - \bar{y}_n$, $X_t = x_t - \bar{x}_{n-1}$ and $\mathcal{E}_t = \varepsilon_t - \bar{\varepsilon}_n$, the resulting (demeaned) regression matrices by $\underline{Y} = (Y'_1, \dots, Y'_n)'$ and $\underline{X} = (X'_0, \dots, X'_{n-1})'$, and the (undemeaned) instrument matrix by $\tilde{Z} = (\tilde{z}'_0, \dots, \tilde{z}'_{n-1})'$. We may obtain invariance to the presence of the intercept μ in the predictive equation by subtracting $\bar{y}_n = \mu + A\bar{x}_{n-1} + \bar{\varepsilon}_n$ from (1) and obtaining the transformed predictive equation

$$Y_t = AX_{t-1} + \mathcal{E}_t. \quad (16)$$

We now proceed with IVX estimation of A from the predictive regression system (16), by considering a two-stage least-squares estimator based on the near-stationary instruments in (5):

$$\tilde{A}_{IVX} = \underline{Y}' \tilde{Z} \left(\underline{X}' \tilde{Z} \right)^{-1} = \sum_{t=1}^n (y_t - \bar{y}_n) \tilde{z}'_{t-1} \left[\sum_{j=1}^n (x_j - \bar{x}_{n-1}) \tilde{z}'_{j-1} \right]^{-1}. \quad (17)$$

Note that the estimator does not involve a demeaned version of the matrix of instruments, as the IVX estimator in (17) is invariant to demeaning \tilde{z}_{t-1} by \bar{z}_{n-1} .

The asymptotic behavior of the normalized and centered IVX estimator in (17) is summarized by Theorem A in the Appendix. The varying persistence levels of the predictor process in (2) and the effect of estimating an intercept in the predictive model (1) become manifest only in the limit distribution of the normalized signal matrix $\underline{X}' \tilde{Z}$. After appropriate centering and

normalization, the $\underline{Y}'\tilde{Z}$ component of the IVX estimator converges in distribution to a Gaussian variate that is independent of the (possibly) random limit in distribution of the signal matrix. As a result, the IVX estimator in (17) follows a mixed Gaussian limit distribution regardless of the degree of persistence of the predictor variable in (2).

The asymptotic mixed normality property of the IVX procedure implies that linear restrictions on the coefficients A generated by the system of predictive Equations (1) can be tested by a standard Wald test based on the IVX estimator for all persistence scenarios conforming to the classes P(i)-P(iv). In particular, we consider a set of linear restrictions

$$H_0 : H \text{vec}(A) = h, \quad (18)$$

where H is a known $q \times mr$ matrix with rank q and h is a known vector. We propose the following IVX-Wald statistic for testing H_0 in (18):

$$W_{IVX} = \left(H \text{vec} \tilde{A}_{IVX} - h \right)' Q_H^{-1} \left(H \text{vec} \tilde{A}_{IVX} - h \right), \quad (19)$$

where \tilde{A}_{IVX} is the IVX estimator in (17),

$$\begin{aligned} Q_H &= H \left[\left(\tilde{Z}' \underline{X} \right)^{-1} \otimes I_m \right] \mathbb{M} \left[\left(\underline{X}' \tilde{Z} \right)^{-1} \otimes I_m \right] H', \\ \mathbb{M} &= \tilde{Z}' \tilde{Z} \otimes \hat{\Sigma}_{\varepsilon\varepsilon} - n \bar{z}_{n-1} \bar{z}'_{n-1} \otimes \hat{\Omega}_{FM}, \end{aligned} \quad (20)$$

$$\hat{\Omega}_{FM} = \hat{\Sigma}_{\varepsilon\varepsilon} - \hat{\Omega}_{\varepsilon u} \hat{\Omega}_{uu}^{-1} \hat{\Omega}'_{\varepsilon u}, \quad (21)$$

and the matrices $\hat{\Sigma}_{\varepsilon\varepsilon}$, $\hat{\Omega}_{\varepsilon u}$ and $\hat{\Omega}_{uu}$ are defined in (13), (14), and (15).

Theorem 1. Consider the model (1)–(3) under Assumption INNOV with instruments \tilde{z}_t defined by (4) and (5). Then the Wald statistic in (19) for testing (18) satisfies

$$W_{IVX} \Rightarrow \chi^2(q) \quad \text{as } n \rightarrow \infty$$

under H_0 , for the following classes of predictor processes x_t in (2):

- (i) P(i)-P(iv) under Assumption INNOV(i),

(ii) P(i)-P(iii) under Assumption INNOV(ii).

The proof of Theorem 1 can be found in the Online Appendix. Theorem 1 establishes the robustness of the IVX-Wald test in (19) to the persistence properties of the predictor process in (2). It shows that the IVX methodology provides a unifying framework of inference in predictive regressions that encompasses the whole range of empirically relevant autoregressive data-generating mechanisms, from stationary processes to purely nonstationary random walks.

The only class of predictor variables not covered by Theorem 1 is that of purely stationary autoregressions P(iv) with conditionally heteroscedastic innovations. This is by no means surprising because, in the above case, the IVX-Wald test statistic is asymptotically equivalent to a standard Wald statistic of the form:

$$W_n = \left(H \text{vec} \hat{A}_{OLS} - h \right)' \left[H \left\{ (\underline{X}' \underline{X})^{-1} \otimes \hat{\Sigma}_{\varepsilon\varepsilon} \right\} H' \right]^{-1} \left(H \text{vec} \hat{A}_{OLS} - h \right),$$

with \hat{A}_{OLS} as the usual OLS estimator. It is well known that, even with a priori knowledge that x_t is a stationary process, W_n will not have a $\chi^2(q)$ limit distribution when the innovation sequence ε_t in (1) is conditionally heteroscedastic because the asymptotic variance of $n^{-1/2} \sum_{t=1}^n (x_{t-1} \otimes \varepsilon_t)$ is given by $\Upsilon = E(x_{t-1} x'_{t-1} \otimes \varepsilon_t \varepsilon'_t)$ and does not factorize to $E(x_{t-1} x'_{t-1}) \otimes \Sigma_{\varepsilon\varepsilon}$ as in the case in which ε_t are conditionally homoscedastic (see Equation (35) in Theorem A). Consequently, the matrix $(\underline{X}' \underline{X})^{-1} \otimes \hat{\Sigma}_{\varepsilon\varepsilon}$ is no longer a consistent estimator of the asymptotic variance of $\text{vec}(\hat{A}_{OLS})$ and W_n will fail to be asymptotically $\chi^2(q)$.

This failure is a characteristic of least-squares regression rather than IVX methodology. It can be rectified by introducing a White-type (1980) of correction in the Wald statistic. In particular, using $\hat{\Upsilon}_n = n^{-1} \sum_{t=1}^n (\tilde{z}_{t-1} \tilde{z}'_{t-1} \otimes \hat{\varepsilon}_t \hat{\varepsilon}'_t)$ as an estimator for Υ , with $\hat{\varepsilon}_t$ being the OLS residuals from (1), and replacing the matrix \mathbb{M} in (20) by $\tilde{\mathbb{M}} = n \hat{\Upsilon}_n - n \bar{z}_{n-1} \bar{z}'_{n-1} \otimes \hat{\Omega}_{FM}$, makes the IVX-Wald statistic in (19) heteroscedasticity-robust even in the P(iv) case.

The robustness of the IVX-Wald statistic to conditional heteroscedasticity for the persistence classes P(i)-P(iii) is a novel result of considerable interest: it depends on establishing the invariance, under Assumptions INNOV(i) and INNOV(ii), of the limit distribution in the central limit theorem for $n^{-(1+\alpha)/2} \sum_{t=1}^n (x_{t-1} \otimes \varepsilon_t)$ for $\alpha \in (0, 1)$; see Lemma B4 of the Online Appendix. Because the IVX instrument \tilde{z}_t behaves asymptotically as a near-stationary process (z_t if $\beta < \alpha$

and x_t if $\beta > \alpha$), Lemma B4 ensures that $\text{vec}(\tilde{A}_{IVX})$ will have the same asymptotic variance under Assumptions INNOV(i) and INNOV(ii) for all $\alpha > 0$. The methods developed in Lemma B4 can be used in a wider context to show that any amount of persistence (even of arbitrarily small order) in the regressor x_t alleviates asymptotically the effect of conditional heteroscedasticity and results to t and Wald statistics with standard limit distributions. This phenomenon becomes manifest in long-horizon predictive regressions when the horizon parameter tends to infinity with the sample size; see Theorem 2(ii) in Section 5.

Removing the finite-sample distortion to the mixed normal limit distribution of the IVX estimator caused by the estimation of the intercept is another subtle issue. The component Q_H in the quadratic form of the Wald statistic in (19) contains a finite-sample correction in the form of a weighted demeaning of the dominating term $\tilde{Z}'\tilde{Z} \otimes \hat{\Sigma}_{\varepsilon\varepsilon}$ of \mathbb{M} in (20) by $n\bar{z}_{n-1}\bar{z}'_{n-1} \otimes \hat{\Omega}_{FM}$. Despite not contributing to the first-order limit theory for W_{IVX} , this correction removes the finite-sample effects of estimating an intercept in (1). As discussed in Remark A(b) of the Appendix, these effects are more prominent for highly persistent regressors that are strongly correlated with the predictive model's innovations ε_t . Weighting the demeaning in (20) by the long-run covariance matrix $\hat{\Omega}_{FM}$ (which appears in Phillips and Hansen's [1990] FM-endogeneity correction for integrated systems) controls the effect of correlation between ε_t and u_t on the remainder term of the Gaussian first-order approximation (see Equation (36) in the Appendix) by the degree of demeaning of the instrument moment matrix $\tilde{Z}'\tilde{Z}$. To obtain better intuition on the nature of the correction in (20), assume for simplicity that $m = r = 1$; then

$$\mathbb{M} = \left[\sum_{t=1}^n \tilde{z}_{t-1}^2 - n\bar{z}_{n-1}^2 (1 - \hat{\rho}_{\varepsilon u}^2) \right] \hat{\Sigma}_{\varepsilon\varepsilon}$$

where $\hat{\rho}_{\varepsilon u} = \hat{\Omega}_{\varepsilon u} / \sqrt{\hat{\Sigma}_{\varepsilon\varepsilon} \hat{\Omega}_{uu}}$ is an estimator of the long-run correlation coefficient between ε_t and u_t . Therefore, the correction in (20) applies a weighting of the demeaning of the $\tilde{Z}'\tilde{Z}$ matrix according to the magnitude of the absolute value of the long-run correlation coefficient $\rho_{\varepsilon u}$, with higher values of $\rho_{\varepsilon u}$ associated with reduced degree of demeaning.

2. Finite-Sample Properties

2.1 Univariate case

This section analyzes the finite-sample performance of the IVX-Wald statistic in (19) by means of an extensive Monte Carlo study and compares it to the performance of the Q-statistic of CY and the $\pi_{0.05}^*$ statistic of Jansson and Moreira (2006; henceforth JM). We run two-sided tests with nominal size 5% for all three statistics under the null hypothesis that the slope coefficient in the predictive regression is zero, that is, $H_0 : A = 0$.

We use the following data-generating process (DGP) for the univariate case, where y_t and x_t are scalars. For $t \in \{1, \dots, n\}$, the innovation sequences $\varepsilon_t \sim NID(0, 1)$ and $e_t \sim NID(0, 1)$ generate the following model:

$$y_t = \mu + Ax_{t-1} + \varepsilon_t, \quad (22)$$

$$x_t = R_n x_{t-1} + u_t, \quad R_n = 1 + C/n, \quad (23)$$

$$u_t = \phi u_{t-1} + e_t. \quad (24)$$

We denote by $\delta = E(\varepsilon_t u_t)$ the contemporaneous correlation coefficient between ε_t and u_t . Simulation results using 10,000 repetitions are presented for values of $C \in \{0, -5, -10, -15, -20, -50\}$, $\delta \in \{-0.95, -0.5, 0, 0.5, 0.95\}$, sample size $n \in \{100, 250, 500, 1,000\}$, and $\phi \in \{0, 0.5\}$. In the Online Appendix, we also present simulation results for $\phi = 0.25$ and $\phi = -0.1$, and additional results for $\delta \in \{-0.75, -0.25, 0.25, 0.75\}$ are available upon request. The system is initialized at $x_0 = 0$. The IVX estimator and the corresponding Wald statistic are invariant to the value of μ , so we opt for $\mu = 0$. We consider the following sequence of local alternatives for power comparisons:

$$A = \frac{b}{n} \sqrt{1 - \delta^2} \text{ for } b \in \{0, 2, 4, \dots, 32, 40, 60, 100\} \quad (25)$$

with $b = 0 \Rightarrow A = 0$ corresponding to the size of each test.

The results regarding the empirical size in the case of no autocorrelation in the predictor's innovation sequence u_t (that is, $\phi = 0$) are presented in Table 1. We observe that for sample sizes $n \geq 250$ the Wald statistic has excellent size control across all values of C and δ . For $n = 100$, it only appears to be slightly oversized when $|\delta| = 0.95$ and $C \in \{0, -5\}$. For the

other combinations of C and δ , the Wald statistic has the correct size. On the other hand, the Q-statistic appears to be undersized for moderate to high values of δ , such as $|\delta| = 0.5$; increasing the sample size does not seem to remedy this problem. Moreover, for autoregressive roots away from unity and very high values of $|\delta|$, the Q-statistic becomes severely oversized; see, e.g., the combinations $R_n = 0.5$ and $|\delta| \in \{0.95, 0.5\}$, as well as $R_n = 0.8$ and $|\delta| = 0.95$. This is a manifestation of the fact that the CY procedure is not valid for predictors with low degree of persistence. Finally, the JM statistic also exhibits severe size distortions. The most striking finding is that it becomes extremely oversized across all degrees of regressor persistence when $|\delta| = 0$. In addition, considering high values of $|\delta|$, such as $|\delta| = 0.95$, and autoregressive roots away from unity, the JM statistic becomes severely oversized too. Its size distortions appear to be minimized when $|\delta| = 0.5$.

–Table 1 here–

We subsequently examine the finite-sample size of these three statistics in the presence of autocorrelation in u_t . Table 2 refers to the case in which u_t is an AR(1) process with root $\phi = 0.5$. We find that the Wald statistic exhibits size very close to the nominal 5% apart from some slight oversizing for $n = 100$, $C \in \{0, -5\}$, and $|\delta| = 0.95$. On the other hand, the Q-statistic has size substantially lower than the nominal 5% for $|\delta| = 0.5$. Moreover, for $C = -50$, $|\delta| = 0.95$, and $n = 100, 250, 500$, the Q-statistic is severely oversized. Regarding the JM statistic, its size distortions are exacerbated in the presence of autocorrelation. The statistic is severely oversized for both high and low values of $|\delta|$ across all degrees of regressor persistence. As in the case of no autocorrelation, the size distortions of this statistic appear to be minimized when $|\delta| = 0.5$.

–Table 2 here–

Next, we examine the power properties of the three statistics. Our simulation study computes power with respect to the local alternatives given in (25) without size adjustment, as there is no oversizing in the proposed Wald statistic. Here, we present results for sample size $n = 250$ and correlations $\delta \in \{-0.95, -0.5, 0\}$, while the corresponding results for $n = 1,000$ are presented in the Online Appendix.⁶ The power plots presented here correspond to the case

⁶In addition, simulation results for $n \in \{100, 500\}$ and $\delta \in \{-0.75, -0.25\}$ are available upon request. The relative performance of the IVX and CY procedures is very similar to the results reported in this section; the Wald statistic dominates the Q-statistic in terms of power, with the exception of the combinations $\delta \in \{-0.95, 0\}$ and $C = 0$, where there is no dominating relationship.

of no autocorrelation in u_t (that is, $\phi = 0$), whereas in the Online Appendix we present the corresponding power plots for $\phi = 0.5$.

Figure 1 presents the power of the three statistics for $n = 250$, $\delta = -0.95$, and all values of C considered. We observe that in the unit root case ($C = 0$), the Wald statistic has higher power than the Q-statistic for alternatives close to the null hypothesis, but this relationship is reversed for alternatives farther away from the null. For all of the other persistence scenarios (that is, for all values of $C < 0$), the Wald statistic dominates the Q-statistic for any choice of local alternative and δ . The distance between the power curves of the two statistics increases in favor of the Wald test as the persistence of the regressor is reduced toward stationarity (that is, as $|C|$ increases). The last panel of Figure 1 for $C = -50$ shows that despite being considerably oversized in this case, the Q-statistic appears to have lower power in comparison to the (correctly sized) Wald test. Moreover, the JM statistic is characterized by a remarkable lack of power, with the exception of the unit root case. For lower degrees of regressor persistence, the power of the JM statistic is approximately equal to its size even for alternatives far away from the null, undermining further its suitability for predictability tests.

–Figure 1 here–

Figure 2 presents power comparisons for $\delta = -0.5$ and $n = 250$. The power of the Wald test uniformly dominates that of the Q-statistic for all persistence scenarios, including that of a unit root regressor ($C = 0$). As before, the dominance of the IVX over the CY procedure increases as the degree of persistence is reduced towards stationarity. In addition, the power of the JM statistic is much lower relative to the other two statistics, especially as we move away from the unit root case.

–Figure 2 here–

Figure 3 presents power comparisons for $\delta = 0$ and $n = 250$. The Q-statistic appears to have higher power relative to the Wald statistic in the unit root case ($C = 0$). However, as the degree of persistence is reduced ($C < 0$), the power of the Wald statistic becomes indistinguishable from the power of the Q-statistic. The lack of power for the JM statistic relative to the Wald and the Q-statistic is evident in this case too. Interestingly, this conclusion holds true even in the cases in which the JM statistic is severely oversized.

2.2 Conditionally heteroscedastic DGP

Recalling that the asymptotic results for the proposed Wald statistic are also valid under conditional heteroscedasticity, we employ a GARCH(1,1) DGP to examine the finite-sample properties of the statistic and compare them with the corresponding properties of the Q-statistic of CY (see the Online Appendix for the DGP specification).

Extensive simulation results are reported in the Online Appendix. We find that the Wald statistic exhibits no size distortion for every parameter combination considered. The Q-statistic exhibits correct size for $\delta = 0$, but it is oversized for the combination $n = 100$, $|\delta| = 0.95$, and $C = -50$, while it is undersized when $|\delta| = 0.5$. With respect to the power of the tests, we find that for $\delta = -0.95$ the Wald statistic dominates the Q-statistic for every degree of regressor persistence considered. The same conclusion is derived for $\delta = -0.5$. For $\delta = 0$, we find that in the unit root case ($C = 0$), the Q-statistic has higher power than the Wald statistic, whereas for all other degrees of regressor persistence ($C < 0$), the two statistics appear to have the same power.

2.3 Additional Monte Carlo results

We additionally examine the robustness of the power properties of the IVX-Wald statistic with respect to the choice of kernel for the estimation of the long-run covariance matrix. In particular, apart from the Bartlett kernel that we use in the benchmark results, we alternatively use (1) the Parzen kernel and (2) the quadratic spectral kernel. Results are reported in the Online Appendix. In most of the cases, we find that the power plots are almost indistinguishable across the three kernels used.

Finally, we examine the robustness of the power properties of the IVX-Wald statistic when alternative lag lengths are used for the Newey-West estimator of the long-run covariance matrix. In particular, apart from the truncation lag $n^{1/3}$ that we use in the benchmark results, we alternatively consider the following truncation lags: (1) $n^{1/4}$ and (2) $n^{1/2}$, where n is the sample size. Overall, the results presented in the Online Appendix show that the choice of truncation lag yields no considerable difference in terms of power.

2.4 Multivariate case

In this section, we examine the finite-sample performance of the Wald statistic in the context of multivariate regressions. We generalize the DGP of Section 2.1 to include more than one predictors. In particular, we use the following DGP:

$$y_t = \mu + Ax_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim NID(0, 1), \quad (26)$$

$$x_t = R_n x_{t-1} + u_t, \quad R_n = I_r + C/n, \quad (27)$$

$$u_t = \Phi u_{t-1} + e_t, \quad \Phi = \text{diag}(\phi_1, \phi_2, \phi_3), \quad e_t \sim N(0, \Theta), \quad (28)$$

$$\Sigma = E(\zeta_t \zeta_t'), \quad \zeta_t = (\varepsilon_t, u_t)'. \quad (29)$$

where x_t is a 3×1 vector that contains three regressors. Each regressor is characterized by a different degree of persistence. In particular, we set $C = \text{diag}(0, -10, -100)$, corresponding to a unit root, a local-to-unity, and a stationary regressor.⁷

To render the examined setup empirically relevant, we use values for Φ and Σ that are estimated from a predictive system with the CRSP S&P 500 log excess returns being the regressand and the earnings-price ratio (unit root), T-bill rate (local-to-unity), and inflation rate (stationary) being the regressors. In particular, correlation set 1 corresponds to the correlation structure of the residuals (δ 's) and autocorrelation coefficients (ϕ 's) that are estimated from monthly data during the full sample period, whereas the corresponding parameters of correlation set 2 are estimated from quarterly data. In addition to these parameters, we also examine the size properties of the Wald test when alternatively $\Phi = 0_{3 \times 3}$ (that is, no autocorrelation), $\Phi = 0.25I_3$, and $\Phi = 0.5I_3$. Finally, we consider sample sizes $n \in \{250, 500, 1,000\}$.

We examine the size properties of four different tests using a 5% significance level. The first one is the joint Wald test (W_{joint}) under the null hypothesis that all three slope coefficients are jointly equal to zero, that is, $H_0 : A = (0, 0, 0)$ in (26). The other three tests refer to the individual significance of each regressor. In particular, W_{UR} corresponds to the Wald test under the null hypothesis that the slope coefficient of the unit root regressor is equal to zero, that is, $H_0 : A_1 = 0$, letting the other two slope coefficients free. Similarly, W_{LTU} corresponds to the Wald test under $H_0 : A_2 = 0$ and $W_{Stationary}$ corresponds to the Wald test under $H_0 : A_3 = 0$.

⁷We would like to thank an anonymous referee for suggesting this setup.

Table 3 presents the finite-sample size of these four Wald tests. For correlation set 1 in the upper panel, we find that the size of the joint Wald test (W_{joint}) is very close to the nominal 5% across all autocorrelation structures examined. With respect to the test of individual significance for the unit root regressor (W_{UR}), we find a slight oversizing, which peaks around 8%. However, this oversizing becomes almost negligible for the test of individual significance for the local-to-unity regressor (W_{LTU}), whereas the corresponding test for the stationary regressor ($W_{Stationary}$) exhibits no size distortion. Examining the size properties using correlation set 2 in the lower panel of Table 3, we find no size distortion across these four tests, regardless of the autocorrelation structure used.

–Table 3 here–

We also examine the power properties of the joint Wald test under the null hypothesis $H_0 : A=(0,0,0)$, as the slope coefficient of each of the three regressors increases. In particular, $Wald_{0.05}^{UR}$ refers to the power of the joint test when, under the alternative, the slope coefficient of the unit root regressor takes nonzero values ($A = \frac{b}{n} (1, 0, 0)$), $Wald_{0.05}^{LTU}$ refers to the power of the joint test when the slope coefficient of the local-to-unity regressor increases ($A = \frac{b}{n} (0, 1, 0)$), whereas $Wald_{0.05}^{Stationary}$ refers to the power of the joint test when the slope coefficient of the stationary regressor increases ($A = \frac{b}{n} (0, 0, 1)$). Local alternatives are derived using $b \in \{0, 2, 4, \dots, 32, 40, 60, 100\}$ with $b = 0$ corresponding to the size of the test, while we consider $n \in \{100, 250, 500, 1, 000\}$.⁸

Figure 4 presents the power plots of the joint Wald statistic using correlation set 1, whereas Figure 5 presents the corresponding power plots using correlation set 2. We find that in every case examined, the joint Wald test has very good power properties, because the rejection rate monotonically increases as the true value of the corresponding slope coefficient increases. This holds true for all sample sizes examined. Moreover, the power of the joint Wald test is remarkably high even for low values of local alternatives for the slope coefficient of the unit root and the local-to-unity regressors.

–Figures 4 and 5 here–

⁸Simulation results for the power properties of the individual significance tests in the presence of multiple regressors are available upon request.

3. Data and Regressors' Degree of Persistence

We implement the proposed methodology to test the predictive ability of commonly used financial variables with respect to excess stock market returns. The examined period is 1927–2012. The employed dataset is an updated version of the one used by Welch and Goyal (2008).⁹ For our benchmark predictability tests, we use monthly and quarterly data, whereas in the Online Appendix we report results for annual data too. Following Welch and Goyal (2008), we use S&P 500 value-weighted log excess returns to proxy for excess market returns. Moreover, we use the following twelve variables as potential predictors: T-bill rate (**tbl**), long-term yield (**lty**), term spread (**tms**), default yield spread (**dfy**), dividend-price ratio (**d/p**), dividend yield (**d/y**), earnings-price ratio (**e/p**), dividend payout ratio (**d/e**), book-to-market value ratio (**b/m**), net equity expansion (**ntis**), inflation rate (**inf**), and consumption-wealth ratio (**cay**). We present the definitions of these variables, as well as a list of prior studies that have examined their predictive ability in the Online Appendix. It should be noted that *cay* is only available at quarterly and annual frequency, starting from 1952 for quarterly and 1945 for annual data.

One of the main advantages of the IVX methodology is that, by virtue of its robustness, it does not require any pretesting to determine the degree of predictors' persistence prior to conducting predictability tests. Pretesting procedures naturally increase the type I error of predictability tests and may well lead to conflicting empirical conclusions. To demonstrate this point, we report for each regressor in Table 4 the least-squares point estimate of the autoregressive root \widehat{R}_n from regression (2) using monthly data as well as the results of four unit root tests that are commonly used as pretests: the Augmented Dickey Fuller (ADF) test, the DF-GLS test by Elliott, Rothenberg, and Stock (1996), the Phillips-Perron (PP) test, and the KPSS test by Kwiatkowski, Phillips, Schmidt and Shin (1992); the lag length for ADF and DF-GLS is determined by the Bayesian information criterion. It is remarkable how close to unity the estimated root is for most of the variables: for *d/y*, *d/p*, and *e/p* the estimated root is exactly equal to unity up to three decimal points. The four pretests agree on the existence of a unit root only for *lty*, *d/y*, and *d/p*. For the remaining variables, the tests yield contradictory results. Even for *inf*, which exhibits a relatively low autoregressive root, the KPSS test would reject the null hypothesis of no unit root at the 5% level.

⁹The dataset updated until December 2012 is sourced from Amit Goyal's Web site: <http://www.hec.unil.ch/agoyal/>.

–Table 4 here–

Table 5 contains the corresponding results for quarterly data, confirming that these variables exhibit a very high degree of persistence, even when they are measured at a lower frequency, and that their autoregressive root is very close to unity, with the exception of *inf*. Interestingly, *cay* also exhibits a very high autoregressive root and the ADF and PP tests would not reject the null hypothesis of unit root. The evidence provided in the Online Appendix using annual data is very similar, though the autoregressive coefficients are somewhat lower for some variables. Overall, neither the conclusions of the pretests nor the estimated autoregressive roots alleviate the uncertainty on the exact degree of persistence of the employed regressors, regardless of the frequency used. This observation, along with type I error considerations, motivates further the use of the proposed IVX econometric framework.

–Table 5 here–

4. Predictability Tests

4.1 Univariate regressions

4.1.1 Monthly data. We firstly examine the individual predictive ability of each of the employed regressors using monthly data. Table 6 contains the results for these univariate regressions using the proposed IVX estimator and the corresponding Wald statistic under the null hypothesis of no predictability. For comparison, we also report: (1) the t-ratio under the standard least-squares approach, (2) the 90% Bonferroni confidence interval from the Q-statistic of CY, and (3) the p -value for the JM statistic. Moreover, we report the correlation coefficient (δ) of the residuals from regressions (1) and (2) as a measure of the regressor’s degree of endogeneity.

–Table 6 here–

Panel A reports the results for the period January 1927–December 2012. Using our test statistic, we find that the null of no predictability can be rejected at the 5% level only when the lagged *e/p*, *b/m*, and *ntis* are used as predictors; *d/y* is significant only at the 10% level. To the contrary, there is no evidence of significant predictive ability for *d/e*, *lty*, *d/p*, *tbl*, *dfy*, *tms*, and *inf* in the full sample period. Comparing our findings with the other test statistics,

there are important differences with respect to which predictors are significant and at which level. Standard least-squares inference indicates that d/y is significant at the 5% and that $ntis$ is significant only at the 10% level. More interestingly, the Q-test of CY fails to report the significance of e/p even at the 10% level. Calculating 95% Bonferroni confidence intervals for the Q-test in unreported results, we find that only $ntis$ is significant at the 5% level. These findings are in line with our simulation results for the size properties of the Q-test, where we documented that it tends to underreject for large sample sizes ($n = 1,000$) and for moderate to high degrees of endogeneity, such as the one estimated for e/p ($\delta = -0.76$). Finally, the JM test does not find e/p or b/m to be significant predictors, whereas it does so for tbl and dfy , which are insignificant according to our test.

Panel B of Table 6 reports the corresponding results for the period after 1952. This subperiod is examined for two reasons. First, the term structure variables (tbl , tms , and lty) are thought to be more informative since the Fed abandoned its policy of pegging the interest rate (1951 Treasury Accord). Moreover, cay becomes available at quarterly frequency during this period. Second, prior studies (see, e.g., CY) have found that the evidence of predictability has weakened in more recent sample periods, and hence it can be attributed to early periods when such patterns were not documented. The proposed testing methodology can shed further light on this conjecture.

In fact, the predictability evidence almost entirely disappears in the post-1952 period. The IVX-Wald test indicates that only inf is significant at the 5% level. Moreover, tbl and tms are significant at the 10% level, supporting the argument that the term structure variables may have become more informative after 1952. Similar is the evidence based on the Q-test of CY, with the main difference that their test additionally finds d/y to be marginally significant at the 10% level. More striking are the differences with least-squares inference, according to which both d/y and d/p are significant at the 10% level, whereas tbl is significant at the 5% level, demonstrating its tendency to overreject the null of no predictability. Using the JM test would also lead to conclusions that are considerably different from ours. Most importantly, this test indicates d/y as a significant predictor, whereas it fails to do so for tbl and inf . Overall, our results support the argument that predictability has considerably weakened, if not disappeared, after 1952.

4.1.2 Quarterly data. We subsequently estimate the univariate predictive regressions using quarterly data and we report the corresponding results in Table 7 for the full sample period (panel A) and the post-1952 period (panel B), respectively. The results are very similar to the ones we derived using monthly data. In particular, the IVX-Wald test indicates that in the full sample period, e/p , b/m , and $ntis$ are again found to be significant predictors at the 5% level, whereas we also report significance for d/p at the 10% level. Standard least-squares inference would point to similar conclusions, with even lower p -values due to the tendency of the t -test to overreject. More striking is the comparison with the inference derived from the Q-test. In particular, the latter fails to find significance for either e/p or b/m even at the 10% level, demonstrating again a tendency to underreject for moderate to high values of δ . The inference derived from the JM test is also very different from ours. In particular, the JM test fails to report significance for e/p and d/p , whereas it indicates d/y and dfy as strongly significant.

–Table 7 here–

For the post-1952 period we find that, according to the IVX-Wald test, only tms out of the previously used variables remains significant at the 10% level. The rest of the tests also show that predictability has overall weakened in this subperiod, but they additionally find some other variables to be significant predictors, at least at the 10% level. The most interesting finding is that cay , which becomes available after 1952, is a highly significant predictor across all tests considered, including our Wald test. This striking finding corroborates the results of Lettau and Ludvigson (2001) for the updated sample period that we examine.

Taken together with the corresponding univariate results for annual data reported in the Online Appendix, the Wald test indicates that there is significant evidence of in-sample predictability for e/p , b/m , and $ntis$ in the full sample period, and weaker evidence for the dividend-based ratios. However, this evidence almost entirely disappears during the post-1952 period, with the exception of some rather weak evidence for the term structure variables (tms and tbl). The only variable that is found to be strongly significant in the post-1952 period is cay .

4.2 Multivariate regressions

The previous section considered univariate predictability tests. However, it is common practice to employ multiple regressors and to assess their joint significance; this approach is informa-

tive for market efficiency tests because predictability should be assessed with respect to the entire information set, not each variable in isolation (see also Cochrane 2011 for a discussion of the multivariate challenge in predictability tests). Moreover, multivariate predictive regressions are widely used in VAR systems for intertemporal asset pricing tests (e.g., Campbell and Vuolteenaho 2004), as well as in conditional performance evaluation studies (e.g., Ferson, Sarkissian, and Simin 2008). Additionally, from a theoretical point of view, recently developed present value models (see, e.g., Ang and Bekaert 2007; Golez 2014) suggest that d/p alone cannot capture the variation in expected stock returns due to stochastic discount rates and/or dividend growth, and hence it should be used jointly with other predictors.

Given the importance of multivariate predictive systems, it is unfortunate that the recent methodological contributions that correct for the bias arising in least-squares inference are developed for univariate regressions only. The notable exception is the iterative procedure of Amihud, Hurvich, and Wang (2009), which is based on the augmented regression method of Amihud and Hurvich (2004) and accommodates multiple regressors in a single-horizon predictive setup under the restriction that the predictors are stationary. Their procedure yields a reduced-bias estimator and the corresponding test statistic is shown to have good size properties, which deteriorate as the persistence of the predictors approaches the nonstationarity boundary.

On the other hand, our instrumental variable approach introduces an easy-to-implement Wald statistic, enabling us to conduct valid inference regardless of the dimensionality of the predictive system and for all known types of regressors' persistence, from strictly stationary to unit root processes, while it is also applicable to long-horizon predictive systems. The proposed Wald test allows us to examine the joint, as well as the individual significance of the regressors used in a multivariate system. In particular, to test their joint significance, we compute the Wald statistic (19) under the null hypothesis that all slope coefficients are equal to zero, that is, $H_0 : A = 0_{1 \times r}$, while the individual significance of each predictor is evaluated under the null hypothesis that the corresponding slope coefficient is equal to zero, that is, $H_0 : A_i = 0$.

We utilize this test to re-examine the predictive ability of certain combinations of regressors that were found to be significant in prior studies, and they are motivated from either a theoretical or an empirical point of view. In particular, we use the following combinations: (1) d/p and tbl (Ang and Bekaert 2007), (2) d/p, tbl, dfy, and tms (Ferson and Schadt 1996), (3) d/p

and b/m (Kothari and Shanken 1997), (4) d/p and d/e (Lamont 1998), and (5) e/p, tms, and b/m (Campbell and Vuolteenaho 2004). Additionally, we follow a general-to-specific statistical approach to come up with the best set of predictors. In particular, starting with a base system that includes d/p, e/p, tbl, tms, dfy and ntis, we eliminate in each estimation round the variable exhibiting the lowest (and insignificant) value of individual Wald statistic. This process is repeated until all remaining variables are individually significant at the 10% level or lower.¹⁰

4.2.1 Monthly data. Table 8 reports the results for monthly data. Panel A contains the results for the full sample period. Interestingly, we find that none of the examined combinations leads to joint significance at the 5% level. Only the combination of e/p, b/m and tms is jointly significant at the 10% level, but none of these predictors' coefficients is individually significant. It is also noteworthy that d/p is individually insignificant in all combinations examined, apart from the case where it is used jointly with d/e. This finding casts more doubt on its predictive ability over short-horizon returns. On the other hand, the general-to-specific approach leads to a rather interesting finding: e/p and tbl are both jointly and individually significant at the 5% level.

–Table 8 here–

Panel B reports the corresponding results for the post-1952 period, leading to very similar conclusions. None of the five combinations considered is found to be jointly significant and d/p is individually insignificant in every case. Only tbl and tms are found to be individually significant in some cases, confirming that term structure variables may be indeed more informative in the post-1952 period. The general-to-specific approach yields again the most interesting result: e/p and tbl are jointly and individually significant during this subperiod too. As a robustness test, we have alternatively included b/m instead of d/p in the base system; unreported results show that we still end up with e/p and tbl being the only two individually and jointly significant predictors in both periods.

4.2.2 Quarterly data. We repeat the previous analysis using quarterly data and we report these results in Table 9. Panel A contains the full sample period results. We find that combinations that include b/m lead to joint significance, but the regressors' coefficients are insignificant.

¹⁰We would like to thank the Editor for suggesting this approach.

Moreover, we find that d/p is individually significant in some combinations, but none of these yields joint significance. On the other hand, according to the general-to-specific approach, e/p , tbl and $ntis$ are both individually and jointly significant.

–Table 9 here–

Panel B reports the corresponding results for the post-1952 period. Interestingly, we find that none of these five combinations yields joint significance. Because cay becomes available in this subperiod, we additionally examine the combination of d/p , d/e and cay , which was considered in Lettau and Ludvigson (2001). In fact, we find that this combination and cay 's coefficient are significant at the 1% level. Moreover, we also include cay in the base system for the general-to-specific approach, given its strong significance in univariate tests. This approach yields a highly significant combination of e/p , tbl , cay and dfy for this subperiod.

Taken together, the multivariate results for monthly and quarterly data show that commonly used combinations of these regressors have limited predictive ability, especially in the post-1952 period. However, a general-to-specific approach indicates that the combination of e/p and tbl is highly significant and robust to the choice of data frequency and the examined period. Finally, these results confirm that cay is a highly significant predictor in the post-1952 period and this significance is not subsumed by other commonly used variables.

5. Long-Horizon Predictive Regressions

The previous tests examined short-horizon predictability using 1-period ahead returns. A related debate in the literature refers to the existence of long-horizon predictability. In particular, a number of studies have found that the predictive ability of certain financial variables becomes stronger as the horizon increases (see, *inter alia*, the surveys of Cochrane 1999; Campbell 2000). On the other hand, some recent studies cast doubt on this prevailing view (see Valkanov 2003; Torous, Valkanov, and Yan 2004; Ang and Bekaert 2007; Boudoukh, Richardson, and Whitelaw 2008; Hjalmarsson 2011). In particular, Ang and Bekaert (2007) find no evidence of long-horizon predictability using standard errors based on the reverse regression approach of Hodrick (1992), which removes the moving average structure in the error term induced by summing returns over long horizons, and hence retains the correct size, as compared with Hansen-Hodrick

(1980) and Newey-West (1987) standard errors that lead to severely oversized test statistics.¹¹ Moreover, Valkanov (2003) and Boudoukh, Richardson, and Whitelaw (2008) show that in the presence of highly persistent regressors, predictability may artificially emerge in standard least-squares regressions as the horizon increases. We contribute to this debate by extending the proposed IVX-Wald test to accommodate long-horizon predictive regressions and conducting the corresponding empirical tests.¹² Section 5.1 develops a long-horizon IVX-Wald test; Section 5.2 examines the finite-sample properties of the newly developed Wald test; and Section 5.3 discusses the corresponding empirical results.

5.1 Long-horizon IVX inference

Long-horizon predictability tests are typically based on estimators derived from regressing the K -period cumulative stock return $y_t(K)$ on a lagged predictor x_{t-1} and an intercept as in the following fitted model:

$$y_t(K) = \mu_f + Ax_{t-1} + \eta_{f,t} \quad t \in \{1, \dots, n - K + 1\}, \quad (30)$$

with $y_t(K) = \sum_{i=0}^{K-1} y_{t+i}$, while the DGP characterizing the true relationship between y_t and x_t continues to be given by (1). For brevity, we introduce the notation $v_t(K) = \sum_{i=0}^{K-1} v_{t+i}$ for any sequence $(v_t)_{t \geq 1}$ and let $n_K = n - K + 1$.

It is clear that the accumulation of predicted variables on the left side of (30) generates additional correlations that are not present in short-horizon regressions and affect the stochastic properties of long-horizon estimators. To fix ideas, assume temporarily that the intercepts μ in (1) and μ_f in (30) are equal to zero. Then the OLS estimator of A from (30) is given by $\hat{A}_{OLS}(K) = \sum_{t=1}^{n_K} y_t(K) x'_{t-1} \left(\sum_{t=1}^{n_K} x_{t-1} x'_{t-1} \right)^{-1}$. Using the DGP (1), it is easy to see that the above OLS estimator is inconsistent for $K > 1$:

$$\hat{A}_{OLS}(K) = \left[A \sum_{t=1}^{n_K} x_{t-1}(K) x'_{t-1} + \sum_{t=1}^{n_K} \varepsilon_t(K) x'_{t-1} \right] \left(\sum_{t=1}^{n_K} x_{t-1} x'_{t-1} \right)^{-1},$$

the inconsistency occurring because $\sum_{t=1}^{n_K} x_{t-1}(K) x'_{t-1}$ has the same order of magnitude as

¹¹The recent study of Wei and Wright (2013) extends the reverse regression approach of Hodrick (1992) to a wider range of null hypotheses even when the predictors are local-to-unity processes.

¹²We would like to thank the Editor for suggesting the extension of IVX methodology to the long-horizon case.

$\sum_{t=1}^{n_K} x_{t-1} x'_{t-1}$ for fixed horizon K and dominates $\sum_{t=1}^{n_K} x_{t-1} x'_{t-1}$ asymptotically when $K \rightarrow \infty$. This imbalance can be easily corrected by modifying the OLS estimator:

$$\hat{A}_{mOLS}(K) = \sum_{t=1}^{n_K} y_t(K) x'_{t-1} \left(\sum_{t=1}^{n_K} x_{t-1}(K) x'_{t-1} \right)^{-1}. \quad (31)$$

While this modified OLS estimator is consistent, the limit distribution of $\hat{A}_{mOLS}(K) - A$ (under suitable normalisation) will not be mixed Gaussian in the case of unit root and local-to-unity regressors. Consequently, inference procedures based on $\hat{A}_{mOLS}(K)$ will not be valid across the range of persistence classes P(i)-P(iv) of Section 1, leading to erroneous empirical conclusions in the case of misspecification of regressor persistence. IVX methodology can be adapted to deliver robust inference in long-horizon predictive regression systems. The key idea is the same as in the short-horizon case: given a consistent least-squares estimator, the IVX estimator is constructed as a feasible instrumental variables estimator that replaces the regressor x_t by the IVX instrument \tilde{z}_t in (31):

$$\hat{A}_{IVX}(K) = \sum_{t=1}^{n_K} y_t(K) \tilde{z}'_{t-1} \left(\sum_{t=1}^{n_K} x_{t-1}(K) \tilde{z}'_{t-1} \right)^{-1}.$$

In the general case in which the intercept terms μ in (1) and μ_f in (30) are nonzero, a standard result on partitioned regression yields that least-squares estimation of A from the regression (30) is equivalent to least-squares estimation of A from the regression:

$$y_t(K) - \bar{y}_{n_K}(K) = A(x_{t-1} - \bar{x}_{n_K-1}) + \vartheta_t \quad t \in \{1, \dots, n_K\}, \quad (32)$$

where $\bar{y}_{n_K}(K) = n_K^{-1} \sum_{t=1}^{n_K} y_t(K)$ and $\bar{x}_{n_K-1}(K) = n_K^{-1} \sum_{t=1}^{n_K} x_{t-1}(K)$ denote the sample means of $y_t(K)$, and $x_{t-1}(K)$ and \bar{y}_{n_K} and \bar{x}_{n_K-1} denote the usual sample means of y_t and x_{t-1} based on the first n_K observations, respectively. We define the data matrices $\underline{X}_{n_K-1} = [x'_0 - \bar{x}'_{n_K-1}, \dots, x'_{n_K-1} - \bar{x}'_{n_K-1}]'$, $\tilde{Z}(K) = [\tilde{z}'_0(K), \dots, \tilde{z}'_{n_K-1}(K)]'$,

$$\begin{aligned} \underline{Y}(K) &= [y'_1(K) - \bar{y}'_{n_K}(K), \dots, y'_{n_K}(K) - \bar{y}'_{n_K}(K)]' \\ \underline{X}(K) &= [x'_0(K) - \bar{x}'_{n_K-1}(K), \dots, x'_{n_K-1}(K) - \bar{x}'_{n_K-1}(K)]' \end{aligned}$$

and $\tilde{Z}_{n_K-1} = [\tilde{z}'_0, \dots, \tilde{z}'_{n_K-1}]'$, where, as before, underlying indicates demeaning. The modified OLS estimator from (30) (equivalently from (32)) can be expressed as:

$$\tilde{A}_{mOLS}(K) = \underline{Y}(K)' \underline{X}_{n_K-1} [\underline{X}(K)' \underline{X}_{n_K-1}]^{-1},$$

and the corresponding IVX estimator of A is given by:

$$\tilde{A}_{IVX}(K) = \underline{Y}(K)' \tilde{Z}_{n_K-1} [\underline{X}(K)' \tilde{Z}_{n_K-1}]^{-1}. \quad (33)$$

The asymptotic behavior of the normalized and centered IVX estimator in (33) is summarized by Theorem B in the Appendix; asymptotic mixed Gaussianity is preserved regardless of the degree of persistence of the predictor variable in (2), as long as the rate of growth of the horizon K is slower than that of the sample size n . This requirement is presented formally below.

Assumption H. The horizon K may be a fixed integer or a sequence $(K_n)_{n \in \mathbb{N}}$ that increases to infinity slower than the sample size n : $K_n/n \rightarrow 0$ as $n \rightarrow \infty$.

As in the short-horizon case, the asymptotic mixed normality property of the long-horizon IVX estimator $\tilde{A}_{IVX}(K)$ implies that the associated IVX-Wald test statistic will have a standard chi-squared limit distribution across the whole range of empirically relevant persistence classes P(i)-P(iv). In particular, we propose the following IVX-Wald statistic for testing the set of linear restrictions (18) in long-horizon predictive regression systems:

$$W_{IVX}(K) = \left[H \text{vec} \tilde{A}_{IVX}(K) - h \right]' Q_{H,K}^{-1} \left[H \text{vec} \tilde{A}_{IVX}(K) - h \right] \quad (34)$$

where

$$\begin{aligned} Q_{H,K} &= H \left[\left(\tilde{Z}'_{n-K} \underline{X}(K) \right)^{-1} \otimes I_m \right] \mathbb{M}_K \left[\left(\underline{X}(K)' \tilde{Z}_{n-K} \right)^{-1} \otimes I_m \right] H' \\ \mathbb{M}_K &= \tilde{Z}(K)' \tilde{Z}(K) \otimes \hat{\Sigma}_{\varepsilon\varepsilon} - n_K \bar{z}_{n_K-1}(K) \tilde{z}'_{n_K-1}(K) \otimes \hat{\Omega}_{FM} \end{aligned}$$

$\bar{z}_{n_K-1}(K) = n_K^{-1} \sum_{t=1}^{n_K} \tilde{z}_{t-1}(K)$ and $\hat{\Omega}_{FM}$ is defined in (21).

Theorem 2. Consider the model (1)–(3) under Assumption INNOV with (9) and H. Then, the IVX-Wald statistic in (34) for testing (18) satisfies

$$\tilde{W}_{IVX}(K) \Rightarrow \chi^2(q) \quad \text{as } n \rightarrow \infty$$

under H_0 for the following classes of predictor processes x_t in (2):

- (i) P(i)-P(iv) under Assumption INNOV(i),
- (ii) P(i)-P(iv) under Assumption INNOV(ii) when $K \rightarrow \infty$,
- (iii) P(i)-P(iii) under Assumption INNOV(ii) when the horizon parameter K is fixed.

Theorem 2 shows that, under Assumption H, the robustness property of IVX methodology extends to long-horizon predictive regressions. Note that when $K = 1$, the long-horizon IVX estimator (33) and the associated IVX-Wald statistic (34) reduce to their short-horizon counterparts (17) and (19), respectively. Note also the robustness that the IVX-Wald statistic exhibits to conditional heteroscedasticity for purely stationary regressors when $K \rightarrow \infty$: this is due to the persistence that the horizon K induces in the predictive regression; see the discussion in the penultimate paragraph of Section 1.

5.2 Finite-sample properties

In this section, we examine the finite-sample properties of the long-horizon Wald statistic in (34) that corresponds to the long-horizon predictive regression in (30). For this Monte Carlo study, we use the DGP specified in (22)–(24) for the univariate case. In particular, we consider the following parameter values: $C \in \{0, -5, -10, -15, -20, -50\}$, $\delta \in \{-0.95, -0.5, 0\}$, $n \in \{100, 500, 1,000\}$, and $\phi = 0$. For sample size $n = 100$, we consider predictive horizons $K = 2, 3, 4, 5$; for $n = 500$, we consider $K = 4, 8, 12, 20$; and for $n = 1,000$, we use $K = 4, 12, 36, 60$. Table 10 presents the finite-sample size of the long-horizon Wald statistic. These simulation results show that the size of the proposed test is remarkably close to the nominal 5% level for all cases considered.

–Table 10 here–

We also examine the power properties of the long-horizon Wald statistic, using local alternatives $A = \frac{b}{n}$ with $b \in \{0, 2, 4, \dots, 32, 40, 60, 100\}$. Power plots for sample size $n = 1,000$ and horizons $K = 12, 36, 60$, as well as for sample size $n = 500$ and horizons $K = 4, 12, 20$, are presented in the Online Appendix.¹³ In sum, these plots show that for all horizons considered, the power of the statistic rapidly increases as the true value of A increases. Moreover, in each case, the power of the statistic decreases as the predictive horizon increases, but this decrease is very small for highly persistent regressors.

5.3 Empirical results

Table 11 reports the results from long-horizon univariate predictability tests using monthly data. In the full sample period (panel A), we find no evidence that predictability becomes stronger as the horizon increases, with the exception of tms. To the contrary, the predictive ability of e/p and b/m weakens, being significant only at the 10% level when we examine horizons longer than 12 and 36 months, respectively. Only tms and ntis are significant at the 5% level when we examine a sixty-month horizon. Regarding d/y and d/p, these are not significant at the 5% level regardless of the examined horizon.¹⁴ In the post-1952 period (panel B), predictability almost entirely disappears, especially for horizons beyond twenty-four months. We find that only d/e becomes significant at long horizons, while tms remains marginally significant at the 10% level.

–Table 11 here–

Table 12 reports the corresponding long-horizon tests using quarterly data. In the full sample period (panel A), predictability becomes weaker as the horizon increases. Interestingly, e/p, b/m, and ntis, which were found to be strongly significant in predicting one-quarter-ahead returns (see Table 7), become less significant as the horizon increases and they are eventually insignificant at the twenty-quarter horizon; d/p remains significant at the 10% level for all horizons considered, whereas tms becomes marginally significant at very long horizons. In the post-1952 period (panel B), there is no evidence of predictability with three exceptions: tms remains significant but only at the 10% level; d/e becomes marginally significant beyond eight quarters; and cay

¹³The corresponding power plots for $n = 100$ are available upon request.

¹⁴To the contrary, in unreported results we find that using Newey-West or Hansen-Hodrick standard errors to calculate least-squares t-ratios, d/y and d/p (as well as most of the other variables), would erroneously appear as highly significant for horizons of twelve months or longer. The findings are similar when we consider quarterly data.

is the only variable that remains significant at the 5% level for all horizons examined. Similar is the pattern of the corresponding results using annual data that are reported in the Online Appendix.

–Table 12 here–

In sum, our evidence is in line with the results of the above cited studies that cast doubt on the ability of commonly used variables to predict stock returns at long horizons, especially in the post-1952 period. We actually find that, if anything, predictability is generally weaker, not stronger, as the horizon increases.

Table 13 presents the results for long-horizon predictability tests with multiple regressors. We present only the combinations of regressors that were found to be both individually and jointly significant under the general-to-specific approach described in Section 4.2 and reported in Tables 8 and 9, using one-month and one-quarter-ahead returns, respectively. Panel A reports the results for monthly data. In the full sample period, we find that while e/p remains individually significant, tbl becomes insignificant as the horizon increases. Their joint predictive ability remains significant but only at the 10% level for horizons beyond twelve months. For the post-1952 period results are more striking: e/p and tbl are neither individually nor jointly significant beyond twelve months.

–Table 13 here–

Using quarterly data in panel B, we get a similar pattern. For the full sample period, only e/p remains individually significant for all the examined horizons, while neither tbl nor $ntis$ are significant for longer than eight-quarter horizons; the joint significance of these three variables becomes weaker as we increase the predictive horizon and eventually disappears at the twenty-quarter horizon. For the post-1952 period, we find that at horizons longer than four quarters, only cay is individually significant at the 5% level, driving the joint significance of the corresponding multivariate system. Overall, our evidence is in broad agreement with the results of Ang and Bekaert (2007), who found that tbl can predict future stock returns within a multivariate setup only at short (less than one-year) horizons.

6. Conclusion

This study revisits the popular issue of stock return predictability via lagged financial variables. We conduct a battery of predictability tests for U.S. stock returns during the 1927–2012 period, proposing a novel methodology, termed as IVX estimation, which is robust to the time-series properties of the employed regressors. The uncertainty regarding the order of integration of these predictive variables has been characterized as a main source of concern for invalid inference, especially in the presence of endogenous regressors (see Stambaugh 1999; CY); the robust methodology we propose successfully addresses this concern. In univariate tests, we find that the earnings-price and book-to-market value ratios as well as net equity expansion are significant predictors of one-period-ahead excess market returns. However, this evidence almost entirely disappears in the post-1952 period. Only the consumption-wealth ratio is found to be strongly significant in this subperiod.

Apart from robustifying inference in predictability tests, this novel methodology presents two additional, particularly attractive features. First, it leads to standard chi-squared inference, and hence the construction of Bonferroni-type confidence intervals is avoided. Such a simplification is mostly welcome for practical purposes, given the large number of predictive regressors that have been employed in prior literature. Second, the IVX estimation methodology is applicable to multivariate systems of both regressors and regressands. This facility allows us to test a wide range of predictability relationships. Most obviously, we can test for the joint ability of a set of regressors to predict stock market returns. While this issue was the main motivation of the early studies in the literature (e.g., Fama and French 1989), most of the recently suggested econometric methodologies have been restricted to setups with a scalar regressor (see Torous, Valkanov, and Yan 2004; CY; JM; Hjalmarsson 2011). Our multivariate tests document that the combination of the earnings-price ratio and T-bill rate is highly significant and robust to the choice of data frequency and examined period.

Interestingly, the proposed testing procedure can be extended to long-horizon predictive regressions. We develop the relevant test statistic, and we show that it exhibits very good finite-sample properties. Using this newly developed statistic, our long-horizon tests document that, if anything, predictability becomes weaker, not stronger, as the horizon increases. Only the consumption-wealth ratio remains strongly significant for all horizons examined. This evidence

is in agreement with the results of recent studies casting doubt on the prevailing view that predictability becomes stronger as the horizon increases (see, *inter alia*, Ang and Bekaert 2007; Boudoukh, Richardson, and Whitelaw 2008).

Concluding, the proposed IVX estimation methodology improves testing in predictive regressions both by extending the range of testable hypotheses and by robustifying inference with respect to misspecification of regressors' persistence. This novel econometric methodology can prove useful for predictability tests in other asset classes too. Successful implementation can shed new light on whether bond yields and exchange rate fluctuations are predictable via publicly available information. Because predictability tests in these asset classes also rely on persistent regressors with uncertain order of integration, this robust methodology can minimize the risk of distorted inference due to incorrect time-series modeling.

Appendix. Asymptotic Mixed-Gaussianity of the IVX Estimator

This Appendix provides a summary and discussion of the asymptotic behavior of the normalized and centered IVX estimator \tilde{A}_{IVX} in (17) and $\tilde{A}_{IVX}(K)$ in (33) arising from short-horizon and long-horizon predictive regressions, respectively. The key property of \tilde{A}_{IVX} and $\tilde{A}_{IVX}(K)$ that ensures robustness of the IVX procedure and a chi-squared limit distribution for the IVX-Wald test statistic is asymptotic mixed normality. Theorem A shows that asymptotic mixed normality applies to all predictors belonging to the classes P(i)-P(iv) of autoregressive processes regardless of their persistence properties. Theorem B shows that the asymptotic mixed normality property of the IVX estimator extends to long-horizon predictive regression systems. We employ the shorthand notation $a \wedge b = \min(a, b)$ and $a \vee b = \max(a, b)$.

Theorem A. Consider the model (1)–(3) under Assumption INNOV with instruments \tilde{z}_t defined by (4) and (5). Let B_u be a r -variate Brownian motion with covariance matrix Ω_{uu} , $J_C(t) = \int_0^t e^{C(t-s)} dB_u(s)$ be an Ornstein-Uhlenbeck process and let

$$\underline{B}_u(t) = B_u(t) - \int_0^1 B_u(t) dt, \quad \underline{J}_C(t) = J_C(t) - \int_0^1 J_C(t) dt$$

denote the demeaned versions of B_u and J_C . The following limit theory as $n \rightarrow \infty$ applies for the estimator \tilde{A}_{IVX} in (17):

- (i) when $\beta < \alpha \wedge 1$, $n^{\frac{1+\beta}{2}} \text{vec}(\tilde{A}_{IVX} - A) \Rightarrow MN\left(0, \left(\tilde{\Psi}_{uu}^{-1}\right)' C_z V_{C_z} C_z \tilde{\Psi}_{uu}^{-1} \otimes \Sigma_{\varepsilon\varepsilon}\right)$
- (ii) when $\alpha \in (0, \beta)$, $n^{\frac{1+\alpha}{2}} \text{vec}(\tilde{A}_{IVX} - A) \Rightarrow N\left(0, V_C^{-1} \otimes \Sigma_{\varepsilon\varepsilon}\right)$
- (iii) when $\alpha = \beta > 0$, $n^{\frac{1+\alpha}{2}} \text{vec}(\tilde{A}_{IVX} - A) \Rightarrow N\left(0, \mathbb{V}^{-1} C^{-1} V_C C^{-1} (\mathbb{V}')^{-1} \otimes \Sigma_{\varepsilon\varepsilon}\right)$
- (iv_a) when $\alpha = 0$, $\sqrt{n} \text{vec}(\tilde{A}_{IVX} - A) \Rightarrow N\left(0, (E x_{0,1} x'_{0,1})^{-1} \otimes \Sigma_{\varepsilon\varepsilon}\right)$ under INNOV(i)
- (iv_b) when $\alpha = 0$, $\sqrt{n} \text{vec}(\tilde{A}_{IVX} - A) \Rightarrow N(0, V_0)$ under INNOV(ii)

where $x_{0,t} = \sum_{j=0}^{\infty} R^j u_{t-j}$ with $R = I_r + C$ is a stationary version of x_t when $\alpha = 0$, the matrices V_C , V_{C_z} , V and V_0 are given by

$$\begin{aligned} V_C &= \int_0^{\infty} e^{rC} \Omega_{uu} e^{rC} dr, \quad V_{C_z} = \int_0^{\infty} e^{rC_z} \Omega_{uu} e^{rC_z} dr, \quad \mathbb{V} = \int_0^{\infty} e^{rC} V_C e^{rC_z} dr, \\ V_0 &= \left([E x_{0,1} x'_{0,1}]^{-1} \otimes I_m \right) E (x_{0,1} x'_{0,1} \otimes \varepsilon_2 \varepsilon'_2) \left([E x_{0,1} x'_{0,1}]^{-1} \otimes I_m \right), \end{aligned} \quad (35)$$

and the random matrix $\tilde{\Psi}_{uu}$ is given by

$$\tilde{\Psi}_{uu} = \begin{cases} \Omega_{uu} + \int_0^1 \underline{B}_u dB'_u & \text{under P(i)} \\ \Omega_{uu} + \int_0^1 \underline{J}_C dJ'_C & \text{under P(ii)} \\ \Omega_{uu} + V_C C & \text{under P(iii)}. \end{cases}$$

The proof of Theorem A can be found in the Online Appendix.

Remarks A.

(1) Theorem A establishes asymptotic mixed normality of the IVX estimator in predictive regression systems the validity of which is invariant to the persistence properties of the generating mechanism of the predictor process x_t . The fact that asymptotic mixed normality extends over the entire range P(i)-P(iv) of autoregression-induced persistence is the key property that ensures robustness of the IVX procedure. The varying rates of convergence and expressions of the (possibly random) limit variance of the IVX estimator along different persistence classes do not affect self-normalized test statistics such as that of the Wald test considered in (19): mixed normality will deliver standard chi-squared asymptotic inference for IVX based self-normalized quadratic forms.

(2) Theorem A shows that the presence of an intercept in the model does not affect the main asymptotic property of IVX estimation, mixed Gaussianity. This, however, is a first-order asymptotic result. In finite samples, the effect of estimating the intercept may become manifest for predictor processes x_t exhibiting high degree of persistence and strong correlation with the innovations ε_t of the predictive Equation (1). In this case, represented by part (i) of Theorem A, the sample moment that drives mixed normality can be written as

$$n^{-\frac{1+\beta}{2}} \underline{\mathcal{E}}' \tilde{Z} = n^{-\frac{1+\beta}{2}} \sum_{t=1}^n \varepsilon_t \tilde{z}'_{t-1} - n^{\frac{1-\beta}{2}} \bar{\varepsilon}_n \bar{z}'_{n-1}.$$

The first term on the right hand side has a $N(0, V_{C_z}^{-1} \otimes \Sigma_{\varepsilon\varepsilon})$ limit distribution which produces the mixed normal limit result for the IVX estimator in part (i) of Theorem A. Using part (i) of Lemma A1 in the Online Appendix, the second term can be analyzed as

follows:

$$n^{\frac{1-\beta}{2}} \bar{\varepsilon}_n \bar{z}'_{n-1} = \frac{-C_z^{-1}}{n^{\frac{1-\beta}{2}} n^{\frac{1-\alpha}{2}}} \left(\frac{1}{\sqrt{n}} \sum_{t=1}^n \varepsilon_t \right) \frac{x'_n}{n^{\alpha/2}} = O_p \left(n^{-\frac{1-\beta}{2}} n^{-\frac{1-\alpha}{2}} \right) \quad (36)$$

We conclude that the term in (36) is asymptotically negligible across the whole range P(i)-P(iv) of predictor processes but its finite-sample contribution depends simultaneously on three factors: the degree of regressor persistence α ; the correlation between innovations ε_t and u_t ; and the choice of β in the instrumentation procedure. The finite-sample impact of the remainder term in (36) is more prominent for highly persistent regressors: persistence of the unit root and local-to-unity type P(i) and P(ii) results to a finite-sample contribution of exact order $O_p \left(n^{-\frac{1-\beta}{2}} \right)$ in (36); the magnitude of this finite-sample contribution declines continuously as the persistence parameter α drives the predictor process towards stationarity and assumes the minimal rate $O_p \left(n^{-1+\beta/2} \right)$ for stationary predictors belonging to the class P(iv). Strong (positive or negative) correlation also exacerbates the finite-sample effect of estimating the intercept in (1): by a simple application of the central limit theorem to (36), it is clear that a unit root predictor $x_n = \sum_{t=1}^n u_t$ induces finite-sample bias of the form $-C_z^{-1} n^{-\frac{1-\beta}{2}} \Omega_{\varepsilon u}$, the magnitude of which depends on the long-run covariance $\Omega_{\varepsilon u}$ between the innovations of (1) and (2), defined in (12). All finite-sample effects (irrespective of their source) are simultaneously removed by the finite-sample correction (20) on the self-normalizing component of the IVX-Wald statistic. This correction employs a weighted demeaning of the IVX instruments by a matrix that depends on $\hat{\Omega}_{\varepsilon u}$ in a way that balances the finite-sample contribution of (36) for all persistence and correlation combinations conforming to P(i)-P(iv) and Assumption INNOV and all admissible choices of the IVX tuning parameter β .

Theorem B. Consider the model (1)–(3) under Assumption INNOV with (9) and Assumption H. The limit distribution as $n \rightarrow \infty$ of the normalized and centered long-horizon IVX estimator in (33) is mixed Gaussian of the following form:

$$(i) \text{ when } K/n^{\alpha \wedge \beta} \rightarrow 0, n^{\frac{1+(\alpha \wedge \beta)}{2}} \text{vec} \left[\tilde{A}_{IVX}(K) - A \right] \Rightarrow MN(0, Q_1 \otimes \Sigma_{\varepsilon \varepsilon}),$$

$$Q_1 = \left(\tilde{\Psi}_{uu}^{-1} \right)' C_z V_{C_z} C_z \tilde{\Psi}_{uu}^{-1} \text{ if } \beta < \alpha; Q_1 = V_C^{-1} \text{ if } \alpha < \beta$$

- (ii) when $K/n^\alpha \rightarrow 0$, $K/n^\beta \rightarrow \infty$, $\sqrt{nK} \text{vec} [\tilde{A}_{IVX}(K) - A] \Rightarrow MN(0, Q_2 \otimes \Sigma_{\varepsilon\varepsilon})$,
 $Q_2 = (\tilde{\Psi}_{uu}^{-1})' \Omega_{uu}^{-1} \tilde{\Psi}_{uu}^{-1}$
- (iii) when $K/n^\alpha \rightarrow \infty$, $K/n^\beta \rightarrow 0$, $\sqrt{n/K} n^\alpha \text{vec} [\tilde{A}_{IVX}(K) - A] \Rightarrow N(0, Q_3 \otimes \Sigma_{\varepsilon\varepsilon})$,
 $Q_3 = CV_C^{-1} C^{-1} \Omega_{uu} C^{-1} V_C^{-1} C$ if $\alpha > 0$; $Q_3 = (G_{x_0, \infty}^{-1})' C^{-1} \Omega_{uu} C^{-1} G_{x_0, \infty}^{-1}$ if $\alpha = 0$
- (iv) when $K/n^{\alpha \vee \beta} \rightarrow \infty$, $n^{1/2 + \alpha - (\alpha \vee \beta)/2} \text{vec} [\tilde{A}_{IVX}(K) - A] \Rightarrow N(0, Q_4 \otimes \Sigma_{\varepsilon\varepsilon})$,
 $Q_4 = 2V_C^{-1}$ if $\beta < \alpha$; $Q_4 = 2CV_C^{-1} C^{-1} V_{C_z} C^{-1} V_C^{-1} C$ if $0 < \alpha < \beta$; if $\alpha = 0$ $Q_4 = 2(G_{x_0, \infty}^{-1})' C^{-1} V_{C_z} C^{-1} G_{x_0, \infty}^{-1}$.

When $\alpha = 0$ and K is fixed,

- (va) $\sqrt{n} \text{vec} [\tilde{A}_{IVX}(K) - A] \Rightarrow N(0, Q_5 \otimes \Sigma_{\varepsilon\varepsilon})$ under INNOV(i),
 $Q_5 = (G_{x_0, K}^{-1})' \sum_{i, j=0}^{K-1} \Gamma_{x_0}(i-j) G_{x_0, K}^{-1}$
- (vb) $\sqrt{n} \text{vec} [\tilde{A}_{IVX}(K) - A] \Rightarrow N\left(0, \left(G_{x_0, K}^{-1'} \otimes I_m\right) W_{0, K} \left(G_{x_0, K}^{-1} \otimes I_m\right)\right)$ under INNOV(ii),
 $W_{0, K} = \sum_{i, j=0}^{K-1} E\left(x_{0, i} x'_{0, j} \otimes \varepsilon_K \varepsilon'_K\right)$

where V_C , V_{C_z} and $\tilde{\Psi}_{uu}$ are defined in Theorem A, $\Gamma_{x_0}(j) = E\left(x_{0, t} x'_{0, t-j}\right)$ is the autocovariance function of the process $x_{0, t}$ defined in Theorem A, and $G_{x_0, K} = \sum_{j=0}^{K-1} \Gamma_{x_0}(j)$.

The proof of Theorem B requires the development of a new limit theory for sample moments arising from long-horizon predictive regressions and joint control of the asymptotic growth rates for n^α , n^β , and K . The details of this asymptotic development are lengthy and highly nontrivial and can be found in Kostakis, Magdalinos, and Stamatogiannis (2014).

References

- Amihud, Y., and C. M. Hurvich. 2004. Predictive regression: A reduced-bias estimation method. *Journal of Financial and Quantitative Analysis* 39:813–41
- Amihud, Y., C. M. Hurvich, and Y. Wang. 2009. Multiple-predictor regressions: Hypothesis testing. *Review of Financial Studies* 22:413–34
- Ang, A., and G. Bekaert. 2007. Stock return predictability: Is it there? *Review of Financial Studies* 20:651–707
- Binsbergen, V. J., and R. Koijen. 2010. Predictive regressions: A present-value approach. *Journal of Finance* 65:1439–71
- Boudoukh, J., M. P. Richardson, and R. F. Whitelaw. 2008. The myth of long-horizon predictability. *Review of Financial Studies* 21:1577–605
- Boussama, F., 2006. Ergodicité des chaînes de Markov à valeurs dans une variété algébrique: application aux modèles GARCH multivariés. *Comptes Rendus de l' Academie des Sciences Paris* 343:275–78
- Campbell, J. Y. 2000. Asset pricing at the millenium. *Journal of Finance* 55:1515–67
- Campbell, J. Y, and R. J. Shiller. 1988. Stock prices, earnings, and expected dividends. *Journal of Finance* 43:661–76
- Campbell, J.Y., and T. Vuolteenaho. 2004. Bad beta, good beta. *American Economic Review* 94:1249–75
- Campbell J. Y., and M. Yogo. 2006. Efficient tests of stock return predictability. *Journal of Financial Economics* 81:27–60
- Cavanagh, C., G. Elliott and J. H. Stock. 1995. Inference in models with nearly integrated regressors. *Econometric Theory* 11:1131–47
- Cochrane, J. 1999. New facts in finance. *Economic Perspectives* 23:36–58
- . 2008. The dog that did not bark: A defense of return predictability. *Review of Financial Studies* 21:1533–75
- . 2011. Presidential address: Discount rates. *Journal of Finance* 66:1047–108
- Elliott, G. 1998. On the robustness of cointegration methods when regressors almost have unit roots. *Econometrica* 66:149–58
- Elliott, G., T. J., Rothenberg, and J. H. Stock. 1996. Efficient tests for an autoregressive unit root. *Econometrica* 64:813–36

- Fama, E. F. 1970. Efficient capital markets: A review of theory and empirical work. *Journal of Finance* 25:383–417
- . 1991. Efficient capital markets: II. *Journal of Finance* 46:1575–617
- Fama, E. F., and K. R. French. 1988. Dividend yields and expected stock returns. *Journal of Financial Economics* 22:3–24
- . 1989. Business conditions and expected returns on stocks and bonds. *Journal of Financial Economics* 25:23–49
- Ferson, W. E., and R. W. Schadt. 1996. Measuring fund strategy and performance in changing economic conditions. *Journal of Finance* 51:425–61
- Ferson, W. E., S. Sarkissian, and T. Simin. 2008. Asset pricing models with conditional betas and alphas: The effects of data snooping and spurious regression. *Journal of Financial and Quantitative Analysis* 43:331–54
- Francq, C., and J. M. Zakoian. 2010. *GARCH models: Structure, statistical inference and financial applications*. Hoboken, NJ: Wiley
- Giraitis, L., P. Kokoszka, and R. Leipus. 2000. Stationary ARCH models: Dependence structure and the central limit theorem. *Econometric Theory* 16:3–22
- Golez, B. 2014. Expected returns and dividend growth rates implied by derivative markets. *Review of Financial Studies* 27:790–822
- Hansen, L., and R. J. Hodrick. 1980. Forward exchange rates as optimal predictors of future spot rates: An econometric analysis. *Journal of Political Economy* 88:829–53
- Hjalmarsson, E. 2011. New methods for inference in long-horizon regressions. *Journal of Financial and Quantitative Analysis* 46:815–39
- Hodrick, R. J. 1992. Dividend yields and expected stock returns: Alternative procedures for inference and measurement. *Review of Financial Studies* 5:357–86
- Jansson, M., and M. J. Moreira. 2006. Optimal inference in regression models with nearly integrated regressors. *Econometrica* 74:681–714
- Keim, D. B., and R. F. Stambaugh. 1986. Predicting returns in the stock and the bond markets. *Journal of Financial Economics* 17:357–90
- Kelly, B., and S. Pruitt. 2013. Market expectations in the cross section of present values. *Journal of Finance* 68:1721–56
- Kostakis, A., T. Magdalinos, and M. P. Stamatogiannis . 2014. IVX limit theory for long-horizon predictive regressions. Unpublished Manuscript, University of Southampton

- Kothari, S., and J. Shanken. 1997. Book-to-market, dividend yield, and expected market returns: A time-series analysis. *Journal of Financial Economics* 44:169–203
- Kwiatkowski, D., P. C. B. Phillips, P. Schmidt, and Y. Shin. 1992. Testing the null hypothesis of stationary against the alternative of a unit root. *Journal of Econometrics* 54:159–78
- Lamont, O. 1998. Earnings and expected returns. *Journal of Finance* 53:1563–87
- Lanne, M. 2002. Testing the predictability of stock returns. *Review of Economics and Statistics* 84:407–15
- Lettau, M., and S. Ludvigson. 2001. Consumption, aggregate wealth, and expected stock returns. *Journal of Finance* 56:815–49
- . 2005. Expected returns and expected dividend growth. *Journal of Financial Economics* 76:583–626
- Lettau, M., and S. van Nieuwerburgh. 2008. Reconciling the return predictability evidence. *Review of Financial Studies* 21:1607–52
- Magdalinos, T., and P. C. B. Phillips. 2009. Limit theory for cointegrated systems with moderately integrated and moderately explosive regressors. *Econometric Theory* 25:482–526
- Newey, W., and K. West. 1987. A simple, positive definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica* 55:703–8
- Phillips, P. C. B. 1987. Time series regression with a unit root. *Econometrica* 55:277–302
- Phillips, P. C. B., and B. E. Hansen. 1990. Statistical inference in instrumental variables regression with I(1) processes. *Review of Economic Studies* 57:99–125
- Phillips, P. C. B., and T. Magdalinos. 2007. Limit theory for Moderate deviations from a unit root. *Journal of Econometrics* 136:115–30
- . 2009. Econometric inference in the vicinity of unity. CoFie Working Paper No. 7, Singapore Management University
- Phillips, P. C. B., and V. Solo. 1992. Asymptotics for linear processes. *Annals of Statistics* 20:971–1001
- Saikkonen, P. 1991. Asymptotically efficient estimation of cointegration regressions. *Econometric Theory* 7:1–21
- Stambaugh, R. F. 1999. Predictive regressions. *Journal of Financial Economics* 54:375–421
- Stock, J.H., and M. W. Watson. 1993. A simple estimator of cointegrating vectors in higher order integrated systems. *Econometrica* 61:783–820

- Torous, W., R. Valkanov, and S. Yan. 2004. On predicting stock returns with nearly integrated explanatory variables. *Journal of Business* 77:937–66
- Valkanov, R. 2003. Long-horizon regressions: Theoretical results and applications. *Journal of Financial Economics* 68:201–32
- Wei, M., and J. H. Wright. 2013. Reverse regressions and long-horizon forecasting. *Journal of Applied Econometrics* 28:353–71
- Welch, I., and A. Goyal. 2008. A comprehensive look at the empirical performance of equity premium prediction. *Review of Financial Studies* 21:1455–508
- White, H. 1980. A heteroskedasticity-consistent covariance matrix estimator and a direct test for heteroskedasticity. *Econometrica* 48:817–39

Table 1**Finite-sample sizes when there is no autocorrelation ($\phi = 0$) in the residuals of the autoregression**

This table presents finite-sample sizes, testing the null hypothesis $H_0 : A = 0$ versus the alternative $H_1 : A \neq 0$ in (22) when there is no autocorrelation in the residuals of the autoregressive equation, that is, $\phi = 0$ in (24). $W_{0.05}$ corresponds to the rejection rate for the Wald statistic, defined in (19), with 5% nominal size; $Q_{0.05}$ corresponds to the rejection rate resulting from the 95% confidence interval for the Campbell and Yogo (2006) Q -test; and $JM_{0.05}$ corresponds to the rejection rate for the $\pi_{0.05}^*$ statistic of Jansson and Moreira (2006).

Results are reported for different degrees of correlation between the residuals of regressions (22) and (23), $\delta = -0.95, -0.5, 0, 0.5,$ and 0.95 , different sample sizes $n = 100, 250, 500,$ and $1,000$, and for different local-to-unity parameters $C = 0, -5, -10, -15, -20,$ and -50 , which in each sample size case correspond to different autoregressive roots (R_n) reported in the third column. The reported results are based on the Monte Carlo simulation described in Section 2.1, and the average rejection rates are calculated over 10,000 repetitions.

n	C	R_n	$\delta = -0.95$			$\delta = -0.50$			$\delta = 0$			$\delta = 0.50$			$\delta = 0.95$		
			$W_{0.05}$	$Q_{0.05}$	$JM_{0.05}$	$W_{0.05}$	$Q_{0.05}$	$JM_{0.05}$	$W_{0.05}$	$Q_{0.05}$	$JM_{0.05}$	$W_{0.05}$	$Q_{0.05}$	$JM_{0.05}$	$W_{0.05}$	$Q_{0.05}$	$JM_{0.05}$
100	0	1.000	0.067	0.055	0.048	0.064	0.044	0.062	0.051	0.050	0.436	0.063	0.042	0.060	0.063	0.054	0.058
	-5	0.950	0.072	0.061	0.046	0.060	0.039	0.046	0.055	0.050	0.192	0.057	0.037	0.052	0.070	0.062	0.044
	-10	0.900	0.066	0.068	0.030	0.060	0.039	0.032	0.059	0.052	0.170	0.056	0.039	0.040	0.065	0.064	0.028
	-20	0.800	0.063	0.088	0.066	0.056	0.044	0.068	0.051	0.045	0.144	0.057	0.042	0.068	0.062	0.085	0.070
	-50	0.500	0.058	0.257	0.150	0.050	0.095	0.058	0.058	0.054	0.148	0.054	0.094	0.048	0.055	0.257	0.148
250	0	1.000	0.060	0.051	0.062	0.053	0.036	0.054	0.050	0.050	0.510	0.057	0.038	0.042	0.057	0.046	0.052
	-5	0.980	0.062	0.047	0.036	0.056	0.034	0.048	0.050	0.050	0.208	0.052	0.031	0.038	0.062	0.046	0.028
	-10	0.960	0.059	0.050	0.042	0.055	0.032	0.052	0.051	0.048	0.158	0.048	0.030	0.036	0.061	0.053	0.042
	-20	0.920	0.057	0.062	0.040	0.050	0.032	0.036	0.052	0.049	0.128	0.054	0.033	0.038	0.059	0.059	0.034
	-50	0.800	0.054	0.169	0.318	0.050	0.050	0.038	0.055	0.052	0.116	0.053	0.054	0.040	0.055	0.166	0.342
500	0	1.000	0.052	0.039	0.042	0.053	0.038	0.046	0.049	0.048	0.582	0.051	0.036	0.072	0.059	0.043	0.048
	-5	0.990	0.062	0.049	0.036	0.051	0.030	0.038	0.051	0.048	0.258	0.052	0.032	0.040	0.064	0.050	0.040
	-10	0.980	0.057	0.044	0.036	0.055	0.031	0.036	0.049	0.049	0.200	0.054	0.033	0.040	0.060	0.047	0.032
	-20	0.960	0.055	0.049	0.050	0.054	0.029	0.042	0.051	0.051	0.178	0.049	0.028	0.048	0.056	0.049	0.054
	-50	0.900	0.052	0.113	0.524	0.052	0.037	0.054	0.048	0.045	0.176	0.051	0.037	0.054	0.054	0.114	0.488
1,000	0	1.000	0.055	0.042	0.038	0.047	0.034	0.038	0.051	0.049	0.646	0.052	0.035	0.032	0.056	0.042	0.046
	-5	0.995	0.059	0.047	0.040	0.051	0.030	0.046	0.052	0.051	0.334	0.055	0.031	0.034	0.061	0.048	0.042
	-10	0.990	0.059	0.046	0.038	0.052	0.030	0.050	0.051	0.048	0.270	0.054	0.032	0.050	0.055	0.046	0.046
	-20	0.980	0.058	0.047	0.042	0.057	0.031	0.034	0.049	0.047	0.222	0.053	0.029	0.040	0.060	0.048	0.036
	-50	0.950	0.052	0.074	0.606	0.050	0.032	0.028	0.049	0.048	0.194	0.049	0.029	0.030	0.056	0.069	0.600

Table 2**Finite-sample sizes with autocorrelation coefficient $\phi = 0.5$ in the residuals of the autoregression**

This table presents finite-sample sizes, testing the null hypothesis $H_0 : A = 0$ versus the alternative $H_1 : A \neq 0$ in (22) when the autocorrelation coefficient in the residuals of the autoregression (23) is $\phi = 0.5$. $W_{0.05}$ corresponds to the rejection rate for the Wald statistic, defined in (19), with 5% nominal size; $Q_{0.05}$ corresponds to the rejection rate resulting from the 95% confidence interval for the Campbell and Yogo (2006) Q -test; and $JM_{0.05}$ corresponds to the rejection rate for the $\pi_{0.05}^*$ statistic of Jansson and Moreira (2006). Results are reported for different degrees of correlation between the residuals of regressions (22) and (23), $\delta = -0.95, -0.5, 0, 0.5$ and 0.95 , different sample sizes $n = 100, 250, 500$, and $1,000$ and for different local-to-unity parameters $C = 0, -5, -10, -15, -20$, and -50 , which in each sample size case correspond to different autoregressive roots (R_n) reported in the third column. The reported results are based on the Monte Carlo simulation described in Section 2.1 and the average rejection rates are calculated over 10,000 repetitions.

n	C	R_n	$\delta = -0.95$			$\delta = -0.50$			$\delta = 0$			$\delta = 0.50$			$\delta = 0.95$		
			$W_{0.05}$	$Q_{0.05}$	$JM_{0.05}$	$W_{0.05}$	$Q_{0.05}$	$JM_{0.05}$	$W_{0.05}$	$Q_{0.05}$	$JM_{0.05}$	$W_{0.05}$	$Q_{0.05}$	$JM_{0.05}$	$W_{0.05}$	$Q_{0.05}$	$JM_{0.05}$
100	0	1.000	0.072	0.054	0.110	0.066	0.044	0.110	0.050	0.051	0.394	0.061	0.039	0.118	0.073	0.054	0.108
	-5	0.950	0.072	0.053	0.148	0.063	0.040	0.056	0.053	0.049	0.162	0.062	0.037	0.050	0.073	0.053	0.136
	-10	0.900	0.068	0.047	0.156	0.060	0.036	0.046	0.054	0.050	0.124	0.061	0.034	0.038	0.071	0.052	0.138
	-20	0.800	0.063	0.059	0.140	0.055	0.032	0.038	0.056	0.051	0.094	0.056	0.033	0.034	0.061	0.056	0.138
	-50	0.500	0.053	0.150	0.134	0.051	0.053	0.042	0.055	0.052	0.100	0.056	0.055	0.046	0.055	0.155	0.112
250	0	1.000	0.064	0.044	0.122	0.055	0.033	0.088	0.051	0.052	0.420	0.054	0.033	0.070	0.060	0.044	0.114
	-5	0.980	0.065	0.046	0.124	0.059	0.033	0.054	0.051	0.048	0.158	0.057	0.034	0.052	0.067	0.045	0.134
	-10	0.960	0.066	0.046	0.118	0.057	0.035	0.044	0.055	0.050	0.108	0.058	0.032	0.038	0.062	0.043	0.116
	-20	0.920	0.054	0.046	0.112	0.056	0.033	0.036	0.049	0.047	0.078	0.058	0.034	0.030	0.056	0.047	0.122
	-50	0.800	0.054	0.150	0.094	0.051	0.044	0.040	0.051	0.048	0.102	0.054	0.046	0.048	0.057	0.144	0.112
500	0	1.000	0.055	0.043	0.070	0.053	0.036	0.052	0.047	0.049	0.410	0.050	0.034	0.072	0.056	0.043	0.088
	-5	0.990	0.064	0.044	0.104	0.056	0.033	0.052	0.052	0.048	0.202	0.056	0.033	0.052	0.062	0.049	0.108
	-10	0.980	0.061	0.044	0.082	0.053	0.032	0.026	0.047	0.044	0.152	0.053	0.030	0.036	0.061	0.044	0.074
	-20	0.960	0.055	0.043	0.114	0.050	0.029	0.040	0.050	0.046	0.136	0.052	0.033	0.042	0.058	0.045	0.102
	-50	0.900	0.051	0.097	0.112	0.049	0.034	0.060	0.056	0.053	0.136	0.050	0.033	0.058	0.057	0.098	0.120
1,000	0	1.000	0.054	0.039	0.066	0.056	0.034	0.044	0.052	0.053	0.468	0.054	0.033	0.044	0.061	0.044	0.096
	-5	0.995	0.065	0.049	0.088	0.057	0.035	0.060	0.053	0.053	0.216	0.054	0.030	0.060	0.063	0.046	0.112
	-10	0.990	0.060	0.047	0.096	0.055	0.031	0.062	0.047	0.045	0.146	0.052	0.032	0.054	0.060	0.046	0.106
	-20	0.980	0.061	0.045	0.100	0.053	0.030	0.040	0.051	0.047	0.124	0.051	0.028	0.042	0.064	0.050	0.104
	-50	0.950	0.052	0.064	0.124	0.052	0.027	0.036	0.053	0.051	0.116	0.053	0.028	0.034	0.053	0.064	0.110

Table 3**Finite-sample sizes for multivariate predictive systems**

This table presents finite-sample sizes for four Wald tests, with nominal size 5%, based on the multivariate predictive system in (26) with three regressors exhibiting different degrees of persistence (unit root, local-to-unity, and stationary), as described in the Monte Carlo simulation in Section 2.4. W_{joint} reports the rejection rate for the joint Wald test, defined in (19), under the null hypothesis $H_0 : A = \mathbf{0}_{1 \times 3}$, that is, that all three coefficients in vector A are equal to zero. W_{UR} reports the corresponding rejection rate for the individual significance of the unit root regressor coefficient, that is, under the null hypothesis $H_0 : A_1 = 0$. W_{LTU} reports the rejection rate for the individual significance of the local-to-unity regressor coefficient, that is, under the null hypothesis $H_0 : A_2 = 0$, whereas $W_{\text{Stationary}}$ reports the rejection rate for the individual significance of the stationary regressor coefficient, that is, under the null hypothesis $H_0 : A_3 = 0$. Results are reported for (1) two sets of correlations (δ 's) between the residuals of regressions (26) and (27), as estimated using S&P 500 value-weighted log excess return (regressand), earnings-price ratio (UR), T-bill rate (LTU), and inflation rate (Stationary) with monthly (correlation set 1) and quarterly (correlation set 2) data for the period 1927–2012, (2) four sets of autocorrelation coefficients in the residuals of autoregressions in (27): $\phi = 0, 0.25, 0.5$, and the corresponding sample estimates for each of the three regressors mentioned above, and (3) different sample sizes: $n=250, 500$, and $1,000$. The average rejection rates are calculated over 10,000 repetitions.

Correlation set 1	n	W_{joint}	W_{UR}	W_{LTU}	$W_{\text{Stationary}}$
$\phi_1 = \phi_2 = \phi_3 = 0$	250	0.052	0.078	0.065	0.057
	500	0.051	0.076	0.060	0.057
	1,000	0.047	0.077	0.065	0.054
$\phi_1 = \phi_2 = \phi_3 = 0.25$	250	0.070	0.076	0.065	0.055
	500	0.064	0.080	0.062	0.053
	1,000	0.063	0.075	0.067	0.049
$\phi_1 = \phi_2 = \phi_3 = 0.5$	250	0.058	0.082	0.067	0.053
	500	0.053	0.079	0.069	0.053
	1,000	0.049	0.080	0.059	0.052
$\phi_1 = 0.28$ $\phi_2 = 0.32$ $\phi_3 = -0.14$	250	0.070	0.084	0.065	0.055
	500	0.064	0.080	0.069	0.052
	1,000	0.067	0.079	0.062	0.053
Correlation set 2	n	W_{joint}	W_1	W_2	W_3
$\phi_1 = \phi_2 = \phi_3 = 0$	250	0.058	0.054	0.058	0.056
	500	0.048	0.053	0.059	0.050
	1,000	0.051	0.054	0.054	0.054
$\phi_1 = \phi_2 = \phi_3 = 0.25$	250	0.057	0.054	0.053	0.058
	500	0.052	0.055	0.050	0.055
	1,000	0.056	0.050	0.053	0.051
$\phi_1 = \phi_2 = \phi_3 = 0.5$	250	0.054	0.058	0.057	0.054
	500	0.049	0.058	0.059	0.047
	1,000	0.052	0.053	0.054	0.052
$\phi_1 = 0.22$ $\phi_2 = -0.1$ $\phi_3 = -0.08$	250	0.058	0.053	0.060	0.052
	500	0.053	0.053	0.052	0.051
	1,000	0.052	0.052	0.049	0.053

Table 4
Unit root tests for predictive regressors, monthly data

This table presents the results of unit root tests for the following list of financial and economic variables defined in Section 3: dividend payout ratio (d/e), long-term yield (lty), dividend yield (d/y), dividend price ratio (d/p), T-bill rate (tbl), earnings price ratio (e/p), book-to-market value ratio (b/m), default yield spread (dfy), net equity expansion (ntis), term spread (tms), and inflation rate (inf). \hat{R}_n corresponds to the least-squares point estimate of the AR(1): $x_t = R_n x_{t-1} + u_t$. ADF stands for the augmented Dickey-Fuller test statistic; DF-GLS refers to the Elliot et al. (1996) Dickey-Fuller-GLS test statistic; PP stands for the Phillips-Perron test statistic; and KPSS refers to the Kwiatkowski et al. (1992) test statistic. The Bayesian information criterion has been used to select the optimal lag length for ADF and DF-GLS test statistics. The sample period is January 1927–December 2012. *, **, and *** imply rejection of the null hypothesis of a unit root (for ADF, DF-GLS, and PP) or stationarity (for KPSS) at 10%, 5%, and 1% level, respectively.

	\hat{R}_n	ADF	DF-GLS	PP	KPSS
Dividend payout ratio	0.999	-5.758***	-5.712***	-4.184***	1.701***
Long-term yield	0.999	-1.286	-1.181	-1.314	1.853***
Dividend yield	1.000	-2.179	-1.448	-2.087	2.502***
Dividend-price ratio	1.000	-2.180	-1.468	-2.149	2.505***
T-bill rate	0.997	-2.238	-2.237**	-2.131	1.313***
Earnings-price ratio	1.000	-3.870***	-3.014***	-3.656***	1.026***
Book-to-market value ratio	0.997	-3.108**	-2.754***	-2.989**	1.384***
Default yield spread	0.993	-3.430**	-3.364***	-3.779***	0.546**
Net equity expansion	0.981	-4.371***	-1.247	-4.592***	1.008***
Term spread	0.985	-5.112***	-3.727***	-4.697***	0.535**
Inflation rate	0.633	-9.161***	-5.257***	-20.531***	0.617**

Table 5
Unit root tests for predictive regressors, quarterly data

This table presents the results of unit root tests for the following list of financial and economic variables defined in Section 3: Dividend payout ratio (d/e), long-term yield (lty), dividend yield (d/y), dividend price ratio (d/p), T-bill rate (tbl), earnings price ratio (e/p), book-to-market value ratio (b/m), default yield spread (dfy), net equity expansion (ntis), term spread (tms), inflation rate (inf), and consumption-wealth ratio (cay). \hat{R}_n corresponds to the least-squares point estimate of the AR(1): $x_t = R_n x_{t-1} + u_t$. ADF stands for the augmented Dickey-Fuller test statistic; DF-GLS refers to the Elliot et al. (1996) Dickey-Fuller-GLS test statistic; PP stands for the Phillips-Perron test statistic; and KPSS refers to the Kwiatkowski et al. (1992) test statistic. The Bayesian information criterion has been used to select the optimal lag length for ADF and DF-GLS test statistics. The sample period is 1927Q1–2012Q4, with the exception of cay, which becomes available at quarterly frequency after 1952. *, **, and *** imply rejection of the null hypothesis of a unit root (for ADF, DF-GLS, and PP) or stationarity (for KPSS) at 10%, 5%, and 1% level, respectively.

	\hat{R}_n	ADF	DF-GLS	PP	KPSS
Dividend payout ratio	0.985	-4.019***	-3.995***	-3.938***	1.288***
Long-term yield	0.997	-1.428	-1.318	-1.213	1.023***
Dividend yield	1.000	-2.159	-1.560	-2.096	1.439***
Dividend-price ratio	1.000	-2.224	-1.619*	-2.284	1.453***
T-bill rate	0.983	-2.141	-2.145**	-2.333	0.765***
Earnings-price ratio	0.999	-4.274***	-2.462**	-3.424**	0.665**
Book-to-market value ratio	0.989	-3.500***	-3.114***	-3.262**	0.800***
Default yield spread	0.971	-3.241**	-3.186***	-4.055***	0.357*
Net equity expansion	0.939	-4.182***	-1.057	-4.654***	0.752***
Term spread	0.944	-4.536***	-2.923***	-5.333***	0.418*
Inflation rate	0.627	-4.364***	-4.366***	-12.360***	0.425*
Consumption-wealth ratio	0.951	-2.408	-2.201**	-2.431	0.232

Table 6
Univariate predictive regressions, monthly data

This table presents the results of univariate predictive regression models, as in Equation (1), during the sample periods January 1927–December 2012 (panel A) and January 1952–December 2012 (panel B). The dependent variable is the monthly S&P 500 value-weighted log excess return and the lagged persistent regressor is each of the following variables defined in Section 3: dividend payout ratio (d/e), long-term yield (lty), dividend yield (d/y), dividend price ratio (d/p), T-bill rate (tbl), earnings price ratio (e/p), book-to-market value ratio (b/m), default yield spread (dfy), net equity expansion (ntis), term spread (tms), and inflation rate (inf). \tilde{A}_{OLS} stands for the least-squares slope coefficient estimated via regression model (1), whereas t_{OLS} is the corresponding t -statistic under the null hypothesis that A is equal to zero (i.e., no predictability). \tilde{A}_{IVX} , defined in (17), stands for the slope coefficient for the predictive regression (16) estimated via the proposed instrumental variable (IVX) approach, whereas IVX-Wald refers to the Wald statistic, defined in Equation (19), under the null hypothesis that the slope coefficient A is equal to zero. δ denotes the correlation coefficient between the residuals of regression models (1) and (2). *, **, and *** imply rejection of the null hypothesis at 10%, 5%, and 1% level, respectively. CY 90% CI stands for the 90% Bonferroni confidence interval for the bias-corrected scaled least-squares slope coefficient of the predictive regression using the Q -test of Campbell and Yogo (2006). Bold indicates rejection of the null hypothesis of no predictability at the 10% level. JM reports the p -value for the $\pi_{0.05}^*$ statistic of Jansson and Moreira (2006) under the null hypothesis of no predictability.

Regressors	\tilde{A}_{OLS}	t_{OLS}	\tilde{A}_{IVX}	IVX-Wald	δ	CY 90% CI		JM
Panel A: January 1927–December 2012								
Dividend payout ratio	-0.0024	-0.46	-0.0033	0.393	-0.067	-0.006	0.003	0.19
Long-term yield	-0.0622	-1.01	-0.0665	1.064	-0.108	-0.007	0.002	0.38
Dividend yield	0.0075	1.97**	0.0081	3.129*	-0.079	0.001	0.014	0.06*
Dividend-price ratio	0.0062	1.63	0.0065	2.031	-0.975	-0.004	0.008	0.32
T-bill rate	-0.0784	-1.40	-0.0761	1.770	-0.062	-0.011	0.001	0.03**
Earnings-price ratio	0.0087	2.13**	0.0088	4.402**	-0.759	-0.003	0.015	0.34
Book-to-market value ratio	0.0148	2.28**	0.0134	4.101**	-0.823	0.001	0.021	0.12
Default yield spread	0.1100	0.45	0.0591	0.058	-0.274	-0.009	0.015	0.03**
Net equity expansion	-0.1355	-1.93*	-0.1720	4.150**	-0.031	-0.026	-0.003	0.01***
Term spread	0.1482	1.13	0.1399	1.095	-0.005	-0.004	0.024	0.15
Inflation rate	-0.3500	-1.07	-0.3555	1.148	0.023	-0.064	0.021	0.35
Panel B: January 1952–December 2012								
Dividend payout ratio	0.0049	0.93	0.0044	0.672	-0.091	-0.003	0.009	0.31
Long-term yield	-0.0725	-1.23	-0.0777	1.396	-0.148	-0.012	0.002	0.16
Dividend yield	0.0075	1.95*	0.0081	1.425	-0.058	0.001	0.014	0.04**
Dividend-price ratio	0.0069	1.79*	0.0072	1.142	-0.986	-0.006	0.005	0.43
T-bill rate	-0.1057	-2.01**	-0.1054	3.537*	-0.126	-0.018	-0.002	0.27
Earnings-price ratio	0.0038	1.04	0.0029	0.588	-0.610	-0.011	0.006	0.46
Book-to-market value ratio	0.0043	0.68	0.0029	0.174	-0.747	-0.007	0.008	0.27
Default yield spread	0.2275	0.65	0.2306	0.389	-0.056	-0.009	0.019	0.46
Net equity expansion	-0.0259	-0.30	-0.0417	0.220	-0.063	-0.016	0.010	0.28
Term spread	0.2071	1.88*	0.2176	3.808*	0.034	0.002	0.038	0.03**
Inflation rate	-1.0501	-2.31**	-1.1057	5.922**	-0.069	-0.130	-0.031	0.15

Table 7**Univariate predictive regressions, quarterly data**

This table presents the results of univariate predictive regression models, as in Equation (1), during the sample period 1927Q1–2012Q4 (panel A) and 1952Q1–2012Q4 (panel B). The dependent variable is the quarterly S&P 500 value-weighted log excess return and the lagged persistent regressor is each of the following variables defined in Section 3: dividend payout ratio (d/e), long-term yield (lty), dividend yield (d/y), dividend price ratio (d/p), T-bill rate (tbl), earnings price ratio (e/p), book-to-market value ratio (b/m), default yield spread (dfy), net equity expansion (ntis), term spread (tms), inflation rate (inf), and consumption-wealth ratio (cay). \tilde{A}_{OLS} stands for the least-squares slope coefficient estimated via regression model (1), whereas t_{OLS} is the corresponding t -statistic under the null hypothesis that A is equal to zero (i.e., no predictability). \tilde{A}_{IVX} , defined in (17), stands for the slope coefficient for the predictive regression (16) estimated via the proposed instrumental variable (IVX) approach, whereas IVX-Wald refers to the Wald statistic, defined in Equation (19), under the null hypothesis that the slope coefficient A is equal to zero. δ denotes the correlation coefficient between the residuals of regression models (1) and (2). *, **, and *** imply rejection of the null hypothesis at 10%, 5% and 1% level respectively. CY 90% CI stands for the 90% Bonferroni confidence interval for the bias-corrected scaled least-squares slope coefficient of the predictive regression using the Q -test of Campbell and Yogo (2006). Bold fonts indicate rejection of the null hypothesis of no predictability at the 10% level. JM reports the p -value for the $\pi_{0.05}^*$ statistic of Jansson and Moreira (2006) under the null hypothesis of no predictability.

Regressors	\tilde{A}_{OLS}	t_{OLS}	\tilde{A}_{IVX}	IVX-Wald	δ	CY 90% CI	JM	
Panel A: 1927Q1–2012Q4								
Dividend payout ratio	-0.0031	-0.18	-0.0053	0.095	-0.138	-0.037	0.020	0.22
Long-term yield	-0.1621	-0.78	-0.1705	0.629	-0.071	-0.022	0.008	0.34
Dividend yield	0.0216	1.69*	0.0232	2.638	0.045	0.001	0.044	0.03**
Dividend-price ratio	0.0230	1.83*	0.0249	2.952*	-0.943	-0.010	0.033	0.35
T-bill rate	-0.2110	-1.13	-0.2032	1.129	-0.029	-0.039	0.008	0.07*
Earnings-price ratio	0.0284	2.10**	0.0289	4.439**	-0.556	-0.002	0.072	0.31
Book-to-market value ratio	0.0610	2.82***	0.0565	6.553**	-0.832	-0.001	0.062	0.10*
Default yield spread	0.6472	0.80	0.5041	0.390	-0.515	-0.026	0.064	0.01**
Net equity expansion	-0.6054	-2.60***	-0.7683	6.596**	0.137	-0.090	-0.022	0.04**
Term spread	0.4245	0.97	0.4007	0.796	-0.005	-0.016	0.076	0.17
Inflation rate	-0.1980	-0.45	-0.1954	0.198	0.033	-0.084	0.061	0.43
Panel B: 1952Q1–2012Q4								
Dividend payout ratio	0.0189	1.13	0.0177	1.097	-0.190	-0.024	0.057	0.50
Long-term yield	-0.1792	-0.93	-0.1881	0.782	-0.095	-0.035	0.009	0.14
Dividend yield	0.0272	2.17**	0.0307	2.235	-0.095	0.004	0.046	0.03**
Dividend-price ratio	0.0237	1.88*	0.0257	1.525	-0.967	-0.016	0.019	0.44
T-bill rate	-0.2835	-1.65*	-0.2806	2.362	-0.073	-0.067	0.001	0.24
Earnings-price ratio	0.0112	0.95	0.0088	0.518	-0.334	-0.028	0.044	0.49
Book-to-market value ratio	0.0200	0.97	0.0171	0.546	-0.793	-0.020	0.028	0.31
Default yield spread	0.6762	0.60	0.6910	0.329	-0.174	-0.041	0.065	0.49
Net equity expansion	-0.0319	-0.11	-0.0718	0.060	-0.034	-0.043	0.344	0.43
Term spread	0.6047	1.68*	0.6349	3.057*	0.040	0.001	0.119	0.05**
Inflation rate	-0.7879	-1.38	-0.8793	2.356	-0.128	-0.193	-0.026	0.15
Consumption-wealth ratio	0.8480	3.38***	0.8746	11.351***	-0.429	0.032	0.110	0.02**

Table 8**Predictive regressions with multiple regressors, monthly data**

This table presents the results of predictive regression models with multiple regressors, as in Equation (1), during the sample periods January 1927–December 2012 (panel A) and January 1952–December 2012 (panel B). In each case, the dependent variable is the monthly S&P 500 value-weighted log excess return and the lagged regressors are combinations of the following variables defined in Section 3: dividend price ratio (d/p), earnings price ratio (e/p), book-to-market value ratio (b/m), dividend payout ratio (d/e), T-bill rate (tbl), default yield spread (dfy), and term spread (tms). \tilde{A}_{IVX} , defined in (17), is the vector containing the slope coefficients with respect to each of the employed variables for the predictive regression (16), estimated via the instrumental variable (IVX) approach. The significance of each individual coefficient is evaluated using the Wald statistic, defined in Equation (19), under the null hypothesis that the corresponding coefficient is equal to zero. Joint Wald refers to the same Wald statistic, under the null hypothesis that all coefficients A are jointly equal to zero. *, **, and *** imply rejection of the null hypothesis at 10%, 5%, and 1% level respectively.

Panel A: January 1927–December 2012								
d/p	e/p	b/m	d/e	tbl	dfy	tms	Joint Wald	Related study/ Model
0.0061	–	–	–	–0.0807	–	–	3.644	Ang and Bekaert (2007)
0.0077	–	–	–	–0.0647	–0.1871	0.0996	4.742	Ferson and Schadt (1996)
–0.0010	–	0.0150	–	–	–	–	4.117	Kothari and Shanken (1997)
0.0091*	–	–	–0.0082	–	–	–	3.655	Lamont (1998)
–	0.0082	0.0053	–	–	–	0.1992	7.321*	Campbell and Vuolteenaho (2004)
–	0.0112**	–	–	–0.1275**	–	–	8.748**	General-to-specific approach
Panel B: January 1952–December 2012								
d/p	e/p	b/m	d/e	tbl	dfy	tms	Joint Wald	Related study/ Model
0.0150	–	–	–	–0.2314**	–	–	4.132	Ang and Bekaert (2007)
0.0130	–	–	–	–0.2044	0.2252	0.0607	7.653	Ferson and Schadt (1996)
0.0237	–	–0.0290	–	–	–	–	2.085	Kothari and Shanken (1997)
0.0067	–	–	0.0025	–	–	–	1.326	Lamont (1998)
–	0.0060	–0.0014	–	–	–	0.2633**	5.420	Campbell and Vuolteenaho (2004)
–	0.0108**	–	–	–0.2113***	–	–	8.160**	General-to-specific approach

Table 9**Predictive regressions with multiple regressors, quarterly data**

This table presents the results of predictive regression models with multiple regressors, as in Equation (1), during the sample periods 1927Q1–2012Q4 (panel A) and 1952Q1–2012Q4 (panel B). In each case, the dependent variable is the quarterly S&P 500 value-weighted log excess return, and the lagged persistent regressors are combinations of the following variables defined in Section 3: dividend price ratio (d/p), earnings price ratio (e/p), book-to-market value ratio (b/m), dividend payout ratio (d/e), T-bill rate (tbl), default yield spread (dfy), term spread (tms), consumption-wealth ratio (cay), and net equity expansion (ntis). \tilde{A}_{IVX} , defined in (17), is the vector containing the slope coefficients with respect to each of the employed variables for the predictive regression (16), estimated via the instrumental variable (IVX) approach. Joint Wald refers to the Wald statistic, defined in Equation (19), under the null hypothesis that all coefficients A are jointly equal to zero. *, **, and *** imply rejection of the null hypothesis at 10%, 5%, and 1% level respectively.

Panel A: 1927Q1–2012Q4									
d/p	e/p	b/m	d/e	tbl	dfy	tms	ntis	Joint Wald	Related study/ Model
0.0240*	–	–	–	–0.2190	–	–	–	3.971	Ang and Bekaert (2007)
0.0267	–	–	–	–0.1731	–0.2871	0.2476	–	4.557	Ferson and Schadt (1996)
–0.0137	–	0.0770	–	–	–	–	–	6.576**	Kothari and Shanken (1997)
0.0321**	–	–	–0.0222	–	–	–	–	4.023	Lamont (1998)
–	0.0160	0.0413	–	–	–	0.5046	–	8.391**	Campbell and Vuolteenaho (2004)
–	0.0361**	–	–	–0.3755*	–	–	–0.6152*	13.469***	General-to-specific approach
Panel B: 1952Q1–2012Q4									
d/p	e/p	b/m	d/e	tbl	dfy	tms	cay	Joint Wald	Related study/ Model
0.0483	–	–	–	–0.6828*	–	–	–	3.745	Ang and Bekaert (2007)
0.0434	–	–	–	–0.5884	0.5073	0.2380	–	6.880	Ferson and Schadt (1996)
0.0706	–	–0.0783	–	–	–	–	–	1.883	Kothari and Shanken (1997)
0.0235	–	–	0.0114	–	–	–	–	1.954	Lamont (1998)
–	0.0089	0.0161	–	–	–	0.7553**	–	4.574	Campbell and Vuolteenaho (2004)
0.0230	–	–	–0.0006	–	–	–	0.7333**	13.199***	Lettau and Ludvigson (2001)
–	0.0390**	–	–	–0.7339***	2.4016**	–	0.9749***	23.985***	General-to-specific approach

Table 10**Finite-sample sizes for long-horizon Wald test**

This table presents finite-sample sizes, derived from K -horizon univariate predictive regressions, as in Equation (30), under the null hypothesis $H_0 : A = 0$ in the DGP (22). $W_{0.05}$ corresponds to the rejection rate for the long-horizon Wald statistic, defined in (34), with 5% nominal size. Results are reported for different degrees of correlation between the residuals of regressions (22) and (23), $\delta = -0.95, -0.5, \text{ and } 0$, different sample sizes $n = 100, 500, \text{ and } 1,000$, different horizons K that are empirically relevant to the corresponding sample size n and different local-to-unity parameters $C = 0, -5, -10, -20, \text{ and } -50$. The reported results are based on the Monte Carlo simulation described in Section 5.2, and the average rejection rates are calculated over 10,000 repetitions.

$n=100$					$n=500$					$n=1,000$				
		$\delta=-0.95$	$\delta=-0.5$	$\delta=0$			$\delta=-0.95$	$\delta=-0.5$	$\delta=0$			$\delta=-0.95$	$\delta=-0.5$	$\delta=0$
C	K	$W_{0.05}$	$W_{0.05}$	$W_{0.05}$	C	K	$W_{0.05}$	$W_{0.05}$	$W_{0.05}$	C	K	$W_{0.05}$	$W_{0.05}$	$W_{0.05}$
0	2	0.067	0.060	0.051	0	4	0.060	0.054	0.050	0	4	0.056	0.055	0.051
	3	0.062	0.062	0.050		8	0.055	0.050	0.051		12	0.053	0.051	0.049
	4	0.059	0.057	0.048		12	0.053	0.050	0.045		36	0.048	0.044	0.048
	5	0.057	0.055	0.047		20	0.050	0.049	0.047		60	0.044	0.042	0.049
-5	2	0.067	0.060	0.053	-5	4	0.060	0.059	0.050	-5	4	0.059	0.056	0.050
	3	0.064	0.060	0.052		8	0.063	0.053	0.053		12	0.060	0.052	0.047
	4	0.062	0.050	0.048		12	0.060	0.052	0.053		36	0.052	0.053	0.045
	5	0.059	0.050	0.048		20	0.057	0.049	0.044		60	0.047	0.043	0.047
-10	2	0.061	0.062	0.050	-10	4	0.059	0.052	0.050	-10	4	0.061	0.049	0.047
	3	0.066	0.057	0.049		8	0.056	0.056	0.050		12	0.054	0.055	0.052
	4	0.059	0.051	0.054		12	0.058	0.054	0.049		36	0.053	0.052	0.048
	5	0.058	0.052	0.047		20	0.053	0.048	0.049		60	0.049	0.044	0.047
-20	2	0.058	0.057	0.055	-20	4	0.057	0.056	0.050	-20	4	0.054	0.051	0.047
	3	0.057	0.052	0.049		8	0.056	0.051	0.050		12	0.057	0.051	0.048
	4	0.063	0.054	0.049		12	0.054	0.052	0.046		36	0.050	0.050	0.048
	5	0.055	0.052	0.052		20	0.054	0.047	0.046		60	0.052	0.049	0.043
-50	2	0.050	0.053	0.059	-50	4	0.052	0.050	0.050	-50	4	0.052	0.052	0.051
	3	0.051	0.055	0.051		8	0.050	0.051	0.050		12	0.048	0.052	0.051
	4	0.050	0.053	0.051		12	0.049	0.048	0.052		36	0.052	0.053	0.047
	5	0.051	0.051	0.053		20	0.051	0.050	0.049		60	0.053	0.050	0.046

Table 11**Long-horizon univariate predictive regressions, monthly data**

This table presents the results of long-horizon univariate predictive regression models, as in Equation (30), during the sample periods January 1927–December 2012 (panel A) and January 1952–December 2012 (panel B), for various horizons (K -mths). The dependent variable is the cumulative S&P 500 value-weighted log excess return from month t to month $t+K-1$, corresponding to a horizon of K months, and the lagged persistent regressor is each of the following variables defined in Section 3: Dividend payout ratio (d/e), long-term yield (lty), dividend yield (d/y), dividend price ratio (d/p), T-bill rate (tbl), earnings price ratio (e/p), book-to-market value ratio (b/m), default yield spread (dfy), net equity expansion (ntis), term spread (tms), and inflation rate (inf). The table reports the long-horizon Wald statistic, defined in Equation (34), under the null hypothesis that the slope coefficient of the long-horizon univariate predictive regression estimated via the proposed instrumental variable (IVX) approach, is equal to zero (i.e., no predictability). *, **, and *** imply rejection of the null hypothesis at 10%, 5%, and 1% level, respectively.

Panel A: January 1927–December 2012											
K -mths	d/e	lty	d/y	d/p	tbl	e/p	b/m	dfy	ntis	tms	inf
4	0.138	0.752	2.322	2.271	1.413	3.978**	4.851**	0.054	4.805**	1.125	0.781
12	0.005	0.195	3.492*	3.230*	0.947	4.538**	5.767**	0.124	9.123***	2.156	0.528
24	0.472	0.061	3.772*	3.782*	0.774	3.335*	4.501**	0.141	8.784***	3.080*	0.022
36	0.803	0.039	3.415*	3.452*	0.918	2.806*	3.866**	0.105	6.816***	5.025**	0.001
48	0.422	0.021	3.150*	3.234*	0.668	3.418*	3.788*	0.222	4.960**	4.642**	0.053
60	0.637	0.024	2.912*	3.018*	0.525	3.044*	2.970*	0.232	4.309**	4.022**	0.057
Panel B: January 1952–December 2012											
K -mths	d/e	lty	d/y	d/p	tbl	e/p	b/m	dfy	ntis	tms	inf
4	1.522	0.821	1.517	1.386	2.483	0.372	0.367	0.866	0.006	3.367*	5.507**
12	1.717	0.133	1.810	1.763	1.406	0.761	0.642	0.549	0.005	4.422**	8.328***
24	4.392**	0.009	1.584	1.639	0.651	0.286	0.241	0.048	0.147	3.494*	3.670*
36	5.779**	0.000	1.269	1.306	0.449	0.203	0.063	0.014	0.119	3.654*	2.400
48	3.317*	0.045	0.901	0.932	0.157	0.467	0.050	0.010	0.040	3.388*	2.297
60	3.856**	0.127	0.883	0.896	0.039	0.541	0.112	0.093	0.001	3.412*	1.311

Table 12**Long-horizon univariate predictive regressions, quarterly data**

This table presents the results of long-horizon univariate predictive regression models, as in Equation (30), during the sample periods 1927Q1–2012Q4 (panel A) and 1952Q1–2012Q4 (panel B), for various horizons (K -qtrs). The dependent variable is the cumulative S&P 500 value-weighted log excess return from quarter t to quarter $t+K-1$, corresponding to a horizon of K quarters, and the lagged persistent regressor is each of the following variables defined in Section 3: dividend payout ratio (d/e), long-term yield (lty), dividend yield (d/y), dividend price ratio (d/p), T-bill rate (tbl), earnings price ratio (e/p), book-to-market value ratio (b/m), default yield spread (dfy), net equity expansion (ntis), term spread (tms), inflation rate (inf), and consumption-wealth ratio (cay). The table reports the long-horizon Wald statistic, defined in Equation (34), under the null hypothesis that the slope coefficient of the long-horizon univariate predictive regression estimated via the proposed instrumental variable (IVX) approach, is equal to zero (i.e., no predictability). *, **, and *** imply rejection of the null hypothesis at 10%, 5%, and 1% level, respectively.

Panel A: 1927Q1–2012Q4												
K -qtrs	d/e	lty	d/y	d/p	tbl	e/p	b/m	dfy	ntis	Tms	inf	
4	0.000	0.173	3.537*	3.362*	0.746	4.221**	5.750**	0.139	7.672***	1.564	0.116	
8	0.424	0.047	3.567*	3.648*	0.614	3.010*	4.207**	0.170	6.135**	2.475	0.021	
12	0.703	0.029	3.190*	3.233*	0.697	2.461	3.428*	0.112	4.466**	3.827*	0.036	
16	0.378	0.017	2.771*	2.954*	0.510	2.906*	3.181*	0.201	3.063*	3.496*	0.083	
20	0.527	0.017	2.562	2.744*	0.408	2.623	2.506	0.203	2.419	3.158*	0.061	
Panel B: 1952Q1–2012Q4												
K -qtrs	d/e	lty	d/y	d/p	tbl	e/p	b/m	dfy	ntis	Tms	inf	Cay
4	1.409	0.132	1.902	1.902	1.201	0.857	0.824	0.391	0.022	3.569*	5.511**	11.022***
8	3.516*	0.005	1.524	1.686	0.530	0.361	0.352	0.030	0.088	2.977*	2.585	8.794***
12	4.865**	0.000	1.269	1.348	0.353	0.244	0.113	0.003	0.079	2.993*	1.784	7.326***
16	2.960*	0.034	0.895	0.961	0.135	0.463	0.062	0.007	0.038	2.809*	1.693	6.048**
20	3.247*	0.112	0.878	0.911	0.034	0.558	0.126	0.069	0.002	2.974*	0.927	5.000**

Table 13**Long-horizon predictive regressions with multiple regressors**

This table presents the results of long-horizon predictive regression models with multiple regressors, as in Equation (30). Panel A contains the results for monthly data, and panel B contains the results for quarterly data. Each panel reports results for the full sample period, 1927–2012, and the subperiod 1952–2012. Results are reported for various horizons (K -mths in panel A; K -qtrs in panel B). In panel A, the dependent variable is the cumulative S&P 500 value-weighted log excess return from month t to month $t+K-1$, corresponding to a horizon of K months. In panel B, the dependent variable is the cumulative S&P 500 value-weighted log excess return from quarter t to quarter $t+K-1$, corresponding to a horizon of K quarters. The lagged persistent regressors are combinations of the following variables: earnings price ratio (e/p), T-bill rate (tbl), default yield spread (dfy), net equity expansion (ntis), and consumption-wealth ratio (cay). The combination of regressors used in each presented case is the one derived from the general-to-specific approach for one-period regressions, as described in Section 4.2 and presented in Tables 8 and 9. The table reports the long-horizon Wald statistic, defined in Equation (34), testing the individual significance of each regressor, that is, under the null hypothesis that the corresponding slope coefficient of the long-horizon regression estimated via the proposed instrumental variable (IVX) approach, is equal to zero. It also reports the corresponding joint Wald statistic testing the joint significance of the employed regressors, that is, under the null hypothesis that all slope coefficients of the long-horizon regression are jointly equal to zero. *, **, and *** imply rejection of the null hypothesis at 10%, 5%, and 1% level, respectively.

Panel A: Monthly data									
Period: January 1927–December 2012					Period: January 1952–December 2012				
K -mths	e/p	tbl	Joint Wald		e/p	tbl	Joint Wald		
4	5.778**	3.894**	7.638**		3.257*	5.666**	5.734*		
12	6.383**	3.166*	7.614**		4.093**	4.986**	5.289*		
24	4.990**	2.124	5.794*		2.273	2.411	2.596		
36	4.599**	1.915	5.383*		2.049	1.885	2.116		
48	4.983**	1.441	5.660*		2.207	1.702	2.216		
60	4.321**	1.039	4.822*		1.814	1.258	1.825		

Panel B: Quarterly data									
Period: 1927Q1–2012Q4					Period: 1952Q1–2012Q4				
K -qtrs	e/p	tbl	ntis	Joint Wald	e/p	tbl	dfy	cay	Joint Wald
4	4.862**	2.922*	4.928**	13.530***	3.741*	6.114**	4.890**	16.786***	23.548***
8	3.500*	2.157	3.988**	10.393**	2.013	3.662*	1.009	11.809***	16.118***
12	3.791*	2.067	2.421	8.296**	1.477	2.683	0.062	6.733***	13.292***
16	4.150**	1.689	1.175	7.300*	1.380	1.854	0.036	3.852**	11.569**
20	3.854**	1.383	0.600	6.102	0.859	0.948	0.045	1.604	10.664**

Figure 1

Power plots for sample size $n=250$ and residuals' correlation coefficient $\delta=-0.95$

This figure shows the rejection rates for tests of the null hypothesis $H_0 : A = 0$ versus the alternative $H_1 : A \neq 0$ in (22) as the true value of A increases. The solid curve ($Wald_{0.05}$) illustrates the rejection rate we get using the Wald test, defined in Equation (19), with 5% nominal size (horizontal line). The dashed curve ($CYQ_{0.05}$) illustrates the rejection rate using the 95% confidence interval of the Campbell and Yogo (2006) Q -test. The dash-dot curve ($JM_{0.05}$) illustrates the rejection rate using the $\pi_{0.05}^*$ statistic of Jansson and Moreira (2006). Each panel corresponds to a different local-to-unity parameter $C= 0, -5, -10, -15, -20,$ and -50 . These rejection rates have been calculated using Monte Carlo simulations described in Section 2.1 with 10,000 repetitions for a sample size of $n=250$, correlation coefficient between the residuals of regressions (22) and (23) $\delta = -0.95$, and for no autocorrelation in the residuals of the autoregressive equation, that is, $\phi = 0$ in (24).




Figure 2

Power plots for sample size $n=250$ and residuals' correlation coefficient $\delta=-0.5$

This figure shows the rejection rates for tests of the null hypothesis $H_0 : A = 0$ versus the alternative $H_1 : A \neq 0$ in (22) as the true value of A increases. The solid curve ($Wald_{0.05}$) illustrates the rejection rate we get using the Wald test, defined in Equation (19), with 5% nominal size (horizontal line). The dashed curve ($CYQ_{0.05}$) illustrates the rejection rate using the 95% confidence interval of the Campbell and Yogo (2006) Q -test. The dash-dot curve ($JM_{0.05}$) illustrates the rejection rate using the $\pi_{0.05}^*$ statistic of Jansson and Moreira (2006). Each panel corresponds to a different local-to-unity parameter $C= 0, -5, -10, -15, -20,$ and -50 . These rejection rates have been calculated using Monte Carlo simulations described in Section 2.1 with 10,000 repetitions for a sample size of $n=250$, correlation coefficient between the residuals of regressions (22) and (23) $\delta=-0.5$, and for no autocorrelation in the residuals of the autoregressive equation, that is, $\phi=0$ in (24).




Figure 3

Power plots for sample size $n=250$ and residuals' correlation coefficient $\delta=0$

This figure shows the rejection rates for tests of the null hypothesis $H_0 : A = 0$ versus the alternative $H_1 : A \neq 0$ in (22) as the true value of A increases. The solid curve ($\text{Wald}_{0.05}$) illustrates the rejection rate we get using the Wald test, defined in Equation (19), with 5% nominal size (horizontal line). The dashed curve ($\text{CYQ}_{0.05}$) illustrates the rejection rate using the 95% confidence interval of the Campbell and Yogo (2006) Q -test. The dash-dot curve ($\text{JM}_{0.05}$) illustrates the rejection rate using the $\pi_{0.05}^*$ statistic of Jansson and Moreira (2006). Each panel corresponds to a different local-to-unity parameter $C = 0, -5, -10, -15, -20,$ and -50 . These rejection rates have been calculated using Monte Carlo simulations described in Section 2.1 with 10,000 repetitions for a sample size of $n=250$, correlation coefficient between the residuals of regressions (22) and (23) $\delta=0$, and for no autocorrelation in the residuals of the autoregressive equation, that is, $\phi = 0$ in (24).




Figure 4

Power plots for joint Wald test with multiple regressors (correlation set 1)

This figure shows the rejection rates for the joint Wald test defined in (19), with 5% nominal size, under the null hypothesis $H_0 : A = 0_{1 \times 3}$, that is, that all three coefficients in vector A are equal to zero, as the true value of each regressor coefficient A_i increases. The joint Wald test is based on the multivariate predictive system in (26), with three regressors exhibiting different degrees of persistence (unit root, local-to-unity, and stationary). The solid curve (Wald^{UR}) illustrates the rejection rate for the joint Wald test as the true value of the unit root regressor coefficient increases. The dashed curve (Wald^{LTU}) illustrates the corresponding rejection rate as the true value of the local-to-unity regressor coefficient increases. The dotted curve (Wald^{Stationary}) illustrates the corresponding rejection rate as the true value of the stationary regressor coefficient increases. These rejection rates have been calculated using Monte Carlo simulations described in Section 2.4 with 10,000 repetitions for different sample sizes: $n=100, 250, 500,$ and $1,000$. The correlation coefficients (δ 's) between the residuals of regressions (26) and (27) are estimated using S&P 500 value-weighted log excess return (regressand), earnings-price ratio (UR), T-bill rate (LTU), and inflation rate (Stationary) with monthly data for the period 1927–2012, that is, correlation set 1. The utilized autocorrelation coefficients (φ 's) for the autoregressions are the corresponding sample estimates for each of the three regressors mentioned above.




Figure 5

Power plots for joint Wald test with multiple regressors (correlation set 2)

This figure shows the rejection rates for the joint Wald test defined in (19), with 5% nominal size, under the null hypothesis $H_0 : A = 0_{1 \times 3}$, that is, that all three coefficients in vector A are equal to zero, as the true value of each regressor coefficient A_i increases. The joint Wald test is based on the multivariate predictive system in (26), with three regressors exhibiting different degrees of persistence (unit root, local-to-unity, and stationary). The solid curve (Wald^{UR}) illustrates the rejection rate for the joint Wald test as the true value of the unit root regressor coefficient increases. The dashed curve (Wald^{LTU}) illustrates the corresponding rejection rate as the true value of the local-to-unity regressor coefficient increases. The dotted curve (Wald^{Stationary}) illustrates the corresponding rejection rate as the true value of the stationary regressor coefficient increases. These rejection rates have been calculated using Monte Carlo simulations described in Section 2.4 with 10,000 repetitions for different sample sizes: $n=100, 250, 500,$ and $1,000$. The correlation coefficients (δ 's) between the residuals of regressions (26) and (27) are estimated using S&P 500 value-weighted log excess return (regressand), earnings-price ratio (UR), T-bill rate (LTU) and inflation rate (Stationary) with quarterly data for the period 1927–2012, that is, correlation set 2. The utilized autocorrelation coefficients (φ 's) for the autoregressions are the corresponding sample estimates for each of the three regressors mentioned above.

