

*Citation for published version:*

Sattar, FA, Darley, DJ, Politano, F, Woodman, TJ, Threadgill, MD & Lloyd, MD 2012, 'Acyl-CoA -proton exchange by -methylacyl-CoA racemase', Biochemical Society Conference, 2012, Cambridge, UK United Kingdom, 1/01/12.

*Publication date:*  
2012

*Document Version*  
Early version, also known as pre-print

[Link to publication](#)

## University of Bath

### Alternative formats

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Unexpected stereoselective exchange of straight-chain fatty acyl-CoA  $\alpha$ -protons by human  $\alpha$ -methylacyl-CoA racemase 1A (P504S)

F. A. Sattar, D. J. Darley, F. Politano, T. J. Woodman, M. D. Threadgill and M. D. Lloyd

Medicinal Chemistry, Department of Pharmacy & Pharmacology, University of Bath, Claverton Down, Bath BA2 7AY, United Kingdom; [M.D.Lloyd@bath.ac.uk](mailto:M.D.Lloyd@bath.ac.uk)

$\alpha$ -Methylacyl-CoA racemase (AMACR; P504S) enzyme levels are increased ~10-fold in prostate and other cancers. The enzyme catalyses chiral inversion of 2-methylacyl-CoA substrates by removal of the C-2 proton followed by non-stereoselective reprotonation resulting in a ~1:1 epimeric mixture of 2*S*- and 2*R*-methyl products. AMACR is localized in peroxisomes and mitochondria. Both organelles contain straight-chain acyl-CoA esters, which are potential substrates for AMACR.

The present study investigates whether straight-chain acyl-CoA esters are substrates for the enzyme. Incubation of decanoyl-CoA with AMACR in the presence of  $^2\text{H}_2\text{O}$  resulted in proton exchange, as judged by  $^1\text{H-NMR}$ . The reaction was reversible, as shown by deuterium loss from 2,2'-[ $^2\text{H}_2$ ]-decanoyl-CoA when incubated with AMACR in  $^1\text{H}_2\text{O}$ -containing buffer. Incubation of 2-[ $^{13}\text{C}_1$ ]-decanoyl-CoA with AMACR in the presence of  $^2\text{H}_2\text{O}$  showed only one proton was exchanged, as judged by the conversion of the substrate  $^{13}\text{C}$ -singlet to a triplet in the NMR spectrum. Unexpectedly, chiral derivatization of incubation products followed by  $^1\text{H}$  NMR analysis showed that the 2*S*-proton was exchanged much more frequently than the 2*R*-proton. Steady-state kinetic analysis of decanoyl-CoA and *S*-2-methyldecanoyl-CoA esters showed that straight-chain acyl-CoA esters are poor substrates for AMACR, as judged by  $k_{\text{cat}}/K_{\text{m}}$  values (114 vs. 37  $\text{M}^{-1} \text{s}^{-1}$ ).

Straight-chain acyl-CoA esters are AMACR substrates *in vitro*, but these results suggest that AMACR does not substantially catalyse proton-exchange of straight-chain acyl-CoA substrates *in vivo*.

219 words

MTG2011/0001041