



Citation for published version:

Denstel, KD, Broyles, ST, Larouche, R, Sarmiento, OL, Barreira, TV, Chaput, J-P, Barreira, TV, Church, TS, Fogelholm, M, Hu, G, Kuriyan, R, Lambert, EV, Maher, C, Maia, J, Matsudo, V, Olds, T, Onywera, V, Standage, M, Tremblay, MS, Tudor-Locke, C, Zhao, P & Katzmarzyk, PT 2015, 'Active school transport and weekday physical activity in 9-11 year old children from 12 countries', *International Journal of Obesity*, vol. 5, no. S2, pp. S100-S106. <https://doi.org/10.1038/ijosup.2015.26>

DOI:

[10.1038/ijosup.2015.26](https://doi.org/10.1038/ijosup.2015.26)

Publication date:

2015

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

Original Article

Active School Transport and Weekday Physical Activity in 9-11 year old Children from 12 Countries

Kara D. Denstel, MPH¹; Stephanie T. Broyles, PhD¹; Richard Larouche, PhD²; Olga L. Sarmiento, MD, PhD³; Tiago V. Barreira, PhD^{1,4}; Jean-Philippe Chaput, PhD²; Timothy S. Church, MD, PhD¹; Mikael Fogelholm, ScD⁵; Gang Hu, MD, PhD¹; Rebecca Kuriyan, PhD⁶; Anura Kurpad, MD, PhD⁶; Estelle V. Lambert, PhD⁷; Carol Maher, PhD⁸; Jose Maia, PhD⁹; Victor Matsudo, MD, PhD¹⁰; Timothy Olds, PhD⁸; Vincent Onywera, PhD¹¹; Martyn Standage, PhD¹²; Mark S. Tremblay, PhD²; Catrine Tudor-Locke, PhD^{1,13}; Pei Zhao, MD¹⁴; and Peter T. Katzmarzyk PhD¹ for the ISCOLE Research Group

¹Pennington Biomedical Research Center, Baton Rouge, LA;

²Children's Hospital of Eastern Ontario Research Institute, Ottawa, Canada;

³School of Medicine, Universidad de los Andes, Bogotá, Colombia;

⁴Exercise Science, University of Syracuse, Syracuse, NY;

⁵Department of Food and Environmental Sciences, University of Helsinki, Helsinki, Finland;

⁶St. Johns Research Institute, Bangalore, India;

⁷Division of Exercise Science and Sports Medicine, Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa;

⁸Alliance for Research in Exercise Nutrition and Activity (ARENA), School of Health Sciences, University of South Australia, Adelaide, Australia;

⁹CIFI²D, Faculdade de Desporto, University of Porto, Porto, Portugal;

¹⁰Centro de Estudos do Laboratório de Aptidão Física de São Caetano do Sul (CELAFISCS), Sao Paulo, Brazil;

¹¹Department of Recreation Management and Exercise Science, Kenyatta University, Nairobi, Kenya;

¹²Department for Health, University of Bath, Bath, United Kingdom;

¹³Department of Kinesiology, University of Massachusetts Amherst, Amherst, United States;

¹⁴Tianjin Women's and Children's Health Center, Tianjin, China

Running Title: Active Transportation and Physical Activity in Children

Address for Correspondence and Reprints: Peter T. Katzmarzyk, PhD, Pennington Biomedical Research Center, 6400 Perkins Road, Baton Rouge, LA 70808-4124; Phone (225) 763-2536; Fax: (225) 763-2927; Email: Peter.Katzmarzyk@pbrc.edu

40 **Abstract**

41 **Background:** Active school transport (AST) may increase the time that children spend in
42 physical activity (PA). This study examined relationships between AST and weekday moderate-
43 to-vigorous physical activity (MVPA), light physical activity (LPA), sedentary time (SED), and
44 total activity during naturally organized time periods (daily, before school, during school, and
45 after school) in a sample of children from 12 countries.

46 **Methods:** The sample included 6 224 children aged 9-11 years. PA and sedentary time were
47 objectively measured using Actigraph accelerometers. AST was self-reported by participants.
48 Multi-level generalized linear and logistic regression statistical models were used to determine
49 associations between PA, SED and AST across and within study sites.

50 **Results:** After adjustment for age, highest parental educational attainment, BMI z-score, and
51 accelerometer wear time, children who engaged in AST accumulated significantly more
52 weekday MVPA during all studied time periods and significantly less time in LPA before school
53 compared to children who used motorized transport to school. AST was unrelated to time spent
54 in sedentary behaviors. Across all study sites, AST was associated with 6.0 min (95% CI: 4.7-
55 7.3; $p < 0.0001$) more of weekday MVPA; however, there was some evidence that this differed
56 across study sites (p for interaction = 0.06). Significant positive associations were identified within
57 7 of 12 study sites, with differences ranging from 4.6 min (95% CI: 0.3-8.9; $p = 0.04$, in Canada)
58 to 10.2 min (95% CI: 5.9-14.4; $p < 0.0001$, in Brazil) more of daily MVPA among children who
59 engaged in AST compared to motorized transport.

60 **Conclusions:** The present study demonstrated that AST was associated with children spending
61 more time engaged in MVPA throughout the day and less time in LPA before school. AST
62 represents a good behavioral target to increase levels of PA in children.

63 **Keywords:** commuting, walking, pediatrics, accelerometer, global

64
65 **Trial Registration:** ClinicalTrials.gov: Identifier NCT01722500.

66 **Introduction**

67 Physical activity (PA) guidelines recommend that children engage in at least 60 min of
68 moderate-to-vigorous physical activity (MVPA) each day to lower their risk of negative health
69 consequences; yet globally, many children do not meet these guidelines.¹⁻³ The journeys to and
70 from school provide an opportunity to establish habitual PA patterns.⁴ Previous research
71 suggests that children who engage in active school transport (AST), such as walking or biking,
72 accumulate more PA and experience better cardiorespiratory, muscular, and metabolic fitness
73 and a lower likelihood of diabetes and obesity than those who travel using motorized transport
74 alternatives including by car or bus.⁵⁻⁸

75 Despite the documented benefits of AST, the prevalence has declined in many countries
76 over recent decades. In the United States (US), the proportion of children who walked or
77 bicycled to school declined from 48% to 13% between 1969 and 2009.⁹ Similar declines have
78 been documented in the United Kingdom (UK)¹⁰, Australia¹¹, Brazil¹², Canada¹³, Kenya¹⁴ and
79 Switzerland.¹⁵ The most common parent-reported barriers to AST include distance from home to
80 school, traffic concerns, and crime.^{16, 17}

81 While a positive association between AST and MVPA among children has been shown,^{4, 13}
82 the majority of evidence has been limited to developed countries. Additionally, few studies have
83 examined associations between AST and light-intensity physical activity (LPA) or sedentary time
84 (SED). Further, little is known about the contribution of AST to PA (MVPA or other intensities)
85 or sedentary time during different naturally organized time periods throughout the segmented
86 school day. It is important to more fully understand how AST contributes to PA of various
87 intensities and SED time globally in order to plan and implement effective interventions,
88 programs, and policies in a variety of settings.

89 Given these research gaps, the aims of this study were to examine: (a) relationships
90 between AST and objectively measured weekday MVPA, LPA, total activity, and SED; (b) how
91 these relationships differ across different time periods of the school day; and (c) associations

92 between children's AST and accumulating an average of at least 60 min per weekday (Monday
93 through Friday) of MVPA in a sample of 9 to 11 year old children from 12 countries ranging in
94 level of socio-economic and human development.

95 **Methods**

96 **Study Design and Participants**

97 The International Study of Childhood Obesity, Lifestyle and the Environment (ISCOLE) is a
98 cross-sectional, multi-national study conducted in sites in 12 countries including Australia,
99 Brazil, Canada, China, Colombia, Finland, India, Kenya, Portugal, South Africa, UK, and US.
100 The ISCOLE protocol was approved by the ethics committee at the Pennington Biomedical
101 Research Center and each participating institution. Written informed consent was obtained from
102 parents or legal guardians, and child assent, if required by the site's local ethics committee, was
103 also obtained before children participated in the study. Data were collected from September
104 2011 through December 2013. Detailed information on the study design and methods can be
105 found elsewhere.¹⁸

106 A total of 7 372 children 9-11 years of age from 256 schools participated in ISCOLE. The
107 present analytical sample includes 6 224 children after excluding participants missing AST
108 (n=62), valid accelerometry (n=798), parental education (n=283), and body mass index (BMI) z-
109 score (n=5). Participants who were excluded in this study did not differ from those included with
110 respect to the proportion in AST, but were more likely to be older, male, have a higher body
111 mass index (BMI), and have parents who achieved less than a college education. The number
112 of participants per site with complete data for each of the variables of interest for this study
113 ranged from 400 in South Africa to 856 in Colombia. The primary sampling frame was schools
114 which were stratified by an indicator of socioeconomic status. Classrooms within the recruited
115 schools were the secondary sampling frame chosen to yield a sample of children with a mean
116 age of 10 years.

117

118 **Measurements**

119 *Accelerometry*

120 The full accelerometer protocol has been previously reported.¹⁹ Briefly, time spent in MVPA,
121 LPA, and SED and total activity counts were obtained from 24-hour accelerometry. Children
122 were encouraged to wear an Actigraph GT3X+ accelerometer (ActiGraph LLC, Pensacola, FL,
123 USA) on a belt around the waist at the right mid-axillary line 24 hours per day for at least 7
124 consecutive days. A valid accelerometer record required ≥ 4 days with ≥ 10 h of waking wear
125 time per day, including at least one weekend day. Sleep and non-wear time were identified
126 using an automated algorithm prior to identification of other activities.^{20, 21} A waking non-wear
127 period began with 20 consecutive minutes of zero activity counts.²⁰ Activity cut points were
128 based on Evenson.²² Total activity was expressed as the number of activity counts per day.
129 MVPA was defined as all activity greater than or equal to 574 activity counts/15 s, LPA was
130 defined as activity from 26 to 573 counts/15 s, and SED was defined as all activity ≤ 25
131 counts/15 s²² not including the sleep period and non-wear time. Minutes spent in MVPA, LPA,
132 and SED and total activity counts were assigned to the before school, during school, and after
133 school time periods using school day schedules provided by each participating school. The
134 before school time period was considered wake time (established using a validated algorithm²⁰)
135 until school start time, during school was defined as the time between school start and end time,
136 and the after school period was considered school end time through the child's bed time (also
137 determined by our validated algorithm²⁰) determined objectively from accelerometry. Only
138 weekday accelerometer data were used in these analyses to more accurately capture the effect
139 of AST on PA.

140 In order to determine whether or not children met PA guidelines, the number of valid
141 weekdays contained in the accelerometer file was determined, and the mean number of MVPA
142 minutes per valid weekday that children accumulated was calculated. Children were then

143 classified as meeting PA guidelines (yes or no) if they obtained a mean of ≥ 60 min of MVPA
144 per weekday over the measurement period.

145 *Active school transport (AST)*

146 AST was self-reported by participants using a diet and lifestyle questionnaire¹⁸ with the
147 question “In the last week you were in school, the main part of your journey to school was by”.
148 Response options included “walking”, “bicycle, roller-blade, skateboard or non-motorized
149 scooter”, “bus, train, tram, underground or boat”, “car, motorcycle or moped”, and “other”. A
150 participant was considered to have engaged in AST if they reported walking, running, bicycle,
151 roller-blades, skateboard, or non-motorized scooter. Motorized transportation response options
152 included bus, train, tram, underground, boat, car, motorcycle, or moped. These options were
153 collapsed into a binomial variable indicating AST or motorized transport to school.

154 *Covariates*

155 Standing height and body weight were measured using standard procedures across study
156 sites.¹⁸ Detailed measurement procedures have been published elsewhere.¹⁸ Body mass index
157 ($[\text{weight}(\text{kg})/\text{height}(\text{m})^2]$; BMI) was derived from the average standing height and weight, and
158 BMI z-score was computed using age- and sex-specific reference values from the World Health
159 Organization (WHO).²³ A demographic questionnaire completed by parents captured the
160 participant’s date of birth, sex, and parents’ educational attainment. Age was computed from the
161 date of birth and date anthropometric measurements were performed. Highest parental
162 educational attainment was created based on the highest education level completed by either
163 parent (did not complete high school, completed high school or some college, completed
164 bachelor’s degree or postgraduate degree).

165 *Statistical analysis*

166 Means and standard deviations were computed for variables by study site and sex for
167 participants with complete data for accelerometry, AST, and covariates. AST was the
168 independent variable in all analyses. Separate multi-level generalized linear mixed models (SAS

169 version 9.4, PROC MIXED) were used to examine the associations between mean daily, before
170 school, during school, and after school MVPA, LPA, SED, and AST (0 = no, 1 = yes). All linear
171 models were stratified by sex and adjusted for mean waking accelerometer wear time, highest
172 parental educational attainment, average BMI z-score, and a site by AST interaction. Study site
173 and school nested within study site were treated as fixed effects to account for the clustering of
174 the data (weekday MVPA intra-class correlation; ICC= .14 at site; ICC=0.26 at school nested
175 within site). Results are presented as adjusted means using the LSMEANS option. The PDIFF
176 option was used to obtain the difference in the number of weekday MVPA minutes obtained by
177 children who used AST versus motorized transport alternatives within each site. Multi-level
178 logistic mixed models (PROC GLIMMIX) with the SLICEDIFFS and ODDSRATIO options
179 specified were used to obtain the within-site odds of children meeting PA guidelines (0 = no, 1=
180 yes) for those who engaged in AST compared to those who used motorized transport. Logistic
181 models were adjusted for mean waking accelerometer wear time, highest parental educational
182 attainment, average BMI z-score, and a site by AST interaction. All data management and
183 statistical analyses were conducted using SAS version 9.4 (SAS Institute, Cary, NC). The
184 significance level was set at $P < 0.05$.

185 **Results**

186 The descriptive characteristics of the study sample, stratified by study site and sex, are
187 provided in Table 1. The mean \pm SD age of the full sample was 10.4 ± 0.6 years and 54.4% of
188 the participants were girls. The prevalence of AST was 43.1% among boys and 41.8% among
189 girls, and this ranged across sites from 7.2% to 76.2% among boys and 3.7% to 81.0% among
190 girls. Approximately 37% of children travelled to school by walking, 5% by bicycle, roller-blade,
191 skateboard, or non-motorized scooter, 22% by bus, train, tram, underground, or boat, 35% by
192 car, motorcycle, or moped, and 1% travelled by some other mode. More detail regarding mode
193 of transportation prevalences by site can be found elsewhere.²⁴ Across sites, the mean time

194 spent in daily MVPA ranged from 50.9 ± 17.6 min/day to 85.8 ± 29.2 min/day among boys and
195 38.9 ± 14.8 min/day to 66.3 ± 21.6 min/day among girls.

196 The adjusted mean minutes of weekday MVPA, LPA, SED, and total activity counts by
197 mode of transport to school are presented in Table 2. After adjustment for age, highest parental
198 educational attainment, BMI z-score, accelerometer wear time, and an AST by site interaction,
199 AST was associated with obtaining more minutes of MVPA and a higher number of total activity
200 counts for boys at all time periods (daily, before, during, and after school) compared to
201 motorized transport alternatives. Results were similar among girls, but the associations
202 apparent during the school time period were not significant. Girls who engaged in AST
203 accumulated less LPA time daily, and also before, during, and after school compared to those
204 who did not. Among boys, daily and before school LPA was significantly lower among those
205 who reported AST compared to those who did not. There were no significant differences in the
206 amounts of SED time.

207 The differences in mean minutes of daily MVPA between children who engaged in AST
208 compared to motorized transport to school adjusted for age, highest parental educational
209 attainment, BMI z-score, accelerometer wear time, and an AST by site interaction (p for
210 interaction=0.06) are presented in Figure 1. Across all study sites, AST was associated with 6.0
211 min (95% CI: 4.7-7.3; $p<0.0001$) more of daily MVPA. The association was significant within 7 of
212 12 ISCOLE study sites, including Brazil, Canada, Finland, Kenya, Portugal, South Africa, and
213 the UK. Among those with a significant association, the effect size of AST on daily MVPA
214 ranged from 4.6 min (95% CI: 0.3-8.9; $p=0.04$) in Canada to 10.2 min (95% CI: 5.9-14.4;
215 $p<0.0001$) in Brazil.

216 The site-specific odds ratios for the association of AST and obtaining a weekday average of
217 60 or more minutes of MVPA adjusted for age, highest parental educational attainment, BMI z-
218 score, accelerometer wear time, and an AST by site interaction are shown in Figure 2. AST was
219 associated with significantly higher odds of obtaining the recommended amount of MVPA

220 across all sites (OR: 1.8; 95% CI: 1.5-2.1) and within 7 of 12 study sites, including Brazil (OR:
221 2.3; 95% CI: 1.4-3.8), Canada (OR:1.7 ; 95% CI: 1.0-2.8), Finland (OR: 2.1; 95% CI: 1.2-3.8),
222 Kenya (OR: 1.9; 95% CI: 1.1-3.4), Portugal (OR: 2.5; 95% CI: 1.6-3.8), South Africa (OR: 2.8;
223 95% CI: 1.4-5.3), and the UK (OR: 2.4; 95% CI: 1.4-3.9).

224 **Discussion**

225 The present study showed that AST was associated with children spending more time
226 engaged in MVPA and less time in LPA throughout the weekday. Overall, children who
227 undertook AST were 1.8 times more likely to meet daily PA guidelines, though this varied across
228 different study sites.

229 Children who practiced AST recorded almost 10% more daily time in MVPA compared to
230 those who did not. A multicenter study of adolescents from 10 European cities reported similar
231 findings in that time spent walking and biking for travel was positively associated with objectively
232 measured MVPA compared to those who reported less time walking and biking for travel.²⁵
233 However, that study did not differentiate active travel to/from school from trips to other locations.
234 Further, a recent review of 49 studies found that 82% of included studies showed a positive
235 association between AST and daily PA levels (e.g., MVPA, energy expenditure expressed as
236 kilocalorie/day, steps, total PA, etc.) as determined by accelerometry, pedometry, or self-report.⁸
237 Additionally, Lee and colleagues²⁶ showed that children who practiced AST accumulated 6.9
238 min of MVPA during the trip contributing approximately 11% to daily time in MVPA. This agrees
239 with our findings that children who engaged in AST obtained 6 additional minutes per day in
240 MVPA. One study we identified examined AST and associations with MVPA at various time
241 periods throughout the day among 10 year old children in England and found that objectively
242 measured MVPA was significantly higher before and after school among boys and before
243 school among girls who participated in AST.²⁷ These results were similar to the present study;
244 however, our findings also suggest significant positive associations between AST and weekday
245 MVPA during school among boys and after school among girls. Further, intervention studies

246 provide causal evidence that introducing AST into children’s daily routine can significantly
247 increase MVPA. For example, in a randomized controlled trial among 149 4th graders, Mendoza
248 and colleagues²⁸ implemented a 5-week walking school bus intervention in which study staff
249 walked the children to and from school. Children in the intervention group significantly increased
250 their daily MVPA levels from 46.6 min/day to 48.8 min/day while control group children
251 significantly decreased their daily MVPA levels from 46.1 min/day to 41.3 min/day.

252 In contrast to our main findings, no association between AST and MVPA was identified for 5
253 ISCOLE study sites including Australia, China, Colombia, India, and the US. This may have
254 been due to the low AST prevalence in India (5.3%) and the US (8.2%) and lack of power to
255 detect a difference. Additionally, the average weekday MVPA time in our sample was high for
256 Australian children irrespective of school transportation mode suggesting that behaviors other
257 than AST may influence their activity levels. Alternatively, both active and motorized travelers in
258 China spent less time than recommended in weekday MVPA (45.9 min/day vs 43.4 min/day)
259 indicating that behavioral interventions in addition to those promoting AST may be needed to
260 increase MVPA. In Colombia, car ownership is low, and the majority of Colombian children in
261 our sample either practices AST (72.6%) or uses a motorized form of public transportation
262 (20.0%). In the present analysis, public transportation (“bus, train, tram, underground or boat”)
263 was categorized as motorized, but this method likely includes some active travel, such as
264 walking to bus stops, which may have resulted in an underestimation of weekday MVPA among
265 Colombian children who use motorized transport to school.

266 Across all 12 study sites, AST was associated with 80% higher odds of children obtaining
267 the recommended 60 min of MVPA on weekdays compared to motorized transport to school.
268 Further, AST was associated with significantly higher odds of children obtaining a weekday
269 average of at least 60 min per day of MVPA in 7 of 12 study sites. This result supports previous
270 studies that have quantified the association between AST and PA guidelines. Although limited to
271 Russian children, Tudor-Locke and colleagues²⁹ found that omitting activity obtained through

272 AST decreased the prevalence of meeting PA guidelines by 17-19% for boys and girls. While in
273 the same direction, our finding was greater in magnitude which may be due to the use of
274 different PA guidelines; we used an average of 60 min per day during the school week as our
275 proxy for meeting PA guidelines while the aforementioned study utilized a different guideline of
276 'at least 150 min per week or 30 min per school day'.³⁰ Additionally, Tudor-Locke²⁹ measured
277 children's PA by parent-proxy report rather than objectively as was done in the present study.

278 In the present study, boys and girls who engaged in AST recorded approximately 7% less
279 time in LPA before school compared to those who used motorized transport to school. No
280 previous studies were found that examined the relationship between LPA time and AST, but our
281 results suggest that children who participate in AST substitute minutes of LPA for MVPA. LPA
282 confers some health benefits to children, especially those overweight or just beginning a PA
283 routine; however, studies suggest greater health benefits are obtained from more intense PA
284 (i.e., MVPA). Carson, et al.³¹ found that each additional hour adolescents spent in objectively
285 measured higher intensity LPA (800 counts/min - <4 METs) was associated with significantly
286 lower diastolic blood pressure and higher high-density lipoprotein (HDL) cholesterol while
287 additional time spent in MVPA was associated with greater cardiometabolic benefits, including
288 lower systolic blood pressure and insulin, smaller waist circumference, and higher insulin
289 sensitivity (HOMA-S%). Additionally, Ekelund and colleagues³² examined associations between
290 various intensities of objectively measured physical activity among children and adolescents
291 from the European Youth Heart Study and found that MVPA was associated with better
292 cardiometabolic risk profiles compared to LPA. Similar results were found in a sample of
293 Spanish adolescents. Objectively measured MVPA was associated with better body
294 composition measures while LPA was not.³³ This suggests that children who engage in AST
295 accumulate additional health benefits by replacing LPA with MVPA.

296 Insight into the relationship between AST and health in the present sample of children was
297 beyond the scope of this study, but has been reported elsewhere.³⁴ Briefly, Sarmiento and

298 colleagues reported that ISCOLE children who engaged in AST had lower odds of obesity
299 (OR=0.72; 95% CI: 0.60-0.87) and lower body fat percentage (β =-0.56 (0.22); p =0.002)
300 compared to those that did not. These findings are in line with previous studies that
301 demonstrated an inverse association between AST and objectively measured overweight and
302 obesity among children.³⁵⁻³⁷ In contrast, a recent review⁸ concluded that the relationship
303 between AST and body composition is inconsistent. Compared to children who engaged in
304 motorized transport to school, 56% of included studies showed no difference in body
305 composition among active travelers, 36% of studies found more favorable body composition
306 among active travelers, and 8% of studies observed a less favorable body composition among
307 active travelers.⁸ Additionally, a systematic review by Lubans and colleagues³⁸ suggested a
308 positive association between bicycling to school and better cardiorespiratory fitness. Finally, it is
309 well accepted that higher levels of MVPA are associated with decreased risk of obesity and
310 better cardiovascular and metabolic risk profiles.³⁹⁻⁴² Therefore, MVPA may be the mechanism
311 by which AST leads to better health and lower odds of obesity, but intervention studies are
312 needed to elucidate causality in these relationships.

313 Finally, we found no evidence of a relationship between AST and time spent in SED
314 behaviors during any period of the day. A study of 9- and 15-year old children in Europe
315 similarly failed to identify a correlation between walking or biking to school and percent of time
316 spent sedentary.⁴³ The null findings between AST and SED in the present study support
317 previous findings that correlates of children's MVPA appear distinct from those of SED.⁴⁴ This is
318 in line with the growing evidence base that SED and MVPA are independent behaviors.⁴⁵⁻⁴⁷
319 Additionally, since children spend such a large amount of time in SED behaviors, it is unlikely
320 that AST alone would be sufficient to significantly reduce SED time. However, Hinckson and
321 colleagues⁴⁸ found that among children who engage in AST, those living in a 1-2 km radius from
322 their school accumulated less SED time than those living closer or further from school. This

323 suggests that future studies should examine the relationship between AST and SED across
324 various home-to-school distances.

325 Our study has several strengths including the large sample size consisting of children from
326 all major world regions and various levels of development. Additionally, the rigorous
327 standardization of the protocol, measurements, and quality control procedures facilitate
328 comparisons across study sites and increase the integrity of our data. Further, ISCOLE was the
329 first study to employ and analyze a 24-hour waist worn accelerometer protocol to objectively
330 measure various intensities of activity, which led to higher wear time and compliance compared
331 to other studies.⁴⁹ Additionally, time-stamped accelerometer data matched to local daily school
332 start and end times enabled the present study to examine the associations between AST and
333 activity levels during natural time periods before school, during school and after school time
334 periods when such behaviors may especially contribute to MVPA, the first to our knowledge to
335 do so in an international sample of children.

336 Our study also has several limitations. First, the cross-sectional study design impedes
337 causal inference. Next, the large number of participants missing valid accelerometer data may
338 have biased our results. However, the observed differences in physical activity between children
339 who engage in AST and motorized transport alternatives are consistent with previous
340 systematic reviews,^{8, 13} so it is unlikely that our results would have been drastically different if all
341 participants had provided valid accelerometry data. Previous studies suggest MVPA from
342 wheeled modes of AST, such as biking, may be underestimated by hip-worn accelerometers. A
343 sensitivity analysis comparing MVPA obtained from wheeled modes (bicycle, roller-blade,
344 skateboard or non-motorized scooter) to all other modes of AST and motorized transport
345 revealed that the number of minutes of weekday MVPA obtained from wheeled modes of AST
346 was significantly lower than that from other modes of AST and did not differ from motorized
347 transport modes (data not shown). This suggests that we may be underestimating the
348 association between weekday MVPA and AST in countries with a higher prevalence of wheeled

349 AST, such as in Finland where nearly 25% of children in our sample bicycled to school.
350 Additionally, several sites had little variability in the prevalence of AST which may have limited
351 our ability to detect significant within-country differences. Further, ISCOLE assessed method of
352 transportation to school, and children who participate in AST both to and from school may
353 accumulate more MVPA than those only using AST to school.⁸ Future work should collect
354 transport mode to and from school to avoid this potential source of bias. Additionally, the
355 literature suggests the distance between a child's home and school may affect both their PA
356 levels and participation in AST.^{9, 50-52} This distance was not included in the present analyses as
357 a potential moderator as it was not collected in ISCOLE. The between-country variation in the
358 relationship between AST and weekday MVPA may be partially explained if the average home-
359 to-school distance varied across ISCOLE sites. Finally, AST was captured by children's self-
360 report which may be unreliable.^{53, 54} Nevertheless, a recent systematic review indicated that
361 child-reported school travel mode showed substantial test-retest reliability and substantial
362 convergent validity with parent reports.⁵⁵

363 This study was the first to show associations between AST and weekday time in MVPA,
364 LPA, SED, and total activity counts in an international sample of 9-11 year old children. It
365 provided evidence that AST is associated with higher levels of objectively measured MVPA
366 throughout the weekday. Future research should seek to elucidate factors that explain variability
367 in MVPA and AST across various countries. Additionally, there is a need for additional
368 prospective and experimental studies to examine whether or not switching from motorized
369 transport to AST increases MVPA as the cross-sectional design is the key limitation of our
370 study. Finally, barriers to AST need to be further explored in order to develop effective
371 interventions aimed at increasing AST prevalence among children.

372

373

374

375 **Acknowledgments**

376 We wish to thank the ISCOLE External Advisory Board and the ISCOLE participants and their
377 families who made this study possible. RL holds a postdoctoral fellowship from the Canadian
378 Institutes of Health Research. A membership list of the ISCOLE Research Group and External
379 Advisory Board is included in Katzmarzyk, Lambert and Church. An Introduction to the
380 International Study of Childhood Obesity, Lifestyle and the Environment (ISCOLE). Int J Obes
381 Suppl. (This Issue).
382 ISCOLE was funded by The Coca-Cola Company. The funder had no role in study design, data
383 collection and analysis, decision to publish, or preparation of the manuscript.

384

385 **Conflicts of Interest**

386 MF has received a research grant from Fazer Finland and has received an honorarium for
387 speaking for Merck. AK has been a member of the Advisory Boards of Dupont and McCain
388 Foods. RK has received a research grant from Abbott Nutrition Research and Development. VM
389 is a member of the Scientific Advisory Board of Actigraph and has received an honorarium for
390 speaking for The Coca-Cola Company. TO has received an honorarium for speaking for The
391 Coca-Cola Company. The authors reported no other potential conflicts of interest.
392

393 **References**

- 394 1. United States Department of Health and Human Services. 2008 Physical Activity
395 Guidelines for Americans. Washington, DC: 2008.
- 396 2. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in
397 the United States measured by accelerometer. *Med Sci Sport Exer* 2008;40(1):181-8.
- 398 3. Fakhouri TH, Hughes JP, V.L. B, Song M, Fulton JE, Ogden CL. Physical activity in U.S.
399 youth aged 12-15 years, 2012. NCHS data brief. 2014;2014(141):1-8.
- 400 4. Tudor-Locke C, Ainsworth B, Popkin B. Active Commuting to School. *Sports Med*
401 2001;31(5):309-13.
- 402 5. Mendoza JA, Watson K, Nguyen N, Cerin E, Baranowski T, Nicklas TA. Active
403 commuting to school and association with physical activity and adiposity among US
404 youth. *J Phys Act Health* 2011;8(4):488-95.
- 405 6. Pizarro AN, Ribeiro JC, Marques EA, Mota J, Santos MP. Is walking to school
406 associated with improved metabolic health? *Int J Behav Nutr Phys Act* 2013;10:12.
- 407 7. Ostergaard L, Kolle E, Steene-Johannessen J, Anderssen SA, Andersen LB. Cross
408 sectional analysis of the association between mode of school transportation and physical
409 fitness in children and adolescents. *Int J Behav Nutr Phys Act* 2013;10:91.
- 410 8. Larouche R, Saunders TJ, Faulkner G, Colley R, Tremblay M. Associations between
411 active school transport and physical activity, body composition, and cardiovascular
412 fitness: a systematic review of 68 studies. *J Phys Act Health* 2014;11(1):206-27.
- 413 9. McDonald NC. Active transportation to school: trends among U.S. schoolchildren, 1969-
414 2001. *Am J Prev Med* 2007;32(6):509-16.
- 415 10. Black C, Collins A, Snell M. Encouraging Walking: The Case of Journey-to-school Trips
416 in Compact Urban Areas. *Urban Studies*. 2001;38(7):1121-41.

- 417 11. Tremblay MS, Gray CE, Akinroye K, Harrington DM, Katzmarzyk PT, Lambert EV, et al.
418 Physical Activity of Children: A Global Matrix of Grades Comparing 15 Countries. *J Phys*
419 *Act Health* 2014;11(Supp 1): S113-S125.
- 420 12. Costa FF, Silva KS, Schmoelz CP, Campos VC, de Assis MA. Longitudinal and cross-
421 sectional changes in active commuting to school among Brazilian schoolchildren. *Prev*
422 *Med* 2012;55(3):212-4.
- 423 13. Faulkner GE, Buliung RN, Flora PK, Fusco C. Active school transport, physical activity
424 levels and body weight of children and youth: a systematic review. *Prev Med*
425 2009;48(1):3-8.
- 426 14. Onywera VO, Adamo KB, Sheel AW, Waudu JN, Boit MK, Tremblay M. Emerging
427 evidence of the physical activity transition in Kenya. *J Phys Act Health* 2012;9(4):554-62.
- 428 15. Grize L, Bringolf-Isler B, Martin E, Braun-Fahrlander C. Trend in active transportation to
429 school among Swiss school children and its associated factors: three cross-sectional
430 surveys 1994, 2000 and 2005. *Int J Behav Nutr Phys Act* 2010;7(1):28.
- 431 16. Hayne CL, Moran PA, Ford MM. Regulating Environments to Reduce Obesity. *J Public*
432 *Health Pol* 2004;25(3/4):391-407.
- 433 17. Mitra R. Independent Mobility and Mode Choice for School Transportation: A Review
434 and Framework for Future Research. *Trans Rev* 2012;33(1):21-43.
- 435 18. Katzmarzyk P, Barreira T, Broyles S, Champagne C, Chaput J-P, Fogelholm M, et al.
436 The International Study of Childhood Obesity, Lifestyle and the Environment (ISCOLE):
437 design and methods. *BMC Public Health* 2013;13(1):900.
- 438 19. Tudor-Locke C, Barreira T, Schuna J, Mire E, Chaput J-P, Fogelholm M, et al. Improving
439 wear time compliance with a 24-hour waist-worn accelerometer protocol in the
440 International Study of Childhood Obesity, Lifestyle and the Environment (ISCOLE). *Int J*
441 *Behav Nutr Phys Act* 2015;12(1):11.

- 442 20. Barreira TV, Schuna JM, Jr., Mire EF, Katzmarzyk PT, Chaput JP, Leduc G, et al.
443 Identifying Children's Nocturnal Sleep Using 24-h Waist Accelerometry. *Med Sci Sport*
444 *Exerc* 2015;47(5):937-43.
- 445 21. Tudor-Locke C, Barreira TV, Schuna JM, Jr., Mire EF, Katzmarzyk PT. Fully automated
446 waist-worn accelerometer algorithm for detecting children's sleep-period time separate
447 from 24-h physical activity or sedentary behaviors. *Appl Physiol Nutr Metab*
448 2014;39(1):53-7.
- 449 22. Evenson KR, Catellier DJ, Gill K, Ondrak KS, McMurray RG. Calibration of two objective
450 measures of physical activity for children. *J Sports Sci* 2008;26(14):1557-65.
- 451 23. Onis Md, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a
452 WHO growth reference for school-aged children and adolescents. *Bulletin of the World*
453 *Health Organization*. 2007;85:660-7.
- 454 24. Larouche R, Sarmiento OL, Broyles ST, Denstel KD, Church TS, Barriera TV, et al. Are
455 the correlates of active school transport context-specific? *Int J Obes* 2015 [this issue].
- 456 25. Chillón P, Ortega FB, Ruiz JR, De Bourdeaudhuij I, Martinez-Gomez D, Vicente-
457 Rodriguez G, et al. Active commuting and physical activity in adolescents from Europe:
458 results from the HELENA study. *Pediatr Exerc Sci* 2011;23(2):207-17.
- 459 26. Lee C, Li L. Demographic, Physical Activity, and Route Characteristics Related to
460 School Transportation: An Exploratory Study. *Am J Health Promot* 2014;28(sp3):S77-
461 S88.
- 462 27. Cooper AR, Page AS, Foster LJ, Qahwaji D. Commuting to school: are children who
463 walk more physically active? *Am J Prev Med* 2003;25(4):273-6.
- 464 28. Mendoza JA, Watson K, Baranowski T, Nicklas TA, Uscanga DK, Hanfling MJ. The
465 Walking School Bus and Children's Physical Activity: A Pilot Cluster Randomized
466 Controlled Trial. *Pediatrics* 2011;128(3):e537-e44.

- 467 29. Tudor-Locke C, Neff LJ, Ainsworth BE, Addy CL, Popkin BM. Omission of active
468 commuting to school and the prevalence of children's health-related physical activity
469 levels: the Russian Longitudinal Monitoring Study. *Child Care Health Dev*
470 2002;28(6):507-12.
- 471 30. Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C, et al. Physical activity
472 and public health. A recommendation from the Centers for Disease Control and
473 Prevention and the American College of Sports Medicine. *JAMA* 1995;273(5):402-7.
- 474 31. Carson V, Ridgers ND, Howard BJ, Winkler EAH, Healy GN, Owen N, et al. Light-
475 Intensity Physical Activity and Cardiometabolic Biomarkers in US Adolescents. *PLoS*
476 *ONE* 2013;8(8):e71417.
- 477 32. Ekelund U, Anderssen SA, Froberg K, Sardinha LB, Andersen LB, Brage S. Independent
478 associations of physical activity and cardiorespiratory fitness with metabolic risk factors
479 in children: the European youth heart study. *Diabetologia* 2007;50(9):1832-40.
- 480 33. Moliner-Urdiales D, Ruiz JR, Ortega FB, Rey-Lopez JP, Vicente-Rodriguez G, Espana-
481 Romero V, et al. Association of objectively assessed physical activity with total and
482 central body fat in Spanish adolescents; The HELENA Study. *Int J Obes*
483 2009;33(10):1126-35.
- 484 34. Sarmiento OL, Lemoine P, Gonzalez SA, Broyles ST, Denstel KD, Larouche R, et al.
485 Relationships between active school transport and adiposity indicators in school age
486 children from low-, middle- and high-income countries. *Int J of Obes* 2015. [this issue]
- 487 35. Li Y, Zhai F, Yang X, Schouten EG, Hu X, He Y, et al. Determinants of childhood
488 overweight and obesity in China. *Br J Nutr* 2007;97(1):210-5.
- 489 36. Ostergaard L, Grontved A, Borrestad LA, Froberg K, Gravesen M, Andersen LB. Cycling
490 to school is associated with lower BMI and lower odds of being overweight or obese in a
491 large population-based study of Danish adolescents. *J Phys Act Health* 2012;9(5):617-
492 25.

- 493 37. Rosenberg DE, Sallis JF, Conway TL, Cain KL, McKenzie TL. Active transportation to
494 school over 2 years in relation to weight status and physical activity. *Obesity*
495 2006;14(10):1771-6.
- 496 38. Lubans DR, Boreham CA, Kelly P, Foster CE. The relationship between active travel to
497 school and health-related fitness in children and adolescents: a systematic review. *Int J*
498 *Behav Nutr Phys Act* 2011;8:5.
- 499 39. Ness AR, Leary SD, Mattocks C, Blair SN, Reilly JJ, Wells J, et al. Objectively Measured
500 Physical Activity and Fat Mass in a Large Cohort of Children. *PLoS Med* 2007;4(3):e97.
- 501 40. Ekelund U, Sardinha LB, Anderssen SA, Harro M, Franks PW, Brage S, et al.
502 Associations between objectively assessed physical activity and indicators of body
503 fatness in 9- to 10-y-old European children: a population-based study from 4 distinct
504 regions in Europe (the European Youth Heart Study). *Am J Clin Nutr* 2004;80(3):584-90.
- 505 41. De Bourdeaudhuij I, Verloigne M, Maes L, Van Lippevelde W, Chinapaw MJM, te Velde
506 SJ, et al. Associations of physical activity and sedentary time with weight and weight
507 status among 10- to 12-year-old boys and girls in Europe: a cluster analysis within the
508 ENERGY project. *Pediatr Obes* 2013;8(5):367-75.
- 509 42. Martinez-Gomez D, Ruiz JR, Ortega FB, Veiga OL, Moliner-Urdiales D, Mauro B, et al.
510 Recommended levels of physical activity to avoid an excess of body fat in European
511 adolescents: the HELENA Study. *Am J Prev Med* 2010;39(3):203-11.
- 512 43. Nilsson A, Bo Andersen L, Ommundsen Y, Froberg K, Sardinha L, Piehl-Aulin K, et al.
513 Correlates of objectively assessed physical activity and sedentary time in children: a
514 cross-sectional study (The European Youth Heart Study). *BMC Public Health*
515 2009;9(1):322.
- 516 44. Larouche R, Faulkner GEJ, Fortier M, Tremblay MS. Active Transportation and
517 Adolescents' Health: The Canadian Health Measures Survey. *Am J Prev Med*
518 2014;46(5):507-15.

- 519 45. Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children
520 and adolescents. *Med Sci Sport Exerc* 2000;32(5):963-75.
- 521 46. Van der Horst K, Paw M, Twisk JW, Van Mechelen W. A brief review on correlates of
522 physical activity and sedentariness in youth. *Med Sci Sport Exerc* 2007;39(8):1241.
- 523 47. Hamilton MT, Healy GN, Dunstan DW, Zderic TW, Owen N. Too Little Exercise and Too
524 Much Sitting: Inactivity Physiology and the Need for New Recommendations on
525 Sedentary Behavior. *Curr Cardiovas Risk Rep* 2008;2(4):292-8.
- 526 48. Hinckson EA, McGrath L, Hopkins W, Oliver M, Badland H, Mavoa S, et al. Distance to
527 School is Associated with Sedentary Time in Children: Findings from the URBAN Study.
528 *Front Public Health* 2014;2.
- 529 49. Tudor-Locke C, Mire E, Dentre K, Barreira T, Schuna J, Zhao P, et al. A model for
530 presenting accelerometer paradata in large studies: ISCOLE. *Int J Behav Nutr Phys Act.*
531 2015;12(1):52.
- 532 50. Faulkner G, Stone M, Buliung R, Wong B, Mitra R. School travel and children's physical
533 activity: a cross-sectional study examining the influence of distance. *BMC Public Health*
534 2013;13(1):1166.
- 535 51. van Sluijs EMF, Fearne VA, Mattocks C, Riddoch C, Griffin SJ, Ness A. The contribution
536 of active travel to children's physical activity levels: Cross-sectional results from the
537 ALSPAC study. *Prev Med* 2009;48(6):519-24.
- 538 52. Panter J, Jones A, Van Sluijs E, Griffin S. The influence of distance to school on the
539 associations between active commuting and physical activity. *Pediatr Exerc Sci*
540 2011;23(1):72-86.
- 541 53. Loprinzi PD, Cardinal BJ. Measuring Children's Physical Activity and Sedentary
542 Behaviors. *J Exerc Sci Fit* 2011;9(1):15-23.
- 543 54. Hussey J, Bell C, Gormley J. The measurement of physical activity in children. *Phys*
544 *Ther Rev* 2007;12(1):52-8.

545 55. Larouche R, Oyeyemi AL, Prista A, Onywera V, Akinroye KK, Tremblay MS. A
546 systematic review of active transportation research in Africa and the psychometric
547 properties of measurement tools for children and youth. *Int J Behav Nutr Phys Act*
548 2014;11:129.

549

550

551

552

553 **Figure Legends**

554 **Figure 1.** The difference in mean school day moderate-to-vigorous physical activity (MVPA)
555 among children who engage in active school transport (AST) compared to motorized transport
556 to school in N=6 224 9-11 year old children. Means are adjusted for sex, BMI z-score, age,
557 highest parental educational attainment, accelerometer waking wear time, site by AST
558 interaction, and school nested within site was treated as a fixed effect in the multilevel analysis.
559 Error bars represent 95% confidence intervals.

560 **Figure 2.** Associations between active school transport (AST) and obtaining at least 60 minutes
561 of moderate-to-vigorous physical activity (MVPA) per school day in N=6 224 9-11 year old
562 children from 12 countries. Odds ratios are adjusted for sex, BMI z-score, age, highest parental
563 educational attainment, accelerometer waking wear time, site by AST interaction, and school
564 nested within site was treated as a fixed effect in the multilevel analysis. Error bars represent
565 95% confidence intervals.

566

567

568

569

570

Table 1. Descriptive characteristics of ISCOLE participants stratified by sex and study site (n=6 224).

Country (Site)	n	Active Transport Yes (%)	Mean (SD)						
			Age (y)	BMI (kg/m ²)	BMI z-score	MVPA ^a (min/day)	LPA ^a (min/day)	SED ^a (min/day)	Total Activity (counts/day)
Boys									
Australia (Adelaide)	220	33.0	10.8 (0.4)	18.6 (2.9)	0.6 (1.1)	79.9 (24.3)	317.6 (48.6)	465.8 (63.9)	564028.9 (132446.3)
Brazil (São Paulo)	217	39.2	10.5 (0.5)	19.9 (4.8)	1.0 (1.5)	74.1 (28.0)	347.4 (53.9)	495.4 (78.5)	536820.8 (147588.2)
Canada (Ottawa)	213	44.1	10.5 (0.4)	18.5 (3.4)	0.6 (1.2)	72.2 (20.8)	318.7 (43.6)	500.8 (66.7)	513246.8 (105426.6)
China (Tianjin)	258	31.4	9.9 (0.5)	19.8 (4.5)	1.1 (1.6)	50.9 (17.6)	307.5 (56.9)	569.1 (72.1)	401917.5 (98427.1)
Colombia (Bogotá)	422	69.4	10.5 (0.6)	17.8 (2.6)	0.3 (1.1)	79.3 (27.4)	344.4 (51.1)	502.5 (71.1)	549247.5 (138015.7)
Finland (Helsinki, Espoo & Vantaa)	218	76.2	10.5 (0.4)	17.6 (2.5)	0.3 (1.1)	85.8 (29.2)	303.0 (44.1)	528.1 (73.0)	578639.7 (156260.4)
India (Bangalore)	249	7.2	10.4 (0.5)	17.6 (3.3)	0.1 (1.5)	62.0 (20.0)	356.1 (51.5)	497.0 (64.1)	489561.4 (104436.9)
Kenya (Nairobi)	230	48.3	10.2 (0.7)	17.1 (2.8)	0.02 (1.2)	78.9 (31.1)	328.6 (56.6)	507.0 (78.6)	546932.8 (147917.5)
Portugal (Porto)	269	29.7	10.4 (0.2)	19.3 (3.3)	1.0 (1.2)	73.6 (24.6)	310.6 (50.8)	539.2 (69.3)	507698.7 (122343.3)
South Africa (Cape Town)	159	52.2	10.3 (0.7)	17.8 (3.3)	0.3 (1.3)	73.6 (25.9)	320.9 (54.0)	495.6 (72.4)	524922.9 (131996.5)
United Kingdom (Bath & NE Somerset)	185	66.0	10.9 (0.4)	18.1 (2.6)	0.4 (1.0)	76.7 (24.8)	289.7 (46.5)	496.1 (63.0)	526362.0 (131745.9)
United States (Baton Rouge)	195	8.7	10.0 (0.6)	18.5 (3.3)	0.7 (1.3)	54.1 (18.0)	318.0 (54.6)	533.9 (65.4)	439918.9 (107853.4)
All Sites	2835	43.1	10.4 (0.6)	18.4 (3.4)	0.5 (1.3)	72.0 (26.8)	324.0 (54.5)	511.8 (74.7)	515841.0 (137393.5)
Girls									
Australia (Adelaide)	258	27.9	10.7 (0.4)	19.0 (3.5)	0.6 (1.1)	59.9 (19.8)	308.4 (49.1)	489.5 (59.2)	473728.8 (119392.7)
Brazil (São Paulo)	228	41.2	10.4 (0.5)	19.7 (4.3)	0.8 (1.3)	49.6 (19.4)	336.3 (55.8)	519.2 (67.5)	418194.0 (108757.1)
Canada (Ottawa)	304	29.9	10.5 (0.4)	18.2 (3.3)	0.3 (1.2)	55.1 (18.3)	308.1 (47.2)	516.3 (63.5)	439736.4 (100951.3)
China (Tianjin)	238	34.0	9.8 (0.4)	18.0 (3.6)	0.3 (1.3)	41.6 (15.1)	281.9 (57.1)	598.8 (72.2)	345893.1 (89348.4)
Colombia (Bogotá)	434	75.6	10.4 (0.6)	17.4 (2.4)	0.1 (1.0)	62.3 (21.5)	333.7 (53.0)	526.4 (70.0)	470448.4 (111298.7)
Finland (Helsinki, Espoo & Vantaa)	252	81.0	10.5 (0.4)	17.9 (2.7)	0.2 (1.0)	66.3 (21.6)	295.9 (46.7)	541.4 (65.3)	486068.3 (131095.2)
India (Bangalore)	297	3.7	10.4 (0.5)	18.2 (3.3)	0.3 (1.3)	38.9 (14.8)	336.1 (53.5)	556.3 (67.0)	370217.0 (89399.1)
Kenya (Nairobi)	266	41.7	10.2 (0.7)	17.3 (3.3)	-0.02 (1.2)	63.0 (30.8)	329.2 (61.5)	521.1 (72.5)	481052.9 (151122.6)
Portugal (Porto)	349	27.5	10.4 (0.3)	19.4 (3.4)	0.8 (1.1)	50.5 (16.6)	301.3 (52.8)	571.3 (60.1)	399559.8 (89138.0)
South Africa (Cape Town)	241	62.7	10.2 (0.7)	18.3 (3.8)	0.4 (1.3)	57.1 (22.6)	319.2 (56.9)	515.5 (64.5)	457427.3 (123594.2)
United Kingdom (Bath & NE Somerset)	243	63.8	10.9 (0.5)	18.7 (3.2)	0.4 (1.1)	59.2 (18.7)	288.3 (47.6)	506.9 (57.3)	444690.1 (100906.2)
United States (Baton Rouge)	279	7.9	9.8 (0.5)	19.0 (4.0)	0.7 (1.3)	42.8 (15.4)	306.9 (53.8)	550.5 (64.4)	385268.2 (100745.5)
All Sites	3389	41.8	10.4 (0.6)	18.4 (3.5)	0.4 (1.2)	54.1 (21.7)	313.2 (55.8)	535.0 (71.3)	431804.9 (118704.3)

BMI: Body Mass Index; MVPA: Moderate-to-Vigorous Physical Activity; LPA: Light Physical Activity; SED: Sedentary Time

^aWeekday (Monday through Friday) average

Table 2. Adjusted mean^a minutes of weekday moderate-to-vigorous physical activity (MVPA), light physical activity (LPA), sedentary time (SED), and total activity counts by active transportation status and gender among 6 224 9-11 year old children.

Active Transportation	Boys		Girls	
	Yes	No	Yes	No
MVPA, min/day	77.1 (75.1-79.1)	70.4 (68.8-72.0)***	58.1 (56.7-59.6)	53.2 (52.0-54.3)***
Before School	8.8 (8.3-9.2)	6.1 (5.7-6.4)***	7.1 (6.8-7.4)	5.0 (4.7-5.2)***
During School	32.3 (31.4-33.2)	30.6 (29.9-31.4)*	22.0 (21.3-22.6)	21.3 (20.8-21.8)
After School	36.2 (34.9-37.6)	33.8 (32.7-34.9)*	29.0 (28.0-30.0)	26.8 (26.1-27.6)**
LPA, min/day	320.7 (316.6-324.9)	326.3 (323.0-329.7)*	310.7 (306.6-314.8)	319.0 (316.0-322.0)**
Before School	35.9 (35.1-36.7)	38.5 (37.9-39.2)***	36.1 (35.3-36.8)	38.8 (38.2-39.4)***
During School	138.6 (136.3-140.9)	140.2 (138.4-142.1)	125.4 (123.1-127.6)	128.5 (126.8-130.1)*
After School	147.9 (146.1-149.7)	147.5 (145.6-149.3)	148.9 (146.7-151.2)	151.7 (150.1-153.4)*
SED, min/day	509.8 (504.7-515.0)	510.9 (506.8-515.1)	533.5 (528.7-538.3)	530.2 (526.7-533.7)
Before School	54.7 (53.7-55.7)	54.8 (54.0-55.6)	56.6 (55.7-57.5)	56.0 (55.3-56.6)
During School	223.7 (221.0-226.5)	223.8 (221.6-226.0)	242.1 (239.5-244.6)	239.6 (237.8-241.5)
After School	230.7 (227.7-233.7)	231.9 (229.5-234.3)	235.3 (232.5-238.1)	234.7 (232.6-236.8)
Total Activity, counts/day	536816 (526375-547258)	513420 (504991-521849)**	449700 (441270-458131)	433500 (427275-439725)*
Before School	58551 (56519-60583)	49463 (47815-51110)***	51308 (49748-52868)	44619 (43465-45773)***
During School	226377 (221540-231214)	220347 (216433-224261)*	173113 (169291-176934)	171169 (168340-173997)
After School	253150 (246040-260260)	244356 (238618-250094)*	224823 (218999-230647)	217264 (212953-221576)*

MVPA: Moderate-to-Vigorous Physical Activity; LPA: Light Physical Activity; SED: Sedentary Time

^aMulti-level models with site and school nested within site treated as fixed effects; adjusted for age, BMI z-score, highest parental education level, mean accelerometer wear time, and site by active transportation interaction.

Data shown as adjusted mean (95% confidence interval).

***p<0.0001; **p<0.001; *p<0.05

Figure 1. The difference in mean school day moderate-to-vigorous physical activity (MVPA) among children who engage in active school transport (AST) compared to motorized transport to school in N=6 224 9-11 year old children. Means are adjusted for sex, BMI z-score, age, highest parental educational attainment, accelerometer waking wear time, site by AST interaction, and school nested within site was treated as a fixed effect in the multilevel analysis. Error bars represent 95% confidence intervals.




Figure 2. Associations between active school transport (AST) and obtaining at least 60 minutes of moderate-to-vigorous physical activity (MVPA) per school day in N=6 224 9-11 year old children from 12 countries. Odds ratios are adjusted for sex, BMI z-score, age, highest parental educational attainment, accelerometer waking wear time, site by AST interaction, and school nested within site was treated as a fixed effect in the multilevel analysis. Error bars represent 95% confidence intervals.

