



Citation for published version:

González-Sánchez, MI, Lee, PT, Guy, RH & Compton, RG 2015, 'In situ detection of salicylate in *Ocimum basilicum* plant leaves via reverse iontophoresis', *Chemical Communications*, vol. 51, no. 92, pp. 16534-16536. <https://doi.org/10.1039/c5cc06909b>

DOI:

[10.1039/c5cc06909b](https://doi.org/10.1039/c5cc06909b)

Publication date:

2015

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

In situ detection of salicylate in *Ocimum Basilicum* plant leaves via reverse iontophoresis

Received 00th January 20xx,
Accepted 00th January 20xx

M. I. Gonzalez-Sanchez^{a,b}, P. T. Lee^a, R. H. Guy^{a,c} and R. G. Compton^{a*}

DOI: 10.1039/x0xx00000x

www.rsc.org/

The quantitative analysis of salicylate provides useful information for the evaluation of metabolic processes in plants. We report a simple, noninvasive method to measure salicylate *in situ* in *Ocimum Basilicum* leaves using reverse iontophoresis in combination with cyclic voltammetry at disposable screen-printed electrodes and the concentration of salicylate in basil leaves was found to be 3 mM.

Salicylates are widely distributed phytohormones in plants, present either as the free acid (SA) (Figure 1) or as its glycosolated or methylated conjugate forms¹. Salicylates participate in plant growth^{2,3}, development^{2,3}, immunity⁴, thermogenesis^{3,5}, flowering⁵, and germination⁶. It follows that the amount of salicylate present in a plant is a reflection of its metabolic and physiological activity, and is responsive, therefore, to biotic and abiotic stress⁵ (for example, to insect feeding⁷, pathogen infection⁵ or environmental changes⁸). The level of endogenous salicylates under non-stress conditions is highly dependent on the specific plant species of interest⁸, ranging from ng to mg per gram of fresh material⁹; as stated, these quantities can increase substantially increase under biotic and abiotic stress⁵.




Figure 1. Chemical structure of SA

^a Department of Chemistry, Physical and Theoretical Chemistry Laboratory, Oxford University, South Parks Road, Oxford, OX1 3QZ, United Kingdom.

^b Department of Physical Chemistry, Castilla-La Mancha University, 02071, Albacete, Spain.

^c Department of Pharmacy & Pharmacology, University of Bath, Bath, BA2 7AY, United Kingdom.

*Corresponding Author: richard.compton@chem.ox.ac.uk

† Electronic Supplementary Information (ESI) available: Experimental materials and procedures. Analytical methods to measure salicylates in leaf extracts. *In vivo* methods to quantify plant metabolites. See DOI: 10.1039/x0xx00000x

Analysis of salicylates in leaf extracts has been reported using chromatography^{10,11}, fluorescence⁶ and electrochemical methods¹² (see Table S1, ESI). While these approaches are quantitative and capable of assessing many samples¹³, sample preparation typically requires plant tissue pulverisation in liquid nitrogen followed by exhaustive extraction with non-aqueous solvents and is non-trivial, expensive, and time-consuming. More relevant, of course, for the study of oxidative stress *in situ* in a relevant agricultural setting would be an *in vivo* determination of plant metabolites in intact leaves^{14,15} (see Table S2, ESI). With respect to salicylate quantification in intact plants, only a few examples have been reported using methods (Table 1) such as colorimetry¹³ and bioluminescence¹⁶. The first *in situ* electrochemical analysis of SA in tomato leaves was performed using a paper-based multi-walled carbon nanotube (MWCNT)/Nafion modified carbon tape electrode¹⁷; however, the measurement required an invasive hole to be punched into the leaf so as to allow SA levels to be determined.

Table 1. Methods to measure SA in plants *in situ*.

Method	Sensor	SA quantity	Plant	Ref
Colorimetric	Titanium dioxide nanoparticles	NA ¹	Tobacco	13
Fluorescence	Acinetobacter sp ADP1	NA ²	Tobacco	16
Electrochemical	MWCNTs/Nafion modified carbon tape electrode	4.2 ng per leaf	Tomato	17

NA: not available.

¹ *In situ* SA labelling prior to its quantification in extracts.

² SA detection after inoculation.

Iontophoresis^{18,19} is a technique that has been primarily used to increase the transport of (primarily) ionised compounds across biological barriers, such as the skin²⁰. The method involves the application, via appropriately positioned (typically reversible Ag/AgCl) electrodes, of a small electric current (≤ 0.05 mA/cm²) that promotes transmembrane transport by electromigration and/or electroosmosis.

Although most often used for transdermal drug delivery²¹, so-called reverse iontophoresis, has been studied as a tool for extraction purposes as well, in particular for drug and glucose monitoring^{20, 22, 23, 24}. In terms of applications in plants, the iontophoretic release of alkaloids from cell suspensions (followed by chromatographic analysis) has been investigated²⁵. However, this process has been never applied on a plant living tissue and constitutes a very promising non-destructive way to extract small plant metabolites.

The aim of this work was to apply iontophoresis coupled with electroanalysis at disposable carbon nanotube screen-printed electrodes to determine the amount of salicylate in intact *Ocimum Basilicum* leaves. Such an approach, to our knowledge, has never been performed on a living plant. Electrochemical analysis using cyclic voltammetry is well suited for this application, the method being cheap, technically straightforward and having good sensitivity. Although electrochemical detection *per se* has used before with several plant species^{17, 26}, the need for a complex pre-treatment protocol to release the target analyte from the sample is clearly disadvantageous as mentioned above. In contrast, the *in situ* method described here to determine salicylate in the leaves of an intact, living plant is facile and non-destructive (Figure 2). The procedure employs a glass iontophoresis cell¹⁸ clipped onto an undamaged and untreated leaf. Then a period of current passage was used to extract materials. Subsequently, cyclic voltammetry on multi-walled carbon nanotube screen printed electrodes (MWCNT-SPE) were used to determine salicylate via its oxidation to the carboxyl-hydroquinone²⁷.




Figure 2. Experimental set-up for the iontophoretic extraction of salicylate from an *Ocimum Basilicum* leaf; (A) real-life image, (B) schematic representation.

Iontophoresis at 0.5 mA applied to the basil leaf for 8 hours, followed by cyclic voltammetry on the solutions present in the electrode chambers, clearly showed that, at pH 7, where salicylic acid ($pK_a = 2.97$ for the carboxylic acid functional group²⁷) is essentially fully and negatively ionised, is extracted to the anode (Figure 3, left panel). The cyclic voltammograms on the iontophoretically-extracted solution in the anode has an obvious oxidation peak at about +0.7 V (vs. Ag), which was absent when no iontophoretic current was applied (i.e., any passive diffusion contribution can be considered negligible).

To verify that the oxidation peak was indeed due to salicylate, the reverse iontophoresis experiment was repeated under identical conditions except that the cathode chamber contained a 10 mM salicylate solution in pH 7.0 phosphate buffer (as opposed to just buffer alone). At the end of 8-hours of current passage, cyclic voltammetry was performed on the anode solution and the results (Figure 3, right panel) shows that the +0.7 V (vs. Ag) oxidation peak was significantly greater. The interpretation of this finding is that salicylate extraction from the leaf is now being augmented by the analyte, which is delivered from the cathode chamber through the leaf into the lower compartment, and then extracted back (through the leaf again) into the anode chamber.




Figure 3. Left panel: Cyclic voltammogram (scan rate = 50 mVs⁻¹) at MWCNT-SPE of the iontophoretically-extracted anode chamber solution after 8-hours with (solid line) and without (dotted line) application of current at 0.5 mA. Right panel: Cyclic voltammograms of the anode solution after 8h of current application in the absence (black) and in the presence (red) of 10 mM SA in the cathode compartment. Scan rate as again 50 mVs⁻¹.

To confirm that an 8-hour extraction period was appropriate for further development of this proof-of-concept study, the reverse iontophoretic extraction of endogenous salicylate (i.e., with no added material to the cathode chamber) followed by cyclic voltammetric detection was repeated following 9 and 10 hours of current passage (Figure 4). It is evident that there is no significant difference between the amounts of salicylate extracted in 8, 9 or 10 hours of iontophoresis (1-way analysis of variance, $\alpha > 0.05$) and that it is apparent that an 8-hour application of 0.5 mA is sufficient to extract essentially all of the analyte in the leaf.




Figure 4. Average salicylate peak current (error bars = standard deviation; $n = 3$) detected by cyclic voltammetry of the anode solution after iontophoresis for different times, with no exogenous addition of salicylate.

Finally, to estimate quantitatively the concentration of salicylate in the leaf, a further series of 8-hour iontophoresis experiments was conducted with different salicylate concentrations (1, 3, 7 and 10 mM) in the cathode chamber, essentially an 'internal' standard addition. Once again, at the end of current passage, cyclic voltammetry was used to detect the analyte in the anode compartment of the iontophoresis cell. The results in Figure 5 show that the salicylate peak current ($I_p / \mu\text{A}$) increases linearly with the added salicylate concentration in the cathode chamber ([SA] mM) with a relationship of $I_p = (0.201 \pm 0.023) + (0.065 \pm 0.005) \times [\text{SA}]$.




Figure 5. Peak oxidation current (mean \pm SD; $n = 3$) of salicylate measured in the anode compartment solution as a function of the salicylate concentration added to the cathode chamber.

Compared to the value of I_p when no salicylate was introduced into the cathode (i.e., about 0.2 μA), it can be seen that the peak current was approximately doubled when the cathode concentration of salicylate was *ca.* (3.1 \pm 0.12) mM. It follows, therefore, that the salicylate concentration in the leaf is approximately 3.0 mM. Although this deduction assumes that the inner and outer surfaces of the leaf offer comparable resistances to salicylate electrotransport, and recognising that there is an additional resistance offered to the salicylate that penetrates from the cathodal chamber. The concentration is likely to reflect the composition of fluid (sap) in the leaf and is significantly different from the composition determined by analysis of the total leaf content, reported in the literature (see Table S1, ESI) as wt/wt.

The successful extraction and proof-of-concept monitoring of endogenous salicylate from the leaves of *Ocimum Basilicum in situ* has been achieved by reverse iontophoresis coupled with cyclic voltammetry. The results of these experiments demonstrate that the concentration of salicylate in the leaf is *ca.* (3.1 \pm 0.12) mM. The method is simple, non-destructive, relatively fast and does not require independent calibration of

any sort. The approach has potential, therefore, as a monitor of plant metabolism, its growth and developmental processes, and its response to a variety of external stressors.

This work was supported in part by the Spanish Government (BFU2013-44095-P, cofounded with FEDER funds, EU). MIGS would like to thank to the Regional Ministry of Education and Science of Castilla-La Mancha for her postdoctoral grant.

Notes and references

1. H. I. Lee, J. León and I. Raskin, *Proc. Natl. Acad. Sci.*, 1995, **92**, 4076-4079.
2. M. Rivas-San Vicente and J. Plasencia, *J. Exp. Bot.*, 2011, **62**, 3321-3338.
3. D. M. A. Dempsey, A. C. Vlot, M. C. Wildermuth and D. F. Klessig, *Arabidopsis*, 2011, **9**, e0156.
4. C. An and Z. Mou, *J. Integr. Plant Biol.*, 2011, **53**, 412-428.
5. C. A. Vlot, A. D. D'Maris and D. F. Klessing, *Annu. Rev. Phytopathol.*, 2009, **47**, 177-206.
6. C. T. DeFraia, E. A. Schmelz and Z. Mou, *Plant Methods*, 2008, **4**, 28-38.
7. R. P. Van Poecke, M. Posthumus and M. Dicke, *J. Chem. Ecol.*, 2001, **27**, 1911-1928.
8. L. V. Kurepin, K. P. Dahal, M. Zaman and R. P. Pharis, *Salicylic Acid*, S. Hayat, A. Ahmad and M. N. Alyemeni, Springer Netherlands, 1st ed., 2013, ch. 5, p. 61-82.
9. H. Bandurska, *Salicylic Acid*, S. Hayat, A. Ahmad and M. N. Alyemeni, Springer Netherlands, 2013, ch. 1, p. 1-14.
10. S. M. Wilbert, L. H. Ericsson and M. P. Gordon, *Anal. Biochem.*, 1998, **257**, 186-194.
11. X. Pan, R. Welti and X. Wang, *Nat. Protocols*, 2010, **5**, 986-992.
12. Z. Wang, F. Wei, S.-Y. Liu, Q. Xu, J.-Y. Huang, X.-Y. Dong, J.-H. Yu, Q. Yang, Y.-D. Zhao and H. Chen, *Talanta*, 2010, **80**, 1277-1281.
13. P.-J. Tseng, C.-Y. Wang, T.-Y. Huang, Y.-Y. Chuang, S.-F. Fu and Y.-W. Lin, *Anal. Methods*, 2014, **6**, 1759-1765.
14. Q.-Q. Ren, X.-J. Yuan, X.-R. Huang, W. Wen, Y.-D. Zhao and W. Chen, *Biosens. Bioelectron.*, 2013, **50**, 318-324.
15. L. Zheng, J. Zhou, B. Li and D. Xing, *Front. Recent Dev. Plant Sci.*, 2015, **6**, 1-8.
16. W. E. Huang, L. Huang, G. M. Preston, M. Naylor, J. P. Carr, Y. Li, A. C. Singer, A. S. Whiteley and H. Wang, *Plant J.*, 2006, **46**, 1073-1083.
17. L.-J. Sun, Q.-M. Feng, Y.-F. Yan, Z.-Q. Pan, X.-H. Li, F.-M. Song, H. Yang, J.-J. Xu, N. Bao and H.-Y. Gu, *Biosens. Bioelectron.*, 2014, **60**, 154-160.
18. P. Glikfeld, C. Cullander, R. S. Hinz and R. H. Guy, *Pharm. Res.*, 1988, **5**, 443-446.
19. P. Glikfeld, R. S. Hinz and R. H. Guy, *Pharm. Res.*, 1989, **6**, 988-990.
20. B. Leboulanger, R. H. Guy and M. B. Delgado-Charro, *Physiol. Meas.*, 2004, **25**, R35-R50.
21. A. Djabri, R. H. Guy and M. B. Delgado-Charro, *Int. J. Pharm.*, 2012, **435**, 27-32.
22. M. B. Delgado-Charro and R. H. Guy, *Pharm. Res.*, 2003, **20**, 1508-1513.

23. G. Rao, P. Glikfeld and R. Guy, *Pharm. Res.*, 1993, **10**, 1751-1755.
24. G. Rao, R. H. Guy, P. Glikfeld, W. R. LaCourse, L. Leung, J. Tamada, R. O. Potts and N. Azimi, *Pharm. Res.*, 1995, **12**, 1869-1873.
25. H. T. Pu, R. Y. K. Yang and F. L. Saus, *Biotechnol. Lett.*, 1989, **11**, 83-86.
26. M. I. Gonzalez-Sanchez, L. Gonzalez-Macia, M. T. Perez-Prior, E. Valero, J. Hancock and A. J. Killard, *Plant, Cell Environ.*, 2013, **36**, 869-878.
27. W.-D. Zhang, B. Xu, Y.-X. Hong, Y.-X. Yu, J.-S. Ye and J.-Q. Zhang, *J. Solid State Electrochem.*, 2010, **14**, 1713-1718.