



*Citation for published version:*

Moschos, SA, Usher, L & Lindsay, MA 2017, 'Clinical potential of oligonucleotide-based therapeutics in the respiratory system', *Pharmacology and Therapeutics*, vol. 169, pp. 83-103.  
<https://doi.org/10.1016/j.pharmthera.2016.10.009>

*DOI:*

[10.1016/j.pharmthera.2016.10.009](https://doi.org/10.1016/j.pharmthera.2016.10.009)

*Publication date:*

2017

*Document Version*

Peer reviewed version

[Link to publication](#)

## University of Bath

### Alternative formats

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **Clinical potential of oligonucleotide-based therapeutics in the respiratory system**

Sterghios A. Moschos<sup>1\*</sup>, Louise Usher<sup>1</sup>, and Mark A. Lindsay<sup>2</sup>

<sup>1</sup>Department of Biomedical Sciences, Faculty of Science and Technology, University of Westminster, 115 New Cavendish Str., London W1W 6UW, United Kingdom and

<sup>2</sup>Department of Pharmacy and Pharmacology, Claverton Down, University of Bath, Bath, BA2 7AY, United Kingdom

\* Corresponding Author. Dr Sterghios A. Moschos. Department of Biomedical Sciences, Faculty of Science and Technology, University of Westminster, 115 New Cavendish Str., London W1W 6UW, UK. +44 (0) 207 911 5000, extension 64153.

[s.moschos@westminster.ac.uk](mailto:s.moschos@westminster.ac.uk)

The discovery of an ever-expanding plethora of coding and non-coding RNAs with nodal and causal roles in the regulation of lung physiology and disease is reinvigorating interest in the clinical utility of the oligonucleotide therapeutic class. This is strongly supported through recent advances in nucleic acids chemistry, synthetic oligonucleotide delivery and viral gene therapy that have succeeded in bringing to market at least three nucleic acid-based drugs. As a consequence, multiple new candidates such as RNA interference modulators, antisense, and splice switching compounds are now progressing through clinical evaluation. Here, manipulation of RNA for the treatment of lung disease is explored, with emphasis on robust pharmacological evidence aligned to the five pillars of drug development: exposure to the appropriate tissue, binding to the desired molecular target, evidence of the expected mode of action, activity in the relevant patient population and commercially viable value proposition.

### **Abbreviations**

AGO, Argonaute; ASO, antisense oligonucleotide; CCR3, C-C chemokine receptor 3; CPP, cell penetrating peptide; COPD, chronic obstructive pulmonary disease; CD, cluster designation; GM-CSF, granulocyte monocyte – colony stimulating factor; IL, interleukin; IL-‘X’R, Interleukin receptor number ‘X’; LNA, locked nucleic acid; miRNA, microRNA; ncRNA, non-coding RNA microRNA; nucleotide, nt; PNA, peptide nucleic acid; PMO, phosphorodiamidate morpholinos; RISC, RNA induced silencing complex; RNAi, RNA interference; RNA-Seq; next generation sequencing of RNA; PPMO, peptide-conjugated phosphorodiamidate morpholinos; siRNA, short (or small) interfering RNA; TLR, toll-like receptor; UTR, untranslated region;  $\beta$ c, common beta chain.

**Keywords**

Oligonucleotide therapeutics, siRNA, miRNA, PNA, PPMO, delivery.

## Table of Contents

<b>1. Introduction .....</b>	<b>5</b>
<b>Modulating RNA homeostasis with oligonucleotide-based therapeutics.....</b>	<b>10</b>
<b>Development of antisense oligonucleotide therapeutics .....</b>	<b>11</b>
<b>Development of siRNA-based therapeutics .....</b>	<b>18</b>
<b>Clinical progress of oligonucleotide therapeutics for lung diseases .....</b>	<b>26</b>
<b>RNAi mediators: ALN-RSV01 and the treatment of RSV infection.....</b>	<b>27</b>
<b>2<sup>nd</sup> Generation ASO therapeutics: Clinical Status.....</b>	<b>30</b>

Commented [SM1]: Ignore this, needs rebuilding

## 1. Introduction

Pharmacologic intervention commonly involves the administration of synthetic small molecules, recombinant proteins or antibody-based therapeutics that target or mimic the action of proteins. However, these approaches can only be employed to target certain classes of proteins (i.e. extracellular protein in the case of antibodies). This has led to emerging interest in targeting proteins at the mRNA level using oligonucleotide-based therapeutics such as antisense and short (or small) interference RNA (siRNA). Since the actions of oligonucleotide therapeutics are mediated through Watson-Crick binding, this approach has the advantage of being able to target all known mRNAs, as well regulatory non-coding RNAs (ncRNA) such as microRNAs (miRNA) (de Hoon, Shin, & Carninci, 2015; Djebali et al., 2012; Kawai et al., 2001). However, in developing these oligonucleotide-based therapeutics, a number of major problems have emerged including poor stability in biological fluids, potential to induce immune response and off-target actions. However, the principal problem is delivery of these large (often > 6 kDa), negatively charged (resulting from the sugar-phosphate backbone) molecules both into the target tissue and then across the plasma membrane (also negatively charged) and into the cell, where they mediate their biological action. Significantly, the delivery problems have meant that much of the effort in this area has focused upon liver disease based on the observation that this organ readily absorbs oligonucleotides following intravenous (IV) administration (Nicklin et al., 1998). Thus, Ionis (previously known as ISIS) Pharmaceuticals has recently obtained regulatory approval in the USA for mipomersan, an antisense oligonucleotide (ASO) that targets apolipoprotein B for the treatment of familial hypercholesterolemia (Santos, Raal, Donovan, & Cromwell, 2015). Nevertheless, the large surface area of the airways and their

accessibility to topical delivery suggests that the respiratory system might also represent a potential target tissue for oligonucleotide-based therapeutics.

### **1.1. Anatomy and physiology of the respiratory system**

Lung architecture (Hasleton, 1972; Lambert, Wilson, Hyatt, & Rodarte, 1982; Lambert, 1989; R. J. Lorenz, 1966) is commonly visualized as a bunch-of-grapes (Mauroy et al., 2015) formed by 23 serial bifurcations from the trachea to the last alveolar duct (generation 22; Fig. 1A) (R. J. Lorenz, 1966). Surface area and volume in adults ranges from 24 - 69 m<sup>2</sup> and 2.16 - 5.23 l, respectively, varying significantly on account of age, height and disease (Hasleton, 1972; Labiris & Dolovich, 2003; Usmani, 2014). Of the ~3 l standard internal lung volume (Hasleton, 1972), 85-90% is occupied by alveoli (lung parenchyma or interstitium) (Ochs, 2014).

One of the key physiological barriers to the delivery of oligonucleotides is the thin layer of pulmonary surfactant and mucus that line lower and upper airways, respectively. The cationic and lipoproteinaceous matrix that constitutes the pulmonary surfactant of the lower airways and alveoli has a number of biological functions including mechanostructural stabilization (Parra & Pérez-Gil, 2015), surface tension reduction to enable gas exchange (Whitsett, Wert, & Weaver, 2015) and as a contributor to immune protection (Han & Mallampalli, 2015). Many of the surfactant lipid components have been successfully synthesized and are used in liposomal and lipid nanoparticle drug formulation. Surfactant proteins (SP) are principally derived from the *SPA*, *SPB*, *SPC* and *SPD* genes, expressed by type II alveolar epithelial cells. These, along with alveolar macrophages, also recycle

surfactant and together participate in innate immune responses to extrinsic inflammatory stimuli of viral (Derscheid & Ackermann, 2013; Hillaire, Haagsman, Osterhaus, Rimmelzwaan, & van Eijk, 2013), bacterial (Chronos, Sever-Chroneos, & Shepherd, 2010; Sender & Stamme, 2014; Whitsett, 2010), fungal (Faro-Trindade et al., 2012; Ledford, Addison, Foster, & Que, 2014; Singh et al., 2015; van de Wetering et al., 2004), parasitic (Blanco, Lugones, Díaz, & Monzote) and particulate nature (Arick, Choi, Kim, & Won, 2015; Ma et al., 2015; Vattanasit et al., 2014). Alveolar macrophages also clear extraneous matter, which may or may not elicit innate and adaptive immune responses (Forbes et al., 2014).

Surfactant genes are also expressed in the larger airways (Whitsett et al., 2015) and even the nose (Gaunsbaek, Kjeldsen, Svane-Knudsen, Henriksen, & Hansen, 2014). However, the upper airways are mainly lined with mucous - a highly complex, anionic, glycoproteinaceous and gelatinous extracellular matrix (S. K. Lai, Wang, Wirtz, & Hanes, 2009) synthesized by goblet cells and submucosal glands that express *MUC5AC* and *MUC5B* and are located throughout the bronchi. The predominant structural component of mucous is large mucin polymer fibers of <10 nm size, that interact to produce a complex physicochemical organization (Shogren, Gerken, & Jentoft, 1989). Its role is the capture and elimination of large particles reaching the upper airways through inhalation. The underlying epithelium contributes to immune surveillance through both immune dampening and pathogen associated molecular pattern recognition systems (Davies, 2014). Removal of particulates is mediated by mucociliary clearance and episodic bronchial smooth muscle contraction, i.e. cough. Materials are propelled into the trachea and then to the gastrointestinal tract for digestion. This mechanobiological phenomenon relies on the microrheology of mucous, which is more elastic in the nasopharynx and distal airways but less so in the intervening



bronchi, and is altered by disease (e.g. increased viscoelasticity due to reduced hydration) (Rubin, 2014). These changes can alter significantly mucociliary clearance rates, bacterial cell interactions with the host, particle motion within the mucous matrix and, consequently, drug delivery mechanics (S. K. Lai et al., 2009).

Respiratory disease is associated with multiple changes in the airways and lungs that can impact upon the delivery and action of oligonucleotide therapeutics. In fibrotic disease thickening of the alveoli reduces the gas diffusion barrier, in many cases preceded or accompanied by surfactant dysfunction. In chronic obstructive pulmonary disease (COPD), alveolar sac volume increases on account of loss of the type II epithelia, in turn reducing the surface area available for gas exchange (Ochs, 2014). Asthmatic patients experience dyspnea as principally airborne allergens (Hurwitz, 1955; Johansson et al., 2008; Romanet-Manent, Charpin, Magnan, Lanteaume, & Vervloet, 2002) trigger constriction of the upper airways (Noble et al., 2014) alongside inflammatory/allergic responses (Erle & Sheppard, 2014), mucous overproduction/increased viscoelasticity (S. K. Lai et al., 2009) and conducting airway/bronchial wall thickening (Olin & Wechsler, 2014). Together, these phenomena reduce the bronchial diameter during an asthmatic attack that can eventually lead to death (Dowell, Lavoie, Solway, & Krishnan, 2014). In cystic fibrosis (CF), genetic mutations (Bombieri et al., 2011) produce solute imbalances in the airways that contribute to mucous hypersecretion, altered mucous motility and dysregulated immune homeostasis, leading to opportunistic infection.

All these indications are associated with exaggerated or disrupted innate and adaptive immune responses. Indeed, recent attention has shifted onto the role of respiratory

epithelia given their importance in inflammation and the exacerbations associated with many chronic lung disease following exposure to inhaled pathogens (Zuo, Lucas, Fortuna, Chuang, & Best, 2015). Interestingly, studies have also indicated that the lung is not sterile, as commonly believed, but populated by numerous microbes (Riiser, 2015). Although sampling approaches and analytical methodologies confound this research (Salter et al., 2014), reports of pulmonary microbiome imbalances have emerged. Whether these are causal or consequential remains to be determined, however, their relevance to immune plasticity during treatment is of therapeutic interest, especially where novel therapeutics may interact with, or affect immune status.

The physiological, mechanistic and molecular differences between pathologies, coupled to the functional differences between species and organizational complexity of the tissue are such that *in vitro* 2D/3D/co-culture system and animal model use is limited, or at least challenging in their translational value (Nichols et al., 2014; Ochs, 2014; J. C. Parker & Townsley, 2008; Reus et al., 2014; Williams & Roman, 2015). This is especially the case when such approaches are used in isolation (Saturni, Contoli, Spanevello, & Papi, 2015), or without consideration of systems-level elements (Dittmar, McIver, Michalak, Garner, & Valdez, 2014; Seok et al., 2013; Takao & Miyakawa, 2015; Tsitsiou et al., 2012).

## **1.2 Pharmacological targeting of selective regions within the respiratory system**

Airway dimensions adhere to fractal branching models (Horsfield, 1990), information essential in engineering aerosolized and dry powder systems that target drugs to the specific location within the airways and lung by inhalation (Denyer & Dyche, 2010; Laube,

2014). Thus, by controlling nebulized solution or dry powder particle size, appropriate deposition targeting can be achieved across the pulmonary tract, with minimal drug loss to the oral/nasal cavity or the environment (Fig. 1B). Drug formulations are presently evaluated for their deposition mechanics through optical particle sizing, time-of-flight spectrometry, cascade impaction and liquid impinger technologies (Denyer & Dyché, 2010; Mitchell, Bauer, Lyapustina, Tougas, & Glaab, 2011; Pu, Kline, Khawaja, Van Liew, & Berry, 2015; Zhu, Haghi, Goud, Young, & Traini, 2015).

## **2. Modulating RNA homeostasis with oligonucleotide-based therapeutics**

A number of classes of oligonucleotide-based therapeutics have been developed although these can be principally divided into i) antisense and ii) RNA interference (RNAi)-based approaches. The first are single-stranded antisense oligonucleotides (ASO) 20-30 nucleotides (nt) in length that can either catalyse target cleavage (via RNase H) or stoichiometrically sequester RNA targets (Fig. 2A, B). Currently, this approach has proved the most effective, with two approved ASOs that are RNase H-active (fomivirsen: approved in 1998; mipomersen: approved in 2012). Antisense-based approaches have also been developed for RNA splicing manipulation, the closest to market being drisapersen, a compound in late stage developed for the treatment of Duchenne muscular dystrophy (Fig. 2C; drisapersen, Biomarin Pharmaceutical (Disterer et al., 2014; Q.-L. Lu, Cirak, & Partridge, 2014; van Deutekom et al., 2007)). Single stranded oligonucleotides in the form of aptamers can also be used as replacements to antibodies. A marketed example is pegaptanib which targets extracellular VEGF and is licensed for use in wet macular degeneration (approved in 2004) (Sundaram, Kurniawan, Byrne, & Wower, 2013). Following the identification of siRNA-

mediated RNAi as a mechanism capable of regulating gene expression at the level of transcription and/or translation, there has also been considerable effort to develop these double stranded RNAs (dsRNA) as potential therapeutic modalities (Fig. 2D) (Moschos, 2013; Sabin, Delás, & Hannon, 2013).

The biggest challenges in the clinical progression of both antisense and RNAi-active drugs are their poor targeting to disease-relevant tissue sites and their inefficient transportation across the plasma membrane into the cytosol, where the pharmacological targets of these drugs reside. In addition, it has been necessary to address a number of other issues including stability, off-target action and immune activation.

## **2.1 Development of antisense oligonucleotide therapeutics**

To simultaneously address the problems of antisense stability and delivery, much of the effort has focused upon the development of modified synthetic oligonucleotides (Bennett, Chiang, Chan, Shoemaker, & Mirabelli, 1992). Specifically, emphasis has been placed on development of chemistries impervious to nuclease degradation, with concomitantly enhanced pharmacokinetic properties, better cellular uptake, improved target affinity, and tailored immunogenicity profiles.

### **2.1.1 First Generation Antisense**

Early studies on RNase H-active ASO evidenced transfection reagents as necessary to elicit appreciable function *in vitro* (Bennett et al., 1992), otherwise micromolar concentrations achieved internalization over a 50 h period through a saturable, cytoplasmic membrane

protein pathway (Loke et al., 1989). The main site of ASO activity was shown to be the nucleus (P L Iversen, Zhu, Meyer, & Zon, 1992). *In vivo* activity, however, was negligible due to the poor stability against nucleases.

The first attempts to resolve this problem involved the use of phosphorothioate backbone modifications, which imparted increased stability in biofluids and improved pharmacokinetics *in vivo* (Fig. 3A, Fig. 4). However, toxicity was observed due to interactions with heparin-binding growth factors and other compounds (Benimetskaya et al., 1997; Fennewald & Rando, 1995; Guvakova, Yakubov, Vlodayevsky, Tonkinson, & Stein, 1995). It was not until many years later that serum albumin was identified as the principle (T. A. Watanabe, Geary, & Levin, 2006), but not the only carrier of phosphorothioate ASO in circulation and into the liver (Bijsterbosch et al., 2000). Uptake is presently understood to involve at least two (Koller et al., 2011), if not more endocytic mechanisms (Juliano, Ming, & Nakagawa, 2012) although this mechanism of action (MAO) is thought to account for only 20% of uptake (Geary et al., 2009). Of note, these antisense were also found to be TLR9 agonists (Krieg et al., 1995; Rutz et al., 2004) (Table 1) and demonstrated a propensity for complement activation (Advani et al., 2005; Henry et al., 1997)

### **2.1.2 Second Generation Antisense**

The second generation of antisense involved a cadre of chemical modifications of the 2' ribose position (Fig. 3B, 4B; mixomers), or the development of entirely novel backbone structures resistant to nuclease activity, such as phosphorodiamidate morpholinos (PMO) and peptide nucleic acids (PNA; Fig. 3A). In many cases, although these modifications greatly increased the binding affinity of the ASO for its target RNA, this often coincided with the loss

of RNaseH inducing activity, as RNase H cleaves 8-12 bases from the 3' end of the ASO (Cerritelli & Crouch, 2009; Crooke, 1999; Vickers & Crooke, 2015; H. Wu, Lima, & Crooke, 1999).

Lack of RNase H induction, however, can be advantageous where the mechanism of action does not require target cleavage, such as exon modulation (splice switching or splice correction) therapy. Interestingly, some mutations in CF patients are indeed amenable to splice-switching therapy as evidenced *in vitro*. Thus, correct exon splicing has been reported in at least two separate studies modeling distinct genetic backgrounds involving aberrant splicing of *CFTR* (Friedman et al., 1999; Igreja, Clarke, Botelho, Marques, & Amaral, 2015). The therapeutic value of this approach is strongly supported by complementary studies which involve the overexpression of splicing factors as opposed to use of splice correction oligonucleotides (Nissim-Rafinia et al., 2004). There is therefore scope for expanding the evidence on the mechanistic utility of splice switching therapies in CF animal models, provided commercially appealing delivery solutions are pursued.

To address the mixomer issue around loss of RNase H activity in 2<sup>nd</sup> generation ASO attempts were made to produce so-called gapmer oligonucleotides (Fig. 4B). These were organized to feature unmodified 8-12 nt DNA sequences at the centre, flanked by modified nucleosides at the 3' and 5' ends. (Cerritelli & Crouch, 2009; Crooke, 1999; Vickers & Crooke, 2015; H. Wu et al., 1999). A number of these 2<sup>nd</sup> generation ASO have been advanced clinically with mixed outcomes (section 3).

PMOs are claimed to have higher affinity and a reduced propensity for off-target RNA binding (J. E. Summerton, 2007), with flagship programs in phase 2 and 3 clinical studies for

the treatment of Duchenne's muscular dystrophy through splicing modulation (eteplirsen (AVI-4658) and SRP-4053; Sarepta Therapeutics). Whilst preclinical data corroborated increased affinity (Q. L. Lu et al., 2005; Tanganyika-de Winter et al., 2012; B. Wu et al., 2010), including in the lung (Q. L. Lu et al., 2005; B. Wu et al., 2010), the claims of increased specificity have been challenged by others exploring off-target binding in depth (Eisen & Smith, 2008; Schulte-Merker & Stainier, 2014). Furthermore, although PMO alone showed delivery *in vitro* and *in vivo*, this can be substantially improved through conjugation with synthetic cell penetrating peptides (CPP) (Moulton et al., 2007) (peptide-conjugated phosphorodiamidate morpholinos (PPMO), or 2<sup>nd</sup> generation PMO's).

PPMOs have been recently suggested to form <90 nm, negatively charged micelles under physiologically relevant conditions (Ezzat et al., 2015), leading to cellular uptake through scavenger receptors (Ezzat et al., 2012, 2015). However, the micromolar critical micelle concentration reported by Ezzat *et al.* is higher than therapeutically relevant concentrations of drug. Although there have been eight clinical trials involving PPMOs, none have involved lung disease. However, preclinical evidence suggests efficacy after topical administration to the lung. Thus, targeting of the respiratory syncytial virus (RSV) by intranasal (IN) dosing in mice demonstrated localized antisense delivery to the bronchial airways and indicated some prophylactic value (S.-H. Lai et al., 2008; Lupfer et al., 2008) if the drug was dosed within a limited window ahead of virus challenge. Therapeutic value, however, was questionable since only a modest, <65% reduction of viral titers was reported (S.-H. Lai et al., 2008). Curiously, the extent of virologic response was comparable to the ~50% reduction of transcript levels for endogenous RNAs when these were targeted by PPMOs (Rajsbaum et al., 2014). Yet efficacious antivirals typically induce acute, multi-log (i.e. >99.9%) drops in

viral target titers. Similar results were reported in piglets (Opriessnig et al., 2011), with RNase H-inactive antisense to RSV inducing a delay, but not elimination of viral infection. Of relevance, development of viral mutational escape (Lupfer et al., 2008) has led to adoption of multi-PPMO strategies and complex dosing regimes (Patrick L Iversen et al., 2012). PPMOs might also be active against bacteria, as a single, acute (5 min), post-infection, IN 0.1 mg dose of PPMO in mice was shown to be protective against respiratory challenge with multidrug resistant *Acinetobacter* by targeting bacterial transcripts (Geller et al., 2013). However, at least some of these activities are class-level bacteriostatic effects from the covalently attached CPPs that were employed as the delivery vehicle (Wesolowski, Alonso, & Altman, 2013). In addition, PMOs also demonstrate both antibacterial and antimalarial activities (Augagneur, Wesolowski, Tae, Altman, & Ben Mamoun, 2012), a finding that further complicates the separation of gene specific from off-target effects (Eisen & Smith, 2008). PPMOs have been also proposed as inhibitors (Francis et al., 2014) of endogenous mediators of RNAi, miRNAs, and miRNA target site protectors (Staton & Giraldez, 2011). More recently, additional backbone modification of the PPMO chemistry has been sought in the so-called PMO*plus* structure (Fig. 3A). This has been evaluated clinically at phase I against lethal filovirus infectious disease (Marburg and Ebola virus), with published data in non-human primates being encouraging with regards to protective capacity (Heald et al., 2014; Patrick L Iversen et al., 2012; Warren et al., 2010, 2015). However, published data lack confirmation of the MOA of PMO*plus* chemistries. Their utility in topical administration to the lung remains unknown.

As with PMOs, PNAs also demonstrated enhanced delivery (Veldhoen, Laufer, & Restle, 2008) when conjugated to simple (Robaczewska et al., 2005; Sazani et al., 2002) and



complex (Cordier et al., 2014) CPP peptides. This includes CPPs used for PMOs (Maekawa et al., 2015) and extends to natural protein transduction domains (Fabani & Gait, 2008; Oh, Ju, & Park, 2009) and small molecule compounds such as triphenylphosphonium (Mehiri et al., 2008) or flavin (Marlin et al., 2012). Yet others have also proposed backbone modification with fluorine (Ellipilli & Ganesh, 2015) or guanidine (Dragulescu-Andrasi et al., 2006) as alternatives to peptide conjugation. With respect to the lung, activity for PNAs has been evidenced only after intraperitoneal injection with CPP-PNAs; however, daily dosing is required with PNA ASOs, whereas phosphorothioate ASOs are compatible with weekly dosing (Sazani et al., 2002).

Although PNA ASO efficacy has been evaluated in various rodent models of systemic disease (Brolin et al., 2015; Fabani & Gait, 2008; Gao et al., 2015; Rembach et al., 2004; Robaczewska et al., 2005), there are no reports examining PNA ASO utility after topical administration to the lung in the absence of any delivery modifications. This would suggest PNA ASO might not be well suited for topical dosing to the airways. Tellingly, whereas Ahn *et al.* proposed topical PNA ASO dosing against respiratory viruses (Ahn et al., 2011), at least one group has reported that a highly complex microparticle formulation is necessary to achieve any degree of efficacy. Thus, the proposed solution is a quaternary system consisting of two polymer blends, CPPs, and the bioactive PNA (Fields et al., 2015). Such approaches bear considerable development and chemistry manufacturing control costs that challenge commercial viability even within orphan disease indications. Unfortunately, evaluation of this solution in a rodent CF model (McNeer et al., 2015), exhibited modest effects. Thus, activity was reported in <0.4% of alveolar epithelia and <1% of macrophages despite particle deposition in both the small and large airways and association with 50-90%

of lung cells (Fields et al., 2015). There remain, therefore, considerable challenges to progressing in the clinic PNA chemistries for lung disease.

### 2.1.3 Third generation Antisense

More recently, 3<sup>rd</sup> generation antisense have been developed that make use of 2'-5' bridging groups (Fig. 3B) resulting in greatly increased stability in biological fluids, as well as higher affinity for their molecular targets. This increased affinity has resulted in the development of shorter antisense, with enhancer delivery (as a result of their smaller size) and more advantageous pharmacokinetic and pharmacodynamic properties (Dirin & Winkler, 2013; Geary, Norris, Yu, & Bennett, 2015). These antisense are also short enough to target miRNAs (anti-miRs) (Elmén, Lindow, Schütz, et al., 2008) and directly inhibit the function of the miRNA and siRNA induced silencing complex (RISC) by preventing engagement of RISC with its RNA targets. The first 2'-5' modification to be described were 'locked' nucleic acid (LNA) (Wengel, 1999) although other structures (e.g. 2'-4' bridges) with similar performance metrics have been developed (Burel et al., 2013; Seth et al., 2009). To date, LNA remains the single 2'-5' bridging modification commercially available for research use. Interestingly, short LNA antisense enter cells *in vitro* without delivery systems i.e. in simple 'naked' saline formulations (Stein et al., 2010). The process has been termed 'gymnosis' from the Greek word for 'naked' (gymnos). This is an alternative and more efficient mechanism to 'free uptake' previously described for 2<sup>nd</sup> generation antisense (Loke et al., 1989). Crucially, activity is principally elicited in the cytosol rather than the nucleus, and possibly via nucleases other than RNase H (Castanotto et al., 2015). It remains to be

elucidated if this is a feature of all 3<sup>rd</sup> generation ASO, LNA-modified ASO alone, or an observation relevant to *in vitro* studies only. Nonetheless, 2,2,7-trimethylguanosine caps could enable nuclear targeting if necessary (Moreno et al., 2009). Presently, phosphorothioate LNA ASO have been shown to associate mainly with the liver and kidney (Straarup et al., 2010) and have been suggested to bind onto their intended pharmacological target to exert the expected MOA in up to phase IIb clinical studies (van der Ree et al., 2014). This has encouraged acquisition of the technology by Roche. However, both precursors and mature miRNA might be targeted by this approach (Gebert et al., 2013), and the methods used to evidence target engagement *in vivo* (Elmén, Lindow, Silahatoglu, et al., 2008) could be simply artifacts of cell association carrying through tissue homogenization and high affinity nucleic acid purification, rather than true target association.

## **2.2. Development of RNAi-based therapeutics**

Historically, RNAi therapeutics were developed on the back of discovery that delivery of exogenous dsRNA sequences of 20-27 nt length or greater, that were fully complementary to practically any part of target mRNAs, resulted in target cleavage and subsequent degradation (Fire et al., 1998). In mammals, >30 nt dsRNA were not thought to be useful in an RNAi manner as such molecules instigate antiviral responses (Bevilacqua & Cech, 1996), through Protein Kinase R engagement (Sledz, Holko, de Veer, Silverman, & Williams, 2003). However, present-day understanding of RNAi involves a considerably larger number of functions and encompasses the endogenous (miRNA) and exogenous (siRNA) mediators of RNAi operating at the transcriptional and post-transcriptional/pre-translational level (Fig. 2D) (Moschos, 2013; Sabin et al., 2013). Interestingly, endogenously expressed long (>30 nt

dsRNA-derived) siRNAs are encountered in reproductive biology, in transposon control and in cancer, operating through DNA methylation (L. Chen, Dahlstrom, Lee, & Rangasamy, 2012; Song et al., 2011; Tam et al., 2008; T. Watanabe et al., 2008; Werner et al., 2014)- a largely overlooked area. The reader is thus encouraged to consider siRNA and miRNA function in the context of RISC bioactivity as opposed to the mechanism of RISC formation as, overall, RISC functions are interchangeable for siRNA and miRNA.

The minimal bioactive complex of RNAi, the RISC complex (Fig. 2D, 5), contains one of four Argonaute proteins found in the mammalian genome, loaded with a short, ~21 nt long RNA strand of either endogenous (miRNA; Fig. 5A) or exogenous (siRNA) origin. RISC is often found in combination with multiple other regulatory proteins and uses Watson-Crick base pairing between the RISC-loaded RNA strand (referred to as the guide or antisense strand) and cellular RNAs to identify RNA targets. Target recognition relies on partial complementarity, often only between bases 2-9 from the 5' end of the guide strand (seed sequence) (Lewis, Shih, Jones-Rhoades, Bartel, & Burge, 2003). Where a transcript contains a sequence complementary to a miRNA seed sequence, the binding site is often referred to as the miRNA response (or recognition) element (MRE) (Leuschner, Ameres, Kueng, & Martinez, 2006; Matranga, Tomari, Shin, Bartel, & Zamore, 2005). These are most commonly found within coding transcript 3' untranslated regions (3'-UTR). However, cases of target recognition that do not involve the seed sequence at all but involve base pairing at the central (Shin et al., 2010) or 3' end of the guide strand have been documented (Sung Wook Chi, Hannon, & Darnell, 2012). The rules of target recognition are not fully described, restricting the utility of computational target prediction. Furthermore, miRNA and siRNA are

subject to 5' and 3' end modification (Azuma-Mukai et al., 2008; Burroughs et al.), which further complicates target prediction and validation.

Detailed descriptions of miRNA genomic organization and biogenesis can be found elsewhere (Moschos, 2013) and are summarized in Figure 3. In conjunction with viral and non-viral gene therapy approaches, miRNA and siRNA expression can be extensively engineered. Structurally, there are minimal differences in mature miRNA and synthetic siRNA structure (Fig. 2D), and siRNA-like synthetic miRNAs (miRNA mimics) are common tools in miRNA complementation / pharmacology studies (Perry et al., 2008). Diverse chemical modifications have been described to direct RISC precursor strand loading (Moschos, 2013) but there is no evidence to suggest AGO protein selection and RISC complex subcellular localization can be manipulated (J. H. Park & Shin, 2015).

As with single stranded ASO, the major issues with the development of RNAi mediators (siRNA and miRNA) as therapeutics has been delivery, stability, off-target actions and induction of immunological responses.

### **2.2.1 Demonstrating On-Target Action of siRNAs**

Of the four human Ago proteins that form RISC, only AGO2 possess an RNA endonuclease, or 'slicer' function, which allows for active target cleavage (F. Li et al., 2007). Slicing of targets occurs on their phosphate backbone opposite positions 10-11 from the 5' end of the guide strand (Leuschner et al., 2006; Matranga et al., 2005), and has been shown to occur both with miRNA and siRNA guide strands (Maniataki & Mourelatos, 2005). Crucially, direct

observation of this cleavage constitutes biochemical evidence of an on-target, AGO2 RISC-mediated pathway and is typically determined using 5' rapid amplification of complementary DNA ends (5'-RACE; qualitative) followed by Sanger sequencing (Soutschek et al., 2004) or next generation sequencing (RACE-Seq; semi-quantitative (Denise et al., 2013)). Global RNA-Seq (Taberner et al., 2013) can report the relative extent of target cleavage (knockdown), assess effects on other transcripts and any impact on splice variants. It is important to note, however, that experimental (Raabe, Tang, Brosius, & Rozhddestvensky, 2014; Sorefan et al., 2012; van Dijk, Jaszczyszyn, & Thermes, 2014) and analytical (Erhard & Zimmer, 2015; X. Liu, Zhang, & Chen, 2015; Yang & Jeong, 2013) bias in deep sequencing may severely skew observations (Lahens et al., 2014).

### **2.2.2 Off-target actions of siRNAs, or documenting miRNA bioactivity.**

It is generally accepted that the majority of off-target actions of siRNAs are mediated through their 'miRNA-like actions' i.e. RISC docking onto the 3'-UTRs of other mRNAs, leading to the unintentional inhibition of their translation (Filipowicz, Bhattacharyya, & Sonenberg, 2008; Jidong Liu, Valencia-Sanchez, Hannon, & Parker, 2005; R. Parker & Sheth, 2007). However, translational inhibition (as opposed to catalytic target cleavage) occurs also when siRNA forms RISC with AGOs 1, 3 and 4 which all lack the slicer function. Non-slicer RISC docking in coding regions (S W Chi, Zang, Mele, & Darnell, 2009; Nelson et al., 2007) may also lead to translational repression (Fang & Rajewsky, 2011; Schnall-Levin et al., 2011; Tay, Zhang, Thomson, Lim, & Rigoutsos, 2008) whilst docking on the 5' UTR may lead to up-regulation of translation (Ørom, Nielsen, & Lund, 2008) (Fig. 2D). These mechanisms can be assessed by determining siRNA guide strand interactions with mRNAs through cross-linking

and immunoprecipitation followed by next generation sequencing (CLIP-Seq) (S W Chi et al., 2009). Notably, this only reports target presence and/or enrichment across the entire gamut of RNAs pulled down through the assay and requires multiple processing (protease, nuclease) steps. To date, a method offering direct MOA evidence remains to be developed. Importantly, reporter constructs (Humphreys et al., 2012) disregard target transcript secondary and tertiary structures and can be misleading, as demonstrated through extensive virological research (Schopman, ter Brake, & Berkhout, 2010). Thus, single point mutations outside MRE / RISC binding sites are adequate to enable virus escape against RNAi mediators.

It is now also well established (Sabin et al., 2013) that mature RISC may translocate back into the nucleus (Bai, Liu, & Laiho, 2014; Castanotto, Lingeman, Riggs, & Rossi, 2009; Földes-Papp et al., 2009; Marcon, Babak, Chua, Hughes, & Moens, 2008; Ohrt et al., 2008; Politz, Zhang, & Pederson, 2006; Weinmann et al., 2009). Functional studies have shown that this can produce transcriptional activation (RNA activation) (L.-C. Li et al., 2006; Matsui, Chu, et al., 2013; Place, Li, Pookot, Noonan, & Dahiya, 2008; Schwartz et al., 2008; Y. Zhang et al., 2014) and repression (Ahlenstiel et al., 2012; Kim, Saetrom, Snøve, & Rossi, 2008; Younger & Corey, 2011), epigenetic remodeling (Ahlenstiel et al., 2012; Kim et al., 2008; L.-C. Li et al., 2006; Morris, Santoso, Turner, Pastori, & Hawkins, 2008; Younger & Corey, 2011; Zardo et al., 2012), miRNA precursor maturation control (Tang et al., 2012), as well as splice switching (Alló et al., 2009; Ameyar-Zazoua et al., 2012; Jing Liu, Hu, & Corey, 2012). Most of these functions involve AGO1 (Kim et al., 2008), but may also include AGO2 (Ohrt et al., 2008; Schraivogel et al., 2015) in a slicer-positive (Gagnon, Li, Chu, Janowski, & Corey, 2014) or negative (Matsui, Chu, et al., 2013) fashion. Functional partners of nuclear RISC include

transcriptional proteins (Matsui, Chu, et al., 2013; Y. Zhang et al., 2014), promoter-associated and natural antisense transcripts (Matsui, Chu, et al., 2013; Morris et al., 2008; Schwartz et al., 2008) or intergenic ncRNA (Matsui, Prakash, & Corey, 2013). Interestingly, some functions appear to integrate the spliceosome, RNA polymerase elongation and chromatin remodeling (Ameyar-Zazoua et al., 2012), and even participate in dsDNA break repair (Wei et al., 2012).

Beyond these functions, viruses have also been shown to encode RNAi mediators and manipulate RNAi homeostasis. Thus, RISC can be used to prevent 5'-3' exonuclease cleavage (Sedano & Sarnow, 2014; Thibault et al., 2015) of RNA viral genomes, promote virus replication (Fan et al., 2015), regulate viral polyprotein translation (Masaki et al., 2015) and organize the viral genome structure (Masaki et al., 2015; Mortimer & Doudna, 2013; Narbus et al., 2011). Whether some of these functions are also relevant to endogenous transcripts remains to be determined. Importantly, as with HIV *in vitro* (Schopman et al., 2010), clinical studies have evidenced mutations external to the RISC binding sites may alter RISC binding capability (Israelow et al., 2014). Yet more recently, components of AGO2 RISC have been isolated from mitochondria (Xiaorong Zhang et al., 2014), although this may not be ubiquitous across cell types (Ro et al., 2013). Mechanistic studies suggest functions in line with other cytosolic RNAi MOA (Jagannathan et al., 2015), although AGO2 RISC translocation within myocyte mitochondria and translational upregulation activity has been proposed (Xiaorong Zhang et al., 2014). Roles in mitochondrial transcription regulation have been so far documented only in plants (Dietrich, Wallet, Iqbal, Gualberto, & Lotfi, 2015), with similar studies lacking in mammals.



### 2.2.3 siRNA and miRNA mediated immune responses

RNAi mediators are now understood to be natural agonists of multiple pathogen associated molecular pattern receptors (Table 1) such as the toll-like receptor (TLR) family members 3 (Cho et al., 2009; Kleinman et al., 2008), 7 and 8 (Hornung et al., 2005; Judge et al., 2005). Their activation drives the release of various proinflammatory cytokines, with diverse kinetics which can vary substantially based on host species, system complexity (*in vitro* vs. *in vivo*) and delivery system type (Broering et al., 2014; Forsbach et al., 2008, 2012; Moschos et al., 2007). Crucially, the effects of at least TLR3 activation are not local: stimulation by an siRNA in the peritoneum can result in remarkable remodeling in tissues as anatomically distant as the eye (Kleinman et al., 2008). This finding contributed substantially to the abandonment of the first-in-class siRNA clinical candidate, bevasiranib. The cytokines presently implicated in TLR-mediated off target effects include alpha interferon, tumor necrosis factor alpha, interleukin (IL) 6, IL-10, beta interferon, interferon-induced protein with tetratricopeptide Repeats 1 and interferon sensitive gene 15, on account of large and small (dinucleotide) sequence motifs (Forsbach et al., 2008; Judge et al., 2005; Robbins et al., 2007; Schlee, Hornung, & Hartmann, 2006) also found in many miRNAs. Indeed, in stark contrast to common misconceptions, it is now confirmed that miRNA are TLR7/8 agonists (at least) that induce inflammatory (Fabbri et al., 2012), pro-apoptotic (W. A. He et al., 2014) and pain responses (C.-K. Park et al., 2014) through exosomal communication pathways (Patton et al., 2015); these functions depend on the miRNA sequence and the recipient cell type, and should come as no surprise to the reader versed in the immunostimulatory potential of synthetic siRNA-like molecules.

Advantageously, nucleoside analogues (Table 1 and Fig. 3B) involving 2' ribose modifications (e.g. 2'-O-methyl, 2'-fluoro and 2'-5' oxygen-bridged 'locked' nucleic acids (LNA)) incorporated in RNAi mediators act as a TLR7/8 antagonists (Robbins et al., 2007; Sarvestani et al., 2015; Sioud, Furset, & Cekaite, 2007), with at least one of these, 2'-O-methyl, altering RISC off-target effect profiles (Fedorov et al., 2006; Jackson et al., 2006). The impact of other modifications on RISC off-target effects has not been publicly disclosed. Although nucleoside suppressors of both TLR3 and TLR7/8 have been described (Karikó, Buckstein, Ni, & Weissman, 2005) (Table 1), their impact on siRNA-mediated TLR3 activation has not been assessed. Presently, reports on the impact of 2-thiouridine on slicer function are contradictory (Prakash, Naik, Sioufi, Bhat, & Swayze, 2009; Sipa et al., 2007). Overall, judicious use of one or two 2' modified nucleosides as formulation adjuncts (e.g. 2'-O-methyl nucleosides, a natural nucleoside) or as siRNA sequence modifications can largely eliminate TLR7/8 activation potential and ablate the pyrimidine-purine specificity of RNase A (i.e. UpA, CpG, etc.) (Turner, Jones, Moschos, Lindsay, & Gait, 2007). Thus, simple changes with minimal impact of strand  $T_m$  can evade TLR7/8 induction and also drive better compound stability and increase confidence of an on-target (i.e. non-inflammatory) MOA. Presently, TLR3 evasion can be engineered only through iterative screening and testing.

Beyond the TLR family, the intracellular pathogen associated molecular pattern sensors 2'-5'-oligoadenylate synthase 1 (OAS1) (Kodym, Kodym, & Story, 2009), double-stranded RNA-activated inhibitor of translation (DAI) (Manche, Green, Schmedt, & Mathews, 1992), melanoma differentiation-associated gene 5 (MDA5) (Kato et al., 2008), and retinoic acid-inducible gene 1 (RIG-I) (Hornung et al., 2006; Marques et al., 2006; Pichlmair et al., 2006)

are also activated by RNAi mediators or their precursors. Fortunately, most are long dsRNA-specific. Among those sensitive to smaller dsRNA OAS1 is specific for the nucleotide motif NNWW(N<sub>9</sub>)WGN (Kodym et al., 2009) whereas RIG-I is inhibited by 3' dinucleotide overhangs typical to miRNAs and endogenous siRNAs (Marques et al., 2006) (Fig. 2) or evaded through lack of 5' triphosphate use on either RISC precursor strand (Kato et al., 2008).

### **3. Clinical progress of oligonucleotide therapeutics for lung disease.**

The original studies on lung-targeted oligonucleotide therapeutics (Moschos, Spinks, Williams, & Lindsay, 2008) expected delivery into lung cells following topical administration, despite historical data suggesting quick absorption into the circulation (Nicklin et al., 1998). This was principally due to the large size (~12 kDa) of the then fashionable siRNA compared to single stranded antisense. Thus, 90-99% reduction of gene expression (i.e. 1-2 logs) was commonly observed in cell culture whilst preclinical studies showed 2-fold (40-60%) reduction of target RNA levels (summarized in (Moschos et al., 2008)). Encouragingly, two co-administered antiviral siRNAs against separate viruses produced a 90-99% reduction in target mRNA and in tandem, alpha interferon production – a possibly advantageous feature for an antiviral RNAi drug (Bitko, Musiyenko, Shulyayeva, & Barik, 2005). Others reported that uptake across the tissue was not uniform for both ASO and siRNA encapsulated in a lipid delivery system (Griesenbach et al., 2006), and CPP conjugation to siRNA also yielded mediocre effects (Moschos et al., 2007). The inconsistencies were interpreted as experimental setup differences and variability between constitutively expressed vs. induced

RNA targets. Based upon these initial observations, a number of oligonucleotide-based therapeutics have been developed/tested in clinical studies.

### **3.1 RNAi mediators: ALN-RSV01 and the treatment of RSV infection.**

Presently, the most clinically advanced RNAi mediator remains Alnylam's ALN-RSV01, a first generation, unmodified siRNA targeted against a conserved region of the nucleoprotein gene encoded in the RNA genome of respiratory syncytial virus (RSV) (DeVincenzo et al., 2008). Dosing the airways was expected to match the respiratory epithelium tropism of the virus (Johnson, Gonzales, Olson, Wright, & Graham, 2007), in a well-defined patient population not catered for by any on-label treatment alternatives (DeVincenzo et al., 2008), and thus likely to achieve commercial success.

#### **3.1.1 Preclinical pharmacology.**

Although mildly immunostimulating (single time point data), 2'-O-methylation of ALN-RSV01 did not affect siRNA IC<sub>50</sub>, suggesting TLR activation in mouse, if any, did not contribute to antiviral responses (Alvarez et al., 2009). On-target RNAi activity *in vivo* by 5'-RACE (Alvarez et al., 2009) further justified no need for use of any nucleoside modifications. Given established *in vivo* pharmacology practice and in line with contemporary studies, drug activity was measured at the tissue level. This involved homogenization of one or more lobes of the lung and reporting virological, RNA or protein assay levels at the whole tissue level. However, direct evidence of tissue loading and retention was not attempted as no systemic absorption was expected (DeVincenzo et al., 2008). It was indeed widely assumed that the large molecular weight of siRNAs (~12.5 kDa) was recalcitrant to systemic exposure.

Rather, their anionic charge should complex with cationic alveolar surfactant (De Backer, Cerrada, Pérez-Gil, De Smedt, & Raemdonck, 2015) to piggy-back through surfactant recycling into alveolar epithelia and macrophage cytosols.

### **3.1.2. Clinical performance and disease biology.**

Curiously, the highest dose (150 mg) tested at phase I resulted in rapid (<10 min) detection of ALN-RSV01 in blood plasma and in the first post-administration urination, at the lower detection limit of the analytical method. This was again interpreted as limited systemic exposure (DeVincenzo et al., 2008) without recourse to more sensitive methods. In an experimental clinical infection model the antiviral effect in the nasal cavity was shown as independent of TLR responses by multivariate regression analysis (DeVincenzo et al., 2010a), but no effort was made to confirm a difference between human and preclinical inflammatory responses. Thus, ALN-RSV01 was suggested to prevent the migration of the infection from the nasopharyngeal epithelium into the lower airways (Hall, 2001), an interesting proposition given RSV was classically considered a disease of the lower airways (Wright et al., 2005).

The progressive onset of symptoms from the upper to the lower airways had been documented in animal models (Gitiban et al., 2005; Richardson et al., 1978) and clinically (Hall, Douglas, & Geiman, 1975, 1976), suggesting the virus might indeed establish in the nares and progressively infect the lower airways over several days. However, no data existed confirming this hypothesis (El Saleeby, Bush, Harrison, Aitken, & Devincenzo, 2011) and the role of adaptive immunity was overlooked. Instead, comparable viral loads were

demonstrated in the upper and lower airways in the mid- to late- stages of infection (Perkins et al., 2005). Crucially, the proposed treatment strategy was not experimentally tested to determine whether antiviral protection across the upper and/or lower airways would be uniquely adequate or jointly necessary. More recent findings indicate RSV might reach the lower airways considerably sooner than clinical symptom onset might otherwise suggest (Rameix-Welti et al., 2014).

The first patient trial sought to test ALN-RSV01 utility in preventing development of RSV-mediated *bronchiolitis obliterans* syndrome, associated inflammation, infection and ultimately tissue rejection in lung transplant patients (Zamora et al., 2011a). In contrast to the previous studies (DeVincenzo et al., 2008, 2010b) the drug was administered as an inhaled, saline-formulated aerosol targeted to the lung only (Gottlieb et al., 2015a). The 0.6 mg/kg dose, ~1/3 of the 150 mg intranasal dose (DeVincenzo et al., 2008), again achieved rapid (<10 min) systemic access at the analytical assay's lower limit of detection. Furthermore, at least one patient not on corticosteroids exhibited elevated cytokine responses (Zamora et al., 2011a). No statistically significant antiviral effect was achieved, but syndrome incidence and associated sequelae reduction was reported. Thus, syndrome incidence was chosen as a sign of efficacy in follow up studies.

In phase IIb, however, the incidence rate of *bronchiolitis obliterans* syndrome was not reduced in RSV-infected individuals in a statistically significant manner, unless strict adherence to the therapeutic protocol was observed (10% incidence with treatment vs 28% incidence with placebo,  $p = 0.025$ ) (Gottlieb et al., 2015b). Notwithstanding the impact on study power engendered through subject removal from analysis on account of protocol

adherence, complete lack of any antiviral effects in oropharyngeal washes and nasal swabs were ascribed to pulmonary targeting of the aerosol formulation. Yet no viral genome load modulation was achieved in the deep lung either: it is unclear if the analytical method used was sensitive to RISC-mediated target cleavage (G. Chen, Kronenberger, Teugels, & De Grève, 2011; Herbert, Coppeters, Lasham, Cao, & Reid, 2011; Holmes, Williams, Chapman, & Cross, 2010; Shepard, Jacobson, & Clark, 2005). It is also unclear if the reported reduction in syndrome incidence was purely on account of ALN-RSV01 treatment: of the patients in the treatment arm of the study, 71% were also receiving pulse steroid treatment vs 58% in the placebo arm.

### **3.2. 2<sup>nd</sup> Generation ASO therapeutics: Clinical Status.**

Parallel to ALN-RSV01, topically administered 2<sup>nd</sup> generation ASO were being clinically investigated for asthma and COPD (Séguin & Ferrari, 2009). These groups also presumed lung retention, measured target modulation at the tissue level and valued disease outcome metrics more than molecular pharmacology. At the time, three clinical candidates were being progressed through to clinical programs: i) AIR645 (previously known as ISIS 369645; 2'-O-methoxyethyl chemistry; sponsored by Altair Therapeutics, an Ionis Pharmaceuticals spin-off) targeting the alpha chain common to IL receptors 4 (IL-4R) and 13 (IL-13R) (Fey et al., 2014), ii) ASM8 (TOP004 and TOP005 dual antisense formulation; 2'-deoxy-2'-fluoro-beta-D-arabinonucleic acid (FANA) chemistry (Fig. 3); sponsored Topigen Pharmaceuticals, later acquired by Pharmaxis) targeting the common beta chain ( $\beta$ c) of the ILR receptors 3, 5 and granulocyte monocyte – colony stimulating factor (GM-CSF) and the C-C chemokine receptor 3 (CCR3), respectively (G M Gauvreau et al., 2011) and iii) ATL1102 (ISIS 107248,

another 2'-O-methoxyethyl; Antisense Therapeutics) targeting the cluster designation 49d subunit of the very late antigen 4 adhesion molecule.

### **3.2.1. Altair AIR645 and the targeting of IL-R4/IL-R13 in asthma.**

The Altair compound was progressed on strong mechanistic rationale around the IL4/IL-13 receptor signaling system in asthma in the context of lung infiltrating, immunoglobulin E-producing, CD4+ T lymphocytes (Chatila, 2004; Karras et al., 2007). Moreover, the group had already explored 2<sup>nd</sup> generation antisense dosing to the mouse lung with other targets (Duan et al., 2005). These early studies indicated dose-dependent antisense retention in the lung but at very low levels (1.1 µg per g tissue 24 hours after a 3 mg/kg estimated dose (Duan et al., 2005)). Aerosol chamber dosing was understandably less efficient than nose-specific aerosol administration (62.4 µg per g tissue from a 3.1 mg/kg dose (Templin et al., 2000)), which also differed from historical intranasal instillation studies (Nicklin et al., 1998). Immunohistochemistry had demonstrated macrophage loading and diffuse association of antisense with lung epithelia cell types (Templin et al., 2000); however, subcellular distribution was not assessed. Nonetheless, Duan *et al.* achieved restriction of multiple asthmatic disease biomarkers efficiently modeled in mice (eosinophilia, mucus production, cytokine production, airway hyper-responsiveness), including ~60% reduction in target mRNA levels in both luminal and draining peribronchial lymph node lymphocytes. However, the mRNA data were not normalized for endogenous housekeeping gene levels (Gorzelnik, Janke, Engeli, & Sharma, 2001), undermining the claim of an on-target mechanism of action. Using a murine-specific version of AIR645, ISIS 231894, Karras *et al.* attempted to address this by measuring the reduction of IL4R alpha chain protein levels (Karras et al., 2007). This involved fluorescent cell sorting of primary lung cell subtypes in mice receiving an estimated



0.01-0.5 mg/kg nose-specific aerosol dose. At best, an ~50% reduction in IL4R alpha chain levels was observed in macrophages and lung epithelia at the highest dose, but it is unclear if multiple comparisons correction was applied in determining statistical significance, or if data distribution profiling was applied to inform linear or logarithmic scale statistical testing. Drug retention within each cell type was also not assessed. Thus, Karras *et al.* reported asthmatic response ablation but the mechanism remained unclear. Interestingly, immunomodulation appeared to be part of the effect of ISIS 231894 as revealed through cytokine profiling in a separate, antiviral study (Ripple *et al.*, 2010). The relevance of these results appears to have been overlooked in the asthma model outputs.

Both AIR645 (primate-specific) and ISIS 231894 were tested in rodents for toxicity and tolerability, but only AIR645 was tested in monkeys (Fey *et al.*, 2014) at 0.4 - 50 mg/kg. This involved classical pulmonary-targeted drug toxicity endpoints and complement activation but excluded cytokines (Advani *et al.*, 2005; Henry *et al.*, 1997). Both antisense and target levels were analyzed at tissue level (Duan *et al.*, 2005; Karras *et al.*, 2007). Target mRNA levels were determined in the tissue (both species) and bronchoalveolar lavage cells (monkeys only), but were expressed with reference to total RNA levels determined by substantially alternative methodologies, an approach known to result in transcript quantification artifacts. No complement activation was observed but up to 24%, dose-dependent, lung weight increases were documented in mice. Macrophage recruitment and enlargement (hystiocytosis; foamy macrophage phenotype) in the parenchyma and tracheobronchial lymph nodes was observed in both species, persisting for 13 weeks. Dose-dependent pro-inflammatory infiltrates were reported only in mice, with hystiocytosis ascribed to 'normal' alveolar macrophage function as defined by Nikula *et al.* (Nikula *et al.*,

2014). Importantly, the Nikula *et al.* article is an expert-level regulatory position paper on the impact of the foamy macrophage induction in response to topically dosed lung therapies. It is worth noting, that Nikula *et al.* make a clear distinction between soluble drugs (i.e. aqueous antisense formulations) and insoluble particles, such as complex formulations and small molecule aggregates (dry powders). Thus, insoluble particles do not rapidly clear from the lung and are removed specifically by macrophages (alveoli) and mucocilliary clearance (bronchial airways). This contrasts soluble antisense and siRNA which rapidly access circulation, for which there is limited clinical data available. The physiological impact of oligonucleotide-induced hystiocytosis requires further elucidation before safety is established as suggested in Karras *et al.*

Fey *et al.* calculated the clearance half-life from the lung at nine days and reported dose-dependent loading in the liver and kidney. The data were in line with parallel studies supporting lower bioavailability for topically dosed 2<sup>nd</sup> generation ASO, but contrasted earlier reports (Nicklin *et al.*, 1998): these differences could be ascribed to the use of 1<sup>st</sup> generation ASOs by Nicklin *et al.* Yet no appreciable change in IL-4R alpha chain mRNA levels was observed. Of note, as stated in Fey *et al.*, the 1000x lower doses of AIR645 used by Karras *et al.* were apparently pharmacologically active, where Fey *et al.* reported no target modulation. The notable discrepancy was ascribed to the low-level target gene expression under naïve conditions in the Fey *et al.* study. In other words, targets expressed at low-levels would appear to be recalcitrant to oligonucleotide-mediated downregulation. Instead, Fey *et al.* postulate that antisense should be used against transcripts undergoing upregulation (e.g. activated pro-inflammatory response genes). Furthermore, antisense efficacy should be measured only when these transcripts are being upregulated, not under

baseline conditions. This contrasts evidence that transcripts expressed at low levels can be knocked down by antisense, provided the tissue is effectively transfected (e.g. apolipoprotein B downregulation in the kidney (Moschos et al., 2011)). It remains to be determined if low-level expression transcripts are malleable targets universally, or in a case-by-case basis. Clinical efficacy data have yet to reach the peer-reviewed public domain for AIR645.

### **3.2.2. Pharmaxis ASM8 and the targeting of CCR3 and the common $\beta$ c of IL-3R, IL-5R, and GM-CSF in asthma.**

The pursuit of a dual formulation consisting of two antisense drugs based on the FANA chemistry (TOP004 and TOP005; ASM8; Fig. 3) targeting two different transcripts was backed by substantial confidence in rationale (Corren, 2012). Thus, haematopoietic myeloid progenitor cells (HMC) are elevated in the circulation the lung tissue and sputum of asthmatics (and in murine models of asthma) under both naïve conditions and after allergen stimulation (Cameron et al., 2000; Dorman et al., 2004; D. S. Robinson et al., 1999; R Sehmi et al., 1996; Sergejeva, Johansson, Malmhäll, & Lötvall, 2004; Southam et al., 2005). HPCs express the receptors CCR3 and IL-5R, and differentiate into eosinophils in response to their activation (Roma Sehmi et al., 2003)(Dorman et al., 2004)(R Sehmi et al., 1997). Eosinophils are a cell type hallmark of allergic responses (Wenzel, 2006), whose concentration in the lung correlates with asthmatic disease severity (Bousquet et al., 1990). In asthmatic disease, HPCs overexpress these receptors (R Sehmi et al., 1997), parallel to the increased levels of these cytokines (Hamid et al., 1991; Humbert et al., 1997; D. Robinson et al., 1993). *Ex vivo*, soluble receptors were shown to attenuate HPC differentiation (Cameron et al., 2000).

Furthermore, clinical and primate studies with injectable monoclonal antibodies against these receptors or their corresponding cytokines, as pioneered by Leckie *et al.* (Leckie *et al.*, 2000), indicated asthma exacerbation management could be achieved by manipulating this immune axis. Indeed, reslizumab (anti-IL-5; Teva Pharmaceutical Industries Ltd.) is presently the third monoclonal antibody on track for market approval in the USA for the treatment of asthma, and the second anti-IL-5 agent after mepolizuman (GlaxoSmithKline Plc.); benralizumab (anti-IL5-R $\alpha$ ; MedImmune LLC.) has also been successful in reducing asthma exacerbations at phase IIb (Castro *et al.*, 2014). Thus, both underlying biology and clinical data contemporary to the development of ASM8 suggested that dual action against CCR3 and IL-5R could potentially achieve broader ablation of asthma exacerbation.

Encouragingly, single antisense studies *in vitro*, in rodents and non-human primates (Z Allakhverdi *et al.*, 2006; Zoufia Allakhverdi, Allam, & Renzi, 2002; Allam & Renzi, 2001; Fortin *et al.*, 2006) were positive, suggesting effective target knockdown at both the RNA and protein level (CCR3 and  $\beta$ c) in tissue and *ex vivo* cell studies, in line with reduced airway eosinophilia and hyper-responsiveness after allergen challenge. However, the gel-based RNA quantification methodology used was poorly quantitative and no explicit MOA results such as 5'-RACE were reported throughout this work. Interestingly, additive effects were observed for the two drugs in rat models, such as reduction of lymphocyte and macrophage recruitment after challenge (Z Allakhverdi *et al.*, 2006). Yet the results obtained in non-human primates was contradictory (Guimond *et al.*, 2008). Thus, as with AIR645, sporadic, but dose-dependent evidence of macrophage accumulation and inflammation was reported, principally at high doses; plasma detection of the drugs was only achieved at the 2.5 mg/kg dose, with follow-on compounds detected also in the liver and kidney at the high

dose only. Curiously, very modest target knockdown was reported (16% for CCR3 and 20% for  $\beta$ c), only in the tracheal tissue and not in the lung parenchyma.

Clinical studies (Gail M Gauvreau et al., 2008) evidenced ASM8 recovery in lung sputum and ablation of the six-fold increase of  $\beta$ c mRNA ( $p = 0.039$ ) induced by allergen challenge. However, the effect on CCR3 mRNA was not statistically significant, nor was that on the protein levels of either target. Similar observations were reported in some, but not all clinical markers of efficacy. Thus, a 46% reduction of sputum eosinophilia did achieve statistical significance, but the trend towards total leukocyte recruitment ablation did not; slightly elevated levels of macrophages were also observed. Dose-dependent, ~40% ablation of the inflammatory response was reproduced in the ensuing dose escalation study (G M Gauvreau et al., 2011), but this also coincided with a dose-dependent increase in sputum macrophages (G M Gauvreau et al., 2011). In later work (Imaoka et al., 2011), ASM8 succeeded in reducing the early (0-2 hours) and late (3-7 hours) asthmatic responses measured as a loss of forced expiratory volume in one second after allergen challenge. However, both phases of the response were ablated only by the high, 8 mg once daily, inhaled dose; the early response was not affected by a 4 mg dose. In these patients eosinophil progenitor recruitment was also significantly reduced in sputum after allergen challenge, accompanied by statistically significant reduction of CCR3<sup>+</sup> and IL-5R<sup>+</sup> HPC numbers, but not total HPC level.

Overall, despite the robust rationale, encouraging pre-clinical data and clinical success with other, injectable therapeutic modalities (antibodies) targeting the same immune signalling axis, the clinical progress of ASM8 was marred by lack of explicit MOA data, inconclusive

bioanalytical results on target RNA and protein levels and mixed clinical outcomes. It is important to note at this point that both ASM8 and its competitor AIR645 have consistently induced macrophage recruitment into the airways, a phenotype not yet cleared as safe with regards to soluble therapeutic compounds.

### **3.3. Third generation ASO.**

The utility of <20 nt LNA ASO was first evaluated in the same study that examined siRNA delivery to the lung at the cell type-specific level (Moschos et al., 2011). Unfortunately, as with the siRNA arm of this work, no activity was observed with LNA ASO in the lung. However, whilst siRNA and phosphodiester LNA ASO were rapidly (<15 min) eliminated in urine, phosphorothioate LNA ASO formed punctate structures within lung epithelia across upper and lower airways. These interactions appeared to be extracellular and membrane specific rather than endocytic, cytosolic or nuclear. Quantifiable amounts of phosphorothioate LNA ASO were reported only in lung macrophages, the liver and the kidney. Crucially, whereas there was no activity in lung macrophages, efficacy was observed in both the kidney and the liver: the latter was found to be indeed comparable to intravenous administration.

These data suggested inhalation might actually be a viable needle-free dosing solution for 3<sup>rd</sup> generation ASO therapies targeting systemic tissues rather than the lung, when the oral bioavailability of 2<sup>nd</sup> generation ASO varies between 2-12% even with the use of uptake enhancers (Nicklin et al., 1998; Raof et al., 2004). Given the differences in ASO structure,

molecular weight (length), target affinity and mismatch propensity between 2<sup>nd</sup> and 3<sup>rd</sup> generation ASO, extrapolation of the LNA findings (Moschos et al., 2011) to 2<sup>nd</sup> generation ASO was actively discouraged. However, comparable doses of the phosphorothioate AIR645 (Fey et al., 2014) and LNA ASO (Moschos et al., 2011) resulted in 10x lower kidney retention of AIR645. Moreover, the low kidney expression of the RNA targeted by the LNA ASO did not prevent ASO activity as suggested for AIR645 (Fey et al., 2014). It is unclear if such limitations are target-specific or ASO-class specific, nor whether prolonged lung dosing of LNA ASO also drive macrophage recruitment.

#### **4. Oligonucleotide Pharmacology: Time for a Reboot?**

##### **4.1 Identifying off-target actions**

The plurality of biological actions mediated via RISC (Fig. 2 and Table 1), and the current observations from clinical development of siRNA and ASO therapeutics have questioned the generic use of this drug class. It is the view of the authors that these issues are also relevant to the laboratory based research using siRNAs, miRNA mimics and, perhaps less so, antisense drugs. Crucially, the vast majority of studies fail to demonstrate on-target MOA and supporting evidence is provided almost universally in an indirect fashion. From a systems pharmacology and toxicology perspective (Cook et al., 2014), confidence in safety requires elimination of unintended, off-target mechanisms throughout the drug development program, using robust and reproducible methods.

Essential in the process is the use of multiple positive and negative controls. Thus, independent verification through alternative tool compounds acting on the same molecular

target or pools such as multiple siRNA targeting the same gene (Hannus et al., 2014) can be employed. Equally, multiple negative controls are also valuable to deconvolute off-target phenomena. Unfortunately, common practice appears to be restricted to one or two (random or habitual) selections (Table 2).

Crucially, it is also important to continuously assess the potential immunomodulatory actions of oligonucleotides and not to restrict these investigations to specific time-points or inflammatory markers. One approach proposes the use of alternating modifications of the passenger strand as a universal tool for ablating TLR7/8 stimulation by siRNA (Hamm et al., 2010). This harkens to the immunostimulatory RNA strand logic of isolating TLR7/8 activation to the passenger strand where this might be mechanistically convenient (Hornung et al., 2005), e.g. in anti-viral therapeutics. It is unclear to what extent this is adequate beyond TLR7/8 activation and within which cell type/tissue context. It is thus not surprising that only two tumor suppressive miRNA analogues, miRNA-16 (TargomiRs; EnGeneIC Ltd.) and miRNA-34 (MRX34; Mirna Therapeutics Inc.), have entered clinical trials to date (Lam, Chow, Zhang, & Leung, 2015). A modified passenger strand drives careful RISC loading of the canonical miRNA-34 guide strand by MRX34 (Daige et al., 2014). Presumably, this also enables evasion of TLR7/8 activation. Conversely, modified nucleotides are used in the 5' ends of both strands of the miRNA-16 mimic TargomiRs (Reid et al., 2013), i.e. within the seed sequence (Fedorov et al., 2006; Jackson et al., 2006). It is unclear at present to what extent the modification strategies adopted with these clinical candidates affect slicer-independent RNAi pathways, successfully evade TLR recognition or impact on guide strand 3' RNA editing common to many miRNAs.

Commented [ML2]: Don't understand this?



Are TLR's 3, 7 and 8 the only concerns of the community? Natural killer cell activation has also been reported with synthetic miRNA analogues both *in vitro* and *in vivo* (S. He et al., 2013). Importantly, this report is unusual as the toll-like receptor implicated, TLR1, is not understood to be a (oligo)nucleotide pattern recognition receptor. Further studies are urgently needed.

One solution proposed to circumvent immune receptor activation has involved the use of corticosteroid inhibitors of TLR signaling (Zamora et al., 2011b). This might be questionable for some indications but can be advantageous in asthma, chronic obstructive pulmonary disease, sarcoidosis and acute lung injury where corticosteroid treatment is recommended, or even pulmonary fibrosis due to habitual use (Xaubet et al., 2013). Corticosteroids and kinase inhibitors may also be of use in deconvoluting off-target pro-inflammatory response induction in drug discovery efforts.

#### **4.2 siRNA therapeutics: The case for next generation *in vivo* pharmacology.**

The preclinical and clinical studies on pulmonary dosing of siRNA showed that topical administration of unmodified compounds could rapidly (<10 min) access circulation and be mildly immunogenic. Success was assessed using clinical outcome metrics and not molecular pharmacology: no 5'-RACE data in man were disclosed. Thus, it remains unknown if topical siRNA administration to the lung in simple formulations such as physiological saline (referred to as 'naked' siRNA) can drive on-target RNAi in man. Notwithstanding the contribution of the immune response by ALN-RSV01 activity, the correct target tissue (nasal and lower airways epithelia) was not fully defined, administration methodology was not

consistent and drug loading was not evidenced directly or followed up adequately given the systemic access outcomes.

Separate preclinical studies sought to determine which cell types within the lung were differentially loaded with oligonucleotide drugs (Moschos et al., 2011), by using cell sorting and confocal microscopy after *in vivo* dosing. Unlike previous efforts (Lomas-Neira, Chung, Wesche, Perl, & Ayala, 2005; Moschos et al., 2007; Perl et al., 2005; Xuchen Zhang et al., 2004) this group sought to quantify drug uptake and activity in specific cell types relevant to lung disease rather than reporting qualitative, low resolution (microscopy), or tissue-level data. In addition, the far-red fluorophore used in these studies, sulphonated Cy5, had an emission wavelength with very limited tissue fluorescence background, and lower lipophilicity ( $\Delta\text{clogD}$ ) to plain Cy5. Indeed, lipophilicity is a feature of many siRNA modifications like cholesterol (similar molecular weight and  $\Delta\text{clogD}$  to Cy5), which induce better cellular uptake of oligonucleotides *in vivo* (Bijsterbosch et al., 2000; C. Lorenz, Hadwiger, John, Vornlocher, & Unverzagt, 2004; Soutschek et al., 2004; Wolfrum et al., 2007). It was thus shown that although 2'-O-methylated siRNA does indeed interact with the lung tissue, it was not subsequently adsorbed, endocytosed or delivered into the cytosol of epithelia, alveolar macrophages or endothelia. No biological activity was observed in any of the examined cell types, even though the study was powered at >94% (n=5) to measure 50% reduction in target levels. Instead, mass spectrometry showed that fully intact, sulphonated Cy5-labelled siRNA was recovered at substantial quantities within the urine of mice within 15 min of intratracheal administration (Moschos et al., 2011). Most convincingly, the colour of the urine was deep blue, i.e. the same naked eye colour of the sulphonated Cy5 dye. Indeed, a similar, unreported observation in the red (Cy3) spectrum

(Moschos et al., 2007) had been previously dismissed as incidental hematuria and remained uninvestigated. Simple allometric scaling on heart rates between species, would translate these findings to siRNA detection in urine ~140 min after administration in man. Although direct comparisons of the compounds tested in mice (Moschos et al., 2007, 2011) to ALN-RSV01 (DeVincenzo et al., 2008; Zamora et al., 2011a) is difficult, rapid (<10 min) circulation access in clinical subjects supports a common absorption and elimination mechanism in both species. As 5'-RACE was not attempted in these studies, it remains unclear if non-quantitative 5'-RACE can report residual RNAi and contribute towards misinterpretation of off-target phenomena as on-target 'slicer' RNAi.

## **5. Concluding remarks and future challenges**

At the time of writing, Alnylam programs involving siRNAs-based approaches to the treatment of genetic and cardiometabolic disease have eclipsed ALN-RSV01 and associated projects, with RSV product development apparently focused in Japan. Elsewhere, attention on RNAi mediator therapeutics for the lung has shifted onto solving the problem of cytosolic delivery (Clark et al., 2013), whereas ongoing clinical trials in the USA are concerned with systemic, non-lung cancer, skin and ocular disorders, almost invariably involving advanced drug delivery systems (Lam et al., 2015). Nevertheless, respiratory medicine is rich in marketed and clinically advanced candidates, some of which are small molecules cleverly designed for topical administration (Jones et al., 2011) and thus are exceptionally competitive commercial propositions against modalities with more complex chemistry, manufacturing and controls requirements.

The studies on the antisense based AIR645, ASM8 and associated compounds indicated effective tissue/cell loading, potentially on-target (mRNA) molecular activity for ASM8 (but not necessarily for AIR645) and effective ablation of asthmatic phenotypes. However, no explicit 5'-RACE MOA data, evidence of sub-cellular localization was evidenced or ASO association with RNase H/intended molecular targets was reported, in line with the so-called three pillars of drug survival (Morgan et al., 2012) and subsequent emerging principles in drug development (Cook et al., 2014). The primary outcome of a phase IIa study for AIR645 (NCT00941577) was not met, and Altair Therapeutics was eventually shut down. In contrast, the efficacy of ASM8 at phase II has not led to the termination of the programme. A follow up phase II study was withdrawn before enrolment in 2012 (NCT01380236), however ASM8 is listed as an active programme by Pharmaxis. Importantly, none of the adverse events observed with ASM8 were definitively drug-related despite the dose-dependent effect on alveolar macrophages, and no complement activation was observed (G M Gauvreau et al., 2011). Thus, without recourse to drug delivery systems, the FANA antisense chemistry remains presently the most promising candidate in realising the clinical potential of oligonucleotide therapeutics for the lung. However, in the absence of explicit 5'-RACE data it is presently uncertain that the action of ASM8 can be ascribed, fully or in part, on an on-target MOA.

In moving forward, it will also be important to take into consideration that both 2<sup>nd</sup> and 3<sup>rd</sup> generation ASO are principally loaded in alveolar macrophages, and administration of 2<sup>nd</sup> generation ASO (at least) promotes macrophage accumulation in the airways. Indeed, concerns around lung macrophage effects have been upgraded to promote use of exploratory inflammatory / toxicology biomarkers (Alton et al., 2012) as encouraged by

Cook *et al.* To date, the activation status of resident and recruited macrophages in the lung and any chronic tissue effects remain unclear (Forbes *et al.*, 2014). Crucially, macrophage recruitment may result in respiratory disease exacerbation (Zaslona *et al.*, 2014), an observation of particular relevance to proposed anti-inflammatory antisense therapeutics with documented macrophage recruitment propensity such as both the Ionis and Pharmaxis chemistries. Moreover, 3<sup>rd</sup> generation ASO have been very recently shown to harbour an off-target class effect that mediates long (>125kb) transcript degradation in an RNase H-dependent manner (Burel *et al.*, 2015) con-currently to on-target MOA. Thus, clinical success for the oligonucleotide therapeutics class in lung disease will pivot on thorough exploratory toxicology studies (Cook *et al.*, 2014; Forbes *et al.*, 2014) *in vitro*, *in vivo* and following chronic exposure trials that effectively compartmentalise on-target value proposition against off-target effect tolerability.


Thus far, the history of oligonucleotide therapeutics shows that success hinges on proactive pursuit of evidence closely aligned with the so-called 5 pillars of drug discovery (Cook *et al.*, 2014). The shortest route to the clinic more often than not hides pharmacological challenges that can lead to costly, late stage failure. This risk can be managed either through documented design approaches (e.g. TLR7/8 avoidance) or systematic screening towards clear, hypothesis-driven and exploratory elimination of toxicological and off-target risks. Key barriers to success have been perpetuated assumptions around adequate drug exposure (pillar 1) and molecular target engagement (pillar 2) extrapolated from tissue loading data, as well as neglect over MOA and off-target effect quantification (pillar 3). This is aligned to the unfortunate continuation of 'dogma' adoption, such as the artificial segregation of the molecular functions of slicer-mediated RNAi and miRNA-mediated RNAi (e.g. (Lorenzer,

Dirin, Winkler, Baumann, & Winkler, 2015), despite calls against this (Moschos, 2013; Sabin et al., 2013). Fortunately, companies focusing on oligonucleotide therapeutics are almost invariably accurate in selecting the appropriate patient population (pillar 4). However, in the absence of orphan indications, as is the case in many lung diseases, the apparent need for delivery solutions makes the commercial potential (pillar 5) of this drug class an exceptionally challenging proposition. The true impact of the foamy alveolar macrophage, and the true need for solutions circumventing its' elicitation is likely to catalyse ultimate success.


**Conflict of Interest Statement**

The authors declare that there are no conflicts of interest.

## Figures



**Figure 1: Anatomy of the lung, airway dynamics and drug delivery.** A) The nasopharynx, trachea and first bronchial bifurcation (upper lung / large airways ( $>2\ \text{mm}$  diameter)) lead to conducting airways ( $<2\ \text{mm}$  diameter: bronchi and bronchioles) that split into alveolar ducts (lower / deep lung or small / peripheral airways) (Labiris & Dolovich, 2003). Alveolar ducts conclude into alveolar sacs wherein gaseous exchange takes place. Bronchi and bronchioles are ringed tangentially by smooth muscle (R. J. Lorenz, 1966), which controls airflow through constriction and relaxation (Gosens & Grainge, 2015), whereas alveolar sacs are surrounded by capillaries, forming a short diffusion gradient for gas exchange. B) Particle dimensions and deposition mechanics by airway compartment (Heyder, 2012; Labiris & Dolovich, 2003); maximal efficiency is achieved with  $1\text{-}3\ \mu\text{m}$  particles, whereas  $<500\ \text{nm}$  particles may diffuse into the alveolar tissue where airflow is low, but are generally exhaled.




**Figure 2: Classes of oligonucleotide drugs used in the lung and their mode of action.**

Oligonucleotide therapeutics (bold black lines) recognize molecular targets (red lines) through Watson-Crick base pairing. (A) ASO form DNA:RNA heteroduplexes in a sense (target) – antisense (drug) manner, recruit RNase H and endonucleotically cleave target RNA to create a novel 5' end (bold), resulting to target degradation and translation inhibition. (B) Alternatively, translation inhibition can be directed by targeting ASOs to translation start sites in transcripts (AUG box), preventing access to eukaryotic translation initiation factors (eIF). Both of these MOAs are active on coding and ncRNA, with stoichiometric inhibition in the latter involving steric hindrances to nucleic acid or ribonucleoprotein complex formation (C) Premature stop codons, cryptic exons and nonsense mutations (etc.) can be 'skipped' by directing spliceosome activity. The clinical implementation in Duchenne's Muscular Dystrophy uses ASOs to stop the spliceosome from including an exon (red boxes) containing a premature stop mutation (vertical grey line), so that the translated protein may retain



some function (Becker phenotype) (Disterer et al., 2014; Q.-L. Lu et al., 2014; van Deutekom et al., 2007). (D) The minimal bioactive complex of human RNAi, the RNA-induced silencing complex (RISC), consists of a ~21 nt strand of RNA in length (black; 'active' or 'guide' strand (Gu et al., 2011; Khvorova, Reynolds, & Jayasena, 2003; Krol et al., 2004)) loaded into one of the four Argonaute (AGO) proteins (Kawamata & Tomari, 2010) from an endogenous (miRNA) or exogenous (siRNA) dsRNA precursor. The reverse complement 'passenger' strand (blue) is discarded. The outcome of target interaction with RISC depends on the AGO component, site of RISC interaction within the target (e.g. coding or untranslated regions (UTR)), degree of complementarity and subcellular complex localization, i.e. the cytosol (blue functions) or the nucleus (RED functions).



**Figure 3: Key backbone and nucleoside modifications used in oligonucleotide therapeutics.** (A) The most popular oligonucleotide modification (orange) of oligonucleotide backbones replaces one oxygen with a sulfur group. Morpholinos (PMO) have an uncharged backbone and six-membered rings replacing riboses that do not impact on solubility, appear to increase affinity (melting temperatures with RNA targets, ~20oC increase per ASO (Summerton & Weller, 1997)). The MOA of PMO is restricted to RNase H-free mechanisms (J. Summerton, 1999) and involves translation inhibition, splice switching and miRNA blockade (Eisen & Smith, 2008). Peptide nucleic acids (PNA) use entirely acyclic backbones conveying generally neutral charge and robust nuclease resistance and improve target affinity (Nielsen, Egholm, Berg, & Buchardt, 1991), evidenced even in siRNA modification (Gong & Desaulniers, 2012). They direct non-RNase H MOA (Uhlmann, 1998) in eukaryotes and prokaryotes (Pooga, Land, Bartfai, & Langel, 2001). (B) Nucleoside modifications have

principally involved substitution of the 2' H (DNA) or OH (RNA) group with more complicated alkyl groups, but also 2'-5' linking groups and 2' arabinose stereoisomers such as the 2'-deoxy-2'-fluoro-beta-D-arabinonucleic acid (2'-fluoroarabino, or FANA) chemistry.

A)

Phosphodiester




Phosphorothioate




B)

2'-modified




C)

2'-5' bicyclic



**Figure 4: The three generations of antisense oligonucleotides.** (A) Uniform modification of the ASO backbone from phosphorothioate (black line) to phosphodiester (green line; first generation) enabled better retention in circulation, significant loading in the liver and kidney and *in vivo* activity without delivery systems (Beltinger et al., 1995; Cossum et al., 1993; Crooke et al., 1996; Nolting et al., 1997) at a small cost of affinity for targets (Summerton & Weller, 1997). (B) Second generation, 2'-modified ASOs retain backbone modifications, are typically 20-30 nt long and are structured either with modifications (orange nucleosides) interspersed in the ASO (mixomers) or as a central 6-8 nucleoside stretch with no ribose modifications and flanked at the 5' and 3' ends with 2'- modified

nucleosides (gapmers). (C) Third generation 2'-5' chemistries (blue nucleosides) impart conformational rigidity that greatly enhances ASO affinity to their targets, increasing melting temperatures by as much as 4-8 °C per modified nucleoside (Kumar et al., 1998). Consequently, shorter ASO (e.g. 16-20 nt) can match or even outperform longer (20-30 nt) 2<sup>nd</sup> generation equivalents: bridged nucleosides force ASO not to tolerate base mismatches with their targets, thereby improving specificity (Valoczi et al., 2004). Short 100% modified octamers (tiny) may block entire miRNA classes with common seed sequences *en masse* (Obad et al., 2011).



**Figure 5: microRNA organization and maturation into bioactive RISC.** (A) miRNA arise from exons (blue boxes) or introns (light blue angled lines). Some short introns can encode miRNA (mirtrons), whereas other miRNA precursors arise from exon-intron junctions, or

span entire short exons (exon-overlap). Yet others are processed out of 5' and 3' UTRs. (B) Endogenous RNAi mediator transcripts typically form stem-loop hairpin structures (primary miRNA) excised by the microprocessor complex in the nucleus into pre-miRNA and actively transported to the cytosol by Exportin 5. Here, the loop is endonucleotically removed by Dicer to form miRNA and then loaded onto AGO to form RISC; guide strand selection is driven by the orientation of the duplex loading onto AGO. Notable updates involve mitochondrial ncRNA-derived miRNA (mito-miRs) (Jagannathan et al., 2015; Ro et al., 2013), as well as AGO-directed small ncRNA (AGO-taxic ncRNA) of distinct genomic origin, but uniquely equipped to select AGO partners in RISC formation (Yamakawa et al., 2014). Small interfering RNA and miRNA replacement therapy can be achieved by gene therapy approaches exploiting any of these genomic organization structures or by providing exogenous, synthetic RNA (e.g. hairpins, duplexes) that engage with the latter processing steps of miRNA biogenesis. Adapted from (Moschos, 2013).

**Table 1:** Innate immune receptors for RNA and DNA oligonucleotide therapeutics.

Receptor	Location	Minimal motif	Oligonucleotide class affected	Evasion solution	Reference(s)
TLR1	Cell surface	Unknown	miRNA analogues	Unknown	He et al. 2013
TLR3	Cell surface	> 19 nt	siRNA, dsRNA	N6-methyladenosine, 2-thiouridine	Cho et al., 2009; Kleinman et al., 2008; Karikó et al., 2005
TLR7/8	Cell surface	UG	siRNA, miRNA, ASO	2' ribose modification, N6-methyladenosine, 2-thiouridine	Hornung et al., 2005; Judge et al., 2005; Fedorov et al. 2006; Jackson et al. 2006
TLR9	Cell surface	unmethylated CpG motifs	DNA ASO	CpG methylation	Krieg et al. 1995; Rutz et al 2004
RIG-I	Cytosolic	blunt duplex ends; 5' triphosphate	siRNA, dsRNA	3' dinucleotide overhangs, no 5' triphosphates	Marques et al., 2006; Kato et al., 2008
MDA5	Cytosolic	> 30 nt	dsRNA	short sequences	Kato et al., 2008
PKR	Cytosolic	> 30 nt	dsRNA	short sequences	Sledz et al., 2003
OAS1	Cytosolic	NNWW(N <sub>9</sub> )WGN	siRNA, miRNA, RNA ASO	Design	Kodym et al., 2009
DAI	Cytosolic	> 30 nt	dsRNA	short sequences	Manche et al., 1992

**Table 2:** Types of negative controls for oligonucleotide therapeutics.

Control type*	Examples	Key features
<b>Scrambled</b>	5'-CUGGUUAGUGGCACUUCGAUU-3' to 5'-CGUGUUAGUGGCACUUCGAUU-3'	<ul style="list-style-type: none"> <li>• impact increases with scrambling amount</li> <li>• may require &gt;1 controls</li> <li>• avoid accidental host gene targeting</li> <li>• scramble 10-11 / central octamer to create slicer / RNase H control</li> </ul>
<b>Mismatch</b>	5'-AUGGUUAGUGGCACUUCGAUU-3' to 5'-ACUGUUAGUGGCACUUCGAUU-3'	<ul style="list-style-type: none"> <li>• seed region placement alters off targets**</li> <li>• slicer / RNase H active site inhibition</li> <li>• &gt;1 controls also recommended</li> </ul>
<b>Reverse</b>	5'-CUGGUUAGUGGCACUUCGAUU-3' to 5'-UUAGCUUCACGGUGAUUGGUC-3'	<ul style="list-style-type: none"> <li>• check for unintended host hits</li> <li>• no complementarity to intended target</li> </ul>
<b>Validated bioactive oligo</b>	Green fluorescent protein, luciferase, <i>Arabidopsis thaliana</i> , <i>C. elegans</i>	<ul style="list-style-type: none"> <li>• target not normally found in host transcriptome</li> <li>• confirm lack of complementarity</li> <li>• confirm no seed sequence in pathway of interest</li> </ul>
<b>Commercial negative control</b>	Research supplier 'universal' negative controls	<ul style="list-style-type: none"> <li>• limited 'negative' validation</li> <li>• unknown modifications</li> <li>• unknown sequence</li> </ul>

\*: Must match design and modification criteria of active oligonucleotide.

\*\* : Relevant to RNAi mediators only.

## References

- Advani, R., Lum, B. L., Fisher, G. A., Halsey, J., Geary, R. S., Holmlund, J. T., ... Sikic, B. I. (2005). A phase I trial of aprinocarsen (ISIS 3521/LY900003), an antisense inhibitor of protein kinase C- $\alpha$  administered as a 24-hour weekly infusion schedule in patients with advanced cancer. *Investigational New Drugs*, 23(5), 467–77. <http://doi.org/10.1007/s10637-005-2906-0>
- Ahlenstiel, C. L., Lim, H. G. W., Cooper, D. A., Ishida, T., Kelleher, A. D., & Suzuki, K. (2012). Direct evidence of nuclear Argonaute distribution during transcriptional silencing links the actin cytoskeleton to nuclear RNAi machinery in human cells. *Nucleic Acids Research*, 40(4), 1579–95. <http://doi.org/10.1093/nar/gkr891>
- Ahn, D.-G., Lee, W., Choi, J.-K., Kim, S.-J., Plant, E. P., Almazán, F., ... Oh, J.-W. (2011). Interference of ribosomal frameshifting by antisense peptide nucleic acids suppresses SARS coronavirus replication. *Antiviral Research*, 91(1), 1–10. <http://doi.org/10.1016/j.antiviral.2011.04.009>
- Allakhverdi, Z., Allam, M., Guimond, A., Ferrari, N., Zemzoumi, K., Séguin, R., ... Renzi, P. M. (2006). Multitargeted approach using antisense oligonucleotides for the treatment of asthma. *Annals of the New York Academy of Sciences*, 1082, 62–73. <http://doi.org/10.1196/annals.1348.047>
- Allakhverdi, Z., Allam, M., & Renzi, P. M. (2002). Inhibition of antigen-induced eosinophilia and airway hyperresponsiveness by antisense oligonucleotides directed against the common beta chain of IL-3, IL-5, GM-CSF receptors in a rat model of allergic asthma. *American Journal of Respiratory and Critical Care Medicine*, 165(7), 1015–21. <http://doi.org/10.1164/ajrccm.165.7.2109095>
- Allam, M., & Renzi, P. M. (2001). Inhibition of GM-CSF/IL-3/IL-5 signaling by antisense oligodeoxynucleotides targeting the common beta chain of their receptors. *Antisense & Nucleic Acid Drug Development*, 11(5), 289–300. <http://doi.org/10.1089/108729001753231678>
- Alló, M., Buggiano, V., Fededa, J. P., Petrillo, E., Schor, I., de la Mata, M., ... Kornblihtt, A. R. (2009). Control of alternative splicing through siRNA-mediated transcriptional gene silencing. *Nature Structural & Molecular Biology*, 16(7), 717–24. <http://doi.org/10.1038/nsmb.1620>
- Alton, E. W., Boushey, H. A., Garn, H., Green, F. H., Hodges, M., Martin, R. J., ... Ferrari, N. (2012). Clinical expert panel on monitoring potential lung toxicity of inhaled oligonucleotides: consensus points and recommendations. *Nucleic Acid Therapeutics*, 22(4), 246–54. <http://doi.org/10.1089/nat.2012.0345>
- Alvarez, R., Elbashir, S., Borland, T., Toudjarska, I., Hadwiger, P., John, M., ... Meyers, R. (2009). RNA Interference-Mediated Silencing of the Respiratory Syncytial Virus Nucleocapsid Defines a Potent Antiviral Strategy. *Antimicrobial Agents and Chemotherapy*, 53(9), 3952–3962. <http://doi.org/10.1128/AAC.00014-09>
- Ameyar-Zazoua, M., Rachez, C., Souidi, M., Robin, P., Fritsch, L., Young, R., ... Harel-



- Bellan, A. (2012). Argonaute proteins couple chromatin silencing to alternative splicing. *Nature Structural & Molecular Biology*, *19*(10), 998–1004. <http://doi.org/10.1038/nsmb.2373>
- Arick, D. Q., Choi, Y. H., Kim, H. C., & Won, Y.-Y. (2015). Effects of nanoparticles on the mechanical functioning of the lung. *Advances in Colloid and Interface Science*, *225*, 218–28. <http://doi.org/10.1016/j.cis.2015.10.002>
- Augagneur, Y., Wesolowski, D., Tae, H. S., Altman, S., & Ben Mamoun, C. (2012). Gene selective mRNA cleavage inhibits the development of *Plasmodium falciparum*. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(16), 6235–40. <http://doi.org/10.1073/pnas.1203516109>
- Azuma-Mukai, A., Oguri, H., Mituyama, T., Qian, Z. R., Asai, K., Siomi, H., & Siomi, M. C. (2008). Characterization of endogenous human Argonautes and their miRNA partners in RNA silencing. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(23), 7964–9. <http://doi.org/10.1073/pnas.0800334105>
- Bai, B., Liu, H., & Laiho, M. (2014). Small RNA expression and deep sequencing analyses of the nucleolus reveal the presence of nucleolus-associated microRNAs. *FEBS Open Bio*, *4*, 441–9. <http://doi.org/10.1016/j.fob.2014.04.010>
- Beltinger, C., Saragovi, H. U., Smith, R. M., LeSauter, L., Shah, N., DeDionisio, L., ... Gewirtz, A. M. (1995). Binding, uptake, and intracellular trafficking of phosphorothioate-modified oligodeoxynucleotides. *The Journal of Clinical Investigation*, *95*(4), 1814–23. <http://doi.org/10.1172/JCI117860>
- Benimetskaya, L., Loike, J. D., Khaled, Z., LoiKe, G., Silverstein, S. C., Cao, L., ... Stein, C. A. (1997). Mac-1 (CD11b/CD18) is an oligodeoxynucleotide-binding protein. *Nature Medicine*, *3*(4), 414–420. <http://doi.org/10.1038/nm0497-414>
- Bennett, C. F., Chiang, M. Y., Chan, H., Shoemaker, J. E., & Mirabelli, C. K. (1992). Cationic lipids enhance cellular uptake and activity of phosphorothioate antisense oligonucleotides. *Molecular Pharmacology*, *41*(6), 1023–33. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/1352033>
- Bevilacqua, P. C., & Cech, T. R. (1996). Minor-groove recognition of double-stranded RNA by the double-stranded RNA-binding domain from the RNA-activated protein kinase PKR. *Biochemistry*, *35*(31), 9983–94. <http://doi.org/10.1021/bi9607259>
- Bijsterbosch, M. K., Rump, E. T., De Vruet, R. L., Dorland, R., van Veghel, R., Tivel, K. L., ... Manoharan, M. (2000). Modulation of plasma protein binding and in vivo liver cell uptake of phosphorothioate oligodeoxynucleotides by cholesterol conjugation. *Nucleic Acids Research*, *28*(14), 2717–25. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=102653&tool=pmcentrez&rendertype=abstract>
- Bitko, V., Musiyenko, A., Shulyayeva, O., & Barik, S. (2005). Inhibition of respiratory viruses by nasally administered siRNA. *Nature Medicine*, *11*(1), 50–5. <http://doi.org/10.1038/nm1164>
- Blanco, O., Lugones, Y., Diaz, E., & Monzote, L. In vitro activity of the clinical pulmonary surfactant Surfactan® against *Leishmania amazonensis*. *Revista Do Instituto de Medicina Tropical de São Paulo*, *53*(4), 235–8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21915470>

- Bombieri, C., Claustres, M., De Boeck, K., Derichs, N., Dodge, J., Girodon, E., ... Ferec, C. (2011). Recommendations for the classification of diseases as CFTR-related disorders. *Journal of Cystic Fibrosis : Official Journal of the European Cystic Fibrosis Society*, 10 Suppl 2, S86–102. [http://doi.org/10.1016/S1569-1993\(11\)60014-3](http://doi.org/10.1016/S1569-1993(11)60014-3)
- Bousquet, J., Chanez, P., Lacoste, J. Y., Barnéon, G., Ghavanian, N., Enander, I., ... Godard, P. (1990). Eosinophilic inflammation in asthma. *The New England Journal of Medicine*, 323(15), 1033–9. <http://doi.org/10.1056/NEJM199010113231505>
- Broering, R., Real, C. I., John, M. J., Jahn-Hofmann, K., Ickenstein, L. M., Kleinehr, K., ... Schlaak, J. F. (2014). Chemical modifications on siRNAs avoid Toll-like-receptor-mediated activation of the hepatic immune system in vivo and in vitro. *International Immunology*, 26(1), 35–46. <http://doi.org/10.1093/intimm/dxt040>
- Brolin, C., Shiraishi, T., Hojman, P., Krag, T. O., Nielsen, P. E., & Gehl, J. (2015). Electroporation Enhanced Effect of Dystrophin Splice Switching PNA Oligomers in Normal and Dystrophic Muscle. *Molecular Therapy. Nucleic Acids*, 4, e267. <http://doi.org/10.1038/mtna.2015.41>
- Burel, S. A., Han, S.-R., Lee, H.-S., Norris, D. A., Lee, B.-S., Machemer, T., ... Henry, S. P. (2013). Preclinical evaluation of the toxicological effects of a novel constrained ethyl modified antisense compound targeting signal transducer and activator of transcription 3 in mice and cynomolgus monkeys. *Nucleic Acid Therapeutics*, 23(3), 213–27. <http://doi.org/10.1089/nat.2013.0422>
- Burel, S. A., Hart, C. E., Cauntay, P., Hsiao, J., Machemer, T., Katz, M., ... Henry, S. P. (2015). Hepatotoxicity of high affinity gapmer antisense oligonucleotides is mediated by RNase H1 dependent promiscuous reduction of very long pre-mRNA transcripts. *Nucleic Acids Research*, gkv1210–. <http://doi.org/10.1093/nar/gkv1210>
- Burroughs, A. M., Ando, Y., de Hoon, M. J. L., Tomaru, Y., Suzuki, H., Hayashizaki, Y., & Daub, C. O. Deep-sequencing of human Argonaute-associated small RNAs provides insight into miRNA sorting and reveals Argonaute association with RNA fragments of diverse origin. *RNA Biology*, 8(1), 158–77. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3127082&tool=pmcentrez&rendertype=abstract>
- Cameron, L., Christodoulopoulos, P., Lavigne, F., Nakamura, Y., Eidelman, D., McEuen, A., ... Hamid, Q. (2000). Evidence for local eosinophil differentiation within allergic nasal mucosa: inhibition with soluble IL-5 receptor. *Journal of Immunology (Baltimore, Md. : 1950)*, 164(3), 1538–45. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10640772>
- Castanotto, D., Lin, M., Kowolik, C., Wang, L., Ren, X.-Q., Soifer, H. S., ... Stein, C. A. (2015). A cytoplasmic pathway for gapmer antisense oligonucleotide-mediated gene silencing in mammalian cells. *Nucleic Acids Research*, 43(19), 9350–61. <http://doi.org/10.1093/nar/gkv964>
- Castanotto, D., Lingeman, R., Riggs, A. D., & Rossi, J. J. (2009). CRM1 mediates nuclear-cytoplasmic shuttling of mature microRNAs. *Proceedings of the National Academy of Sciences of the United States of America*, 106(51), 21655–9. <http://doi.org/10.1073/pnas.0912384106>
- Castro, M., Wenzel, S. E., Bleecker, E. R., Pizzichini, E., Kuna, P., Busse, W. W., ... Raible, D. G. (2014). Benralizumab, an anti-interleukin 5 receptor  $\alpha$  monoclonal antibody,

- versus placebo for uncontrolled eosinophilic asthma: a phase 2b randomised dose-ranging study. *The Lancet. Respiratory Medicine*, 2(11), 879–90.  
[http://doi.org/10.1016/S2213-2600\(14\)70201-2](http://doi.org/10.1016/S2213-2600(14)70201-2)
- Cerritelli, S. M., & Crouch, R. J. (2009). Ribonuclease H: the enzymes in eukaryotes. *The FEBS Journal*, 276(6), 1494–505. <http://doi.org/10.1111/j.1742-4658.2009.06908.x>
- Chatila, T. A. (2004). Interleukin-4 receptor signaling pathways in asthma pathogenesis. *Trends in Molecular Medicine*, 10(10), 493–9.  
<http://doi.org/10.1016/j.molmed.2004.08.004>
- Chen, G., Kronenberger, P., Teugels, E., & De Grève, J. (2011). Influence of RT-qPCR primer position on EGFR interference efficacy in lung cancer cells. *Biological Procedures Online*, 13, 1. <http://doi.org/10.1186/1480-9222-13-1>
- Chen, L., Dahlstrom, J. E., Lee, S.-H., & Rangasamy, D. (2012). Naturally occurring endo-siRNA silences LINE-1 retrotransposons in human cells through DNA methylation. *Epigenetics*, 7(7), 758–71. <http://doi.org/10.4161/epi.20706>
- Chi, S. W., Hannon, G. J., & Darnell, R. B. (2012). An alternative mode of microRNA target recognition. *Nature Structural & Molecular Biology*, 19(3), 321–7.  
<http://doi.org/10.1038/nsmb.2230>
- Chi, S. W., Zang, J. B., Mele, A., & Darnell, R. B. (2009). Argonaute HITS-CLIP decodes microRNA-mRNA interaction maps. *Nature*, 460(7254), 479–486.  
<http://doi.org/10.1038/nature08170>
- Cho, W. G., Albuquerque, R. J. C., Kleinman, M. E., Tarallo, V., Greco, A., Nozaki, M., ... Ambati, J. (2009). Small interfering RNA-induced TLR3 activation inhibits blood and lymphatic vessel growth. *Proceedings of the National Academy of Sciences of the United States of America*, 106(17), 7137–42. <http://doi.org/10.1073/pnas.0812317106>
- Chroneos, Z. C., Sever-Chroneos, Z., & Shepherd, V. L. (2010). Pulmonary surfactant: an immunological perspective. *Cellular Physiology and Biochemistry: International Journal of Experimental Cellular Physiology, Biochemistry, and Pharmacology*, 25(1), 13–26. <http://doi.org/10.1159/000272047>
- Clark, K. L., Hughes, S. A., Bulsara, P., Coates, J., Moores, K., Parry, J., ... Edbrooke, M. R. (2013). Pharmacological Characterization of a Novel ENaC $\alpha$  siRNA (GSK2225745) With Potential for the Treatment of Cystic Fibrosis. *Molecular Therapy. Nucleic Acids*, 2, e65. <http://doi.org/10.1038/mtna.2012.57>
- Cook, D., Brown, D., Alexander, R., March, R., Morgan, P., Satterthwaite, G., & Pangalos, M. N. (2014). Lessons learned from the fate of AstraZeneca's drug pipeline: a five-dimensional framework. *Nature Reviews. Drug Discovery*, 13(6), 419–31.  
<http://doi.org/10.1038/nrd4309>
- Cordier, C., Boutimah, F., Bourdeloux, M., Dupuy, F., Met, E., Alberti, P., ... Saison-Behmoaras, T. E. (2014). Delivery of antisense peptide nucleic acids to cells by conjugation with small arginine-rich cell-penetrating peptide (R/W)9. *PloS One*, 9(8), e104999. <http://doi.org/10.1371/journal.pone.0104999>
- Corren, J. (2012). Inhibition of Interleukin-5 for the Treatment of Eosinophilic Diseases. *Discovery Medicine*, 13(71), 305–312. Retrieved from <http://www.discoverymedicine.com/Jonathan-Corren/2012/04/24/inhibition-of-interleukin-5-for-the-treatment-of-eosinophilic-diseases/>

- Cossum, P. A., Sasmor, H., Dellinger, D., Truong, L., Cummins, L., Owens, S. R., ... Crooke, S. (1993). Disposition of the 14C-labeled phosphorothioate oligonucleotide ISIS 2105 after intravenous administration to rats. *The Journal of Pharmacology and Experimental Therapeutics*, 267(3), 1181–90. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8166890>
- Crooke, S. T. (1999). Molecular mechanisms of action of antisense drugs. *Biochimica et Biophysica Acta*, 1489(1), 31–44. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10806995>
- Crooke, S. T., Graham, M. J., Zuckerman, J. E., Brooks, D., Conklin, B. S., Cummins, L. L., ... Griffey, R. H. (1996). Pharmacokinetic properties of several novel oligonucleotide analogs in mice. *The Journal of Pharmacology and Experimental Therapeutics*, 277(2), 923–37. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8627575>
- Daige, C. L., Wiggins, J. F., Priddy, L., Nelligan-Davis, T., Zhao, J., & Brown, D. (2014). Systemic delivery of a miR34a mimic as a potential therapeutic for liver cancer. *Molecular Cancer Therapeutics*, 13(10), 2352–60. <http://doi.org/10.1158/1535-7163.MCT-14-0209>
- Davies, D. E. (2014). Epithelial barrier function and immunity in asthma. *Annals of the American Thoracic Society*, 11 Suppl 5, S244–51. <http://doi.org/10.1513/AnnalsATS.201407-304AW>
- De Backer, L., Cerrada, A., Pérez-Gil, J., De Smedt, S. C., & Raemdonck, K. (2015). Bio-inspired materials in drug delivery: Exploring the role of pulmonary surfactant in siRNA inhalation therapy. *Journal of Controlled Release : Official Journal of the Controlled Release Society*. <http://doi.org/10.1016/j.jconrel.2015.09.004>
- de Hoon, M., Shin, J. W., & Carninci, P. (2015). Paradigm shifts in genomics through the FANTOM projects. *Mammalian Genome : Official Journal of the International Mammalian Genome Society*, 26(9-10), 391–402. <http://doi.org/10.1007/s00335-015-9593-8>
- Denise, H., Moschos, S. A., Sidders, B., Burden, F., Perkins, H., Carter, N., ... Corbau, R. (2013). Deep sequencing insights in therapeutic shRNA processing and siRNA target cleavage precision. *Molecular Therapy Nucleic Acids*, in press.
- Denyer, J., & Dyche, T. (2010). The Adaptive Aerosol Delivery (AAD) technology: Past, present, and future. *Journal of Aerosol Medicine and Pulmonary Drug Delivery*, 23 Suppl 1, S1–10. <http://doi.org/10.1089/jamp.2009.0791>
- Derscheid, R. J., & Ackermann, M. R. (2013). The innate immune system of the perinatal lung and responses to respiratory syncytial virus infection. *Veterinary Pathology*, 50(5), 827–41. <http://doi.org/10.1177/0300985813480216>
- DeVincenzo, J., Cehelsky, J. E., Alvarez, R., Elbashir, S., Harborth, J., Toudjarska, I., ... Meyers, R. (2008). Evaluation of the safety, tolerability and pharmacokinetics of ALN-RSV01, a novel RNAi antiviral therapeutic directed against respiratory syncytial virus (RSV). *Antiviral Research*, 77(3), 225–31. <http://doi.org/10.1016/j.antiviral.2007.11.009>
- DeVincenzo, J., Lambkin-Williams, R., Wilkinson, T., Cehelsky, J., Nochur, S., Walsh, E., ... Vaishnav, A. (2010a). A randomized, double-blind, placebo-controlled study of an RNAi-based therapy directed against respiratory syncytial virus. *Proceedings of the National Academy of Sciences of the United States of America*, 107(19), 8800–5. <http://doi.org/10.1073/pnas.0912186107>

- DeVincenzo, J., Lambkin-Williams, R., Wilkinson, T., Cehelsky, J., Nochur, S., Walsh, E., ... Vaishnav, A. (2010b). A randomized, double-blind, placebo-controlled study of an RNAi-based therapy directed against respiratory syncytial virus. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(19), 8800–5. <http://doi.org/10.1073/pnas.0912186107>
- Dietrich, A., Wallet, C., Iqbal, R. K., Gualberto, J. M., & Lotfi, F. (2015). Organellar non-coding RNAs: Emerging regulation mechanisms. *Biochimie*, *117*, 48–62. <http://doi.org/10.1016/j.biochi.2015.06.027>
- Dirin, M., & Winkler, J. (2013). Influence of diverse chemical modifications on the ADME characteristics and toxicology of antisense oligonucleotides. *Expert Opinion on Biological Therapy*, *13*(6), 875–88. <http://doi.org/10.1517/14712598.2013.774366>
- Disterer, P., Kryczka, A., Liu, Y., Badi, Y. E., Wong, J. J., Owen, J. S., & Khoo, B. (2014). Development of therapeutic splice-switching oligonucleotides. *Human Gene Therapy*, *25*(7), 587–98. <http://doi.org/10.1089/hum.2013.234>
- Dittmar, W. J., McIver, L., Michalak, P., Garner, H. R., & Valdez, G. (2014). EvoCor: a platform for predicting functionally related genes using phylogenetic and expression profiles. *Nucleic Acids Research*, *42*(Web Server issue), W72–5. <http://doi.org/10.1093/nar/gku442>
- Djebali, S., Davis, C. A., Merkel, A., Dobin, A., Lassmann, T., Mortazavi, A., ... Gingeras, T. R. (2012). Landscape of transcription in human cells. *Nature*, *489*(7414), 101–8. <http://doi.org/10.1038/nature11233>
- Dorman, S. C., Efthimiadis, A., Babirad, I., Watson, R. M., Denburg, J. A., Hargreave, F. E., ... Sehmi, R. (2004). Sputum CD34+IL-5Ralpha+ cells increase after allergen: evidence for in situ eosinophilopoiesis. *American Journal of Respiratory and Critical Care Medicine*, *169*(5), 573–7. <http://doi.org/10.1164/rccm.200307-1004OC>
- Dowell, M. L., Lavoie, T. L., Solway, J., & Krishnan, R. (2014). Airway smooth muscle: a potential target for asthma therapy. *Current Opinion in Pulmonary Medicine*, *20*(1), 66–72. <http://doi.org/10.1097/MCP.000000000000011>
- Dragulescu-Andrasi, A., Rapireddy, S., He, G., Bhattacharya, B., Hyldig-Nielsen, J. J., Zon, G., & Ly, D. H. (2006). Cell-permeable peptide nucleic acid designed to bind to the 5'-untranslated region of E-cadherin transcript induces potent and sequence-specific antisense effects. *Journal of the American Chemical Society*, *128*(50), 16104–12. <http://doi.org/10.1021/ja063383v>
- Duan, W., Chan, J. H. P., McKay, K., Crosby, J. R., Choo, H. H., Leung, B. P., ... Wong, W. S. F. (2005). Inhaled p38alpha mitogen-activated protein kinase antisense oligonucleotide attenuates asthma in mice. *American Journal of Respiratory and Critical Care Medicine*, *171*(6), 571–8. <http://doi.org/10.1164/rccm.200408-1006OC>
- Eisen, J. S., & Smith, J. C. (2008). Controlling morpholino experiments: don't stop making antisense. *Development (Cambridge, England)*, *135*(10), 1735–43. <http://doi.org/10.1242/dev.001115>
- El Saleeby, C. M., Bush, A. J., Harrison, L. M., Aitken, J. A., & DeVincenzo, J. P. (2011). Respiratory syncytial virus load, viral dynamics, and disease severity in previously healthy naturally infected children. *The Journal of Infectious Diseases*, *204*(7), 996–1002. <http://doi.org/10.1093/infdis/jir494>

- Ellipilli, S., & Ganesh, K. N. (2015). Fluorous Peptide Nucleic Acids: PNA Analogues with Fluorine in Backbone ( $\gamma$ -CF<sub>2</sub>-app-PNA) Enhance Cellular Uptake. *The Journal of Organic Chemistry*, 80(18), 9185–91. <http://doi.org/10.1021/acs.joc.5b01614>
- Elmén, J., Lindow, M., Schütz, S., Lawrence, M., Petri, A., Obad, S., ... Kauppinen, S. (2008). LNA-mediated microRNA silencing in non-human primates. *Nature*, 452(7189), 896–9. <http://doi.org/10.1038/nature06783>
- Elmén, J., Lindow, M., Silahtaroglu, A., Bak, M., Christensen, M., Lind-Thomsen, A., ... Kauppinen, S. (2008). Antagonism of microRNA-122 in mice by systemically administered LNA-antimiR leads to up-regulation of a large set of predicted target mRNAs in the liver. *Nucleic Acids Research*, 36(4), 1153–62. <http://doi.org/10.1093/nar/gkm1113>
- Erhard, F., & Zimmer, R. (2015). Count ratio model reveals bias affecting NGS fold changes. *Nucleic Acids Research*, 43(20), e136. <http://doi.org/10.1093/nar/gkv696>
- Erle, D. J., & Sheppard, D. (2014). The cell biology of asthma. *The Journal of Cell Biology*, 205(5), 621–31. <http://doi.org/10.1083/jcb.201401050>
- Ezzat, K., Aoki, Y., Koo, T., McClorey, G., Benner, L., Coenen-Stass, A., ... Wood, M. J. A. (2015). Self-Assembly into Nanoparticles Is Essential for Receptor Mediated Uptake of Therapeutic Antisense Oligonucleotides. *Nano Letters*, 15(7), 4364–73. <http://doi.org/10.1021/acs.nanolett.5b00490>
- Ezzat, K., Helmfors, H., Tudoran, O., Juks, C., Lindberg, S., Padari, K., ... Langel, U. (2012). Scavenger receptor-mediated uptake of cell-penetrating peptide nanocomplexes with oligonucleotides. *FASEB Journal : Official Publication of the Federation of American Societies for Experimental Biology*, 26(3), 1172–80. <http://doi.org/10.1096/fj.11-191536>
- Fabani, M. M., & Gait, M. J. (2008). miR-122 targeting with LNA/2'-O-methyl oligonucleotide mixmers, peptide nucleic acids (PNA), and PNA-peptide conjugates. *RNA (New York, N.Y.)*, 14(2), 336–46. <http://doi.org/10.1261/rna.844108>
- Fabbri, M., Paone, A., Calore, F., Galli, R., Gaudio, E., Santhanam, R., ... Croce, C. M. (2012). MicroRNAs bind to Toll-like receptors to induce prometastatic inflammatory response. *Proceedings of the National Academy of Sciences of the United States of America*, 109(31), E2110–6. <http://doi.org/10.1073/pnas.1209414109>
- Fan, B., Sutandy, F. X. R., Syu, G.-D., Middleton, S., Yi, G., Lu, K.-Y., ... Kao, C. C. (2015). Heterogeneous ribonucleoprotein K (hnRNP K) binds miR-122, a mature liver-specific microRNA required for hepatitis C virus replication. *Molecular & Cellular Proteomics : MCP*. <http://doi.org/10.1074/mcp.M115.050344>
- Fang, Z., & Rajewsky, N. (2011). The impact of miRNA target sites in coding sequences and in 3'UTRs. *PLoS One*, 6(3), e18067. <http://doi.org/10.1371/journal.pone.0018067>
- Faro-Trindade, I., Willment, J. A., Kerrigan, A. M., Redelinghuys, P., Hadebe, S., Reid, D. M., ... Brown, G. D. (2012). Characterisation of innate fungal recognition in the lung. *PLoS One*, 7(4), e35675. <http://doi.org/10.1371/journal.pone.0035675>
- Fedorov, Y., Anderson, E. M., Birmingham, A., Reynolds, A., Karpilow, J., Robinson, K., ... Khvorova, A. (2006). Off-target effects by siRNA can induce toxic phenotype. *RNA (New York, N.Y.)*, 12(7), 1188–96. <http://doi.org/10.1261/rna.28106>

- Fennewald, S. M., & Rando, R. F. (1995). Inhibition of High Affinity Basic Fibroblast Growth Factor Binding by Oligonucleotides. *Journal of Biological Chemistry*, 270(37), 21718–21721. <http://doi.org/10.1074/jbc.270.37.21718>
- Fey, R. A., Templin, M. V., McDonald, J. D., Yu, R. Z., Hutt, J. A., Gigliotti, A. P., ... Reed, M. D. (2014). Local and systemic tolerability of a 2'-O-methoxyethyl antisense oligonucleotide targeting interleukin-4 receptor- $\alpha$  delivery by inhalation in mouse and monkey. *Inhalation Toxicology*. Retrieved from <http://www.tandfonline.com/doi/abs/10.3109/08958378.2014.907587>
- Fields, R. J., Quijano, E., McNeer, N. A., Caputo, C., Bahal, R., Anandalingam, K., ... Saltzman, W. M. (2015). Modified poly(lactic-co-glycolic acid) nanoparticles for enhanced cellular uptake and gene editing in the lung. *Advanced Healthcare Materials*, 4(3), 361–6. <http://doi.org/10.1002/adhm.201400355>
- Filipowicz, W., Bhattacharyya, S. N., & Sonenberg, N. (2008). Mechanisms of post-transcriptional regulation by microRNAs: are the answers in sight? *Nature Reviews. Genetics*, 9(2), 102–14. <http://doi.org/10.1038/nrg2290>
- Fire, A., Xu, S., Montgomery, M. K., Kostas, S. A., Driver, S. E., & Mello, C. C. (1998). Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*, 391(6669), 806–11. <http://doi.org/10.1038/35888>
- Földes-Papp, Z., König, K., Studier, H., Bückle, R., Breunig, H. G., Uchugonova, A., & Kostner, G. M. (2009). Trafficking of mature miRNA-122 into the nucleus of live liver cells. *Current Pharmaceutical Biotechnology*, 10(6), 569–78. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19619125>
- Forbes, B., O'Lone, R., Allen, P. P., Cahn, A., Clarke, C., Collinge, M., ... Wolfreys, A. (2014). Challenges for inhaled drug discovery and development: Induced alveolar macrophage responses. *Advanced Drug Delivery Reviews*, 71, 15–33. <http://doi.org/10.1016/j.addr.2014.02.001>
- Forsbach, A., Müller, C., Montino, C., Kritzler, A., Curdt, R., Benahmed, A., ... Vollmer, J. (2012). Impact of delivery systems on siRNA immune activation and RNA interference. *Immunology Letters*, 141(2), 169–80. <http://doi.org/10.1016/j.imlet.2011.10.001>
- Forsbach, A., Nemorin, J.-G., Montino, C., Müller, C., Samulowitz, U., Vicari, A. P., ... Vollmer, J. (2008). Identification of RNA sequence motifs stimulating sequence-specific TLR8-dependent immune responses. *Journal of Immunology (Baltimore, Md. : 1950)*, 180(6), 3729–38. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18322178>
- Fortin, M., Ferrari, N., Higgins, M.-E., Séguin, S., Allam, M., Allakhverdi, Z., ... Renzi, P. M. (2006). Effects of antisense oligodeoxynucleotides targeting CCR3 on the airway response to antigen in rats. *Oligonucleotides*, 16(3), 203–12. <http://doi.org/10.1089/oli.2006.16.203>
- Francis, H., McDaniel, K., Han, Y., Liu, X., Kennedy, L., Yang, F., ... Meng, F. (2014). Regulation of the extrinsic apoptotic pathway by microRNA-21 in alcoholic liver injury. *The Journal of Biological Chemistry*, 289(40), 27526–39. <http://doi.org/10.1074/jbc.M114.602383>
- Friedman, K. J., Kole, J., Cohn, J. A., Knowles, M. R., Silverman, L. M., & Kole, R. (1999). Correction of aberrant splicing of the cystic fibrosis transmembrane conductance regulator (CFTR) gene by antisense oligonucleotides. *The Journal of Biological Chemistry*, 274(51), 36193–9. Retrieved from

<http://www.ncbi.nlm.nih.gov/pubmed/10593905>

- Gagnon, K. T., Li, L., Chu, Y., Janowski, B. A., & Corey, D. R. (2014). RNAi factors are present and active in human cell nuclei. *Cell Reports*, 6(1), 211–21. <http://doi.org/10.1016/j.celrep.2013.12.013>
- Gao, X., Shen, X., Dong, X., Ran, N., Han, G., Cao, L., ... Yin, H. (2015). Peptide Nucleic Acid Promotes Systemic Dystrophin Expression and Functional Rescue in Dystrophin-deficient mdx Mice. *Molecular Therapy. Nucleic Acids*, 4, e255. <http://doi.org/10.1038/mtna.2015.27>
- Gaunsbaek, M. Q., Kjeldsen, A. D., Svane-Knudsen, V., Henriksen, M. L., & Hansen, S. (2014). Surfactant proteins A, B, C and D in the human nasal airway: associated with mucosal glands and ciliated epithelium but absent in fluid-phase secretions and mucus. *ORL; Journal for Oto-Rhino-Laryngology and Its Related Specialties*, 76(5), 288–301. <http://doi.org/10.1159/000369143>
- Gauvreau, G. M., Boulet, L. P., Cockcroft, D. W., Baatjes, A., Cote, J., Deschesnes, F., ... O'Byrne, P. M. (2008). Antisense therapy against CCR3 and the common beta chain attenuates allergen-induced eosinophilic responses. *American Journal of Respiratory and Critical Care Medicine*, 177(9), 952–8. <http://doi.org/10.1164/rccm.200708-1251OC>
- Gauvreau, G. M., Pageau, R., Séguin, R., Carballo, D., Gauthier, J., D'Anjou, H., ... Renzi, P. M. (2011). Dose-response effects of TPI ASM8 in asthmatics after allergen. *Allergy*, 66(9), 1242–8. <http://doi.org/10.1111/j.1398-9995.2011.02638.x>
- Geary, R. S., Norris, D., Yu, R., & Bennett, C. F. (2015). Pharmacokinetics, biodistribution and cell uptake of antisense oligonucleotides. *Advanced Drug Delivery Reviews*, 87, 46–51. <http://doi.org/10.1016/j.addr.2015.01.008>
- Geary, R. S., Wancewicz, E., Matson, J., Pearce, M., Siwkowski, A., Swayze, E., & Bennett, F. (2009). Effect of dose and plasma concentration on liver uptake and pharmacologic activity of a 2'-methoxyethyl modified chimeric antisense oligonucleotide targeting PTEN. *Biochemical Pharmacology*, 78(3), 284–91. <http://doi.org/10.1016/j.bcp.2009.04.013>
- Gebert, L. F. R., Rebhan, M. A. E., Crivelli, S. E. M., Denzler, R., Stoffel, M., & Hall, J. (2013). Miravirsin (SPC3649) can inhibit the biogenesis of miR-122. *Nucleic Acids Research*, 42(1), 609–621. <http://doi.org/10.1093/nar/gkt852>
- Geller, B. L., Marshall-Batty, K., Schnell, F. J., McKnight, M. M., Iversen, P. L., & Greenberg, D. E. (2013). Gene-silencing antisense oligomers inhibit acinetobacter growth in vitro and in vivo. *The Journal of Infectious Diseases*, 208(10), 1553–60. <http://doi.org/10.1093/infdis/jit460>
- Gitiban, N., Jurcisek, J. A., Harris, R. H., Mertz, S. E., Durbin, R. K., Bakaletz, L. O., & Durbin, J. E. (2005). Chinchilla and murine models of upper respiratory tract infections with respiratory syncytial virus. *Journal of Virology*, 79(10), 6035–42. <http://doi.org/10.1128/JVI.79.10.6035-6042.2005>
- Gong, W., & Desaulniers, J.-P. (2012). Gene-silencing properties of siRNAs that contain internal amide-bond linkages. *Bioorganic & Medicinal Chemistry Letters*, 22(22), 6934–7. <http://doi.org/10.1016/j.bmcl.2012.09.009>
- Gorzelnik, K., Janke, J., Engeli, S., & Sharma, A. M. (2001). Validation of endogenous



controls for gene expression studies in human adipocytes and preadipocytes. *Hormone and Metabolic Research = Hormon- Und Stoffwechselforschung = Hormones et Métabolisme*, 33(10), 625–7. <http://doi.org/10.1055/s-2001-17911>

Gosens, R., & Grainge, C. (2015). Bronchoconstriction and airway biology: potential impact and therapeutic opportunities. *Chest*, 147(3), 798–803. <http://doi.org/10.1378/chest.14-1142>

Gottlieb, J., Zamora, M. R., Hodges, T., Musk, A. W., Sommerwerk, U., Dilling, D., ... Glanville, A. R. (2015a). ALN-RSV01 for prevention of bronchiolitis obliterans syndrome after respiratory syncytial virus infection in lung transplant recipients. *The Journal of Heart and Lung Transplantation : The Official Publication of the International Society for Heart Transplantation*. <http://doi.org/10.1016/j.healun.2015.08.012>

Gottlieb, J., Zamora, M. R., Hodges, T., Musk, A. W., Sommerwerk, U., Dilling, D., ... Glanville, A. R. (2015b). ALN-RSV01 for prevention of bronchiolitis obliterans syndrome after respiratory syncytial virus infection in lung transplant recipients. *The Journal of Heart and Lung Transplantation : The Official Publication of the International Society for Heart Transplantation*. <http://doi.org/10.1016/j.healun.2015.08.012>

Griesenbach, U., Kitson, C., Escudero Garcia, S., Farley, R., Singh, C., Somerton, L., ... Alton, E. W. F. W. (2006). Inefficient cationic lipid-mediated siRNA and antisense oligonucleotide transfer to airway epithelial cells in vivo. *Respiratory Research*, 7, 26. <http://doi.org/10.1186/1465-9921-7-26>

Gu, S., Jin, L., Zhang, F., Huang, Y., Grimm, D., Rossi, J. J., & Kay, M. A. (2011). Thermodynamic stability of small hairpin RNAs highly influences the loading process of different mammalian Argonautes. *Proceedings of the National Academy of Sciences of the United States of America*, 108(22), 9208–13. <http://doi.org/10.1073/pnas.1018023108>

Guimond, A., Viau, E., Aubé, P., Renzi, P. M., Paquet, L., & Ferrari, N. (2008). Advantageous toxicity profile of inhaled antisense oligonucleotides following chronic dosing in non-human primates. *Pulmonary Pharmacology & Therapeutics*, 21(6), 845–54. <http://doi.org/10.1016/j.pupt.2008.08.001>

Guvakova, M. A., Yakubov, L. A., Vlodaysky, I., Tonkinson, J. L., & Stein, C. A. (1995). Phosphorothioate Oligodeoxynucleotides Bind to Basic Fibroblast Growth Factor, Inhibit Its Binding to Cell Surface Receptors, and Remove It from Low Affinity Binding Sites on Extracellular Matrix. *Journal of Biological Chemistry*, 270(6), 2620–2627. <http://doi.org/10.1074/jbc.270.6.2620>

Hall, C. B. (2001). Respiratory syncytial virus and parainfluenza virus. *The New England Journal of Medicine*, 344(25), 1917–28. <http://doi.org/10.1056/NEJM200106213442507>

Hall, C. B., Douglas, R. G., & Geiman, J. M. (1975). Quantitative shedding patterns of respiratory syncytial virus in infants. *The Journal of Infectious Diseases*, 132(2), 151–6. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/808581>

Hall, C. B., Douglas, R. G., & Geiman, J. M. (1976). Respiratory syncytial virus infections in infants: quantitation and duration of shedding. *The Journal of Pediatrics*, 89(1), 11–5. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/180274>

Hamid, Q., Azzawi, M., Ying, S., Moqbel, R., Wardlaw, A. J., Corrigan, C. J., ... Jeffery, P.

- K. (1991). Expression of mRNA for interleukin-5 in mucosal bronchial biopsies from asthma. *The Journal of Clinical Investigation*, 87(5), 1541–6. <http://doi.org/10.1172/JCI115166>
- Hamm, S., Latz, E., Hangel, D., Müller, T., Yu, P., Golenbock, D., ... Bauer, S. (2010). Alternating 2'-O-ribose methylation is a universal approach for generating non-stimulatory siRNA by acting as TLR7 antagonist. *Immunobiology*, 215(7), 559–69. <http://doi.org/10.1016/j.imbio.2009.09.003>
- Han, S., & Mallampalli, R. K. (2015). The Role of Surfactant in Lung Disease and Host Defense against Pulmonary Infections. *Annals of the American Thoracic Society*, 12(5), 765–74. <http://doi.org/10.1513/AnnalsATS.201411-507FR>
- Hannus, M., Beitzinger, M., Engelmann, J. C., Weickert, M.-T., Spang, R., Hannus, S., & Meister, G. (2014). siPools: highly complex but accurately defined siRNA pools eliminate off-target effects. *Nucleic Acids Research*, 42(12), 8049–61. <http://doi.org/10.1093/nar/gku480>
- Hasleton, P. S. (1972). The internal surface area of the adult human lung. *Journal of Anatomy*, 112(Pt 3), 391–400. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1271180&tool=pmcentrez&rendertype=abstract>
- He, S., Chu, J., Wu, L.-C., Mao, H., Peng, Y., Alvarez-Breckenridge, C. A., ... Yu, J. (2013). MicroRNAs activate natural killer cells through Toll-like receptor signaling. *Blood*, 121(23), 4663–71. <http://doi.org/10.1182/blood-2012-07-441360>
- He, W. A., Calore, F., Londhe, P., Canella, A., Guttridge, D. C., & Croce, C. M. (2014). Microvesicles containing miRNAs promote muscle cell death in cancer cachexia via TLR7. *Proceedings of the National Academy of Sciences of the United States of America*, 111(12), 4525–9. <http://doi.org/10.1073/pnas.1402714111>
- Heald, A. E., Iversen, P. L., Saoud, J. B., Sazani, P., Charleston, J. S., Axtelle, T., ... Kaye, E. (2014). Safety and pharmacokinetic profiles of phosphorodiamidate morpholino oligomers with activity against ebola virus and marburg virus: results of two single-ascending-dose studies. *Antimicrobial Agents and Chemotherapy*, 58(11), 6639–47. <http://doi.org/10.1128/AAC.03442-14>
- Henry, S. P., Giclas, P. C., Leeds, J., Pangburn, M., Auletta, C., Levin, A. A., & Kornbrust, D. J. (1997). Activation of the alternative pathway of complement by a phosphorothioate oligonucleotide: potential mechanism of action. *The Journal of Pharmacology and Experimental Therapeutics*, 281(2), 810–6. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9152389>
- Herbert, M., Coppieters, N., Lasham, A., Cao, H., & Reid, G. (2011). The importance of RT-qPCR primer design for the detection of siRNA-mediated mRNA silencing. *BMC Research Notes*, 4(1), 148. <http://doi.org/10.1186/1756-0500-4-148>
- Heyder, J. (2012). Deposition of Inhaled Particles in the Human Respiratory Tract and Consequences for Regional Targeting in Respiratory Drug Delivery. *Proceedings of the American Thoracic Society*. Retrieved from <http://www.atsjournals.org/doi/full/10.1513/pats.200409-046TA#.VoGe7ZOLQUE>
- Hillaire, M. L. B., Haagsman, H. P., Osterhaus, A. D. M. E., Rimmelzwaan, G. F., & van Eijk, M. (2013). Pulmonary surfactant protein D in first-line innate defence against influenza A virus infections. *Journal of Innate Immunity*, 5(3), 197–208.

<http://doi.org/10.1159/000346374>

- Holmes, K., Williams, C. M., Chapman, E. A., & Cross, M. J. (2010). Detection of siRNA induced mRNA silencing by RT-qPCR: considerations for experimental design. *BMC Research Notes*, 3, 53. <http://doi.org/10.1186/1756-0500-3-53>
- Hornung, V., Ellegast, J., Kim, S., Brzózka, K., Jung, A., Kato, H., ... Hartmann, G. (2006). 5'-Triphosphate RNA is the ligand for RIG-I. *Science (New York, N.Y.)*, 314(5801), 994–7. <http://doi.org/10.1126/science.1132505>
- Hornung, V., Guenther-Biller, M., Bourquin, C., Ablasser, A., Schlee, M., Uematsu, S., ... Hartmann, G. (2005). Sequence-specific potent induction of IFN-alpha by short interfering RNA in plasmacytoid dendritic cells through TLR7. *Nature Medicine*, 11(3), 263–70. <http://doi.org/10.1038/nm1191>
- Horsfield, K. (1990). Diameters, generations, and orders of branches in the bronchial tree. *Journal of Applied Physiology (Bethesda, Md. : 1985)*, 68(2), 457–61. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/2318756>
- Humbert, M., Corrigan, C. J., Kimmitt, P., Till, S. J., Kay, A. B., & Durham, S. R. (1997). Relationship between IL-4 and IL-5 mRNA expression and disease severity in atopic asthma. *American Journal of Respiratory and Critical Care Medicine*, 156(3 Pt 1), 704–8. <http://doi.org/10.1164/ajrccm.156.3.9610033>
- Humphreys, D. T., Hynes, C. J., Patel, H. R., Wei, G. H., Cannon, L., Fatkin, D., ... Preiss, T. (2012). Complexity of murine cardiomyocyte miRNA biogenesis, sequence variant expression and function. *PLoS One*, 7(2), e30933. <http://doi.org/10.1371/journal.pone.0030933>
- HURWITZ, S. H. (1955). Nonallergic asthma; differential diagnosis and treatment. *California Medicine*, 83(2), 61–7. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1532783&tool=pmcentrez&rendertype=abstract>
- Igreja, S., Clarke, L. A., Botelho, H. M., Marques, L., & Amaral, M. D. (2015). Correction of a Cystic Fibrosis Splicing Mutation by Antisense Oligonucleotides. *Human Mutation*. <http://doi.org/10.1002/humu.22931>
- Imaoka, H., Campbell, H., Babirad, I., Watson, R. M., Mistry, M., Sehmi, R., & Gauvreau, G. M. (2011). TPI ASM8 reduces eosinophil progenitors in sputum after allergen challenge. *Clinical and Experimental Allergy : Journal of the British Society for Allergy and Clinical Immunology*, 41(12), 1740–6. <http://doi.org/10.1111/j.1365-2222.2011.03816.x>
- Israelow, B., Mullokandov, G., Agudo, J., Sourisseau, M., Bashir, A., Maldonado, A. Y., ... Evans, M. J. (2014). Hepatitis C virus genetics affects miR-122 requirements and response to miR-122 inhibitors. *Nature Communications*, 5, 5408. <http://doi.org/10.1038/ncomms6408>
- Iversen, P. L., Warren, T. K., Wells, J. B., Garza, N. L., Mourich, D. V., Welch, L. S., ... Bavari, S. (2012). Discovery and early development of AVI-7537 and AVI-7288 for the treatment of Ebola virus and Marburg virus infections. *Viruses*, 4(11), 2806–30. <http://doi.org/10.3390/v4112806>
- Iversen, P. L., Zhu, S., Meyer, A., & Zon, G. (1992). Cellular uptake and subcellular distribution of phosphorothioate oligonucleotides into cultured cells. *Antisense Research*

*and Development*, 2(3), 211–22. Retrieved from  
<http://www.ncbi.nlm.nih.gov/pubmed/1490072>

- Jackson, A. L., Burchard, J., Leake, D., Reynolds, A., Schelter, J., Guo, J., ... Linsley, P. S. (2006). Position-specific chemical modification of siRNAs reduces “off-target” transcript silencing. *RNA (New York, N.Y.)*, 12(7), 1197–205.  
<http://doi.org/10.1261/rna.30706>
- Jagannathan, R., Thapa, D., Nichols, C. E., Shepherd, D. L., Stricker, J. C., Croston, T. L., ... Hollander, J. M. (2015). Translational Regulation of the Mitochondrial Genome Following Redistribution of Mitochondrial MicroRNA (MitomiR) in the Diabetic Heart. *Circulation. Cardiovascular Genetics*.  
<http://doi.org/10.1161/CIRCGENETICS.115.001067>
- Johansson, S. G. O., Hourihane, J. O., Bousquet, J., Bruijnzeel-Koomen, C., Dreborg, S., Haahtela, T., ... Wüthrich, B. (2008). A revised nomenclature for allergy: An EAACI position statement from the EAACI nomenclature task force. *Allergy*, 56(9), 813–824.  
<http://doi.org/10.1111/j.1398-9995.2001.00002.x-i1>
- Johnson, J. E., Gonzales, R. A., Olson, S. J., Wright, P. F., & Graham, B. S. (2007). The histopathology of fatal untreated human respiratory syncytial virus infection. *Modern Pathology: An Official Journal of the United States and Canadian Academy of Pathology, Inc*, 20(1), 108–19. <http://doi.org/10.1038/modpathol.3800725>
- Jones, L. H., Baldock, H., Bunnage, M. E., Burrows, J., Clarke, N., Coghlan, M., ... Price, D. A. (2011). Inhalation by design: dual pharmacology  $\beta$ -2 agonists/M3 antagonists for the treatment of COPD. *Bioorganic & Medicinal Chemistry Letters*, 21(9), 2759–63.  
<http://doi.org/10.1016/j.bmcl.2010.10.132>
- Judge, A. D., Sood, V., Shaw, J. R., Fang, D., McClintock, K., & MacLachlan, I. (2005). Sequence-dependent stimulation of the mammalian innate immune response by synthetic siRNA. *Nature Biotechnology*, 23(4), 457–62. <http://doi.org/10.1038/nbt1081>
- Juliano, R. L., Ming, X., & Nakagawa, O. (2012). Cellular uptake and intracellular trafficking of antisense and siRNA oligonucleotides. *Bioconjugate Chemistry*, 23(2), 147–57.  
<http://doi.org/10.1021/bc200377d>
- Karikó, K., Buckstein, M., Ni, H., & Weissman, D. (2005). Suppression of RNA recognition by Toll-like receptors: the impact of nucleoside modification and the evolutionary origin of RNA. *Immunity*, 23(2), 165–75. <http://doi.org/10.1016/j.immuni.2005.06.008>
- Karras, J. G., Crosby, J. R., Guha, M., Tung, D., Miller, D. A., Gaarde, W. A., ... Gregory, S. A. (2007). Anti-inflammatory activity of inhaled IL-4 receptor-alpha antisense oligonucleotide in mice. *American Journal of Respiratory Cell and Molecular Biology*, 36(3), 276–85. <http://doi.org/10.1165/rcmb.2005-0456OC>
- Kato, H., Takeuchi, O., Mikamo-Satoh, E., Hirai, R., Kawai, T., Matsushita, K., ... Akira, S. (2008). Length-dependent recognition of double-stranded ribonucleic acids by retinoic acid-inducible gene-I and melanoma differentiation-associated gene 5. *The Journal of Experimental Medicine*, 205(7), 1601–10. <http://doi.org/10.1084/jem.20080091>
- Kawai, J., Shinagawa, A., Shibata, K., Yoshino, M., Itoh, M., Ishii, Y., ... Hayashizaki, Y. (2001). Functional annotation of a full-length mouse cDNA collection. *Nature*, 409(6821), 685–90. <http://doi.org/10.1038/35055500>
- Kawamata, T., & Tomari, Y. (2010). Making RISC. *Trends in Biochemical Sciences*, 35(7),

368–76. <http://doi.org/10.1016/j.tibs.2010.03.009>

- Khvorova, A., Reynolds, A., & Jayasena, S. D. (2003). Functional siRNAs and miRNAs exhibit strand bias. *Cell*, *115*(2), 209–216. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14567918>
- Kim, D. H., Saetrom, P., Snøve, O., & Rossi, J. J. (2008). MicroRNA-directed transcriptional gene silencing in mammalian cells. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(42), 16230–5. <http://doi.org/10.1073/pnas.0808830105>
- Kleinman, M. E., Yamada, K., Takeda, A., Chandrasekaran, V., Nozaki, M., Baffi, J. Z., ... Ambati, J. (2008). Sequence- and target-independent angiogenesis suppression by siRNA via TLR3. *Nature*, *452*(7187), 591–7. <http://doi.org/10.1038/nature06765>
- Kodym, R., Kodym, E., & Story, M. D. (2009). 2'-5'-Oligoadenylate synthetase is activated by a specific RNA sequence motif. *Biochemical and Biophysical Research Communications*, *388*(2), 317–22. <http://doi.org/10.1016/j.bbrc.2009.07.167>
- Koller, E., Vincent, T. M., Chappell, A., De, S., Manoharan, M., & Bennett, C. F. (2011). Mechanisms of single-stranded phosphorothioate modified antisense oligonucleotide accumulation in hepatocytes. *Nucleic Acids Research*, *39*(11), 4795–807. <http://doi.org/10.1093/nar/gkr089>
- Krieg, A. M., Yi, A.-K., Matson, S., Waldschmidt, T. J., Bishop, G. A., Teasdale, R., ... Klinman, D. M. (1995). CpG motifs in bacterial DNA trigger direct B-cell activation. *Nature*, *374*(6522), 546–549. <http://doi.org/10.1038/374546a0>
- Krol, J., Sobczak, K., Wilczynska, U., Drath, M., Jasinska, A., Kaczynska, D., & Krzyzosiak, W. J. (2004). Structural features of microRNA (miRNA) precursors and their relevance to miRNA biogenesis and small interfering RNA/short hairpin RNA design. *J Biol Chem*, *279*(40), 42230–42239. <http://doi.org/10.1074/jbc.M404931200>
- Kumar, R., Singh, S. K., Koshkin, A. A., Rajwanshi, V. K., Meldgaard, M., & Wengel, J. (1998). The first analogues of LNA (locked nucleic acids): phosphorothioate-LNA and 2'-thio-LNA. *Bioorganic & Medicinal Chemistry Letters*, *8*(16), 2219–22. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9873516>
- Labiris, N. R., & Dolovich, M. B. (2003). Pulmonary drug delivery. Part I: physiological factors affecting therapeutic effectiveness of aerosolized medications. *British Journal of Clinical Pharmacology*, *56*(6), 588–99. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1884307&tool=pmcentrez&rendertype=abstract>
- Lahens, N. F., Kavakli, I. H., Zhang, R., Hayer, K., Black, M. B., Dueck, H., ... Hogenesch, J. B. (2014). IVT-seq reveals extreme bias in RNA sequencing. *Genome Biology*, *15*(6), R86. <http://doi.org/10.1186/gb-2014-15-6-r86>
- Lai, S. K., Wang, Y.-Y., Wirtz, D., & Hanes, J. (2009). Micro- and macrorheology of mucus. *Advanced Drug Delivery Reviews*, *61*(2), 86–100. <http://doi.org/10.1016/j.addr.2008.09.012>
- Lai, S.-H., Stein, D. A., Guerrero-Plata, A., Liao, S.-L., Ivanciuc, T., Hong, C., ... Garofalo, R. P. (2008). Inhibition of respiratory syncytial virus infections with morpholino oligomers in cell cultures and in mice. *Molecular Therapy: The Journal of the American Society of Gene Therapy*, *16*(6), 1120–8. <http://doi.org/10.1038/mt.2008.81>

- Lam, J. K. W., Chow, M. Y. T., Zhang, Y., & Leung, S. W. S. (2015). siRNA Versus miRNA as Therapeutics for Gene Silencing. *Molecular Therapy. Nucleic Acids*, 4, e252. <http://doi.org/10.1038/mtna.2015.23>
- Lambert, R. K. (1989). A new computational model for expiratory flow from nonhomogeneous human lungs. *Journal of Biomechanical Engineering*, 111(3), 200–5. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/2779184>
- Lambert, R. K., Wilson, T. A., Hyatt, R. E., & Rodarte, J. R. (1982). A computational model for expiratory flow. *Journal of Applied Physiology: Respiratory, Environmental and Exercise Physiology*, 52(1), 44–56. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7061277>
- Laube, B. L. (2014). The expanding role of aerosols in systemic drug delivery, gene therapy and vaccination: an update. *Translational Respiratory Medicine*, 2, 3. <http://doi.org/10.1186/2213-0802-2-3>
- Leckie, M. J., ten Brinke, A., Khan, J., Diamant, Z., O'Connor, B. J., Walls, C. M., ... Barnes, P. J. (2000). Effects of an interleukin-5 blocking monoclonal antibody on eosinophils, airway hyper-responsiveness, and the late asthmatic response. *Lancet (London, England)*, 356(9248), 2144–8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11191542>
- Ledford, J. G., Addison, K. J., Foster, M. W., & Que, L. G. (2014). Eosinophil-associated lung diseases. A cry for surfactant proteins A and D help? *American Journal of Respiratory Cell and Molecular Biology*, 51(5), 604–14. <http://doi.org/10.1165/rcmb.2014-0095TR>
- Leuschner, P. J., Ameres, S. L., Kueng, S., & Martinez, J. (2006). Cleavage of the siRNA passenger strand during RISC assembly in human cells. *EMBO Rep*, 7(3), 314–320. <http://doi.org/10.1038/sj.embor.7400637>
- Lewis, B. P., Shih, I., Jones-Rhoades, M. W., Bartel, D. P., & Burge, C. B. (2003). Prediction of mammalian microRNA targets. *Cell*, 115(7), 787–98. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14697198>
- Li, F., Pallan, P. S., Maier, M. A., Rajeev, K. G., Mathieu, S. L., Kreutz, C., ... Egli, M. (2007). Crystal structure, stability and in vitro RNAi activity of oligoribonucleotides containing the ribo-difluorotoluy nucleotide: insights into substrate requirements by the human RISC Ago2 enzyme. *Nucleic Acids Research*, 35(19), 6424–38. <http://doi.org/10.1093/nar/gkm664>
- Li, L.-C., Okino, S. T., Zhao, H., Pookot, D., Place, R. F., Urakami, S., ... Dahiya, R. (2006). Small dsRNAs induce transcriptional activation in human cells. *Proceedings of the National Academy of Sciences of the United States of America*, 103(46), 17337–42. <http://doi.org/10.1073/pnas.0607015103>
- Liu, J., Hu, J., & Corey, D. R. (2012). Expanding the action of duplex RNAs into the nucleus: redirecting alternative splicing. *Nucleic Acids Research*, 40(3), 1240–50. <http://doi.org/10.1093/nar/gkr780>
- Liu, J., Valencia-Sanchez, M. A., Hannon, G. J., & Parker, R. (2005). MicroRNA-dependent localization of targeted mRNAs to mammalian P-bodies. *Nature Cell Biology*, 7(7), 719–23. <http://doi.org/10.1038/ncb1274>
- Liu, X., Zhang, L., & Chen, S. (2015). Modeling Exon-Specific Bias Distribution Improves

the Analysis of RNA-Seq Data. *PLOS ONE*, *10*(10), e0140032.  
<http://doi.org/10.1371/journal.pone.0140032>

- Loke, S. L., Stein, C. A., Zhang, X. H., Mori, K., Nakanishi, M., Subasinghe, C., ... Neckers, L. M. (1989). Characterization of oligonucleotide transport into living cells. *Proceedings of the National Academy of Sciences of the United States of America*, *86*(10), 3474–8. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=287160&tool=pmcentrez&rendertype=abstract>
- Lomas-Neira, J. L., Chung, C.-S., Wesche, D. E., Perl, M., & Ayala, A. (2005). In vivo gene silencing (with siRNA) of pulmonary expression of MIP-2 versus KC results in divergent effects on hemorrhage-induced, neutrophil-mediated septic acute lung injury. *Journal of Leukocyte Biology*, *77*(6), 846–53. <http://doi.org/10.1189/jlb.1004617>
- Lorenz, C., Hadwiger, P., John, M., Vornlocher, H.-P., & Unverzagt, C. (2004). Steroid and lipid conjugates of siRNAs to enhance cellular uptake and gene silencing in liver cells. *Bioorganic & Medicinal Chemistry Letters*, *14*(19), 4975–7. <http://doi.org/10.1016/j.bmcl.2004.07.018>
- Lorenz, R. J. (1966). Weibel, E. R.: Morphometry of the Human Lung. Springer Verlag, Berlin-Göttingen-Heidelberg 1963; 151 S., 109 Abb., DM 36,-. *Biometrische Zeitschrift*, *8*(1-2), 143–144. <http://doi.org/10.1002/bimj.19660080155>
- Lorenzer, C., Dirin, M., Winkler, A.-M., Baumann, V., & Winkler, J. (2015). Going beyond the liver: progress and challenges of targeted delivery of siRNA therapeutics. *Journal of Controlled Release: Official Journal of the Controlled Release Society*, *203*, 1–15. <http://doi.org/10.1016/j.jconrel.2015.02.003>
- Lu, Q. L., Rabinowitz, A., Chen, Y. C., Yokota, T., Yin, H., Alter, J., ... Partridge, T. (2005). Systemic delivery of antisense oligoribonucleotide restores dystrophin expression in body-wide skeletal muscles. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(1), 198–203. <http://doi.org/10.1073/pnas.0406700102>
- Lu, Q.-L., Cirak, S., & Partridge, T. (2014). What Can We Learn From Clinical Trials of Exon Skipping for DMD? *Molecular Therapy. Nucleic Acids*, *3*, e152. <http://doi.org/10.1038/mtna.2014.6>
- Lupfer, C., Stein, D. A., Mourich, D. V., Tepper, S. E., Iversen, P. L., & Pastey, M. (2008). Inhibition of influenza A H3N8 virus infections in mice by morpholino oligomers. *Archives of Virology*, *153*(5), 929–37. <http://doi.org/10.1007/s00705-008-0067-0>
- Ma, J., Mercer, R. R., Barger, M., Schwegler-Berry, D., Cohen, J. M., Demokritou, P., & Castranova, V. (2015). Effects of amorphous silica coating on cerium oxide nanoparticles induced pulmonary responses. *Toxicology and Applied Pharmacology*, *288*(1), 63–73. <http://doi.org/10.1016/j.taap.2015.07.012>
- Maekawa, K., Azuma, M., Okuno, Y., Tsukamoto, T., Nishiguchi, K., Setsukinai, K.-I., ... Rokushima, M. (2015). Antisense peptide nucleic acid-peptide conjugates for functional analyses of genes in *Pseudomonas aeruginosa*. *Bioorganic & Medicinal Chemistry*, *23*(22), 7234–7239. <http://doi.org/10.1016/j.bmc.2015.10.020>
- Manche, L., Green, S. R., Schmedt, C., & Mathews, M. B. (1992). Interactions between double-stranded RNA regulators and the protein kinase DA1. *Molecular and Cellular Biology*, *12*(11), 5238–48. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=360457&tool=pmcentrez&rendertype=abstract>

endertype=abstract

- Maniataki, E., & Mourelatos, Z. (2005). A human, ATP-independent, RISC assembly machine fueled by pre-miRNA. *Genes & Development*, *19*(24), 2979–90. <http://doi.org/10.1101/gad.1384005>
- Marcon, E., Babak, T., Chua, G., Hughes, T., & Moens, P. B. (2008). miRNA and piRNA localization in the male mammalian meiotic nucleus. *Chromosome Research : An International Journal on the Molecular, Supramolecular and Evolutionary Aspects of Chromosome Biology*, *16*(2), 243–60. <http://doi.org/10.1007/s10577-007-1190-6>
- Marlin, F., Simon, P., Bonneau, S., Alberti, P., Cordier, C., Boix, C., ... Giovannangeli, C. (2012). Flavin conjugates for delivery of peptide nucleic acids. *Chembiochem : A European Journal of Chemical Biology*, *13*(17), 2593–8. <http://doi.org/10.1002/cbic.201200505>
- Marques, J. T., Devosse, T., Wang, D., Zamanian-Daryoush, M., Serbinowski, P., Hartmann, R., ... Williams, B. R. G. (2006). A structural basis for discriminating between self and nonself double-stranded RNAs in mammalian cells. *Nature Biotechnology*, *24*(5), 559–65. <http://doi.org/10.1038/nbt1205>
- Masaki, T., Arend, K. C., Li, Y., Yamane, D., McGivern, D. R., Kato, T., ... Lemon, S. M. (2015). miR-122 stimulates hepatitis C virus RNA synthesis by altering the balance of viral RNAs engaged in replication versus translation. *Cell Host & Microbe*, *17*(2), 217–28. <http://doi.org/10.1016/j.chom.2014.12.014>
- Matranga, C., Tomari, Y., Shin, C., Bartel, D. P., & Zamore, P. D. (2005). Passenger-strand cleavage facilitates assembly of siRNA into Ago2-containing RNAi enzyme complexes. *Cell*, *123*(4), 607–620. <http://doi.org/10.1016/j.cell.2005.08.044>
- Matsui, M., Chu, Y., Zhang, H., Gagnon, K. T., Shaikh, S., Kuchimanchi, S., ... Janowski, B. A. (2013). Promoter RNA links transcriptional regulation of inflammatory pathway genes. *Nucleic Acids Research*, *41*(22), 10086–109. <http://doi.org/10.1093/nar/gkt777>
- Matsui, M., Prakash, T. P., & Corey, D. R. (2013). Transcriptional silencing by single-stranded RNAs targeting a noncoding RNA that overlaps a gene promoter. *ACS Chemical Biology*, *8*(1), 122–6. <http://doi.org/10.1021/cb300490j>
- Mauroy, B., Flaud, P., Pelca, D., Fausser, C., Merckx, J., & Mitchell, B. R. (2015). Toward the modeling of mucus draining from human lung: role of airways deformation on air-mucus interaction. *Frontiers in Physiology*, *6*, 214. <http://doi.org/10.3389/fphys.2015.00214>
- McNeer, N. A., Anandalingam, K., Fields, R. J., Caputo, C., Kopic, S., Gupta, A., ... Egan, M. E. (2015). Nanoparticles that deliver triplex-forming peptide nucleic acid molecules correct F508del CFTR in airway epithelium. *Nature Communications*, *6*, 6952. <http://doi.org/10.1038/ncomms7952>
- Mehiri, M., Upert, G., Tripathi, S., Di Giorgio, A., Condom, R., Pandey, V. N., & Patino, N. (2008). An efficient biodelivery system for antisense polyamide nucleic acid (PNA). *Oligonucleotides*, *18*(3), 245–56. <http://doi.org/10.1089/oli.2008.0126>
- Mitchell, J., Bauer, R., Lyapustina, S., Tougas, T., & Glaab, V. (2011). Non-impactor-based methods for sizing of aerosols emitted from orally inhaled and nasal drug products (OINDPs). *AAPS PharmSciTech*, *12*(3), 965–988. <http://doi.org/10.1208/s12249-011-9662-6>



- Moreno, P. M. D., Wenska, M., Lundin, K. E., Wrangle, O., Strömberg, R., & Smith, C. I. E. (2009). A synthetic snRNA m3G-CAP enhances nuclear delivery of exogenous proteins and nucleic acids. *Nucleic Acids Research*, *37*(6), 1925–35. <http://doi.org/10.1093/nar/gkp048>
- Morgan, P., Van Der Graaf, P. H., Arrowsmith, J., Feltner, D. E., Drummond, K. S., Wegner, C. D., & Street, S. D. A. (2012). Can the flow of medicines be improved? Fundamental pharmacokinetic and pharmacological principles toward improving Phase II survival. *Drug Discovery Today*, *17*(9-10), 419–24. <http://doi.org/10.1016/j.drudis.2011.12.020>
- Morris, K. V., Santoso, S., Turner, A.-M., Pastori, C., & Hawkins, P. G. (2008). Bidirectional Transcription Directs Both Transcriptional Gene Activation and Suppression in Human Cells. *PLoS Genetics*, *4*(11), e1000258. <http://doi.org/10.1371/journal.pgen.1000258>
- Mortimer, S. A., & Doudna, J. A. (2013). Unconventional miR-122 binding stabilizes the HCV genome by forming a trimolecular RNA structure. *Nucleic Acids Research*, *41*(7), 4230–40. <http://doi.org/10.1093/nar/gkt075>
- Moschos, S. A. (2013). MicroRNA Biotherapeutics: Key Challenges from a Drug Development Perspective. In L. Jones & A. J. McKnight (Eds.), *Biotherapeutics: Recent Developments using Chemical and Molecular Biology* (pp. 176–223). London: Royal Society of Chemistry Publishing.
- Moschos, S. A., Frick, M., Taylor, B., Turnpenny, P., Graves, H., Spink, K. G., ... Yeadon, M. (2011). Uptake, efficacy, and systemic distribution of naked, inhaled short interfering RNA (siRNA) and locked nucleic acid (LNA) antisense. *Mol Ther*, *19*(12), 2163–2168. <http://doi.org/10.1038/mt.2011.206>
- Moschos, S. A., Jones, S. W., Perry, M. M., Williams, A. E., Erjefalt, J. S., Turner, J. J., ... Lindsay, M. A. (2007). Lung delivery studies using siRNA conjugated to TAT(48-60) and penetratin reveal peptide induced reduction in gene expression and induction of innate immunity. *Bioconjug Chem*, *18*(5), 1450–1459. <http://doi.org/10.1021/bc070077d>
- Moschos, S. A., Spinks, K., Williams, A. E., & Lindsay, M. A. (2008). Targeting the lung using siRNA and antisense based oligonucleotides. *Curr Pharm Des*, *14*(34), 3620–3627. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19075738>
- Moulton, H. M., Fletcher, S., Neuman, B. W., McClorey, G., Stein, D. A., Abes, S., ... Iversen, P. L. (2007). Cell-penetrating peptide-morpholino conjugates alter pre-mRNA splicing of DMD (Duchenne muscular dystrophy) and inhibit murine coronavirus replication in vivo. *Biochemical Society Transactions*, *35*(Pt 4), 826–8. <http://doi.org/10.1042/BST0350826>
- Narbus, C. M., Israelow, B., Sourisseau, M., Michta, M. L., Hopcraft, S. E., Zeiner, G. M., & Evans, M. J. (2011). HepG2 cells expressing microRNA miR-122 support the entire hepatitis C virus life cycle. *Journal of Virology*, *85*(22), 12087–92. <http://doi.org/10.1128/JVI.05843-11>
- Nelson, P. T., De Planell-Saguer, M., Lamprinaki, S., Kiriakidou, M., Zhang, P., O'Doherty, U., & Mourelatos, Z. (2007). A novel monoclonal antibody against human Argonaute proteins reveals unexpected characteristics of miRNAs in human blood cells. *RNA (New York, N.Y.)*, *13*(10), 1787–92. <http://doi.org/10.1261/rna.646007>
- Nichols, J. E., Niles, J. A., Vega, S. P., Argueta, L. B., Eastaway, A., & Cortiella, J. (2014). Modeling the lung: Design and development of tissue engineered macro- and micro-physiologic lung models for research use. *Experimental Biology and Medicine*, *239*(9),

1135–1169. <http://doi.org/10.1177/1535370214536679>

- Nicklin, P. L., Bayley, D., Giddings, J., Craig, S. J., Cummins, L. L., Hastewell, J. G., & Phillips, J. A. (1998). Pulmonary bioavailability of a phosphorothioate oligonucleotide (CGP 64128A): comparison with other delivery routes. *Pharm Res*, *15*(4), 583–591. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9587955>
- Nielsen, P. E., Egholm, M., Berg, R. H., & Buchardt, O. (1991). Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide. *Science (New York, N.Y.)*, *254*(5037), 1497–500. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/1962210>
- Nikula, K. J., McCartney, J. E., McGovern, T., Miller, G. K., Odin, M., Pino, M. V., & Reed, M. D. (2014). STP position paper: interpreting the significance of increased alveolar macrophages in rodents following inhalation of pharmaceutical materials. *Toxicologic Pathology*, *42*(3), 472–86. <http://doi.org/10.1177/0192623313507003>
- Nissim-Rafinia, M., Aviram, M., Randell, S. H., Shushi, L., Ozeri, E., Chiba-Falek, O., ... Kerem, B. (2004). Restoration of the cystic fibrosis transmembrane conductance regulator function by splicing modulation. *EMBO Reports*, *5*(11), 1071–7. <http://doi.org/10.1038/sj.embor.7400273>
- Noble, P. B., Pascoe, C. D., Lan, B., Ito, S., Kistemaker, L. E. M., Tatler, A. L., ... West, A. R. (2014). Airway smooth muscle in asthma: linking contraction and mechanotransduction to disease pathogenesis and remodelling. *Pulmonary Pharmacology & Therapeutics*, *29*(2), 96–107. <http://doi.org/10.1016/j.pupt.2014.07.005>
- Nolting, A., DeLong, R. K., Fisher, M. H., Wickstrom, E., Pollack, G. M., Juliano, R. L., & Brouwer, K. L. (1997). Hepatic distribution and clearance of antisense oligonucleotides in the isolated perfused rat liver. *Pharmaceutical Research*, *14*(4), 516–21. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9144742>
- Obad, S., dos Santos, C. O., Petri, A., Heidenblad, M., Broom, O., Ruse, C., ... Kauppinen, S. (2011). Silencing of microRNA families by seed-targeting tiny LNAs. *Nature Genetics*, *43*(4), 371–8. <http://doi.org/10.1038/ng.786>
- Ochs, M. (2014). Estimating structural alterations in animal models of lung emphysema. Is there a gold standard? *Annals of Anatomy = Anatomischer Anzeiger : Official Organ of the Anatomische Gesellschaft*, *196*(1), 26–33. <http://doi.org/10.1016/j.aanat.2013.10.004>
- Oh, S. Y., Ju, Y., & Park, H. (2009). A highly effective and long-lasting inhibition of miRNAs with PNA-based antisense oligonucleotides. *Molecules and Cells*, *28*(4), 341–5. <http://doi.org/10.1007/s10059-009-0134-8>
- Ohr, T., Mütze, J., Staroske, W., Weinmann, L., Höck, J., Crell, K., ... Schwille, P. (2008). Fluorescence correlation spectroscopy and fluorescence cross-correlation spectroscopy reveal the cytoplasmic origination of loaded nuclear RISC in vivo in human cells. *Nucleic Acids Research*, *36*(20), 6439–49. <http://doi.org/10.1093/nar/gkn693>
- Olin, J. T., & Wechsler, M. E. (2014). Asthma: pathogenesis and novel drugs for treatment. *BMJ (Clinical Research Ed.)*, *349*(nov24\_8), g5517. <http://doi.org/10.1136/bmj.g5517>
- Opriessnig, T., Patel, D., Wang, R., Halbur, P. G., Meng, X.-J., Stein, D. A., & Zhang, Y.-J. (2011). Inhibition of porcine reproductive and respiratory syndrome virus infection in piglets by a peptide-conjugated morpholino oligomer. *Antiviral Research*, *91*(1), 36–42.

<http://doi.org/10.1016/j.antiviral.2011.04.012>

- Ørom, U. A., Nielsen, F. C., & Lund, A. H. (2008). MicroRNA-10a binds the 5'UTR of ribosomal protein mRNAs and enhances their translation. *Molecular Cell*, *30*(4), 460–71. <http://doi.org/10.1016/j.molcel.2008.05.001>
- Park, C.-K., Xu, Z.-Z., Berta, T., Han, Q., Chen, G., Liu, X.-J., & Ji, R.-R. (2014). Extracellular microRNAs activate nociceptor neurons to elicit pain via TLR7 and TRPA1. *Neuron*, *82*(1), 47–54. <http://doi.org/10.1016/j.neuron.2014.02.011>
- Park, J. H., & Shin, C. (2015). Slicer-independent mechanism drives small-RNA strand separation during human RISC assembly. *Nucleic Acids Research*, *43*(19), 9418–33. <http://doi.org/10.1093/nar/gkv937>
- Parker, J. C., & Townsley, M. I. (2008). Physiological determinants of the pulmonary filtration coefficient. *American Journal of Physiology. Lung Cellular and Molecular Physiology*, *295*(2), L235–7. <http://doi.org/10.1152/ajplung.00064.2008>
- Parker, R., & Sheth, U. (2007). P bodies and the control of mRNA translation and degradation. *Molecular Cell*, *25*(5), 635–46. <http://doi.org/10.1016/j.molcel.2007.02.011>
- Parra, E., & Pérez-Gil, J. (2015). Composition, structure and mechanical properties define performance of pulmonary surfactant membranes and films. *Chemistry and Physics of Lipids*, *185*, 153–75. <http://doi.org/10.1016/j.chemphyslip.2014.09.002>
- Patton, J. G., Franklin, J. L., Weaver, A. M., Vickers, K., Zhang, B., Coffey, R. J., ... McManus, M. T. (2015). Biogenesis, delivery, and function of extracellular RNA. *Journal of Extracellular Vesicles*, *4*, 27494. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4553266&tool=pmcentrez&rendertype=abstract>
- Perkins, S. M., Webb, D. L., Torrance, S. A., El Saleeby, C., Harrison, L. M., Aitken, J. A., ... DeVincenzo, J. P. (2005). Comparison of a real-time reverse transcriptase PCR assay and a culture technique for quantitative assessment of viral load in children naturally infected with respiratory syncytial virus. *Journal of Clinical Microbiology*, *43*(5), 2356–62. <http://doi.org/10.1128/JCM.43.5.2356-2362.2005>
- Perl, M., Chung, C.-S., Lomas-Neira, J., Rachel, T.-M., Biffl, W. L., Cioffi, W. G., & Ayala, A. (2005). Silencing of Fas, but not caspase-8, in lung epithelial cells ameliorates pulmonary apoptosis, inflammation, and neutrophil influx after hemorrhagic shock and sepsis. *The American Journal of Pathology*, *167*(6), 1545–59. [http://doi.org/10.1016/S0002-9440\(10\)61240-0](http://doi.org/10.1016/S0002-9440(10)61240-0)
- Perry, M. M., Moschos, S. A., Williams, A. E., Shepherd, N. J., Larner-Svensson, H. M., & Lindsay, M. A. (2008). Rapid changes in microRNA-146a expression negatively regulate the IL-1beta-induced inflammatory response in human lung alveolar epithelial cells. *J Immunol*, *180*(8), 5689–5698. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18390754>
- Pichlmair, A., Schulz, O., Tan, C. P., Näsund, T. I., Liljeström, P., Weber, F., & Reis e Sousa, C. (2006). RIG-I-mediated antiviral responses to single-stranded RNA bearing 5'-phosphates. *Science (New York, N.Y.)*, *314*(5801), 997–1001. <http://doi.org/10.1126/science.1132998>
- Place, R. F., Li, L.-C., Pookot, D., Noonan, E. J., & Dahiya, R. (2008). MicroRNA-373 induces expression of genes with complementary promoter sequences. *Proceedings of*

*the National Academy of Sciences of the United States of America*, 105(5), 1608–13.  
<http://doi.org/10.1073/pnas.0707594105>

- Politz, J. C. R., Zhang, F., & Pederson, T. (2006). MicroRNA-206 colocalizes with ribosome-rich regions in both the nucleolus and cytoplasm of rat myogenic cells. *Proceedings of the National Academy of Sciences of the United States of America*, 103(50), 18957–62.  
<http://doi.org/10.1073/pnas.0609466103>
- Pooga, M., Land, T., Bartfai, T., & Langel, U. (2001). PNA oligomers as tools for specific modulation of gene expression. *Biomolecular Engineering*, 17(6), 183–92. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11337277>
- Prakash, T. P., Naik, N., Sioufi, N., Bhat, B., & Swayze, E. E. (2009). Activity of siRNAs with 2-thio-2'-O-methyluridine modification in mammalian cells. *Nucleosides, Nucleotides & Nucleic Acids*, 28(10), 902–10.  
<http://doi.org/10.1080/15257770903316145>
- Pu, Y., Kline, L. C., Khawaja, N., Van Liew, M., & Berry, J. (2015). Comparison of optical particle sizing and cascade impaction for measuring the particle size of a suspension metered dose inhaler. *Drug Development and Industrial Pharmacy*, 41(5), 737–43.  
<http://doi.org/10.3109/03639045.2014.900079>
- Raabe, C. A., Tang, T.-H., Brosius, J., & Rozhdestvensky, T. S. (2014). Biases in small RNA deep sequencing data. *Nucleic Acids Research*, 42(3), 1414–26.  
<http://doi.org/10.1093/nar/gkt1021>
- Rajsbaum, R., Versteeg, G. A., Schmid, S., Maestre, A. M., Belicha-Villanueva, A., Martínez-Romero, C., ... García-Sastre, A. (2014). Unanchored K48-linked polyubiquitin synthesized by the E3-ubiquitin ligase TRIM6 stimulates the interferon-IKKe kinase-mediated antiviral response. *Immunity*, 40(6), 880–95.  
<http://doi.org/10.1016/j.immuni.2014.04.018>
- Rameix-Welti, M.-A., Le Goffic, R., Hervé, P.-L., Sourimant, J., Rémot, A., Riffault, S., ... Eléouët, J.-F. (2014). Visualizing the replication of respiratory syncytial virus in cells and in living mice. *Nature Communications*, 5, 5104.  
<http://doi.org/10.1038/ncomms6104>
- Raouf, A. A., Chiu, P., Ramtoola, Z., Cumming, I. K., Teng, C., Weinbach, S. P., ... Geary, R. S. (2004). Oral bioavailability and multiple dose tolerability of an antisense oligonucleotide tablet formulated with sodium caprate. *J Pharm Sci*, 93(6), 1431–1439.  
<http://doi.org/10.1002/jps.20051>
- Reid, G., Pel, M. E., Kirschner, M. B., Cheng, Y. Y., Mugridge, N., Weiss, J., ... van Zandwijk, N. (2013). Restoring expression of miR-16: a novel approach to therapy for malignant pleural mesothelioma. *Annals of Oncology: Official Journal of the European Society for Medical Oncology / ESMO*, 24(12), 3128–35.  
<http://doi.org/10.1093/annonc/mdt412>
- Rembach, A., Turner, B. J., Bruce, S., Cheah, I. K., Scott, R. L., Lopes, E. C., ... Cheema, S. S. (2004). Antisense peptide nucleic acid targeting GluR3 delays disease onset and progression in the SOD1 G93A mouse model of familial ALS. *Journal of Neuroscience Research*, 77(4), 573–82. <http://doi.org/10.1002/jnr.20191>
- Reus, A. A., Maas, W. J. M., Jansen, H. T., Constant, S., Staal, Y. C. M., van Triel, J. J., & Kuper, C. F. (2014). Feasibility of a 3D human airway epithelial model to study respiratory absorption. *Toxicology in Vitro: An International Journal Published in*

*Association with BIBRA*, 28(2), 258–64. <http://doi.org/10.1016/j.tiv.2013.10.025>

- Richardson, L. S., Belshe, R. B., Sly, D. L., London, W. T., Prevar, D. A., Camargo, E., & Chanock, R. M. (1978). Experimental respiratory syncytial virus pneumonia in cebus monkeys. *Journal of Medical Virology*, 2(1), 45–59. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/210254>
- Riiser, A. (2015). The human microbiome, asthma, and allergy. *Allergy, Asthma, and Clinical Immunology: Official Journal of the Canadian Society of Allergy and Clinical Immunology*, 11, 35. <http://doi.org/10.1186/s13223-015-0102-0>
- Ripple, M. J., You, D., Honnegowda, S., Giaimo, J. D., Sewell, A. B., Becnel, D. M., & Cormier, S. A. (2010). Immunomodulation with IL-4R alpha antisense oligonucleotide prevents respiratory syncytial virus-mediated pulmonary disease. *Journal of Immunology (Baltimore, Md. : 1950)*, 185(8), 4804–11. <http://doi.org/10.4049/jimmunol.1000484>
- Ro, S., Ma, H.-Y., Park, C., Ortogero, N., Song, R., Hennig, G. W., ... Yan, W. (2013). The mitochondrial genome encodes abundant small noncoding RNAs. *Cell Research*, 23(6), 759–74. <http://doi.org/10.1038/cr.2013.37>
- Robaczewska, M., Narayan, R., Seigner, B., Schorr, O., Thermet, A., Podhajaska, A. J., ... Cova, L. (2005). Sequence-specific inhibition of duck hepatitis B virus reverse transcription by peptide nucleic acids (PNA). *Journal of Hepatology*, 42(2), 180–7. <http://doi.org/10.1016/j.jhep.2004.10.010>
- Robbins, M., Judge, A., Liang, L., McClintock, K., Yaworski, E., & MacLachlan, I. (2007). 2'-O-methyl-modified RNAs act as TLR7 antagonists. *Molecular Therapy: The Journal of the American Society of Gene Therapy*, 15(9), 1663–9. <http://doi.org/10.1038/sj.mt.6300240>
- Robinson, D., Hamid, Q., Bentley, A., Ying, S., Kay, A. B., & Durham, S. R. (1993). Activation of CD4+ T cells, increased TH2-type cytokine mRNA expression, and eosinophil recruitment in bronchoalveolar lavage after allergen inhalation challenge in patients with atopic asthma. *The Journal of Allergy and Clinical Immunology*, 92(2), 313–24. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8349942>
- Robinson, D. S., Damia, R., Zeibecoglou, K., Molet, S., North, J., Yamada, T., ... Hamid, Q. (1999). CD34(+)/interleukin-5Ralpha messenger RNA+ cells in the bronchial mucosa in asthma: potential airway eosinophil progenitors. *American Journal of Respiratory Cell and Molecular Biology*, 20(1), 9–13. <http://doi.org/10.1165/ajrcmb.20.1.3449>
- Romanet-Manent, S., Charpin, D., Magnan, A., Lanteaume, A., & Vervloet, D. (2002). Allergic vs nonallergic asthma: what makes the difference? *Allergy*, 57(7), 607–13. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12100301>
- Rubin, B. K. (2014). Secretion properties, clearance, and therapy in airway disease. *Translational Respiratory Medicine*, 2(1), 6. <http://doi.org/10.1186/2213-0802-2-6>
- Rutz, M., Metzger, J., Gellert, T., Lippa, P., Lipford, G. B., Wagner, H., & Bauer, S. (2004). Toll-like receptor 9 binds single-stranded CpG-DNA in a sequence- and pH-dependent manner. *European Journal of Immunology*, 34(9), 2541–50. <http://doi.org/10.1002/eji.200425218>
- Sabin, L. R., Delás, M. J., & Hannon, G. J. (2013). Dogma Derailed: The Many Influences of RNA on the Genome. *Molecular Cell*, 49(5), 783–794.

<http://doi.org/10.1016/j.molcel.2013.02.010>

- Salter, S. J., Cox, M. J., Turek, E. M., Calus, S. T., Cookson, W. O., Moffatt, M. F., ... Walker, A. W. (2014). Reagent and laboratory contamination can critically impact sequence-based microbiome analyses. *BMC Biology*, *12*(1), 87. <http://doi.org/10.1186/s12915-014-0087-z>
- Santos, R. D., Raal, F. J., Donovan, J. M., & Cromwell, W. C. (2015). Mipomersen preferentially reduces small low-density lipoprotein particle number in patients with hypercholesterolemia. *Journal of Clinical Lipidology*, *9*(2), 201–209. <http://doi.org/10.1016/j.jacl.2014.12.008>
- Sarvestani, S. T., Stunden, H. J., Behlke, M. A., Forster, S. C., McCoy, C. E., Tate, M. D., ... Gantier, M. P. (2015). Sequence-dependent off-target inhibition of TLR7/8 sensing by synthetic microRNA inhibitors. *Nucleic Acids Research*, *43*(2), 1177–88. <http://doi.org/10.1093/nar/gku1343>
- Saturni, S., Contoli, M., Spanevello, A., & Papi, A. (2015). Models of Respiratory Infections: Virus-Induced Asthma Exacerbations and Beyond. *Allergy, Asthma & Immunology Research*, *7*(6), 525–33. <http://doi.org/10.4168/air.2015.7.6.525>
- Sazani, P., Gemignani, F., Kang, S.-H., Maier, M. A., Manoharan, M., Persmark, M., ... Kole, R. (2002). Systemically delivered antisense oligomers upregulate gene expression in mouse tissues. *Nature Biotechnology*, *20*(12), 1228–33. <http://doi.org/10.1038/nbt759>
- Schlee, M., Hornung, V., & Hartmann, G. (2006). siRNA and isRNA: two edges of one sword. *Molecular Therapy : The Journal of the American Society of Gene Therapy*, *14*(4), 463–70. <http://doi.org/10.1016/j.yth.2006.06.001>
- Schnall-Levin, M., Rissland, O. S., Johnston, W. K., Perrimon, N., Bartel, D. P., & Berger, B. (2011). Unusually effective microRNA targeting within repeat-rich coding regions of mammalian mRNAs. *Genome Research*, *21*(9), 1395–403. <http://doi.org/10.1101/gr.121210.111>
- Schopman, N. C. T., ter Brake, O., & Berkhout, B. (2010). Anticipating and blocking HIV-1 escape by second generation antiviral shRNAs. *Retrovirology*, *7*, 52. <http://doi.org/10.1186/1742-4690-7-52>
- Schraivogel, D., Schindler, S. G., Danner, J., Kremmer, E., Pfaff, J., Hannus, S., ... Meister, G. (2015). Importin-β facilitates nuclear import of human GW proteins and balances cytoplasmic gene silencing protein levels. *Nucleic Acids Research*, *43*(15), 7447–61. <http://doi.org/10.1093/nar/gkv705>
- Schulte-Merker, S., & Stainier, D. Y. R. (2014). Out with the old, in with the new: reassessing morpholino knockdowns in light of genome editing technology. *Development (Cambridge, England)*, *141*(16), 3103–4. <http://doi.org/10.1242/dev.112003>
- Schwartz, J. C., Younger, S. T., Nguyen, N.-B., Hardy, D. B., Monia, B. P., Corey, D. R., & Janowski, B. A. (2008). Antisense transcripts are targets for activating small RNAs. *Nature Structural & Molecular Biology*, *15*(8), 842–8. <http://doi.org/10.1038/nsmb.1444>
- Sedano, C. D., & Sarnow, P. (2014). Hepatitis C virus subverts liver-specific miR-122 to protect the viral genome from exoribonuclease Xrn2. *Cell Host & Microbe*, *16*(2), 257–64. <http://doi.org/10.1016/j.chom.2014.07.006>

- Séguin, R. M., & Ferrari, N. (2009). Emerging oligonucleotide therapies for asthma and chronic obstructive pulmonary disease. *Expert Opinion on Investigational Drugs*, *18*(10), 1505–17. <http://doi.org/10.1517/13543780903179294>
- Sehmi, R., Dorman, S., Baatjes, A., Watson, R., Foley, R., Ying, S., ... Denburg, J. A. (2003). Allergen-induced fluctuation in CC chemokine receptor 3 expression on bone marrow CD34+ cells from asthmatic subjects: significance for mobilization of haemopoietic progenitor cells in allergic inflammation. *Immunology*, *109*(4), 536–46. Retrieved from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1782995&tool=pmcentrez&rendertype=abstract>
- Sehmi, R., Howie, K., Sutherland, D. R., Schragge, W., O'Byrne, P. M., & Denburg, J. A. (1996). Increased levels of CD34+ hemopoietic progenitor cells in atopic subjects. *American Journal of Respiratory Cell and Molecular Biology*, *15*(5), 645–55. <http://doi.org/10.1165/ajrcmb.15.5.8918371>
- Sehmi, R., Wood, L. J., Watson, R., Foley, R., Hamid, Q., O'Byrne, P. M., & Denburg, J. A. (1997). Allergen-induced increases in IL-5 receptor alpha-subunit expression on bone marrow-derived CD34+ cells from asthmatic subjects. A novel marker of progenitor cell commitment towards eosinophilic differentiation. *The Journal of Clinical Investigation*, *100*(10), 2466–75. <http://doi.org/10.1172/JCI119789>
- Sender, V., & Stamme, C. (2014). Lung cell-specific modulation of LPS-induced TLR4 receptor and adaptor localization. *Communicative & Integrative Biology*, *7*, e29053. <http://doi.org/10.4161/cib.29053>
- Seok, J., Warren, H. S., Cuenca, A. G., Mindrinos, M. N., Baker, H. V., Xu, W., ... Tompkins, R. G. (2013). Genomic responses in mouse models poorly mimic human inflammatory diseases. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(9), 3507–12. <http://doi.org/10.1073/pnas.1222878110>
- Sergejeva, S., Johansson, A.-K., Malmhäll, C., & Lötvall, J. (2004). Allergen exposure-induced differences in CD34+ cell phenotype: relationship to eosinophilopoietic responses in different compartments. *Blood*, *103*(4), 1270–7. <http://doi.org/10.1182/blood-2003-05-1618>
- Seth, P. P., Siwkowski, A., Allerson, C. R., Vasquez, G., Lee, S., Prakash, T. P., ... Swayze, E. E. (2009). Short antisense oligonucleotides with novel 2'-4' conformationally restricted nucleoside analogues show improved potency without increased toxicity in animals. *Journal of Medicinal Chemistry*, *52*(1), 10–3. <http://doi.org/10.1021/jm801294h>
- Shepard, A. R., Jacobson, N., & Clark, A. F. (2005). Importance of quantitative PCR primer location for short interfering RNA efficacy determination. *Analytical Biochemistry*, *344*(2), 287–8. <http://doi.org/10.1016/j.ab.2005.06.005>
- Shin, C., Nam, J.-W., Farh, K. K.-H., Chiang, H. R., Shkumatava, A., & Bartel, D. P. (2010). Expanding the microRNA targeting code: functional sites with centered pairing. *Molecular Cell*, *38*(6), 789–802. <http://doi.org/10.1016/j.molcel.2010.06.005>
- Shogren, R., Gerken, T. A., & Jentoft, N. (1989). Role of glycosylation on the conformation and chain dimensions of O-linked glycoproteins: light-scattering studies of ovine submaxillary mucin. *Biochemistry*, *28*(13), 5525–36. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/2775721>

- Singh, M., Mahajan, L., Chaudhary, N., Kaur, S., Madan, T., & Sarma, P. U. (2015). Murine models of Aspergillosis: Role of collectins in host defense. *Indian Journal of Experimental Biology*, 53(11), 691–700. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/26669011>
- Sioud, M., Furset, G., & Cekaite, L. (2007). Suppression of immunostimulatory siRNA-driven innate immune activation by 2'-modified RNAs. *Biochemical and Biophysical Research Communications*, 361(1), 122–6. <http://doi.org/10.1016/j.bbrc.2007.06.177>
- Sipa, K., Sochacka, E., Kazmierczak-Baranska, J., Maszewska, M., Janicka, M., Nowak, G., & Nawrot, B. (2007). Effect of base modifications on structure, thermodynamic stability, and gene silencing activity of short interfering RNA. *RNA (New York, N.Y.)*, 13(8), 1301–16. <http://doi.org/10.1261/rna.538907>
- Sledz, C. A., Holko, M., de Veer, M. J., Silverman, R. H., & Williams, B. R. G. (2003). Activation of the interferon system by short-interfering RNAs. *Nature Cell Biology*, 5(9), 834–9. <http://doi.org/10.1038/ncb1038>
- Song, R., Hennig, G. W., Wu, Q., Jose, C., Zheng, H., & Yan, W. (2011). Male germ cells express abundant endogenous siRNAs. *Proceedings of the National Academy of Sciences of the United States of America*, 108(32), 13159–64. <http://doi.org/10.1073/pnas.1108567108>
- Sorefan, K., Pais, H., Hall, A. E., Kozomara, A., Griffiths-Jones, S., Moulton, V., & Dalmay, T. (2012). Reducing ligation bias of small RNAs in libraries for next generation sequencing. *Silence*, 3(1), 4. <http://doi.org/10.1186/1758-907X-3-4>
- Southam, D. S., Widmer, N., Ellis, R., Hirota, J. A., Inman, M. D., & Sehmi, R. (2005). Increased eosinophil-lineage committed progenitors in the lung of allergen-challenged mice. *The Journal of Allergy and Clinical Immunology*, 115(1), 95–102. <http://doi.org/10.1016/j.jaci.2004.09.022>
- Soutschek, J., Akinc, A., Bramlage, B., Charisse, K., Constien, R., Donoghue, M., ... Vornlocher, H.-P. (2004). Therapeutic silencing of an endogenous gene by systemic administration of modified siRNAs. *Nature*, 432(7014), 173–8. <http://doi.org/10.1038/nature03121>
- Staton, A. A., & Giraldez, A. J. (2011). Use of target protector morpholinos to analyze the physiological roles of specific miRNA-mRNA pairs in vivo. *Nature Protocols*, 6(12), 2035–49. <http://doi.org/10.1038/nprot.2011.423>
- Stein, C. A., Hansen, J. B., Lai, J., Wu, S., Voskresenskiy, A., Høg, A., ... Koch, T. (2010). Efficient gene silencing by delivery of locked nucleic acid antisense oligonucleotides, unassisted by transfection reagents. *Nucleic Acids Research*, 38(1), e3. <http://doi.org/10.1093/nar/gkp841>
- Straarup, E. M., Fisker, N., Hedtjærn, M., Lindholm, M. W., Rosenbohm, C., Aarup, V., ... Koch, T. (2010). Short locked nucleic acid antisense oligonucleotides potently reduce apolipoprotein B mRNA and serum cholesterol in mice and non-human primates. *Nucleic Acids Research*, 38(20), 7100–11. <http://doi.org/10.1093/nar/gkq457>
- Summerton, J. (1999). Morpholino antisense oligomers: the case for an RNase H-independent structural type. *Biochimica et Biophysica Acta (BBA) - Gene Structure and Expression*, 1489(1), 141–158. [http://doi.org/10.1016/S0167-4781\(99\)00150-5](http://doi.org/10.1016/S0167-4781(99)00150-5)
- Summerton, J. E. (2007). Morpholino, siRNA, and S-DNA compared: impact of structure



and mechanism of action on off-target effects and sequence specificity. *Current Topics in Medicinal Chemistry*, 7(7), 651–60. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17430206>

- SUMMERTON, J., & WELLER, D. (1997). Morpholino Antisense Oligomers: Design, Preparation, and Properties. *Antisense and Nucleic Acid Drug Development*, 7(3), 187–195. <http://doi.org/10.1089/oli.1.1997.7.187>
- Sundaram, P., Kurniawan, H., Byrne, M. E., & Wower, J. (2013). Therapeutic RNA aptamers in clinical trials. *European Journal of Pharmaceutical Sciences : Official Journal of the European Federation for Pharmaceutical Sciences*, 48(1-2), 259–71. <http://doi.org/10.1016/j.ejps.2012.10.014>
- Taberero, J., Shapiro, G. I., LoRusso, P. M., Cervantes, A., Schwartz, G. K., Weiss, G. J., ... Burris, H. A. (2013). First-in-humans trial of an RNA interference therapeutic targeting VEGF and KSP in cancer patients with liver involvement. *Cancer Discovery*, 3(4), 406–17. <http://doi.org/10.1158/2159-8290.CD-12-0429>
- Takao, K., & Miyakawa, T. (2015). Genomic responses in mouse models greatly mimic human inflammatory diseases. *Proceedings of the National Academy of Sciences of the United States of America*, 112(4), 1167–72. <http://doi.org/10.1073/pnas.1401965111>
- Tam, O. H., Aravin, A. A., Stein, P., Girard, A., Murchison, E. P., Cheloufi, S., ... Hannon, G. J. (2008). Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature*, 453(7194), 534–8. <http://doi.org/10.1038/nature06904>
- Tang, R., Li, L., Zhu, D., Hou, D., Cao, T., Gu, H., ... Zen, K. (2012). Mouse miRNA-709 directly regulates miRNA-15a/16-1 biogenesis at the posttranscriptional level in the nucleus: evidence for a microRNA hierarchy system. *Cell Research*, 22(3), 504–15. <http://doi.org/10.1038/cr.2011.137>
- Tanganyika-de Winter, C. L., Heemskerk, H., Karnaoukh, T. G., van Putten, M., de Kimpe, S. J., van Deutekom, J., & Aartsma-Rus, A. (2012). Long-term Exon Skipping Studies With 2'-O-Methyl Phosphorothioate Antisense Oligonucleotides in Dystrophic Mouse Models. *Molecular Therapy. Nucleic Acids*, 1, e44. <http://doi.org/10.1038/mtna.2012.38>
- Tay, Y., Zhang, J., Thomson, A. M., Lim, B., & Rigoutsos, I. (2008). MicroRNAs to Nanog, Oct4 and Sox2 coding regions modulate embryonic stem cell differentiation. *Nature*, 455(7216), 1124–8. <http://doi.org/10.1038/nature07299>
- Templin, M. V., Levin, A. A., Graham, M. J., Åberg, P. M., Axelsson, B. I., Butler, M., ... Bennett, C. F. (2000). Pharmacokinetic and Toxicity Profile of a Phosphorothioate Oligonucleotide Following Inhalation Delivery to Lung in Mice. *Antisense and Nucleic Acid Drug Development*, 10(5), 359–368. <http://doi.org/10.1089/oli.1.2000.10.359>
- Thibault, P. A., Huys, A., Amador-Cañizares, Y., Gailius, J. E., Pinel, D. E., & Wilson, J. A. (2015). Regulation of Hepatitis C Virus Genome Replication by Xrn1 and MicroRNA-122 Binding to Individual Sites in the 5' Untranslated Region. *Journal of Virology*, 89(12), 6294–311. <http://doi.org/10.1128/JVI.03631-14>
- Tsitsiou, E., Williams, A. E., Moschos, S. A., Patel, K., Rossios, C., Jiang, X., ... Lindsay, M. A. (2012). Transcriptome analysis shows activation of circulating CD8+ T cells in patients with severe asthma. *J Allergy Clin Immunol*, 129(1), 95–103. [http://doi.org/S0091-6749\(11\)01310-8](http://doi.org/S0091-6749(11)01310-8) [pii]10.1016/j.jaci.2011.08.011
- Turner, J. J., Jones, S. W., Moschos, S. A., Lindsay, M. A., & Gait, M. J. (2007). MALDI-

- TOF mass spectral analysis of siRNA degradation in serum confirms an RNase A-like activity. *Mol Biosyst*, 3(1), 43–50. <http://doi.org/10.1039/b611612d>
- Uhlmann, E. (1998). Peptide nucleic acids (PNA) and PNA-DNA chimeras: from high binding affinity towards biological function. *Biological Chemistry*, 379(8-9), 1045–52. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9792437>
- Usmani, O. S. (2014). Small airways dysfunction in asthma: evaluation and management to improve asthma control. *Allergy, Asthma & Immunology Research*, 6(5), 376–88. <http://doi.org/10.4168/aaair.2014.6.5.376>
- Valoczi, A., Hornyik, C., Varga, N., Burgyan, J., Kauppinen, S., & Havelda, Z. (2004). Sensitive and specific detection of microRNAs by northern blot analysis using LNA-modified oligonucleotide probes. *Nucleic Acids Res*, 32(22), e175. <http://doi.org/10.1093/nar/gnh171>
- van de Wetering, J. K., van Remoortere, A., Vaandrager, A. B., Batenburg, J. J., van Golde, L. M. G., Hokke, C. H., & van Hellemond, J. J. (2004). Surfactant protein D binding to terminal alpha1-3-linked fucose residues and to *Schistosoma mansoni*. *American Journal of Respiratory Cell and Molecular Biology*, 31(5), 565–72. <http://doi.org/10.1165/rcmb.2004-0105OC>
- van der Ree, M. H., van der Meer, A. J., de Bruijne, J., Maan, R., van Vliet, A., Welzel, T. M., ... Reesink, H. W. (2014). Long-term safety and efficacy of microRNA-targeted therapy in chronic hepatitis C patients. *Antiviral Research*, 111, 53–9. <http://doi.org/10.1016/j.antiviral.2014.08.015>
- van Deutekom, J. C., Janson, A. A., Ginjaar, I. B., Frankhuizen, W. S., Aartsma-Rus, A., Bremmer-Bout, M., ... van Ommen, G.-J. B. (2007). Local dystrophin restoration with antisense oligonucleotide PRO051. *The New England Journal of Medicine*, 357(26), 2677–86. <http://doi.org/10.1056/NEJMoa073108>
- van Dijk, E. L., Jaszczyszyn, Y., & Thermes, C. (2014). Library preparation methods for next-generation sequencing: tone down the bias. *Experimental Cell Research*, 322(1), 12–20. <http://doi.org/10.1016/j.yexcr.2014.01.008>
- Vattanasit, U., Navasumrit, P., Khadka, M. B., Kanitwithayanun, J., Promvijit, J., Autrup, H., & Ruchirawat, M. (2014). Oxidative DNA damage and inflammatory responses in cultured human cells and in humans exposed to traffic-related particles. *International Journal of Hygiene and Environmental Health*, 217(1), 23–33. <http://doi.org/10.1016/j.ijheh.2013.03.002>
- Veldhoen, S., Laufer, S. D., & Restle, T. (2008). Recent developments in peptide-based nucleic acid delivery. *International Journal of Molecular Sciences*, 9(7), 1276–320. <http://doi.org/10.3390/ijms9071276>
- Vickers, T. A., & Croke, S. T. (2015). The rates of the major steps in the molecular mechanism of RNase H1-dependent antisense oligonucleotide induced degradation of RNA. *Nucleic Acids Research*, 43(18), 8955–63. <http://doi.org/10.1093/nar/gkv920>
- Warren, T. K., Warfield, K. L., Wells, J., Swenson, D. L., Donner, K. S., Van Tongeren, S. A., ... Bavari, S. (2010). Advanced antisense therapies for postexposure protection against lethal filovirus infections. *Nature Medicine*, 16(9), 991–4. <http://doi.org/10.1038/nm.2202>
- Warren, T. K., Whitehouse, C. A., Wells, J., Welch, L., Heald, A. E., Charleston, J. S., ...

- Bavari, S. (2015). A single phosphorodiamidate morpholino oligomer targeting VP24 protects rhesus monkeys against lethal Ebola virus infection. *mBio*, 6(1). <http://doi.org/10.1128/mBio.02344-14>
- Watanabe, T. A., Geary, R. S., & Levin, A. A. (2006). Plasma protein binding of an antisense oligonucleotide targeting human ICAM-1 (ISIS 2302). *Oligonucleotides*, 16(2), 169–80. <http://doi.org/10.1089/oli.2006.16.169>
- Watanabe, T., Totoki, Y., Toyoda, A., Kaneda, M., Kuramochi-Miyagawa, S., Obata, Y., ... Sasaki, H. (2008). Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature*, 453(7194), 539–43. <http://doi.org/10.1038/nature06908>
- Wei, W., Ba, Z., Gao, M., Wu, Y., Ma, Y., Amiard, S., ... Qi, Y. (2012). A role for small RNAs in DNA double-strand break repair. *Cell*, 149(1), 101–12. <http://doi.org/10.1016/j.cell.2012.03.002>
- Weinmann, L., Höck, J., Ivacevic, T., Ohrt, T., Mütze, J., Schwille, P., ... Meister, G. (2009). Importin 8 is a gene silencing factor that targets argonaute proteins to distinct mRNAs. *Cell*, 136(3), 496–507. <http://doi.org/10.1016/j.cell.2008.12.023>
- Wengel, J. (1999). Synthesis of 3'-C- and 4'-C-Branched Oligodeoxynucleotides and the Development of Locked Nucleic Acid (LNA). *Accounts of Chemical Research*, 32(4), 301–310. <http://doi.org/10.1021/ar980051p>
- Wenzel, S. E. (2006). Asthma: defining of the persistent adult phenotypes. *Lancet (London, England)*, 368(9537), 804–13. [http://doi.org/10.1016/S0140-6736\(06\)69290-8](http://doi.org/10.1016/S0140-6736(06)69290-8)
- Werner, A., Cockell, S., Falconer, J., Carlile, M., Alnumeir, S., & Robinson, J. (2014). Contribution of natural antisense transcription to an endogenous siRNA signature in human cells. *BMC Genomics*, 15, 19. <http://doi.org/10.1186/1471-2164-15-19>
- Wesolowski, D., Alonso, D., & Altman, S. (2013). Combined effect of a peptide-morpholino oligonucleotide conjugate and a cell-penetrating peptide as an antibiotic. *Proceedings of the National Academy of Sciences of the United States of America*, 110(21), 8686–9. <http://doi.org/10.1073/pnas.1306911110>
- Whitsett, J. A. (2010). Review: The intersection of surfactant homeostasis and innate host defense of the lung: lessons from newborn infants. *Innate Immunity*, 16(3), 138–42. <http://doi.org/10.1177/1753425910366879>
- Whitsett, J. A., Wert, S. E., & Weaver, T. E. (2015). Diseases of pulmonary surfactant homeostasis. *Annual Review of Pathology*, 10, 371–93. <http://doi.org/10.1146/annurev-pathol-012513-104644>
- Williams, K., & Roman, J. (2015). Studying human respiratory disease in animals - Role of induced and naturally-occurring models. *The Journal of Pathology*, 238(2), 220–32. <http://doi.org/10.1002/path.4658>
- Wolfrum, C., Shi, S., Jayaprakash, K. N., Jayaraman, M., Wang, G., Pandey, R. K., ... Stoffel, M. (2007). Mechanisms and optimization of in vivo delivery of lipophilic siRNAs. *Nature Biotechnology*, 25(10), 1149–57. <http://doi.org/10.1038/nbt1339>
- Wright, P. F., Ikizler, M. R., Gonzales, R. A., Carroll, K. N., Johnson, J. E., & Werkhaven, J. A. (2005). Growth of respiratory syncytial virus in primary epithelial cells from the human respiratory tract. *Journal of Virology*, 79(13), 8651–4.

<http://doi.org/10.1128/JVI.79.13.8651-8654.2005>

- Wu, B., Lu, P., Benrashid, E., Malik, S., Ashar, J., Doran, T. J., & Lu, Q. L. (2010). Dose-dependent restoration of dystrophin expression in cardiac muscle of dystrophic mice by systemically delivered morpholino. *Gene Therapy*, *17*(1), 132–40. <http://doi.org/10.1038/gt.2009.120>
- Wu, H., Lima, W. F., & Crooke, S. T. (1999). Properties of cloned and expressed human RNase H1. *The Journal of Biological Chemistry*, *274*(40), 28270–8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10497183>
- Xaubet, A., Ancochea, J., Bollo, E., Fernández-Fabrellas, E., Franquet, T., Molina-Molina, M., ... Serrano-Mollar, A. (2013). Guidelines for the Diagnosis and Treatment of Idiopathic Pulmonary Fibrosis. *Archivos de Bronconeumología (English Edition)*, *49*(8), 343–353. <http://doi.org/10.1016/j.arbr.2013.06.003>
- Yamakawa, N., Okuyama, K., Ogata, J., Kanai, A., Helwak, A., Takamatsu, M., ... Kotani, A. (2014). Novel functional small RNAs are selectively loaded onto mammalian Ago1. *Nucleic Acids Research*, *42*(8), 5289–301. <http://doi.org/10.1093/nar/gku137>
- Yang, T. Y., & Jeong, S. (2013). Grouped False-Discovery Rate for Removing the Gene-Set-Level Bias of RNA-seq. *Evolutionary Bioinformatics Online*, *9*, 467–78. <http://doi.org/10.4137/EBO.S13099>
- Younger, S. T., & Corey, D. R. (2011). Transcriptional gene silencing in mammalian cells by miRNA mimics that target gene promoters. *Nucleic Acids Research*, *39*(13), 5682–91. <http://doi.org/10.1093/nar/gkr155>
- Zamora, M. R., Budev, M., Rolfé, M., Gottlieb, J., Humar, A., Devincenzo, J., ... Glanville, A. R. (2011a). RNA interference therapy in lung transplant patients infected with respiratory syncytial virus. *American Journal of Respiratory and Critical Care Medicine*, *183*(4), 531–8. <http://doi.org/10.1164/rccm.201003-0422OC>
- Zamora, M. R., Budev, M., Rolfé, M., Gottlieb, J., Humar, A., Devincenzo, J., ... Glanville, A. R. (2011b). RNA interference therapy in lung transplant patients infected with respiratory syncytial virus. *American Journal of Respiratory and Critical Care Medicine*, *183*(4), 531–8. <http://doi.org/10.1164/rccm.201003-0422OC>
- Zardo, G., Ciolfi, A., Vian, L., Billi, M., Racanicchi, S., Grignani, F., & Nervi, C. (2012). Transcriptional targeting by microRNA-polycomb complexes: a novel route in cell fate determination. *Cell Cycle (Georgetown, Tex.)*, *11*(19), 3543–9. <http://doi.org/10.4161/cc.21468>
- Zasłona, Z., Przybranowski, S., Wilke, C., van Rooijen, N., Teitz-Tennenbaum, S., Osterholzer, J. J., ... Peters-Golden, M. (2014). Resident alveolar macrophages suppress, whereas recruited monocytes promote, allergic lung inflammation in murine models of asthma. *Journal of Immunology (Baltimore, Md. : 1950)*, *193*(8), 4245–53. <http://doi.org/10.4049/jimmunol.1400580>
- Zhang, X., Shan, P., Jiang, D., Noble, P. W., Abraham, N. G., Kappas, A., & Lee, P. J. (2004). Small interfering RNA targeting heme oxygenase-1 enhances ischemia-reperfusion-induced lung apoptosis. *The Journal of Biological Chemistry*, *279*(11), 10677–84. <http://doi.org/10.1074/jbc.M312941200>
- Zhang, X., Zuo, X., Yang, B., Li, Z., Xue, Y., Zhou, Y., ... Fu, X.-D. (2014). MicroRNA directly enhances mitochondrial translation during muscle differentiation. *Cell*, *158*(3),

607–19. <http://doi.org/10.1016/j.cell.2014.05.047>

Zhang, Y., Fan, M., Zhang, X., Huang, F., Wu, K., Zhang, J., ... Zhang, H. (2014). Cellular microRNAs up-regulate transcription via interaction with promoter TATA-box motifs. *RNA (New York, N.Y.)*, *20*(12), 1878–89. <http://doi.org/10.1261/rna.045633.114>

Zhu, B., Haghi, M., Goud, M., Young, P. M., & Traini, D. (2015). The formulation of a pressurized metered dose inhaler containing theophylline for inhalation. *European Journal of Pharmaceutical Sciences : Official Journal of the European Federation for Pharmaceutical Sciences*, *76*, 68–72. <http://doi.org/10.1016/j.ejps.2015.04.016>

Zuo, L., Lucas, K., Fortuna, C. A., Chuang, C.-C., & Best, T. M. (2015). Molecular Regulation of Toll-like Receptors in Asthma and COPD. *Frontiers in Physiology*, *6*, 312. <http://doi.org/10.3389/fphys.2015.00312>