



*Citation for published version:*

Morgan, B 2017, 'lattice\_mc: A Python Lattice-Gas Monte Carlo Module', *The Journal of Open Source Software*, vol. 2, no. 13, 00247. <https://doi.org/10.5281/zenodo.546262>, <https://doi.org/10.5281/zenodo.582402>, <https://doi.org/10.21105/joss.00247>

*DOI:*

[10.5281/zenodo.546262](https://doi.org/10.5281/zenodo.546262)

[10.5281/zenodo.582402](https://doi.org/10.5281/zenodo.582402)

[10.21105/joss.00247](https://doi.org/10.21105/joss.00247)

*Publication date:*

2017

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication](#)

*Publisher Rights*

Unspecified

## University of Bath

### Alternative formats

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# lattice\_mc: A Python Lattice-Gas Monte Carlo Module

Benjamin J. Morgan<sup>1</sup>

<sup>1</sup>Department of Chemistry, University of Bath, Bath, BA2 7AY, United Kingdom.

13 April 2017

**Paper DOI:** <http://dx.doi.org/10.21105/joss.00247>

**Software Repository:** [https://github.com/bjmorgan/lattice\\_mc](https://github.com/bjmorgan/lattice_mc)

**Software Archive:** <http://dx.doi.org/10.5281/zenodo.582402>

## Summary

`lattice_mc` is Python module for performing (kinetic) lattice-gas Monte Carlo (LGMC) simulations of ionic transport in solid electrolytes.

In solid electrolytes, ionic motion is typically effected by a series of discrete “jumps” where ions move between adjacent lattice sites (Catlow 1983). For dilute mobile ions, ionic trajectories are random walks, and simple analytical expressions exist that relate macroscopic transport coefficients, i.e. diffusion coefficients and ionic conductivities, to the microscopic jump frequency of individual ions (Howard and Lidiard 1964; Harding 1989). Practical solid electrolytes have high mobile ion concentrations, with significant interparticle interactions producing deviations from the dilute limit random walk behaviour. In general, the *quantitative* effect of interparticle interactions cannot be determined analytically. As an alternative, numerical simulations, such as lattice-gas Monte Carlo methods, can be used to directly calculate these relationships. Lattice-gas Monte Carlo methods are particularly suited to studying how varying properties across broad classes of materials give quantitative differences in macroscopic ionic transport, and can be used to understand the different transport properties of materials with, for example, different crystal structures or mobile ion stoichiometries.

`lattice_mc` has been written to allow materials scientists and solid-state chemists model how the microscopic physics of solid electrolytes (crystal structure, stoichiometry, interaction models) determine macroscopic transport behaviour (diffusion and ionic conduction), with the goal of understanding the factors that make different materials more or less useful for specific applications (e.g. Li-ion batteries or fuel cells).

The code allows the programmatic construction of simple lattices (presently implemented are square, honeycomb, and cubic lattices). Lattices with arbitrary geometries can be constructed from a file format that defines the lattice sites and their connectivity, allowing models based on crystallographic data. Calculated properties include tracer and “jump” diffusion coefficients; where the latter is proportional to the mobility (and hence the conductivity for charged particles) (Van der Ven, Bhattacharya, and Belak 2013); and tracer (single particle) and collective correlation factors,  $f$  and  $f_I$  (Murch 1982). The simplest interaction model is for “non-interacting” particles, where the only restriction is volume exclusion (two particles cannot simultaneously occupy a single site) (Kutner 1981). Additional interaction models include nearest-neighbour repulsion and on-site energies for inequivalent sites. Simulations are performed using an efficient rejection-free Monte Carlo scheme (Voter 2007).

## Acknowledgements

BJM acknowledges support from the Royal Society (UF130329).

## References

- Catlow, C. Richard A. 1983. “Static Lattice Simulation of Structure and Transport in Superionic Conductors.” *Sol. Stat. Ionics* 8: 89–107.
- Harding, J. F. 1989. “Defects and Transport in Ionic Solids.” In *Ionic Solids at High Temperatures*, edited by A. M. Stoneham. World Scientific. doi:10.1142/9789814503228\_0003.
- Howard, R. E., and A. B. Lidiard. 1964. “Matter Transport in Solids.” *Rep Prog Phys.* 27: 161–240.
- Kutner, R. 1981. “Chemical Diffusion in the Lattice Gas of Non-Interacting Particles.” *Phys. Lett.* 81A (4): 239–40.
- Murch, G. E. 1982. “The Haven Ratio in Fast Ionic Conductors.” *Sol. Stat. Ionics* 7 (3): 177–98.
- Van der Ven, Anton, Jishnu Bhattacharya, and Anna A. Belak. 2013. “Understanding Li Diffusion in Li-Intercalation Compounds.” *Acc. Chem. Res.* 46 (5): 1216–25.
- Voter, Arthur F. 2007. “Introduction to the Kinetic Monte Carlo Method.” In *Radiation Effects in Solids*, edited by Kurt E. Sickafus, Eugene A. Kotomin, and Blas P. Uberuaga, 1–23. Dordrecht: Springer Netherlands. doi:10.1007/978-1-4020-5295-8\_1.