



Citation for published version:

Loades, M, Coetzee, B, du Toit, S & Kagee, A 2018, '...But I'm still tired': The experience of fatigue among South African adolescents receiving antiretroviral therapy.', *Aids Care-Psychological and Socio-Medical Aspects of Aids/Hiv*, vol. 30, no. 6, pp. 746-750. <https://doi.org/10.1080/09540121.2017.1394433>

DOI:

[10.1080/09540121.2017.1394433](https://doi.org/10.1080/09540121.2017.1394433)

Publication date:

2018

Document Version

Peer reviewed version

[Link to publication](#)

This is an Accepted Manuscript of an article published by Taylor & Francis in *AIDS Care* on 25 October 2017, available online: <http://www.tandfonline.com/10.1080/09540121.2017.1394433>.

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Title: ‘...But I’m still tired’: The experience of fatigue among South African adolescents receiving antiretroviral therapy

Authors: Loades, M.E¹., Coetzee, B²., Du Toit, S²., & Kagee, A²

Affiliations

Department of Psychology, University of Bath, England

Department of Psychology, Stellenbosch University, South Africa

Corresponding Author

Dr Maria Loades, Department of Psychology, University of Bath, Claverton Down, Bath BA2 7AY, England; m.e.loades@bath.ac.uk

Acknowledgements

We would like to acknowledge Stellenbosch Universities Sub-Committee A, as well as the University of Bath’s Faculty of Humanities and Social Sciences & Department of Psychology for funding this research.

Dr Loades is funded by the National Institute for Health Research (Doctoral Research Fellowship, DRF-2016-09-021). This report is independent research. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, The National Institute for Health Research or the Department of Health.

Title: ‘...But I’m still tired’: The experience of fatigue among South African adolescents receiving antiretroviral therapy

Abstract

In adults with HIV, fatigue is a common and disabling symptom associated with suboptimal adherence to antiretroviral therapy (ART), poor health and well-being, reduced economic productivity, and increased health care resource use. Fatigue among adolescents with HIV is relatively unexplored. We recruited 14 adolescents with HIV receiving ART in South Africa whom we interviewed about fatigue. We used thematic analysis to analyse the data.

Participants reported difficulties in maintaining concentration at school and resorting to self-care strategies such as napping during school hours. Adolescents also described actively avoiding activities thought to lead to fatigue and spoke of being socially excluded by their peers. The findings demonstrate that adolescents experienced symptoms consistent with a definition of fatigue, which had implications for their academic and social lives.

Keywords: HIV, antiretroviral therapy, adolescents, fatigue, qualitative

Introduction

Fatigue refers to decreased capacity for activity due to an imbalance in the availability of resources needed to perform that activity. “Chronic fatigue”, which is ongoing physical and/or mental exhaustion and tiredness that persists for at least one month (Voss, Portillo, Holzemer, & Dodd, 2007) is commonly reported among specific chronic illness populations, including rheumatoid arthritis (Varni, Burwinkle, & Szer, 2004), chronic pain (Gold, Mahrer, Yee, & Palermo, 2009), neurological conditions (Nutini, Karczewski, & Capoor, 2009) and multiple sclerosis (Banwell et al., 2007).

Among adults with HIV, fatigue is a common and disabling symptom that is associated with poorer treatment adherence (Gay et al., 2011), greater functional impairment, and lower survival rates (Justice et al., 1999), and with poorer health, lower work productivity, and increased health care resource use (DiBonaventura, Gupta, Cho, & Mrus, 2012).

To our knowledge, no research has been conducted on fatigue in adolescents living with HIV. We explored the lived experience of fatigue among a sample of South African adolescents receiving antiretroviral treatment (ART).

Method

Setting and participants

Participants were HIV-infected adolescents aged 11-18, attending ART clinics in South

Africa. Participants had to be competent in either English or Afrikaans, and not require urgent medical attention; or have severe mental health problems (e.g. psychosis).

Procedure

We received approval from the Stellenbosch University Health Research Ethics Committee, and permission from relevant health authority and clinic managers.

Interviews were audio recorded, transcribed and anonymised, and translated into English where necessary. Thematic analysis was used to analyse the data (Braun & Clarke, 2006). Themes emerged from the data, with no themes specified a priori.

Measures

Interviews were guided by an interview schedule (appendix 1) exploring how fatigue impacted participants' day-to-day activities, and how they managed and coped with fatigue. The Chalder Fatigue Scale (CFS) (Chalder et al., 1993) measured fatigue severity, and the Revised Children's Anxiety and Depression Scale (RCADS) depression and generalised anxiety subscales measured mood and anxiety (Chorpita, Yim, Moffitt, Umemoto, & Francis, 2000).

Results

Fourteen participants were interviewed (table 1). Using a cut-off score of ≥ 4 (Cella & Chalder, 2010), bimodal scoring on the CFS identified 12 of the 14 adolescents as fatigued.

Participants reported sleep disturbance, unrefreshing sleep, pain, poor concentration and/or memory, and a lack of energy which were not ameliorated by sleep. We identified three overarching themes namely, the impact of fatigue on daily functioning, strategies to

overcome excessive tiredness, and adolescents' understanding and experience of their illness.

Impact on Functioning

Participants avoided vigorous physical activities such as sports as a result of their fatigue. For example, Cassandra stated "*I don't feel like doing the sport anymore because it just makes me more tired than I already am*".

Fatigue impacted on functioning and productivity in school and participants reported sleeping at their desks. Lisa stated "... *sometimes when my teacher is teaching I am very tired and I sleep in the class so my teacher will shout at me....*"

Due to poor concentration and memory and a general lack of energy, adolescents struggled with the demands of school.

When I'm doing my homework then I will become tired and just close the books and think that I will finish them in the evening or the next morning. (Brenda).

Fatigue influenced participants' social interactions. Lisa said "*When I play with my friends I feel tired and I just want to sleep, sometimes*". Participants felt that their peers did not understand their tiredness. For example, Cassandra reported that "*They can't understand why I'm tired every time.*"

Adolescents stated that their fatigue resulted in social exclusion, which for some generated difficult feelings including anger. For example, Cassandra, who wanted to sleep rather than to socialise stated:

When I'm tired ... I just want to be alone and sleep. Sometimes my friends come to my house and then I can't sleep and then they want to play there and then I get very angry.

Participants spoke about sadness and mood fluctuations, for example *"I feel so happy sometimes, sometimes sad"* (Derek). Mood was not consistently seen as directly linked to tiredness, and participants could separate tiredness from low mood:

...my mood is not like my tiredness.... My mood only comes now and then. But my tiredness stays. (Brenda)

Participants lacked motivation as a result of their tiredness. Lisa stated *"I feel tired, I don't want to do some things, I just want to rest"*.

What helps?

Helpful strategies included the support of friends and family, rest, and activities which adolescents purposefully engaged in to stay awake.

Short naps in the afternoons, and sometimes during school hours were helpful. Christine said *"I stay in class during break and sleep. When the bell rings I wake up again."*

Activities like cooking, eating, walking, exercising, playing and reading provided stimulation to remain alert.

Me and my sister would always go for a walk but not too far from our home, just walk around a bit.” (Cassandra)

Motivation to persist through tiredness was enhanced by support from friends, teachers and family members. Friends relieved some of the fatigue through engagement in conversation and encouragement to play. Brenda stated, *“They are like normal friends. When I tell them that I am tired they will tell me that I can go and lay down or they will do something to wake me up or so”*.

Illness perceptions

Some participants attributed their fatigue to exertion and activity: for example, *“Because on the Monday we always do a lot of work at school so then I feel tired on a Tuesday”* (Faith).

Other participants attributed their fatigue to hunger, or to medication: *“I think it’s the pills because after I drink the pills it keeps me awake and then I feel tired during the day”* (Cassandra), but not all agreed:

I don’t think it’s my medication, because I take my medication at night and it’s usually during the day that I feel so tired. (Justin)

Participants had difficulty making sense of their tiredness, which for some was distressing, for example:

When it started, I was at the school and I sleep in school, ...and my teacher shouted at me and I cried because I didn’t know what was wrong me. (Lisa)

Discussion

Adolescents with HIV experienced symptoms consistent with a definition of abnormal fatigue (Aaronson et al., 1999), which impacted on their quality of life, most notably on their school performance and their social lives.

Given that adolescents with HIV are cognitively worse off than those without HIV (Laughton, Cornell, Boivin, & Van Rie, 2013; Sherr, Croome, Castaneda, Bradshaw, & Romero, 2014) it is essential that these additional apparently debilitating symptoms of fatigue are identified early on and remediated for where possible.

Our participants withdrew from social life and abandoned sports, a social activity crucial for forming and maintaining relationships with peers. Prolonged avoidance of activity may escalate depressed mood and perpetuate feelings of anxiety when confronted with activity again. Fatigue is also associated with a loss of motivation to take part in activities, a common consequence of tiredness experienced by adults with HIV (Jenkin, Koch, & Kralik, 2006), and paediatric populations with fatigue (Parslow et al., 2017).

Adolescents spoke of deliberate exclusion from social activities by their peers and perceived this an inability of their peers to understand their experiences of fatigue. Repeated instances of social exclusion led to anger and perpetuated a cycle of negative feelings and thoughts which are consistent with other studies (Power, Badley, French, Wall, & Hawker, 2008; Stormorken, Jason, & Kirkevold, 2015).

Similar to other populations, adolescents had strategies that helped manage their tiredness, such as rest and moderate activity (Jenkin et al., 2006) and social support (Mburu et al., 2014).

Expressing experiences of fatigue, which is a complex and inherently subjective concept,

requires clear articulation of the problem (Jenkin et al., 2006), and may be difficult for adolescents. Although our participants did not attribute their fatigue to a single cause, they were able to distinguish it from tiredness or sleepiness following taking their medication.

Strengths and limitations

Our data provide insights into the complexity of fatigue experiences in this population and the language used to articulate these experiences. However, we only included adolescents who were able to speak English and Afrikaans. The study design does not enable us to disentangle the complex relationships between fatigue and mood, although participants did seem to view these as separable, and those with high levels of fatigue did not necessarily also score more highly on depression or anxiety.

Conclusions

Fatigue is a widespread and disabling problem among adults with HIV, and may be also be salient among adolescent ART users. Further research may include investigations of the prevalence and correlates of fatigue among adolescents living with HIV (including mood and anxiety), the efficacy of psychosocial interventions aimed at ameliorating fatigue, and the role of fatigue in ART adherence. Comparing the experience of fatigue in adolescents with HIV to other chronic disease populations, to mental health patients, and to healthy individuals would further the understanding of the similarities and differences between groups.

Table 1. Sample characteristics of adolescents

Pseudo nym	Age	Gender	Ethnicity	Chalder Fatigue Scale Score (bimodal)	RCADS Depression Subscale Score	RCADS Generalised Anxiety Subscale Score
Lisa	14	Female	African	9	17	15
James	12	Male	Coloured	3	2	5
John	13	Male	Coloured	1	5	4
Jenny	18	Female	Black	4	3	15
Victor	16	Male	Black	6	9	6
Derek	13	Male	Black	6	7	10
Justin	17	Male	Coloured	5	7	1
Brenda	15	Female	Black	5	12	5
Christi ne	12	Female	Black	7	2	6
Violet	14	Female	Black	4	4	1
Steven	16	Male	Black	11	12	5
Lucy	13	Female	Coloured	7	0	5
Cassan dra	17	Female	Black	8	16	12
Faith	18	Female	Black	9	7	11

Appendix 1: Semi-structured interview topic guide

The Lived Experience of Fatigue in Adolescents receiving medication: A qualitative study.

TOPIC GUIDE

To explore the lived experience of fatigue in adolescents receiving medication and coping styles, daily routines and sleep, and their perception of the causes

These questions will be used as prompts to ensure all important areas are covered.

Welcome, introduction, confidentiality. Discuss consent, sign form or check continues to be happy with consent.

1. Tell me how fatigue/tiredness/exhaustion affects your life.

Prompts: how does fatigue/tiredness/exhaustion interfere with your daily life? School? Work? Social life? Family?

2. How do you do things differently day-to-day because of the fatigue?

Prompts: Are there things that you can't do because you are fatigued? What are they? Things you would do if you didn't feel so fatigued/tired?

3. What do you think is making you fatigued/tired?

4. How does fatigue/tiredness/exhaustion make you feel?

Prompts: Sad? Frustrated? Worried? If so, why?

5. What do you do to help yourself to feel better when you are feeling fatigued/tired/exhausted?

Prompts: What makes it worse? What makes it better?

6. Tell me about your sleep. Do you have problems with sleeping?

Prompts: How is getting to sleep? And can you stay asleep or do you wake up often?

How is waking up? Do you feel rested when you wake up?

7. Tell me about your mood.

Prompts: how does your mood affect your daily life? School? Work? Social life?

Family? How do you do things differently day-to-day because of your mood?

8. Do you think that fatigue/tiredness/exhaustion, sleep and mood are connected for you? If so, how?

Prompts: what comes first for you? Fatigue/bad sleep/bad or low mood?

9. Do you have anything else you would like to add? Do you have any questions?

References

- Aaronson, L. S., Teel, C. S., Cassmeyer, V., Neuberger, G. B., Pallikkathayil, L., Pierce, J., . . . Wingate, A. (1999). Defining and measuring fatigue. *Image: the journal of nursing scholarship*, 31(1), 45-50.
- Banwell, B., Krupp, L., Kennedy, J., Tellier, R., Tenenbaum, S., Ness, J., . . . Waubant, E. (2007). Clinical features and viral serologies in children with multiple sclerosis: a multinational observational study. *The Lancet Neurology*, 6(9), 773-781.
- Braun, V., & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative research in psychology*, 3(2), 77-101.
- Cella, M., & Chalder, T. (2010). Measuring fatigue in clinical and community settings. *J Psychosom Res*, 69(1), 17-22.
- Chalder, T., Berelowitz, G., Pawlikowska, T., Watts, L., Wessely, S., Wright, D., & Wallace, E. (1993). Development of a fatigue scale. *J Psychosom Res*, 37(2), 147-153.
- Chorpita, B. F., Yim, L., Moffitt, C., Umemoto, L. A., & Francis, S. E. (2000). Assessment of symptoms of DSM-IV anxiety and depression in children: a revised child anxiety and depression scale. *Behav Res Ther*, 38(8), 835-855.
- DiBonaventura, M. d., Gupta, S., Cho, M., & Mrus, J. (2012). The association of HIV/AIDS treatment side effects with health status, work productivity, and resource use. *AIDS care*, 24(6), 744-755.
- Gay, C., Portillo, C. J., Kelly, R., Coggins, T., Davis, H., Aouizerat, B. E., . . . Lee, K. A. (2011). Self-reported medication adherence and symptom experience in adults with HIV. *Journal of the Association of Nurses in AIDS Care*, 22(4), 257-268.
- Gold, J. I., Mahrer, N. E., Yee, J., & Palermo, T. M. (2009). Pain, fatigue and health-related quality of life in children and adolescents with chronic pain. *Clin J Pain*, 25(5), 407.
- Jenkin, P., Koch, T., & Kralik, D. (2006). The experience of fatigue for adults living with HIV. *J Clin Nurs*, 15(9), 1123-1131. doi:10.1111/j.1365-2702.2006.01343.x
- Justice, A. C., Rabeneck, L., Hays, R. D., Wu, A. W., Bozzette, S. A., & Group, O. C. o. t. A. C. T. (1999). Sensitivity, specificity, reliability, and clinical validity of provider-reported symptoms: a comparison with self-reported symptoms. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 21(2), 126-133.
- Laughton, B., Cornell, M., Boivin, M., & Van Rie, A. (2013). Neurodevelopment in perinatally HIV-infected children: a concern for adolescence. *J Int AIDS Soc*, 16(1).
- Mburu, G., Hodgson, I., Kalibala, S., Haamujompa, C., Cataldo, F., Lowenthal, E. D., & Ross, D. (2014). Adolescent HIV disclosure in Zambia: barriers, facilitators and outcomes. *J Int AIDS Soc*, 17(1).
- Nutini, M., Karczewski, M., & Capoor, J. (2009). Fatigue in children with neurologic impairments. *Physical Medicine and Rehabilitation Clinics of North America*, 20(2), 339-346.
- Parslow, R. M., Harris, S., Broughton, J., Alattas, A., Crawley, E., Haywood, K., & Shaw, A. (2017). Children's experiences of chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME): a systematic review and meta-ethnography of qualitative studies. *BMJ Open*, 7(1), e012633. doi:10.1136/bmjopen-2016-012633
- Power, J. D., Badley, E. M., French, M. R., Wall, A. J., & Hawker, G. A. (2008). Fatigue in osteoarthritis: a qualitative study. *BMC Musculoskelet Disord*, 9, 63. doi:10.1186/1471-2474-9-63
- Sherr, L., Croome, N., Castaneda, K. P., Bradshaw, K., & Romero, R. H. (2014). Developmental challenges in HIV infected children—An updated systematic review. *Children and Youth Services Review*, 45, 74-89.
- Stormorken, E., Jason, L. A., & Kirkevold, M. (2015). Fatigue in adults with post-infectious fatigue syndrome: a qualitative content analysis. *BMC Nurs*, 14, 64. doi:10.1186/s12912-015-0115-5

- Varni, J. W., Burwinkle, T. M., & Szer, I. S. (2004). The PedsQL Multidimensional Fatigue Scale in pediatric rheumatology: reliability and validity. *The Journal of Rheumatology*, 31(12), 2494-2500.
- Voss, J. G., Portillo, C. J., Holzemer, W. L., & Dodd, M. J. (2007). Symptom Cluster of Fatigue and Depression in HIV/AIDS. *Journal of Prevention & Intervention in the Community*, 33(1-2), 19-34. doi:10.1300/J005v33n01_03