



Citation for published version:

Cowles, M & Hogg, L 2019, 'An Experimental Investigation into the Effect of State-Anxiety on State-Paranoia in People Experiencing Psychosis', *Behavioural and Cognitive Psychotherapy*, vol. 47, no. 1, pp. 52-66.
<https://doi.org/10.1017/S1352465818000401>

DOI:

[10.1017/S1352465818000401](https://doi.org/10.1017/S1352465818000401)

Publication date:

2019

Document Version

Peer reviewed version

[Link to publication](#)

This article is forthcoming in the journal *Behavioural and Cognitive Psychotherapy* doi.org/10.1017/S1352465818000401. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © Cambridge University Press.

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract

Background: There is strong evidence to suggest that anxiety is associated with paranoia in clinical and non-clinical samples. However, no research to date has directly manipulated anxiety to investigate if state-anxiety has a causal role in state-paranoia in clinical populations.

Aims: To investigate whether an anxious-mood induction leads to greater paranoia than a neutral-mood induction in people experiencing psychosis and paranoia and, if so, whether this is predicted by anxiety over and above other variables.

Method: 22 participants with a psychosis-spectrum diagnosis took part in a two condition cross-over experimental design. Participants underwent a neutral-mood and an anxious-mood induction with levels of state-anxiety, state-affect and state-paranoia being measured before and after each condition.

Results: State-paranoia was significantly higher after the anxious condition compared to the neutral condition. State-anxiety and state-paranoia were significant predictors of levels of state-paranoia after the anxious condition. When both predictors were included in a regression model, only negative-affect was a significant predictor of state-paranoia after the anxious condition. There were a number of methodological limitations.

Conclusions: State-anxiety and negative-affect may both be involved in the maintenance of paranoia in clinical populations, as predicted by cognitive models. Negative-affect may be the strongest predictor of state-paranoia in clinical populations. Interventions that seek to reduce state-affect may be beneficial in managing state-paranoia. Further research is warranted to explore the suggested clinical and theoretical implications of these findings.

Keywords: paranoia; anxiety; psychosis; schizophrenia; affect

Introduction

Paranoid delusions can be a particularly distressing experience that are reported by up to 70% of people with a first episode of psychosis (Freeman & Garety, 2014), as well as by people with other diagnoses and non-clinical populations (Bebbington et al., 2013).

The Role of Anxiety in Paranoid Delusions

Paranoid ideation is the unfounded belief that harm is, or will be, purposefully inflicted on the individual (Freeman & Garety, 2000). Given that anxiety is commonly understood as threat-anticipation, and disorders of anxiety as chronic overestimations of threat (American Psychiatric Association, 2013; Butler, Fennell, & Hackmann, 2010; Wells, 1997), it makes intuitive sense that delusions involving anticipation of harm would be influenced by, and in turn influence, anxiety. Anxiety is given a key role in the threat-anticipation model of the formation and maintenance of paranoid delusions (Freeman, 2007; Freeman, Garety, Kuipers, Fowler, & Bebbington, 2002).

The research base supporting the role of anxiety in paranoia is substantial. Anxiety has been found to be associated with paranoia in the general population (Freeman et al., 2011), high-risk groups (Salokangas et al., 2016) and clinical groups (Bosanac, Mancuso, & Castle, 2016; Huppert & Smith, 2005). Both anxiety and worry, a key cognitive component of anxiety, were found to have a dose-response relationship to paranoia in a study involving low paranoid, high non-clinically paranoid and clinically paranoid groups (Freeman, Pugh, Vorontsova, Antley, & Slater, 2010). An experience sampling study found that increases in anxiety predicted the onset of paranoid episodes in people with and without clinical paranoia (Thewissen et al., 2011) and anxiety has been found to predict the persistence of paranoia in high-risk groups (Salokangas et al., 2016). In sum, there is good evidence that anxiety is associated with paranoia.

However, a systematic review (Hartley, Barrowclough, & Haddock, 2013) concluded that, whilst there is convincing evidence that anxiety and depression are associated with psychotic experiences, there is a need for research that explores causal relationships between these affective states and specific psychotic experiences. An experimental investigation in a non-clinical population found that an anxious-mood induction led to significantly greater state-paranoia than a neutral-mood induction and that this was predicted by anxiety levels after the mood induction over and above other mood states (Lincoln, Lange, Burau, Exner, & Moritz,

1
2
3 2010). Furthermore, it was demonstrated that the anxious-mood induction increased paranoia
4 only for those with higher baseline paranoia, adding support to the stress-vulnerability model
5 inherent in cognitive models. However, these results may not be the same in people who have
6 already developed clinically significant paranoid delusions; this remains to be investigated.
7
8
9

10 Research that helps to elucidate which factors are involved in the formation and
11 maintenance of paranoia is crucial to indicate useful targets for prevention and treatment.
12 Preliminary studies into treatments for anxiety in people with paranoia have had promising
13 outcomes. A cognitive-behavioural intervention targeting anxiety and reasoning biases was found
14 to reduce anxiety and paranoia in people with anxiety disorders; this effect was greater for those
15 with higher baseline paranoia (Giusti et al., 2017). Pilot studies into brief CBT interventions for
16 worry have found that when worry is reduced, there are significant reductions in paranoid
17 ideation and paranoia-related distress compared to treatment as usual in people with paranoid
18 delusions and psychosis (Foster, Startup, Potts, & Freeman, 2010; Freeman et al., 2015). Further
19 studies that clarify the role of anxiety and other factors in paranoia are necessary to develop and
20 refine theoretical models and clinical interventions. Specifically, research is needed to explore if
21 there is a causal role for anxiety in paranoid thinking and if this interacts with other factors
22 (Freeman, 2007; Hartley et al., 2013).
23
24
25
26
27
28
29
30
31
32
33

34 **Aims of This Paper**

35
36 No study to date has directly manipulated anxiety to see if it has an effect on paranoia in
37 people experiencing such beliefs in the context of clinical psychosis. This study sought to
38 investigate whether an anxious-mood induction lead to greater paranoia than a neutral-mood
39 induction in people suffering with psychosis and paranoia and, if so, whether this was predicted
40 by anxiety over and above other variables.
41
42
43
44

45 It was predicted that: (1) the anxious condition would lead to more state-paranoia than the
46 neutral condition; (2) state-paranoia after the anxious condition would be predicted by state-
47 anxiety; (3) state-anxiety would predict state-paranoia to a greater extent than other affective
48 states; and (4) higher trait-paranoia would predict higher state-paranoia after the anxious
49 condition.
50
51
52
53
54
55
56
57
58
59
60

Method

Ethical approval was obtained from the NHS Health Research Authority (project ID: 201330) prior to conducting this study. Participants gave informed consent before taking part in the study and were fully debriefed at the end of the study.

Participants

Participants were recruited from community mental health teams in the South West of England. Inclusion criteria were: (a) primary psychosis-spectrum diagnosis (ICD-10 classification F20-F29); (b) experiencing paranoia with at least 50% conviction; (c) aged 18-65; (d) not acutely unwell at present; (e) sufficient English language and cognitive ability to take part in the study; and (f) being supported by a mental health team. Twenty-seven people were initially recruited and five of these were excluded: three due to insufficient levels of paranoia, and two as the neutral memory they chose to recall provoked intense emotion and therefore were not, as required by the research questions, neutral. The remaining 22 participants were aged between 20 to 60 years-old ($M = 32.82$; $SD = 10.72$) with 9 being female and 13 being male. The time since first recorded psychotic episode ranged from 0.5 to 27 years ($N = 22$; $M = 9.13$; $SD = 9.22$) and the most common psychiatric diagnosis was paranoid schizophrenia (10 participants).

Measures

Paranoia Checklist (PC: Freeman, Garety, Bebbington, Smith, et al., 2005). The PC was used to measure paranoia conviction. This is an 18-item self-report measure with statements relating to paranoid ideation. For the purpose of this study only the conviction sub-scale (18-items) was used; respondents use a 5-point rating scale (0 = ‘do not believe it’ to 4 = ‘absolutely believe it’). The scale has been found to have excellent internal reliability (Freeman, Garety, Bebbington, Smith, et al., 2005). Responses were summed with higher scores indicating more paranoia conviction (range: 0 – 72). Cronbach’s alpha was .88 in this study, indicating good reliability.

Psychotic Symptoms Rating Scale – section B (PSYRATS-B: Haddock, McCarron, Tarrier, & Faragher, 1999). The PSYRATS-B was used to measure degree of trait paranoia. This scale consists of six clinician-administered questions rated on a 5-point rating scale from 0 (none) to 4 (extreme). The scores are summed with high scores indicating greater severity of delusions; in this study only paranoid delusions were enquired about (range: 0-24). The scale has been found to have excellent reliability (Drake, Haddock, Tarrier, Bentall, & Lewis, 2007).

Brief State Measure. (BSM) The BSM was used to measure fluctuations in state-anxiety, affect, paranoia and paranoia-related distress. This is an 11-item self-report measure designed specifically for this study to measure variables of interest in a short timeframe.

All statements on the BSM were presented on a 10cm visual analogue scale (VAS) with participants drawing a line where they felt that each statement applied to them *right now* (anchors: ‘not at all’, ‘extremely’). The order that the questions were presented in was different each time.

The items were derived from existing measures and previous research studies:

- *State-paranoia* was measured using the 3-item change sensitive version of the PC, which has been found to be a valid and change-sensitive scale for clinically paranoid populations ($\alpha = .74$; Schlier, Moritz, & Lincoln, 2016). Items were: “I have to be on my guard against others”, “Strangers and friends look at me critically”, and “People are trying to make me upset”.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
- Measures of *positive-affect*, *negative-affect*, and *anxiety* were derived from a reliable measure of state-affect ($\alpha = .81$ to $.94$; Stemmler, Heldmann, Pauls, & Scherer, 2001). The measure was modified similarly to other studies in this area (Ascone, Sundag, Schlier, & Lincoln, 2016; Lincoln et al., 2010). The following words were presented after the statement ‘I feel...’:
 - *State-anxiety sub-scale* – ‘anxious’, ‘calm’ (reversed), ‘relaxed’ (reversed)
 - *State-negative-affect sub-scale* – ‘ashamed’, ‘sad’, ‘irritated’
 - *State-positive-affect sub-scale* – ‘happy’
- For the anxiety and negative-affect scales the mean score from the three items was used; higher scores indicated higher affect (range: 0 – 10).
- *Paranoia-related distress* was measured with the item, ‘I am distressed by thoughts of others wishing me harm’.

26 **Materials**

27
28
29
30
31
32
33
34
35

The Brief Relaxation Script was used before each condition. This was based on previous studies looking at paranoia (e.g. Ascone et al., 2016; Lincoln, Hohenhaus, & Hartmann, 2013) and directs the individual to breathe calmly and relax into their chair with eyes closed.

36
37
38
39
40
41
42
43
44

The Imagery Recall Script was used to enhance remembering of the chosen event. This included prompts to elicit *in-sensu* memory recall (e.g. what can you see/hear/feel? What are you thinking/feeling?). This was based on previous studies looking at paranoia (e.g. Ascone et al., 2016; Lincoln et al., 2013).

45
46
47
48
49
50
51
52

The Compassionate-Colour Imagery Script was used to reduce any distress and elicit soothing feelings. This was based on compassion-focused therapy techniques and existing imagery scripts (e.g. Gilbert, 2010) and entails the individual imagining a colour they associate with compassion.

53
54
55
56
57

All scripts are available upon request from the corresponding author.

Design and Procedure

A two condition crossover experimental design was used. Participants were randomised to either the neutral or anxious condition first, and completed the other condition after a break.

Participants were then asked to recall a recent (within the last two months) event that caused them to feel anxious, but that they did not consider to be traumatic, and an event that they considered to be emotionally neutral, such as a regular daily activity. Once an appropriate event had been identified (that did not directly involve paranoia), participants were guided through the brief relaxation script followed by the in-sensu memory recall script for the first condition until the image was rated as at least 60% vivid. The BSM was completed before and after the memory recall in each condition. Following this, participants were guided through the compassionate colour imagery exercise to reduce any distress caused. There was a minimum 20-minute break before the procedure was repeated for second condition. Please see Figure 2 for a diagram of the study protocol which was completed twice, once for each condition, by each participant.

-----FIGURE 1 ABOUT HERE-----

Statistical Analysis

The effect of the anxious condition compared to the neutral condition on each variable of interest (paranoia, distress, anxiety, negative- and positive-affect) was explored using a 2x2 repeated-measures analysis of variance (ANOVA) with within-subjects factors 'time' (pre, post) and 'condition' (anxious, neutral). Significant interactions between time and condition were furthered analysed using paired t-tests to analyse which variables were significantly different between the two conditions. Once the impact of the anxious versus the neutral condition had been identified, regression analyses were used to assess whether state-anxiety predicted state-paranoia and whether this was to a greater extent than prediction by negative-affect. Simple linear regression was used to test if trait-paranoia predicted state-paranoia after the anxious condition. The analysis strategy was carried out using SPSS version 23.

A priori power analyses (calculated with G*Power software, version 3.1.5 [Faul, Erdfelder, Buchner, & Lang, 2009]) showed that a sample size of 21 would be needed to detect a significant difference from pre to post the anxiety condition using a paired samples t-test (based on $p = .05$, power = .8, and effect size = .66 [based on Lincoln et al. (2010)]).

Results

Parametric Assumptions Check

Data was checked for skewness, kurtosis, normality of distribution and homogeneity of variance. All data was adequately normally distributed apart from paranoia after the anxious condition. An extreme outlier that was more than three standard deviations below the mean for this variable was excluded from further statistical analyses, resulting in adequately normally distributed data for this variable.

Order Effects Check

A mixed-model 2 x 2 ANOVA with between-subjects factor 'randomisation order' (anxious first, neutral first) and within-subjects factor 'time' (pre, post) was carried out with each variable of interest as the dependent variable. The results were examined for significant interactions between randomisation order and time to see if there were order-effects, despite counterbalancing. No significant interactions were found for any variable apart from paranoia distress in the neutral condition where there was a significant time x randomization interaction, $F(1, 20) = 5.05, p = .036$. This shows that paranoia distress ratings were different in the neutral condition depending on which condition participants were randomised to experience first. Independent t-tests showed that paranoia distress was the same before the neutral condition for the anxious first ($M = 3.55; SD = 3.05$) and neutral first ($M = 3.55; SD = 3.58$) participants; $t(20) = .0, p = 1$. However, after the neutral condition, people who were randomised to have the anxious condition first had significantly less paranoia distress ($M = 1.31; SD = 2$) than those who had the neutral condition first ($M = 4.15; SD = 3.36$); $t(16.28) = 2.41, p = .028$. The neutral condition may have reduced paranoia distress in those who were already familiar with the protocol. Due to these carryover-effects paranoia distress was not included in further analyses.

Manipulation Check

Results of the repeated measures ANOVAs, and t-tests investigating significant interactions are shown in Table 1.

There were no significant differences between conditions at baseline (T1). Anxiety, negative-affect, and paranoia were significantly higher, and positive-affect significantly lower,

1
2
3 after the anxious condition compared to the neutral condition (T2). This confirmed that the
4 anxious-condition was effective at increasing anxiety. Figure 2 provides a visual depiction of
5 how anxiety and paranoia ratings changed from pre-to-post each condition.
6
7
8
9

10 -----TABLE 1 ABOUT HERE-----

11
12
13 -----FIGURE 2 ABOUT HERE-----

14 15 16 17 **Hypothesis One: Effect of Anxious Condition on Paranoia**

18 To check if these differences were due to changes from pre-to-post the anxious or the
19 neutral condition, paired t-tests were conducted (Table 2).
20
21

22 -----TABLE 2 ABOUT HERE-----

23
24
25 The results showed that there was a significant increase in anxiety, negative-affect and
26 paranoia, and a significant decrease in positive-affect, from pre-to-post the anxious-condition.
27 There was also a significant decrease in anxiety and paranoia from pre-to-post the neutral
28 condition. This confirms hypothesis one, the anxious-condition led to more state-paranoia than
29 the neutral condition.
30
31
32
33

34 Although the anxious condition increased anxiety, it also significantly increased negative-
35 affect and reduced positive-affect. For this reason, the relationship between these variables and
36 levels of paranoia after the anxious condition was explored.
37
38
39
40

41 **Hypothesis Two: State-Anxiety will Predict State-Paranoia**

42 Simple linear regression showed that state-anxiety after the anxious condition was a
43 significant predictor of state-paranoia after the anxious condition, $R^2 = .31$, $F(1, 19) = 8.41$, p
44 $= .009$, (Table 3). This supports hypothesis two, state-anxiety was a significant predictor of state-
45 paranoia.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Hypothesis Three: State-Anxiety as a Unique Predictor of State-Paranoia

Negative-affect after the anxious condition was a significant predictor of state-paranoia after the anxious condition, $R^2 = .39$, $F(1, 19) = 12.28$, $p = .002$. Positive-affect was not a significant predictor, $R^2 = .17$, $F(1, 19) = 4.01$, $p = .06$.

Multiple regression (Method: ENTER) was used to ascertain if anxiety still significantly predicted paranoia when the variance accounted for by negative-affect was taken into account (Table 3). Together, the two variables predicted 48% of the variance in paranoia, $R^2 = .48$, $F(2, 18) = 8.38$, $p = .003$. However anxiety no longer significantly predicted paranoia when negative-affect was included, $\beta = .34$, $t = 1.77$, $p = .094$, but negative-affect remained a significant predictor, $\beta = .47$, $t = 2.47$, $p = .024$. Hypothesis three was not supported; negative-affect, rather than anxiety, was a unique predictor of paranoia after the anxiety induction.

-----TABLE 3 ABOUT HERE-----

Hypothesis Four: The Relationship between Trait-Paranoia and State-Paranoia

Regression analyses showed trait paranoia as measured by the PSYRATS-B was a significant predictor of state-paranoia after the anxious-mood condition. Trait-paranoia did not predict state-paranoia before or after the neutral-mood condition or before the anxious-mood condition (Table 4). This supports hypothesis four, suggesting that people with higher trait-paranoia are more sensitive to the impact of fluctuations in affect on levels of state-paranoia.

-----TABLE 4 ABOUT HERE-----

Discussion

This experimental study set out to investigate the role of state-anxiety in state-paranoia in people experiencing paranoid delusions in the context of a psychosis-spectrum diagnosis. To the authors' knowledge, it was the first study to attempt to directly manipulate anxiety in people with paranoia in the context of clinical psychosis.

The first hypothesis; that the anxious-mood induction would lead to greater state-paranoia than the neutral-mood induction, was supported. The second hypothesis; that state-paranoia would be predicted by state-anxiety after the anxious condition, was supported. The third hypothesis, that state-anxiety would predict state-paranoia to a greater extent than other affective states after the anxious condition, was not supported. The anxious-mood induction

1
2
3 significantly increased anxiety but also significantly increased negative-affect and decreased
4 positive-affect. Results showed that whilst level of anxiety predicted paranoia, negative-affect
5 was also a significant predictor of paranoia, over and above anxiety. The fourth hypothesis, that
6 higher trait-paranoia would predict higher state-paranoia after the anxious-mood induction, was
7 supported.
8
9
10
11
12

13 **Implications**

14 **Theory and Research**

15
16
17 This study confirmed findings that affect is linked to paranoia in clinically paranoid
18 samples (Bosanac et al., 2016; Drake et al., 2004; Freeman et al., 2010; Hartley et al., 2013;
19 Thewissen et al., 2011). By directly manipulating anxiety, this study has demonstrated that
20 anxiety appears to be causally implicated in paranoia, rather than being a consequence of it.
21
22
23

24 The study question was ‘does state-anxiety predict state-paranoia *to a greater extent than*
25 *other factors*’ and this study found the answer to be, ‘no’. Negative affect was a unique
26 significant predictor of paranoia when included in a model with anxiety. However, the threat-
27 anticipation model of paranoia posits that anxiety interacts with negative-affect, reasoning biases,
28 safety behaviours, anomalous experiences and previous life experiences (Freeman, 2007),
29 similarly to other cognitive models (e.g. Chadwick, Birchwood, & Trower, 1996; Morrison,
30 2001). It may be that anxiety is necessary for state-paranoia to be triggered but that negative
31 affect, at least in populations experiencing clinically significant paranoia, can play a more crucial
32 role.
33
34
35
36
37
38

39 Lincoln and colleague’s (2010) contradictory finding, that anxiety *was* a unique predictor
40 of paranoia when included in a model with negative affect, was found with a non-clinical
41 population. It is feasible that anxiety is a unique predictor for state-paranoia in non-clinical
42 populations but that once paranoia has reached clinically relevant levels, it is maintained in a
43 different way. For example, feeling state-negative-affect may be less triggering of state-paranoia
44 for someone who has not faced the stigma, shame and depression often inherent in having a
45 diagnosis of psychosis or seeking help. Interestingly, the measure of affect used in this study
46 included shame, which may have contributed to the strong links found between negative affect
47 and paranoia. Indeed, high shame has been found to increase vulnerability to paranoia in the face
48 of stressful situations (Johnson, Jones, Lin, Wood, Heinze & Jackson, 2014) and decrease sense
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 of social safeness (Castilho, Pinto, Viegas, Carvalho, Madeira & Martins, 2017) in clinical
4 populations. Clinically paranoid individuals have been found to experience significantly higher
5 levels of negative cognitions about the self, such as low self-esteem and low self-compassion
6 (Collett, Pugh, Waite, & Freeman, 2016).”
7
8
9

10 The finding that trait-paranoia was predictive of state-paranoia after the anxious-mood
11 induction fits with other research showing that anxiety has a greater impact on psychotic
12 phenomena in people with higher baseline symptomology (Kesting, Bredenpohl, Klenke,
13 Westermann, & Lincoln, 2013; Lincoln et al., 2010; Lincoln, Peter, Schäfer, & Moritz, 2009).
14 The finding that baseline paranoia *did not* predict state-paranoia at any time point other than after
15 the anxious-condition supports the threat-anticipation model of paranoia, showing that a higher
16 level of trait paranoia does not mean higher levels of state-paranoia unless other relevant factors
17 are present (e.g. changes in affect).
18
19
20
21
22
23
24

25 **Clinical**

26 Although more rigorous research is needed to corroborate that state-anxiety and state-
27 negative-affect play a causal role in state-paranoia in clinical samples, the results of this study
28 imply that interventions for emotion regulation could be beneficial. There is emerging evidence
29 that interventions for anxiety and worry can help to reduce trait paranoia (Foster et al., 2010;
30 Freeman et al., 2015). Based on the findings of this study, it may be that basic emotion regulation
31 skills could be useful as tools to manage state-paranoia in clinical populations not currently in
32 crisis. For example, diaphragmatic ‘belly’ breathing, attention training, progressive muscle
33 relaxation and compassionate imagery are all simple techniques that could be delivered at a low
34 cost by most healthcare professionals to provide basic coping mechanisms for this population.
35
36
37
38
39
40
41
42

43 Interestingly, the neutral-mood induction in this study led to significant decreases in
44 paranoia and anxiety. This is in line with findings that imagining a neutral object in detail can
45 lead to reductions in paranoia and negative-affect in clinically paranoid populations (Ascone et
46 al., 2016). It is likely that the attentional capacity needed to conjure up imagery in sensory detail
47 allows attention to be diverted from other potentially distressing stimuli. Distraction has been
48 found to effectively reduce state-anxiety in people with a psychosis-spectrum diagnosis
49 (Grezellschak, Lincoln, & Westermann, 2015). Further research could look to simple
50
51
52
53
54
55
56
57
58
59
60

1
2
3 intervention strategies that involve attentional capacity to see if these are found to be effective at
4 reducing negative-affect and paranoia in clinical settings.
5
6
7

8 **Limitations**

9

10 A crossover design with counter-balancing was used with both conditions being
11 conducted on the same day, which risked carry-over effects. Counterbalancing controlled this for
12 most variables, but order-effects were present for paranoia-distress, meaning that this variable of
13 interest could not be further explored. Future studies would ideally replicate with an independent
14 groups design and a larger sample size.
15
16
17

18 The anxious-memory recall manipulation was chosen as other studies have found this to
19 lead to increases in anxiety in this population (Ascone et al., 2016; Lincoln et al., 2010). As it
20 was a memory that the participants had already experienced and coped with, it was felt to be a
21 task that this population could undergo with minimal chance of adverse effects and that would
22 mimic 'real-world' anxiety. However, given the semi-ideographic nature of such a manipulation,
23 it was not possible to ensure that the condition was matched across participants. Chosen
24 memories may have involved affect other than anxiety, as indicated by the observed increases in
25 negative-affect and decreases in positive-affect.
26
27
28
29
30
31
32
33

34 **Future Directions**

35

36 Further research is needed to delineate the unique and interacting effects of different
37 types of affect, and other factors such as reasoning biases, predisposition to anomalous
38 experiences, safety behaviours and previous life experiences, in the maintenance of clinical
39 paranoia. Based on the findings of this study it could be particularly interesting to look at shame,
40 anger and sadness as individual components of negative affect rather than measuring them as one
41 construct.
42
43
44
45

46 Much of the experimental research to date has been conducted in non-clinical groups.
47 Experimental studies in clinically paranoid populations can help to advance theories of paranoia
48 formation and maintenance, and inform interventions. It would be particularly interesting to
49 further explore triggers of state-paranoia to elucidate possible targets for developing coping
50 strategies. Brief interventions for state-paranoia may present a more targeted and efficient
51 alternative to current CBT approaches for delusions and be less demanding on services.
52
53
54
55
56
57
58
59
60

1
2
3 The conflicting findings of this study in a clinical population with Lincoln and
4 colleague's (2010) findings in a non-clinical population requires further investigation. Studies
5 that seek to untangle the relationship between negative-affect, anxiety and paranoia in both at-
6 risk and clinical groups can help highlight which factors can be most usefully targeted to help
7 with the prevention and management of paranoia. Future studies would benefit from addressing
8 the methodological limitations outlined in this paper.
9
10
11
12
13
14

15 **Conclusions**

16
17 In summary, this was the first study to directly manipulate anxiety in order to explore its
18 impact on state-paranoia in a population of people experiencing clinically significant psychosis.
19 This study found that state-paranoia was higher after an anxious-mood induction and that this
20 was predicted by levels of state-anxiety and negative-affect. Negative-affect was a unique
21 predictor when both were taken into account. Trait paranoia was also shown to predict levels of
22 state-paranoia after the anxious mood-induction, indicating that vulnerability to paranoia makes
23 it more likely that changes in affective states will trigger state-paranoia. Although these findings
24 need to be interpreted with caution due to methodological limitations, they highlight some
25 interesting avenues for further research.
26
27
28
29
30
31
32
33

34 **Acknowledgments**

35
36 We would like to acknowledge all of the participants who took part in this study and the
37 services who helped us to recruit, particularly Dr Kate Chapman who acted as a field supervisor.
38 Many thanks also to Dr Emma Griffiths, Dr Angus MacBeth and the reviewers at Behavioural
39 and Cognitive Psychotherapy who offered invaluable feedback on earlier manuscripts.
40
41
42
43

44 **Conflict of Interest**

45 None
46
47
48

49 **Financial Support**

50 This research was conducted as part of a Doctorate in Clinical Psychology at the
51 University of Bath.
52
53
54
55
56
57
58
59
60

References

- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*: American Psychiatric Pub.
- Ascone, L., Sundag, J., Schlier, B., & Lincoln, T. M. (2016). Feasibility and Effects of a Brief Compassion-Focused Imagery Intervention in Psychotic Patients with Paranoid Ideation: A Randomized Experimental Pilot Study. *Clinical Psychology & Psychotherapy, in press*. doi:10.1002/cpp.2003
- Bebbington, P. E., McBride, O., Steel, C., Kuipers, E., Radovanovic, M., Brugha, T., . . . Freeman, D. (2013). The structure of paranoia in the general population. *British Journal of Psychiatry, 202*, 419-427. doi:10.1192/bjp.bp.112.119032
- Bosanac, P., Mancuso, S. G., & Castle, D. J. (2016). Anxiety Symptoms in Psychotic Disorders: Results from the Second Australian National Mental Health Survey. *Clinica Schizophrenia & Related Psychoses, 10*(2), 93-100. doi:10.3371/1935-1232-10.2.93
- Butler, G., Fennell, M., & Hackmann, A. (2010). *Cognitive-Behavioral Therapy for Anxiety Disorders: Mastering Clinical Challenges (Guides to Individualized Evidence-based Treatment)*: Guilford Press.
- Castilho, P., Pinto, A. M., Viegas, R., Carvalho, S., Madeira, N., & Martins, M. J. (2017). External Shame as a Mediator between Paranoia and Social Safeness in Psychosis, *Clinical Psychology*. doi:10.1111/cp.12136
- Chadwick, P. D., Birchwood, M. J., & Trower, P. (1996). *Cognitive Therapy for Delusions Voices and Paranoia*. Oxford, England: Wiley.
- Collett, N. Pugh, K., Waite, F., & Freeman, D. (2016). Negative cognitions about the self in patients with persecutory delusions: An empirical study of self-compassion, self-stigma, schematic beliefs, self-esteem, fear of madness, and suicidal ideation. *Psychiatry Research, 239*. 79-84. <http://dx.doi.org/10.1016/j.psychres.2016.02.043>
- Drake, R., Haddock, G., Tarrier, N., Bentall, R., & Lewis, S. (2007). The Psychotic Symptom Rating Scales (PSYRATS): Their usefulness and properties in first episode psychosis. *Schizophrenia Research, 89*(Issues 1–3), 119–122. doi:10.1016/j.schres.2006.04.024
- Drake, R. J., Pickles, A., Bentall, R. P., Kinderman, P., Haddock, G., Tarrier, N., & Lewis, S. W. (2004). The evolution of insight, paranoia and depression during early schizophrenia. *Psychological Medicine, 34*(2), 285-292.
- Faul, F., Erdfelder, E., Buchner, A., & Lang, A. G. (2009). Statistical power analyses using G*Power 3.1: tests for correlation and regression analyses. *Behavior Research Methods, 41*(4), 1149-1160. doi:10.3758/brm.41.4.1149
- Foster, C., Startup, H., Potts, L., & Freeman, D. (2010). A randomised controlled trial of a worry intervention for individuals with persistent persecutory delusions. *Journal of Behavior Therapy and Experimental Psychiatry, 41*(1), 45-51. doi:10.1016/j.jbtep.2009.09.001
- Freeman, D. (2007). Suspicious minds: the psychology of persecutory delusions. *Clinical Psychology Review, 27*(4), 425-457. doi:10.1016/j.cpr.2006.10.004
- Freeman, D., Dunn, G., Startup, H., Pugh, K., Cordwell, J., Mander, H., . . . Kingdon, D. (2015). Effects of cognitive behaviour therapy for worry on persecutory delusions in patients with psychosis (WIT): a parallel, single-blind, randomised controlled trial with a mediation analysis. *The Lancet, 2*(4), 305–313. doi:10.1016/S2215-0366(15)00039-5
- Freeman, D., & Garety, P. (2000). Comments on the content of persecutory delusions: Does the definition need clarification? *British Journal of Clinical Psychology, 39*(4), 407-414. doi:10.1348/014466500163400
- Freeman, D., & Garety, P. (2014). Advances in understanding and treating persecutory delusions: a review. *Social Psychiatry and Psychiatric Epidemiology, 49*(8), 1179-1189. doi:10.1007/s00127-014-0928-7

- 1
2
3 Freeman, D., Garety, P., Bebbington, P., Smith, B., Rollinson, R., Fowler, D., . . . Dunn, G. (2005).
4 Psychological investigation of the structure of paranoia in a non-clinical population. *The British*
5 *Journal of Psychiatry*, 186(5), 427-435. doi:10.1192/bjp.186.5.427
- 6 Freeman, D., Garety, P. A., Bebbington, P., Slater, M., Kuipers, E., Fowler, D., . . . Dunn, G. (2005). The
7 psychology of persecutory ideation II: a virtual reality experimental study. *Journal of Nervous*
8 *and Mental Disorders*, 193(5), 309-315.
- 9 Freeman, D., Garety, P. A., Kuipers, E., Fowler, D., & Bebbington, P. E. (2002). A cognitive model of
10 persecutory delusions. *British Journal of Clinical Psychology*, 41(Pt 4), 331-347.
- 11 Freeman, D., McManus, S., Brugha, T., Meltzer, H., Jenkins, R., & Bebbington, P. (2011). Concomitants
12 of paranoia in the general population. *Psychological Medicine*, 41(5), 923-936.
13 doi:10.1017/S0033291710001546
- 14 Freeman, D., Pugh, K., Vorontsova, N., Antley, A., & Slater, M. (2010). Testing the Continuum of
15 Delusional Beliefs: An Experimental Study Using Virtual Reality. *Journal of Abnormal*
16 *Psychology*, 119(1), 83-92. doi:10.1037/a0017514
- 17 Gilbert, P. (2010). *Compassion Focused Therapy*: Routledge.
- 18 Giusti, L., Ussorio, D., Salza, A., Malavolta, M., Aggio, A., Bianchini, V., . . . Roncone, R. (2017).
19 Preliminary study of effects on paranoia ideation and jumping to conclusions in the context of
20 group treatment of anxiety disorders in young people. *Early Intervention in Psychiatry*, ePud
21 ahead of print. doi:10.1111/eip.12415
- 22 Grezellschak, S., Lincoln, T. M., & Westermann, S. (2015). Cognitive emotion regulation in patients with
23 schizophrenia: Evidence for effective reappraisal and distraction. *Psychiatry Reserach*, 229(1-2),
24 434-439. doi:10.1016/j.psychres.2015.05.103
- 25 Haddock, G., McCarron, J., Tarrrier, N., & Faragher, E. B. (1999). Scales to measure dimensions of
26 hallucinations and delusions: the psychotic symptom rating scales (PSYRATS). *Psychological*
27 *Medicine*, 29(4), 879-889.
- 28 Hartley, S., Barrowclough, C., & Haddock, G. (2013). Anxiety and depression in psychosis: a systematic
29 review of associations with positive psychotic symptoms. *Acta Psychiatrica Scandinavica*,
30 128(5), 327-346. doi:10.1111/acps.12080
- 31 Huppert, J. D., & Smith, T. E. (2005). Anxiety and schizophrenia: the interaction of subtypes of anxiety
32 and psychotic symptoms. *CNS Spectrums*, 10(9), 721-731.
- 33 Johnson, J., Jones, C., Lin, A., Wood, S., Heinze, K., & Jackson, C. (2014). Shame amplifies the
34 association between stressful life events and paranoia amongst young adults using mental health
35 services: Implications for understanding risk and psychological resilience. *Psychiatry Research*,
36 220 (1-2). 217-225. doi:10.1016/j.psychres.2014.07.022
- 37 Kesting, M. L., Bredenpohl, M., Klenke, J., Westermann, S., & Lincoln, T. M. (2013). The impact of
38 social stress on self-esteem and paranoid ideation. *Journal of Behavior Therapy and*
39 *Experimental Psychiatry*, 44(1), 122-128. doi:10.1016/j.jbtep.2012.07.010
- 40 Lincoln, T. M., Hohenhaus, F., & Hartmann, M. (2013). Can Paranoid Thoughts be Reduced by Targeting
41 Negative Emotions and Self-Esteem? An Experimental Investigation of a Brief Compassion-
42 Focused Intervention. *Cognitive Therapy Research*, 37, 390-402. doi:10.1007/s10608-012-9470-7
- 43 Lincoln, T. M., Lange, J., Burau, J., Exner, C., & Moritz, S. (2010). The effect of state anxiety on
44 paranoid ideation and jumping to conclusions. An experimental investigation. *Schizophrenia*
45 *Bulletin*, 36(6), 1140-1148. doi:10.1093/schbul/sbp029
- 46 Lincoln, T. M., Peter, N., Schäfer, M., & Moritz, S. (2009). Impact of stress on paranoia: an experimental
47 investigation of moderators and mediators. *Psychol Med*, 39(7), 1129-1139.
48 doi:10.1017/S0033291708004613
- 49 Morrison, A. P. (2001). The interpretation of intrusions in psychosis: an integrative cognitive approach to
50 hallucinations and delusions. *Behavioural and Cognitive Psychotherapy*, 29, 257-276.
51 doi:doi:10.1017/S1352465801003010
- 52 Salokangas, R. K. R., Schultze-Lutter, F., Hietala, J., Heinimaa, M., From, T., Ilonen, T., . . . Ruhrmann,
53 S. (2016). Depression predicts persistence of paranoia in clinical high-risk patients to psychosis:
54
55
56
57
58
59
60

- 1
2
3 results of the EPOS project | SpringerLink. *Social Psychiatry and Psychiatric Epidemiology*,
4 51(2), 247-257. doi:10.1007/s00127-015-1160-9
5 Schlier, B., Moritz, S., & Lincoln, T. M. (2016). Measuring fluctuations in paranoia: Validity and
6 psychometric properties of brief state versions of the Paranoia Checklist. *Psychiatry Research*,
7 241, 323-332. doi:10.1016/j.psychres.2016.05.002
8 Stemmler, G., Heldmann, M., Pauls, C. A., & Scherer, T. (2001). Constraints for emotion specificity in
9 fear and anger: The context counts. *Psychophysiology*, 38(2), 275-291. doi:10.1111/1469-
10 8986.3820275
11 Thewissen, V., Bentall, R. P., Oorschot, M., J, A. C., van Lierop, T., van Os, J., & Myin-Germeys, I.
12 (2011). Emotions, self-esteem, and paranoid episodes: an experience sampling study. *The British*
13 *Journal of Clinical Psychology*, 50(2), 178-195. doi:10.1348/014466510x508677
14 Wells, A. (1997). *Cognitive Therapy of Anxiety Disorders: A Practice Manual And Conceptual Guide*:
15 JW.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Flow Diagram of Study Procedure with Time Estimates. (Note: Participants completed this procedure twice –order of recalling neutral and anxious memory were counterbalanced. Sections in dark grey were not analysed as part of this paper)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Mean Anxiety and Paranoia Ratings at each Time Point. (N1 = pre-neutral condition; N2 = post-neutral condition; A1 = pre-anxious condition; A2 = post-anxious condition)

Table 1

2 (Time) x 2 (Condition) Repeated Measures ANOVA Results and Post-hoc t-test Results.

Variable	Time	Condition	Time x condition	t test T1 (N vs A)	t test T2 (N vs A)
Anxiety	$F(1, 21) = 21.55,$ $p < .001^{***}$	$F(1, 21) = 39.39,$ $p < .001^{***}$	$F(1, 21) = 66.29,$ $p < .001^{***}$	$t(21) = -.33;$ $p = .75$	$t(21) = -10.8;$ $p < .001^{***}$
Positive-Affect	$F(1, 21) = 10.63,$ $p = .004^{**}$	$F(1, 21) = 33.7,$ $p < .001^{***}$	$F(1, 21) = 20.78,$ $p < .001^{***}$	$t(21) = 1.24;$ $p = .23$	$t(21) = 6.66;$ $p < .001^{***}$
Negative-Affect	$F(1, 21) = 20.65,$ $p < .001^{***}$	$F(1, 21) = 17.01,$ $p < .001^{***}$	$F(1, 21) = 39.24,$ $p < .001^{***}$	$t(21) = -.26;$ $p = .8$	$t(21) = -6.95;$ $p < .001^{***}$
Paranoia	$F(1, 20) = 4.25,$ $p = .052^*$	$F(1, 20) = 14.31,$ $p = .001^{**}$	$F(1, 20) = 18.73,$ $p < .001^{***}$	$t(21) = -.42;$ $p = .67$	$t(20) = -5.56;$ $p < .001^{***}$

Note: T1=pre-condition; T2=post-condition; N vs A=neutral versus anxious; *=significant at .05 level; **=significant at .01 level; ***=significant at .001 level

Table 2

Paired t-test Results for Pre-to-Post Condition Scores

Variable Condition	Pre - Post Rating: M (SD)	Difference: M (SD)	N	<i>t</i>	<i>p</i>	<i>d</i>
Paranoia						
<i>Neutral</i>	13.35 (8.83) – 10.12 (8.33)	3.23 (6.3)	21	2.35	.029*	0.37
<i>Anxious</i>	14.24 (8.38) – 22.19 (6.42)	-7.94 (9.23)	21	-3.95	.001***	1.07
Anxiety						
<i>Neutral</i>	4.09 (2.76) – 2.96 (2.2)	1.13 (1.88)	22	2.81	.011**	0.45
<i>Anxious</i>	4.28 (2.57) – 8.26 (1.72)	-3.98 (2.22)	22	-8.39	<.001***	1.82
Negative						
<i>Neutral</i>	3.25 (2.64) – 2.92 (1.36)	.32 (1.89)	22	.8	.43	0.16
<i>Anxious</i>	3.4 (2.65) – 6.81 (2.65)	-3.41 (2.32)	22	-6.88	<.001***	1.29
Positive						
<i>Neutral</i>	5.02 (3.18) – 5.6 (3.07)	-.58 (2.65)	22	-1.03	.315	0.19
<i>Anxious</i>	4.46 (2.77) – 1.81 (2)	2.65 (1.7)	22	7.32	<.001***	1.1

*Note: M=mean; SD=standard deviation; *=significant at .05 level; **=significant at .01 level; ***=significant at .001 level. Effect size indicated by Cohen's *d*, where 0.2 = small, 0.5 = medium, and 0.8 = large.*

Table 3

Regression Results for Predictors of Paranoia post the Anxiety Induction (n = 21)

Predictor variable	<i>b</i>	Std Error	β	<i>p</i>	R^2	f^2
Constant	3.8	6.45		< .001***		
AnxietyT2A	2.19	.75	.55	.009**	.31	.45
Constant	9.99	3.66		< .001***		
NegativeAffectT2A	1.72	.49	.63	.002**	.39	.64
Constant	24.55	1.76		< .001***		
PositiveAffectT2A	-1.43	.71	-.42	.06	.17	.2
Constant	1.81	5.79		.007**		
AnxietyT2A	1.33	.75	.34	.094	.48	.92
NegativeAffectT2A	1.29	.52	.47	.024*		

*Note: 'T2A'=post anxiety condition; 'Std'= standard; *=significant at .05 level;*

***=significant at .01 level*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 4

Regression Results for Trait-Paranoia as a Predictor of State-Paranoia (n = 21)

Dependent Variable	Predictor Variable	<i>b</i>	Standard Error	β	<i>p</i>	R^2	f^2
State-Anxiety N1	Constant	-.24	10.82		.98		
	PSYRATS-B	.77	.64	.26	.24	.07	0.08
State-Anxiety N2	Constant	-2.27	10.03		.82		
	PSYRATS-B	.71	.59	.26	.24	.07	0.08
State-Anxiety A1	Constant	-.01	10.32		1		
	PSYRATS-B	.81	.61	.29	.2	.08	0.09
State-Anxiety A2	Constant	1.43	8.59		.87		
	PSYRATS-B	1.18	.5	.46	.03*	.21	0.27

*Note: N = neutral condition; A = anxious condition; 1 = pre; 2 = post; * = significant at the .05 level*