

Citation for published version:

Chrzanowski-Smith, OJ, Edinburgh, RM, Betts, JA, Stokes, KA & Gonzalez, JT 2018, 'Evaluation of a graded exercise test to determine peak fat oxidation in individuals with low cardiorespiratory fitness: Estimating maximal capacity for fat oxidation', *Applied Physiology Nutrition and Metabolism*, vol. 43, no. 12, pp. 1288-1297. <https://doi.org/10.1139/apnm-2018-0098>

DOI:

[10.1139/apnm-2018-0098](https://doi.org/10.1139/apnm-2018-0098)

Publication date:

2018

Document Version

Peer reviewed version

[Link to publication](#)

Forthcoming in *Applied Physiology, Nutrition and Metabolism*, at NRC Research Press (Canadian Science Publishing). © The Authors.

All data supporting this study is openly available from the University of Bath Research Data Archive at <https://doi.org/10.15125/BATH-00706>.

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

TITLE: Evaluation of a graded exercise test to determine peak fat oxidation in individuals with low cardiorespiratory fitness.

AUTHORS: Oliver J. Chrzanowski-Smith¹, Robert M. Edinburgh¹, James A. Betts PhD¹, Keith A. Stokes PhD¹, Javier T. Gonzalez PhD¹.

CORRESPONDENCE: Dr Javier T. Gonzalez

Department for Health

University of Bath

Bath, BA2 7AY

United Kingdom

Tel: +44-1225-385-518

Email: J.T.Gonzalez@bath.ac.uk

AFFILIATIONS: ¹ Department for Health, University of Bath, Bath, BA2 7AY, UK

O.J.Chrzanowski-Smith@bath.ac.uk

R.M.Edinburgh@bath.ac.uk

J.Betts@bath.ac.uk

K.Stokes@bath.ac.uk

J.T.Gonzalez@bath.ac.uk

RUNNING HEAD: Estimating maximal capacity for fat oxidation

ABSTRACT

The maximal capacity to utilise fat (peak fat oxidation [PFO]) may have implications for health and ultra-endurance performance, and is commonly determined by incremental exercise tests employing 3-minute stages. However, 3-minute stages may be insufficient to attain steady-state gas kinetics, compromising test validity. We assessed whether 4-minute stages produce steady-state gas exchange and reliable PFO in adults with $\dot{V}O_{2\text{peak}} < 40 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$. Fifteen participants (9F) completed a graded test to determine PFO and the intensity this occurred at (FAT_{MAX}). Three short continuous exercise sessions (SCE) were then completed in a randomised order, involving completion of the graded test to the stage: 1) preceding; 2) equal to ($\text{SCE}_{\text{equal}}$); or 3) after, PFO was attained, where participants then continued to cycle for 10-minutes at that respective intensity. Expired gases were sampled at minutes 3-4, 5-6, 7-8 and 9-10. Individual data showed steady-state gas exchange was achieved within 4-minutes during $\text{SCE}_{\text{equal}}$. Mean fat oxidation rates were not different across time within $\text{SCE}_{\text{equal}}$, nor compared to the graded test at FAT_{MAX} (both $p > 0.05$). However, the graded test displayed poor surrogate validity ($\text{SCE}_{\text{equal}}$ minute 3-4 vs 5-6, 7-8 and 9-10) and day-to-day reliability (minute 3-4 $\text{SCE}_{\text{equal}}$ vs graded test) to determine PFO, as evident by correlations (range: 0.47 – 0.83), and typical errors and 95% limits of agreement (range: 0.03 – 0.05 and ± 0.09 – 0.15 $\text{g}\cdot\text{min}^{-1}$, respectively). In conclusion, intra-individual variation in PFO is substantial despite four-minute stages establishing steady-state gas exchange in individuals with low fitness. Individual assessment of PFO may require multiple assessments.

KEY WORDS: Peak Fat Oxidation, FAT_{MAX} , Exercise Metabolism, Gas Exchange Kinetics, Low Cardiorespiratory Fitness

INTRODUCTION

There is emerging interest into the importance of peak fat oxidation during exercise (PFO; a whole-body measure of the 'maximal' capacity to utilise fat) and identifying the exercise intensity that elicits PFO (FAT_{MAX}) (Dandanell et al. 2017a; Isacco et al. 2014; Mogensen et al. 2009; Robinson et al. 2015; Romain et al. 2012; Rosenkilde et al. 2010). For example, PFO may represent a relatively non-invasive and easily measureable determinant of endurance exercise performance (Frandsen et al. 2017), metabolic health (Rosenkilde et al, 2010) and long-term weight loss and/or maintenance (Dandanell et al. 2017a). However, the valid determination of PFO is paramount to allow an appropriate interpretation of the outcomes of such studies, and to infer any role in sports performance or metabolic disease.

Early studies characterised fat oxidation during exercise over few stages, sometimes only *two* (Friedlander et al. 1998; Sidossis et al. 1997; Steffan et al. 1999; Thompson et al. 1998), *three* (Bircher and Knechtle 2004; Romijn et al. 2000; Romijn et al. 1993; van Loon et al. 2001) or *four* (Bergman and Brooks 1999) progressive exercise intensities, limiting the resolution to detect PFO. Attempting to address this poor resolution, Achten and colleagues (2002) developed a practical, one-off graded protocol (GE_{35/3}) in 2001. The test comprised three-minute stages commencing at 90 Watts (W) and progressively increasing by 35 W increments until volitional exhaustion. The protocol is widely advocated as a reliable and valid method to determine FAT_{MAX} in trained cyclists (Achten et al. 2002).

Slight variations of the GE_{35/3} protocol have since been widely employed to determine PFO and FAT_{MAX} in various populations. However, even within each standardised protocol, substantial inter-individual variability is apparent (0.18–1.01 g·min⁻¹ for PFO and 25-77 % $\dot{V}O_{2\max}$ for FAT_{MAX}) (Achten et al. 2002; Venables et al. 2005). Variations in the graded protocol typically involve alterations in work intensities, with early stage durations (i.e. three-minutes) held constant (Ara et al. 2011; Astorino et al. 2017; Blaize et al. 2014; Dandanell et al. 2017a; Dandanell et al. 2017b; Larsen et al. 2009; Nordby et al. 2015; Robinson et al. 2015; Rosenkilde et al. 2010; Venables et al. 2005). However, the delayed attainment of steady-state gaseous exchange (oxygen consumption and carbon dioxide production) in individuals with low vs high cardiorespiratory fitness (Poole and Jones 2012) raises the question as to whether three-minute stages are sufficient to produce valid and reliable estimates of PFO, or whether longer stages are required in populations with lower cardiorespiratory fitness (Bordenave et al. 2007; Brun et al. 2011).

To date, only one study has investigated the day-to-day reliability and surrogate validity of a graded test to exhaustion in determining PFO and FAT_{MAX} in individuals with low cardiorespiratory fitness (Dandanell et al. 2017b). In sixteen obese adults (8 females), the graded protocol produced PFO values that were not in close agreement (e.g. 95 % limits of agreement \pm 0.17 g·min⁻¹) with estimates derived from three short continuous exercise bouts at three exercise intensities that were not workload matched *per se* to the graded test (10-minutes at 35 %, 50 % and 65 % $\dot{V}O_{2\max}$). The substantial intra-individual (i.e. day-to-day) variation in PFO between the protocols indicates further optimisation of

graded exercise tests is required to improve their validity (Dandanell et al. 2017b). Importantly, neither Dandanell and co-workers (2017b) nor others (Achten et al. 2002; Bordenave et al. 2007) have systematically evaluated whether steady-state gaseous exchange is attained with short-stage graded exercise protocols. Thus, the substantial intra-individual variability (Dandanell et al. 2017b) and reported overestimation of fat oxidation in graded protocols utilising three- vs six-minute stages (Bordenave et al. 2007), may be due to non-steady-state gas kinetics.

Therefore, the main aims of this study were to develop a graded exercise protocol employing four-minute initial stages to: (i) observe whether steady-state gaseous exchange is attained within four-minutes; and to assess the graded test's (ii) surrogate validity (by comparing four-minute stages to six-, eight- and ten-minute stages); and (iii) day-to-day reliability, to determine PFO in individuals with low cardiorespiratory fitness. Four-minute stages were selected to '*optimise*' the graded test (i.e. increase the likelihood that steady-state gas exchange would be achieved in the initial stages, in an attempt to improve the validity and reliability of determining PFO). The hypotheses were (i) that steady-state gaseous exchange would be achieved within four-minutes and (ii) the graded protocol would provide valid and reliable estimates of PFO.

MATERIALS AND METHODS

Participants

Following written informed consent, sixteen adults (10 females) with low cardiorespiratory fitness ($\dot{V}O_{2\text{peak}} < 40 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) were recruited and completed the

study protocol between March and July 2017 (Supplementary Material S1a¹). Exclusion criteria included; age < 18 or > 65 yrs; current or history of cardio-pulmonary, metabolic or musculoskeletal disease; a body mass index < 18.9 or > 35 kg·m⁻²; breastfeeding or was / potentially pregnant; or any conditions or concurrent behaviour (including medication) that may pose undue personal risk to the participant or introduce bias to the study. Menstrual cycle and use of oral contraceptives was recorded but not controlled. One participant was excluded from the analyses due to hyperventilation during expired gas collection in all exercise trials (RER > 1.00), leaving a total sample of $n = 15$ (9 females). Participant characteristics are presented in Table 1.

Experimental design

A cross-over design was adopted where participants completed four exercise trials (one graded exercise test and three short continuous exercise sessions) in a laboratory at the University of Bath, UK. Trials were completed at a similar time (± 1 h within participant) in the morning (0600 – 1100 h) after an overnight fast (11 ± 1 h) and all within 21 days. Over the 24-h preceding each trial, participants were asked to: a) abstain from alcohol and strenuous physical activity; and b) replicate their dietary intake and physical activity. Additionally, participants maintained their habitual lifestyle throughout their involvement in the study. These criteria were confirmed by verbal questioning. All trials (within-subject) were performed under similar laboratory conditions (ambient temperature, humidity and barometric pressure; all $p > 0.05$; data not shown) where *ad-libitum* water intake and use of fans was permitted. The study was performed in accordance with the

¹ Supplementary data are available with the article through the journal Web site at <http://nrcresearchpress.com/doi/suppl/10.1139/apnm-2018-0098>.

1964 Declaration of Helsinki and was approved by the Research Ethics Approval Committee for Health at the University of Bath (REF: EP 16/17 141).

Graded exercise test

Participants first completed an incremental graded cycling test on a mechanically-braked cycle ergometer (Monark Peak Bike Ergomedic 894E, Varberg, Sweden). Upon arrival to the laboratory, anthropometric measurements were taken before placement of a heart rate monitor and self-selected ergometer set-up (seat and handlebar height were recorded to standardise for subsequent sessions). Participants then completed the graded test which comprised: a) four-minute stages for the first seven stages and b) two-minute stages from the eighth stage onwards. Initial power output was set at ~40 W and increased by ~25 W over the next six stages and by ~50 W from stage seven onwards. Participants exercised up until volitional exhaustion. The two-phase test was developed upon unpublished observations of pilot testing and adopted to 1) increase the sensitivity of the graded test for determining PFO in the early stages and 2) to facilitate the achievement of $\dot{V}O_2$ peak and attainment of volitional exhaustion quickly to reduce the confounding effects of fatigue. Expired gases were collected in the final minute of the first seven stages and upon the participant's signal of one-minute remaining before volitional exhaustion. The mouthpiece was removed in between stages and provided one-minute before each gas collection to ensure adequate 'flushing' of ambient air. Expired gas samples were analysed immediately after each collection. Heart rate (HR) via telemetry (Polar RS400 Heart Rate Monitor, Kempele, Finland) and ratings of perceived exertion (RPE) (Borg 1973) were recorded during each expired gas sample. The graded test was used to determine:

- a) Peak fat oxidation [$\text{g}\cdot\text{min}^{-1}$; the stage with the highest recorded fat oxidation value (measured values approach)];
- b) FAT_{MAX} (expressed as absolute $\dot{V}\text{O}_2$, W and HR at stage eliciting PFO and as a % of $\dot{V}\text{O}_{2\text{peak}}$, PPO and HR_{MAX});
- c) Peak power output (W; power output of the last completed stage, plus the fraction of time in the final non-completed stage, multiplied by the Watt increment of that stage);
- d) An estimate of peak oxygen consumption [$\dot{V}\text{O}_{2\text{peak}}$; $\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$; the upper functional limit (i.e. highest $\dot{V}\text{O}_2$ value) recorded in the single graded exercise test (Green and Askew, 2018)].

Short continuous exercise sessions (SCE)

Participants then completed three short continuous exercise sessions (SCE) in a randomised order on separate days (randomisation list created by O.C-S on www.randomizer.org/ Date: 24/02/2017). Upon arrival, anthropometric measurements were taken before participants completed the graded test protocol up until either the stage: 1) preceding (SCE_{pre}); 2) equal to ($\text{SCE}_{\text{equal}}$); or 3) after (SCE_{post}), PFO was previously determined, where participants then continued to cycle for a further ten-minutes at that respective intensity. If PFO was measured in the first stage of the graded test, SCE_{pre} power output was lowered to ~ 30 W ($n = 13$), where participants exercised for ten-minutes at this absolute intensity. One-minute expired gas samples were collected during the 10-

minute continuous period at minutes 3-4, 5-6, 7-8 and 9-10. Participants were provided the mouthpiece one-minute before the first collection and this was then kept in for remainder of the session. Heart rate and RPE were collected during each expired gas sample. Expired gas samples were analysed immediately at the end of the session in order of collection.

Anthropometric measurements

All anthropometric measurements were performed according to International Standards for Anthropometric Assessment 2007 (Marfell-Jones et al. 2007). Body stature was measured to the nearest 0.1 cm using a wall-mounted stadiometer (Holtain Ltd, Pembrokeshire, UK). Body mass (to nearest 0.1 kg) and body fat % (bioelectrical impedance analysis to nearest 0.1 %) were measured using electronic weighing scales (BC-543 Monitor, Tanita, Tokyo, Japan). Waist and hip circumference (cm) were additionally measured in the SCE sessions to the nearest 0.1 cm using a non-elastic measuring tape (SECA 201, Hamburg, Germany).

Metabolic measurements

Expired gas samples were collected into 100-150 L Douglas bags (Cranlea and Hans Rudolph, Birmingham, UK) via a mouthpiece connected to a two-way, T shaped non-rebreathing valve (Model 2700, Hans Rudolph Inc, Kansas City, USA) and falconia tubing (Hans Rudolph Inc, Kansas City, USA). Concentrations of O₂ and CO₂ were measured in a known volume of each sample via paramagnetic and infrared transducers, respectively (Mini MP 5200, Servomex Group Ltd., Crowborough, East Sussex, UK) and until values were stable. The sensor was calibrated to a two-point low and high calibration

of known gas concentrations (Low: 99.998 % Nitrogen, 0 % O₂ and CO₂; High: Balance nitrogen mix, 20.06 % O₂, 8.11 % CO₂) (BOC Industrial Gases, Linde AG, Munich, Germany). Concurrent measurements of inspired air composition were made during collection of each expired gas sample to adjust for changes in ambient O₂ and CO₂ concentrations (Betts and Thompson 2012). Indirect calorimetry was used to determine: $\dot{V}O_2$ (L·min⁻¹); $\dot{V}CO_2$ (L·min⁻¹); $\dot{V}E$ (L·min⁻¹); respiratory exchange ratio (RER); and rate of fat oxidation [g·min⁻¹; estimated by stoichiometric equations (Frayn 1983) assuming urinary nitrogen excretion was negligible].

Statistical analysis

Normality and equal variance were explored via the Shapiro-Wilk test and visual inspection of histograms, Q-Q plots, P-P plots and scatterplots on raw data and residuals of comparisons. Assumption of normality and equal variance was assumed unless otherwise stated and sensitivity analyses were performed where applicable. The attainment of steady-state gas exchange was visually inspected and confirmed as described in Supplementary Material S1b². A range of *a priori* statistical analysis tests were performed to evaluate the graded test in determining primarily PFO as advocated (Atkinson and Nevill 1998). To test for systematic differences in metabolic variables within each SCE session and *versus* the graded test at FAT_{MAX}, one-way repeated measures ANOVAs [time of expired gas sample as within-subject factor; irrespective of normality due to robustness of ANOVA (Maxwell 1990)] and paired sample *t*-tests or Wilcoxon Signed-Rank tests were performed, respectively. *Post-hoc* Bonferroni corrected *t*-tests

² Supplementary data are available with the article through the journal Web site at <http://nrcresearchpress.com/doi/suppl/10.1139/apnm-2018-0098>.

were performed where significant main effects of time were detected. The surrogate validity of four-minute stages (i.e. minute 3-4) to estimate fat oxidation was further explored within SCE_{pre} and SCE_{equal} using Pearson correlation coefficient (r), Intra-Class Correlation [ICC; 3,1 absolute agreement, single measures (Shrout and Fleiss 1979)], Within-subject Coefficient of Variation (CV; root mean square method), Typical Error [TE; SD of difference between scores / $\sqrt{2}$ (Hopkins 2015)] and Bland-Altman plot with mean difference (bias) and 95 % Limits of Agreement (LoA) (Bland and Altman 1986). These tests were not performed at the stage after FAT_{MAX} due to systematic differences in fat oxidation at this exercise intensity (data reported below). The day-to-day reliability of peak fat oxidation was also further explored by comparing fat oxidation estimates in the graded test at FAT_{MAX} (minute 3-4) vs SCE_{equal} minute 3-4 through the aforementioned intra-individual variation statistical tests. Mean difference in fat oxidation was calculated within SCE by subtraction of the minute 3-4 value from the value obtained at each 'respective time point', while comparisons between SCE_{equal} and the graded test were calculated through subtraction of the graded test value from the value obtained at SCE_{equal} minute 3-4. Individual data are plotted on all graphs. A maximum sample size of $n = 15$ was available for statistical analyses (see Supplementary Material S1c³ for description on sample sizes of analyses). Descriptive and statistical analyses were run on Microsoft Excel (2013) and IBM SPSS statistics version 22 for windows (IBM, New York, USA) and graphs were created on Graph Pad Prism 7 software (La Jolla, Calif., USA). Data are presented as means \pm 95 % confidence intervals unless otherwise stated and statistical significance was accepted at $p \leq 0.05$.

³ Supplementary data are available with the article through the journal Web site at <http://nrcresearchpress.com/doi/suppl/10.1139/apnm-2018-0098>.

RESULTS

Gas exchange kinetics and surrogate validity of fat oxidation

There were no systematic differences detected across the four time points during SCE_{pre}, SCE_{equal} or SCE_{post}, for $\dot{V}O_2$ (Fig. 1a, 1c, 1g; F (3, 42) = 0.489, p = 0.692; F (3, 36) = 0.405, p = 0.750; F (3, 42) = 1.905, p = 0.143 respectively), $\dot{V}CO_2$ (Fig. 1b, 1d, 1f; F (3, 42) = 0.393, p = 0.759; F (3, 36) = 0.910, p = 0.446; F (3, 42) = 0.456, p = 0.714 respectively) or $\dot{V}E$ (Table 2). Visual inspection shows no clear within-subject fluctuation in $\dot{V}O_2$ and $\dot{V}CO_2$ across time points during SCE_{pre} and SCE_{equal} in all but one participant, indicating steady-state gas kinetics were typically achieved by minute 3-4 (Fig. 1a – 1f).

There were no systematic differences detected across the four time points during SCE_{pre} or SCE_{equal} for fat oxidation (Fig. 2a – 2b; F (3, 42) = 0.191, p = 0.902 and F (3, 36) = 2.782, p = 0.055, respectively), or RER (Table 2). However, systematic differences were found for fat oxidation (Fig. 2c; F (3, 42) = 8.069, p < 0.001) and RER (Table 2) during SCE_{post}. As depicted in Fig. 2c, fat oxidation was lower during minute 3-4 and 5-6 vs minute 9-10 (p = 0.010 and 0.002, respectively), whilst RER was higher during minute 3-4 and 5-6 vs minute 9-10 (Table 2; p = 0.016 and 0.004, respectively).

Pearson correlations and ICCs between PFO determined at minute 3-4 vs 'longer stage durations' in SCE_{equal} were high to excellent (ranged between 0.61 to 0.83), but coincided with large 95 % CIs (Table 3). There were large within-subject CVs and reasonably small typical errors [range between 19 to 25 % and 0.03 to 0.04 g·min⁻¹,

respectively (Table 3)] and the Bland-Altman analysis of agreements revealed a small bias (range of -0.01 to -0.04 g·min⁻¹) and moderately large 95 % LoAs [range of ± 0.09 to 0.12 g·min⁻¹ (Fig. 3a – 3c)], in the determination of fat oxidation at minute 3-4 vs longer stages at SCE_{equal}. Reliability and intra-individual variation of fat oxidation at SCE_{pre} is presented in Table 3 and Supplementary Figure S1⁴.

Day-to-day reliability

There were no systematic differences in $\dot{V}O_2$ (Fig. 1g; $T(11) = -0.167$, $p = 0.870$), $\dot{V}CO_2$ (Fig. 1h; $Z = -0.78$, $p = 0.937$) or $\dot{V}E$ (Supplementary Table S1⁴) between the graded protocol and SCE minute 3-4 at the stage equal to FAT_{MAX}. Similarly, visual inspection shows little within-subject fluctuation in $\dot{V}O_2$ and $\dot{V}CO_2$ in all but one participant (Fig. 1g and 1h).

No systematic differences were found in fat oxidation (Fig. 2d; $T(11) = 0.954$, $p = 0.361$), or RER (Supplementary Table S1⁴) between the graded test and SCE minute 3-4 at the stage equal to FAT_{MAX}.

The Pearson correlation and ICC of the graded test vs SCE minute 3-4 at the stage equal to FAT_{MAX} to determine PFO were fair with large 95 % CIs apparent [0.48 (95 % CI: -0.14 – 0.82) and 0.47 (95 % CI: -0.10 – 0.81), respectively]. There was a large within-subject CV [23 % (95 % CI: 6 – 32 %)] and a moderately small typical error [0.05 g·min⁻¹ (95 % CI: 0.04 – 0.09 g·min⁻¹)], whilst Bland-Altman analysis of agreement revealed a

⁴ Supplementary data are available with the article through the journal Web site at <http://nrcresearchpress.com/doi/suppl/10.1139/apnm-2018-0098>.

small mean bias ($-0.02 \text{ g}\cdot\text{min}^{-1}$) and a considerably large 95 % LoA [$\pm 0.15 \text{ g}\cdot\text{min}^{-1}$ (Fig. 3d)], in the determination of PFO between the graded test and SCE minute 3-4 at FAT_{MAX} .

Sensitivity analysis performed on an outlier revealed no meaningful effect nor altered interpretation of $\text{SCE}_{\text{equal}}$ and graded test minute 3-4 vs SCE minute 3-4 at FAT_{MAX} analyses, except for Pearson correlation and ICC on the latter analysis ($r = 0.48$ vs 0.15 and $\text{ICC} = 0.47$ vs 0.14 , respectively). Furthermore, outcomes of the SCE analyses were independent of the statistical analysis employed (Friedman vs repeated measures ANOVA; data not shown).

DISCUSSION

The capacity to oxidise fat may represent a determinant of metabolic health and ultra-endurance performance. Acknowledging the recent call to '*optimise*' graded exercise tests for accurate determination of PFO and FAT_{MAX} (Dandanell et al. 2017b), this is the first study to examine the surrogate validity and day-to-day reliability of a graded test employing four-minute stages to determine PFO in individuals with low cardiorespiratory fitness. The novel data provide evidence that steady-state gas exchange is attained within four-minutes at intensities equal to or below FAT_{MAX} . However, large intra-individual variability was apparent when determining PFO with four-minute stages both when compared to longer stage durations *and* across two separate days. This indicates that graded tests employing four-minute stages have poor surrogate validity and day-to-day reliability, respectively. Consequently, multiple testing is likely required to confidently assess PFO in individuals.

Incremental (graded) exercise tests are frequently used in a wide range of research studies to estimate PFO. However, there is currently no clear consensus (or justification) as to the most appropriate stage duration of these tests. We showed that steady-state gas exchange was achieved within four-minutes as there were no systematic differences alongside only minor fluctuations in the individual time course of gas kinetic variables during SCE_{equal} (Fig. 1). The attainment of steady-state gas exchange is a key requirement for the use of indirect calorimetry to accurately assess substrate metabolism (Macfarlane 2017; Shephard and Aoyagi 2012). This suggests that despite previous uncertainty over utilising stage durations under six-minutes (Bordenave et al. 2007), graded exercise tests with as short as four-minute stages can be legitimately recommended for use in healthy populations with low cardiorespiratory fitness.

The current study is the first to comprehensively explore the surrogate validity of graded exercise tests employing various stage lengths (4-, 6-, 8- and 10-minute stages) in the determination of PFO. We observed no systematic differences in fat oxidation rates estimated across the various stage lengths at FAT_{MAX} (i.e. during SCE_{equal} ; Fig. 2b). This is somewhat in contrast to Bordenave et al. (2007) who reported three-minute stages 'over-estimated' fat oxidation rates by $\sim 0.02 \text{ g}\cdot\text{min}^{-1}$ when compared to six-minute stages. However, in this previous study differences in fat oxidation rates were pooled across 4 x 6-minute submaximal exercise intensities and potential carry-over effects (i.e. previous stages influencing data from subsequent stages) were not controlled for, potentially limiting the interpretation of these results. The potential carry over effect of previous stages is also pertinent to previous reliability and validity investigations of graded tests

(Achten et al. 2002; Dandanell et al. 2017*b*). Whilst numerically we report a slightly higher 'over-estimation' of PFO with four-minute vs six-minute stages ($-0.03 \text{ g}\cdot\text{min}^{-1}$), better agreement in PFO estimates were evident when stage durations of eight- or ten-minutes were compared to four-minute stages [-0.02 and $-0.01 \text{ g}\cdot\text{min}^{-1}$, respectively (Fig. 2*b*)]. Collectively, this suggests that stage lengths between four- to ten-minutes provide equally useful group level estimates of PFO.

Despite no evident systematic differences, we found considerable intra-individual variance in the determination of PFO when four-minute stages were compared to longer stage durations (i.e. graded tests with four-minute stages have poor surrogate validity). For example, the wide 95 % CIs surrounding the Pearson and Intra-Class correlation coefficients indicates large intra-individual variability exists when assessing PFO. This is further supported by the large within-subject CVs (Table 3) and 95 % LoAs. The latter suggest that estimates of PFO made with four-minute stages may vary by ± 0.09 to $0.12 \text{ g}\cdot\text{min}^{-1}$ when compared to longer stages (Fig. 3*a* – 3*c*). This corresponds to the finding of Bordenave and colleagues (2007) who report a 95 % LoA of ± 0.06 to $0.10 \text{ g}\cdot\text{min}^{-1}$ when three-minute stages were compared to six-minute stages. Notwithstanding that group level estimates of PFO are similar, researchers should acknowledge the large intra-individual variation in the determination of PFO with different stage durations. Thus, as graded tests currently stand, repeated testing with different stage lengths may be required to more confidently assess PFO.

The present study also assessed the day-to-day reliability of the graded test to determine PFO. We found no systematic difference at the group level when estimating

PFO with four-minute stages across two days at the same absolute power output (Fig. 2d). Alternatively, Dandanell and co-workers (2017b) reported a statistically significant systematic difference in the determination of PFO between a graded test employing three-minute stages and three ten-minute bouts at non-equivalent exercise intensities in sixteen adults (8 females) with obesity and low cardiorespiratory fitness ($0.04 \text{ g}\cdot\text{min}^{-1}$, $p = 0.03$). This slight contrast in finding may derive from methodological discrepancies such as the use of three- vs four-minute stages, as it is still unknown whether three-minute stages are sufficient to establish steady-state gas exchange in the population we studied. Unfortunately, the present study was unable to explore the comparison between three- vs four-minute stages due to the Douglas bag method preventing immediate collection of back-to-back samples. Notably, graded tests employing three-minute stages have reported larger group mean systematic differences in $\dot{V}O_2$, $\dot{V}CO_2$ and $\dot{V}E$ when exercising at FAT_{MAX} on separate days (Achten et al. 2002; Dandanell et al. 2017b). Collectively, this suggests graded tests with three-minute stages may systematically produce greater variability in gas exchange kinetics and PFO estimates, advocating the use of four-minute stages in future studies.

Despite this, we observed large intra-individual variability in the day-to-day reliability of PFO (Fig. 3d and Table 3). Similarly large limits of agreement have previously been reported ($\pm 0.13 \text{ g}\cdot\text{min}^{-1}$) alongside a lower within-subject CV [11 % (95 % CI: 8 – 15 %)] in individuals with low cardiorespiratory fitness (Dandanell et al. 2017b). Moreover, corresponding high CVs (range of 23 to 26 %) and even greater 95 % LoAs (range of ± 0.24 to $0.26 \text{ g}\cdot\text{min}^{-1}$) have also been reported in PFO despite small systematic differences (range of $0.00 - 0.01 \text{ g}\cdot\text{min}^{-1}$) between two identical sub-maximal cycling protocols utilising

five-minute stages in moderately trained males (Croci et al. 2014). Our results support and build upon this previous work by using the still considered gold standard method for expired gas sampling and analysis [i.e. the Douglas bag technique (Shephard 2017)] over the more frequently used breath-by-breath metabolic cart systems. We also adjusted for inspired air composition in substrate utilisation calculations (Betts and Thompson 2012) which is important due to (the largely overlooked) changes in gas composition of both atmospheric air and ambient laboratory air. Taken collectively, these data all suggest that there is poor day-to-day reliability at the individual level in the determination of PFO across a range of graded test protocols.

Numerous factors may contribute to the large intra-individual variation observed in the determination of PFO with graded exercise tests. While exercise intensity and duration are primary modulators of fat oxidation during exercise (Romijn et al. 1993; van Loon et al. 2001), nutrient status (Gonzalez et al. 2013), dietary macronutrient composition (Fletcher et al. 2017; Støa et al. 2016) and both acute and chronic physical activity levels (Venables et al. 2005) are also contributing factors. Standardising diet and physical activity for at least 48-h prior to testing is likely important for more reliable estimates of fat oxidation during exercise (Støa et al. 2016). However, prior research (including the present study) has rarely objectively ensured strict pre-trial standardisation of either diet or physical activity for ≥ 48 -h (Astorino and Schubert 2018). While this may partially account for the day-to-day variability between protocols, similar intra-individual variability was apparent across different stage durations [i.e. within SCE_{pre} and SCE_{equal} (Fig. 3)] with metabolic estimates exposed to the same 'standardisation'. Nevertheless, although

challenging to implement, greater standardisation may be integral to obtaining reliable estimates of PFO and should be further explored.

Like many previous studies, the current graded exercise protocols were based on absolute watt increments. Consequently, each stage would represent a different percentage of exercise capacity between participants (e.g. % $\dot{V}O_{2\text{peak}}$) and thus elicit potentially divergent physiological responses. This may not only contribute to the variability in fat oxidation estimates, particularly as high intra-individual day-to-day variability in the metabolic milieu has previously been reported at FAT_{MAX} (Dandanell et al. 2017b), but may account for the systematic unstable fat oxidation rates prevalent during SCE_{post} (Fig. 2c). At intensities equal to or above lactate threshold, attainment of steady-state gas exchange may be delayed (for ~15 minutes or more) due to contribution of the $\dot{V}O_2$ slow component and $\dot{V}CO_2$ from non-aerobic sources (e.g. buffering of $[H^+]$ ions by bicarbonate) (Rossiter 2011). It is plausible that the exercise intensity elicited by the stage after FAT_{MAX} may have coincided or been above the lactate threshold for some individuals (Achten and Jeukendrup 2004; Michallet et al. 2008). An alternative approach would be to characterise the relative exercise intensities of each stage using the lactate threshold. However, this would require an additional testing day (and standardisation procedure), losing the one-off, time-effective practicality element of such graded tests. Our data also suggest caution should be applied when using more complex analytical approaches (i.e. curve modelling from multiple exercise intensities), which utilise estimates above FAT_{MAX} , to characterise fat oxidation kinetics and PFO (Cheneviere et al. 2009; Croci et al. 2014).

In this work it should be acknowledged that cardiorespiratory fitness was estimated via $\dot{V}O_{2peak}$ which while is an acceptable marker (Green and Askew, 2018) may underestimate participants 'true' maximal cardiorespiratory fitness (Poole and Jones, 2017). Additionally, PFO was recorded in the first stage of the graded test for thirteen participants. Consequently, PFO could have occurred at a lower intensity than the initial 40 W stage in the graded test with SCE_{equal} also potentially not representing 'FAT_{MAX}'. However, at the group level fat oxidation rates were lower in SCE_{pre} than SCE_{equal} (Fig. 2). Additionally, the considerable intra-individual day-to-day variation in PFO and FAT_{MAX} reported here and previously (Crocì et al. 2014; Dandanell et al. 2017b), suggests large overlap in fat oxidation rates may exist between coinciding exercise intensities (Achten et al. 2004). The overlap in fat oxidation rates evident between SCE_{pre} and SCE_{equal} further suggest PFO could have occurred below 30 W. Acknowledging this shortcoming, practitioners may wish to commence graded tests at lower workloads than 30 / 40 W to increase the resolution of capturing PFO and FAT_{MAX} in individuals with low cardiorespiratory fitness.

Collectively, the above demonstrates that future work should seek to further optimise graded tests to more accurately capture and determine PFO. The practical relevance of this is reiterated by exercise training induced increases in PFO [e.g. + 0.12 to 0.22 g·min⁻¹ (Mogensen et al. 2009; Nordby et al. 2015)] that are within the realm of reported intra-individual variability. Consequently, to confidently ascribe any effects to an intervention, observed increases in PFO may need to exceed the reported intra-individual variability in PFO from the most similar protocol employed (Crocì et al. 2014; Dandanell et al. 2017b; De Souza Silveira et al. 2016; Marzouki et al. 2014). That noted, only once

PFO has been quantified reliably can the question of what may constitute a meaningful change or difference in PFO for health and performance be addressed.

Conclusion

In summary, the present study shows for the first time that graded tests utilising four-minute stages are sufficient to establish steady-state gaseous exchange and group level peak fat oxidation rates compared to longer stage durations (\leq 10-minutes) in individuals with low levels of cardiorespiratory fitness. However, our data show that poor surrogate validity and substantial day-to-day variation are apparent when determining PFO. Consequently, graded tests with four-minute stages are advocated, but until further 'optimisation' is performed (particularly with a focus on strict pre-test physical activity and dietary standardisation), repeated assessments are prudent to more confidently and precisely determine 'true' peak fat oxidation rates.

ACKNOWLEDGMENTS

All participants are thanked for their involvement in the study. Dr Mark Thomas, Pippa Heath, Esther Punter, Max Simms and Aaron Hengist (Department for Health, University of Bath) are thanked for their help in the study.

AUTHORSHIPS

The study was designed by O.J.C-S., R.M.E., and J.T.G. Data collection was performed by O.J.C-S, and data analysis and interpretation assisted by J.A.B., and J.T.G. The manuscript was written by O.J.C-S. All authors revised the manuscript and approved the final article.

DECLARATIONS OF FUNDING SOURCES

Funding was provided by the Department for Health and University Research Studentship from the Graduate School, Faculty of Humanities and Social Sciences at the University of Bath. The funds were not involved in any part of the study except for financial and equipment support.

CONFLICTS OF INTEREST

The authors declare no conflict of interests.

REFERENCES

- Achten, J., Gleeson, M., and Jeukendrup, A. E. 2002. Determination of the exercise intensity that elicits maximal fat oxidation. *Med. Sci. Sports Exerc.* 34(1): 92-97. PMID: 11782653.
- Achten, J., and Jeukendrup, A. E. 2004. Relation between plasma lactate concentration and fat oxidation rates over a wide range of exercise intensities. *Int. J. Sports. Med.* 25(1): 32-37. doi: 10.1055/s-2003-45231. PMID: 14750010.
- Ara, I., Larsen, S., Stallknecht, B., Guerra, B., Morales-Alamo, D., Andersen, J. L., et al. 2011. Normal mitochondrial function and increased fat oxidation capacity in leg and arm muscles in obese humans. *Int. J. Obes. (Lond.)*. 35(1): 99-108. doi: 10.1038/ijo.2010.123. PMID: 20548301.
- Astorino, T. A., Edmunds, R. M., Clark, A., Gallant, R., King, L., Ordille, G. M., et al. 2017. Change in maximal fat oxidation in response to different regimes of periodized high-intensity interval training (HIIT). *Eur. J. Appl. Physiol.* 117(4): 745-755. doi: 10.1007/s00421-017-3535-y. PMID: 28251399.
- Astorino, T. A., and Schubert, M. M. 2018. Changes in fat oxidation in response to various regimes of high intensity interval training (HIIT). *Eur. J. Appl. Physiol.* 118(1): 51-63. doi: 10.1007/s00421-017-3756-0. PMID: 29124325.
- Atkinson, G., and Nevill, A. M. 1998. Statistical methods for assessing measurement error (reliability) in variables relevant to sports medicine. *Sports Med.* 26(4): 217-238. PMID: 9820922
- Bergman, B. C., and Brooks, G. A. 1999. Respiratory gas-exchange ratios during graded exercise in fed and fasted trained and untrained men. *J. Appl. Physiol.* (1985). 86(2): 479-487. doi: 10.1152/jappl.1999.86.2.479. PMID: 9931180.

- Betts, J. A., and Thompson, D. A. 2012. Thinking outside the Bag (Not Necessarily outside the Lab). *Med. Sci. Sports. Exerc.* 44(10): 2040. doi: 10.1249/MSS.0b013e318264526f. PMID: 22986475.
- Bircher, S., and Knechtle, B. 2004. Relationship between fat oxidation and lactate threshold in athletes and obese women and men. *J. Sports Sci. Med.* 3(3): 174-181. PMID: 24482595.
- Blaize, A. N., Potteiger, J. A., Claytor, R. P., and Noe, D. A. 2014. Body fat has no effect on the maximal fat oxidation rate in young, normal, and overweight women. *J. Strength Cond. Res.* 28(8): 2121-2126. doi: 10.1519/JSC.0000000000000512. PMID: 24796985.
- Bland, J. M., and Altman, D. G. 1986. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet*, 1(8476): 307-310. PMID: 2868172.
- Bordenave, S., Flavier, S., Fédou, C., Brun, J. F., and Mercier, J. 2007. Exercise calorimetry in sedentary patients: procedures based on short 3 min steps underestimate carbohydrate oxidation and overestimate lipid oxidation. *Diabetes Metab.* 33(5): 379-384. doi: 10.1016/j.diabet.2007.04.003. PMID: 17936665.
- Borg, G. A. 1973. Perceived exertion: a note on "history" and methods. *Med. Sci. Sports.* 5(2): 90-93. PMID: 4721012.
- Brun, J. F., Romain, A. J., and Mercier, J. 2011. Maximal lipid oxidation during exercise (Lipox max): From physiological measurements to clinical applications. Facts and uncertainties. *Science & Sports*, 26(2): 57-71. doi : 10.1016/j.scispo.2011.02.001.

- Cheneviere, X., Malatesta, D., Peters, E. M., and Borrani, F. 2009. A mathematical model to describe fat oxidation kinetics during graded exercise. *Med. Sci. Sports Exerc.* 41(8): 1615-1625. doi: 10.1249/MSS.0b013e31819e2f91. PMID: 19568198.
- Compher, C., Frankenfield, D., Keim, N., Roth-Yousey, L., and Evidence Analysis Working Group. 2006. *J. Am. Diet. Assoc.* 106(6): 881-903. doi: 10.1016/j.jada.2006.02.009. PMID: 16720129.
- Croci, I., Borrani, F., Byrne, N. M., Wood, R. E., Hickman, I. J., Chenevière, X., et al. 2014. Reproducibility of Fat max and Fat Oxidation Rates during Exercise in Recreationally Trained Males. *PloS one*, 9(6): e97930. doi: 10.1371/journal.pone.0097930. PMID: 24886715.
- Dandanell, S., Husted, K., Amdisen, S., Vigelsø, A., Dela, F., Larsen, S., et al. 2017a. Influence of maximal fat oxidation on long-term weight loss maintenance in humans. *J. Appl. Physiol.* (1985). 123(1): 267-274. doi: 10.1152/jappphysiol.00270.2017. PMID: 28546468.
- Dandanell, S., Præst, C. B., Søndergård, S. D., Skovborg, C., Dela, F., Larsen, S., et al. 2017b. Determination of the exercise intensity that elicits maximal fat oxidation in individuals with obesity. *Appl. Physiol. Nutr. Metab.* 42(4): 405-412. doi: 10.1139/apnm-2016-0518. PMID: 28177732.
- De Souza Silveira, R., Carlsohn, A., Langen, G., Mayer, F., and Scharhag-Rosenberger, F. 2016. Reliability and day-to-day variability of peak fat oxidation during treadmill ergometry. *J. Int. Soc. Sports Nutr.* 13(4): 1-8. doi: 10.1186/s12970-016-0115-1. PMID: 26816497.
- Fletcher, G., Eves, F. F., Glover, E. I., Robinson, S. L., Vernooij, C. A., Thompson, J. L., et al. 2017. Dietary intake is independently associated with the maximal capacity

- for fat oxidation during exercise. *Am. J. Clin. Nutr.* 105(4): 864-872. doi: 10.3945/ajcn.116.133520. PMID: 28251936.
- Frandsen, J., Vest, S. D., Larsen, S., Dela, F., and Helge, J. W. 2017. Maximal Fat Oxidation is Related to Performance in an Ironman Triathlon. *Int. J. Sports Med.* 38(13): 975-982. doi: 10.1055/s-0043-117178. PMID: 29050040.
- Frayn, K. N. 1983. Calculation of substrate oxidation rates in vivo from gaseous exchange. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* 55(2): 628-634. doi: 10.1152/jappl.1983.55.2.628. PMID: 6618956.
- Friedlander, A. L., Casazza, G. A., Horning, M. A., Buddinger, T. F., and Brooks, G. A. 1998. Effects of exercise intensity and training on lipid metabolism in young women. *Am. J. Physiol.* 275(5 Pt 1): E853-E863. PMID: 9815006.
- Green, S., and Askew, C. D. 2018. $\dot{V}O_2$ peak is an acceptable estimate of cardiorespiratory fitness but not $\dot{V}O_2$ max. *J. Appl. Physiol.* (1985). doi: 10.1152/japplphysiol.00850.2017. PMID: 29420148
- Gonzalez, J. T., Veasey, R. C., Rumbold, P. L., and Stevenson, E. J. 2013. Breakfast and exercise contingently affect postprandial metabolism and energy balance in physically active males. *Br. J. Nutr.* 110(4): 721-732. doi: 10.1017/S0007114512005582. PMID: 23340006.
- Hopkins, W. G. 2015. Measures of reliability in sports medicine and science. *Sportscience.* 19: 36-44. sportsci.org/2015/ValidRely.htm.
- Isacco, L., Thivel, D., Duclos, M., Aucouturier, J., and Boisseau, N. 2014. Effects of adipose tissue distribution on maximum lipid oxidation rate during exercise in normal-weight women. *Diabetes Metab.* 40(3): 215-219. doi: 10.1016/j.diabet.2014.02.006. PMID: 24698815.

- Larsen, S., Ara, I., Rabøl, R., Andersen, J. L., Boushel, R., Dela, F., et al. 2009. Are substrate use during exercise and mitochondrial respiratory capacity decreased in arm and leg muscle in type 2 diabetes? *Diabetologia*, 52(7): 1400-1408. doi: 10.1007/s00125-009-1353-4. PMID: 19396425.
- Macfarlane, D. J. 2017. Open-circuit respirometry: a historical review of portable gas analysis systems. *Eur. J. Appl. Physiol.* 117(12): 1369-2386. doi: 10.1007/s00421-017-3716-8. PMID: 29043499.
- Marfell-Jones, M., Olds, T., Stewart, A., and Carter, L. 2007. *International Standards for Anthropometric Assessment* (2006 ed.). Potchefstroom: The International Society for the Advancement of Kinanthropometry.
- Marzouki, H., Farhani, Z., Gmada, N., Tabka, Z., Shephard, R. J., and Bouhlef, E. 2014. Relative and absolute reliability of the crossover and maximum fat oxidation points during treadmill running. *Science and Sports*, 29(6): e107-e114. doi: 10.1016/j.scispo.2014.07.013.
- Maxwell, S. E. 1990. *Designing experiments and analyzing data : a model comparison perspective*. Belmont, Ca.: Belmont, Ca.: Wadsworth.
- Michallet, A. S., Tonini, J., Regnier, J., Guinot, M., Favre-Juvin, A., Bricout, V., et al. 2008. Methodological aspects of crossover and maximum fat-oxidation rate point determination. *Diabetes Metab.* 34(5): 514-523. doi: 10.1016/j.diabet.2008.04.004. PMID: 18823806.
- Mogensen, M., Vind, B. F., Højlund, K., Beck-Nielsen, H., and Sahlin, K. 2009. Maximal lipid oxidation in patients with type 2 diabetes is normal and shows an adequate increase in response to aerobic training. *Diabetes Obes. Metab.* 11(9): 874-883. doi: 10.1111/j.1463-1326.2009.01063.x. PMID: 19531056.

- Nordby, P., Rosenkilde, M., Ploug, T., Westh, K., Feigh, M., Nielsen, N. B., et al. 2015. Independent effects of endurance training and weight loss on peak fat oxidation in moderately overweight men: a randomized controlled trial. *J. Appl. Physiol.* (1985). 118(7): 803-810. doi: 10.1152/jappphysiol.00715.2014. PMID: 25614598.
- Poole, D. C., and Jones, A. M. 2012. Oxygen uptake kinetics. *Compr. Physiol.* 2(2): 933-996. doi: 10.1002/cphy.c100072. PMID: 23798293.
- Poole, D. C., and Jones, A. M. 2017. Measurement of the maximum oxygen uptake $\dot{V}O_{2max}$: $\dot{V}O_{2peak}$ is no longer acceptable. *J. Appl. Physiol.* (1985). 122(4): 997-1002. doi: 10.1152/jappphysiol.01063.2016. PMID: 28153947.
- Robinson, S. L., Hattersley, J., Frost, G. S., Chambers, E. S., and Wallis, G. A. 2015. Maximal fat oxidation during exercise is positively associated with 24-hour fat oxidation and insulin sensitivity in young, healthy men. *J. Appl. Physiol.* (1985). 118(11): 1415-1422. doi: 10.1152/jappphysiol.00058.2015. PMID: 25814634.
- Romain, A. J., Carayol, M., Desplan, M., Fedou, C., Ninot, G., Mercier, J., et al. 2012. Physical activity targeted at maximal lipid oxidation: a meta-analysis. *J. Nutr. Metab.* 2012: 285395. doi: 10.1155/2012/285395. PMID: 22928092.
- Romijn, J. A., Coyle, E. F., Sidossis, L. S., Gastaldelli, A., Horowitz, J. F., Endert, E., et al. 1993. Regulation of endogenous fat and carbohydrate metabolism in relation to exercise intensity and duration. *Am. J. Physiol.* 265(3 Pt 1): E380-391. doi: 10.1152/ajpendo.1993.265.3.E380. PMID: 8214047.
- Romijn, J. A., Coyle, E. F., Sidossis, L. S., Rosenblatt, J., and Wolfe, R. R. 2000. Substrate metabolism during different exercise intensities in endurance-trained women. *J. Appl. Physiol.* (1985). 88(5): 1707-1714. doi: 10.1152/jappl.2000.88.5.1707. PMID: 10797133.

- Rosenkilde, M., Nordby, P., Nielsen, L. B., Stallknecht, B. M., and Helge, J. W. 2010. Fat oxidation at rest predicts peak fat oxidation during exercise and metabolic phenotype in overweight men. *Int. J. Obes. (Lond.)*. 34(5): 871-877. doi: 10.1038/ijo.2010.11. PMID: 20157319.
- Rossiter, H. B. 2011. Exercise: Kinetic considerations for gas exchange. *Compr. Physiol.* 1(1): 203-244. doi: 10.1002/cphy.c090010. PMID: 23737170.
- Shepard, R. J., and Aoyagi, Y. 2012. Measurement of human energy expenditure, with particular reference to field studies: an historical perspective. *Eur. J. Appl. Physiol.* 112(8): 2785-815. doi: 10.1007/s00421-011-2268-6. PMID: 22160180.
- Shephard, R. J. 2017. Open-circuit respirometry: a brief historical review of the use of Douglas bags and chemical analyzers. *Eur. J. Appl. Physiol.* 117(3): 381-387. doi: 10.1007/s00421-017-3556-6. PMID: 28210818.
- Shrout, P. E., and Fleiss, J. L. 1979. Intraclass correlations: uses in assessing rater reliability. *Psychol. Bull.* 86(2): 420-428. PMID: 18839484.
- Sidossis, L. S., Gastaldelli, A., Klein, S., and Wolfe, R. R. 1997. Regulation of plasma fatty acid oxidation during low- and high-intensity exercise. *Am. J. Physiol.* 272(6 Pt 1): E1065-E1070. doi: 10.1152/ajpendo.1997.272.6.E1065. PMID: 9227453.
- Steffan, H. G., Elliott, W., Miller, W. C., and Fernhall, B. 1999. Substrate utilization during submaximal exercise in obese and normal-weight women. *Eur. J. Appl. Physiol. Occup. Physiol.* 80(3): 233-239. PMID: 10453926.
- Støa, E. M., Nyhus, L. K., Børresen, S. C., Nygaard, C., Hovet, Å. M., Bratland-Sanda, S., et al. 2016. Day to day variability in fat oxidation and the effect after only 1 day of change in diet composition. *Appl. Physiol. Nutr. Metab.* 41(4): 397-404. doi: 10.1139/apnm-2015-0334. PMID: 26960444.

- Thompson, D. L., Townsend, K. M., Boughey, R., Patterson, K., and Bassett, D. R. Jr. 1998. Substrate use during and following moderate- and low-intensity exercise: Implications for weight control. *Eur. J. Appl. Physiol. Occup. Physiol.* 78(1): 43-49. doi: 10.1007/s004210050385. PMID: 9660155.
- van Loon, L. J., Greenhaff, P. L., Constantin-Teodosiu, D., Saris, W. H., and Wagenmakers, A. J. 2001. The effects of increasing exercise intensity on muscle fuel utilisation in humans. *J. Physiol.* 536(Pt 1): 295-304. doi: 10.1111/j.1469-7793.2001.00295.x. PMID: 11579177.
- Venables, M. C., Achten, J., and Jeukendrup, A. E. 2005. Determinants of fat oxidation during exercise in healthy men and women: a cross-sectional study. *J. Appl. Physiol.* (1985). 98(1): 160-167. doi: 10.1152/jappphysiol.00662.2003. PMID: 15333616.

TABLES

Table 1. Participant Characteristics

	Total Sample	Females	Males
Sex (Females/Males)	9/6	-	-
Age, yrs	33.9 (27.4 – 40.4)	32.7 (22.9 – 42.4)	35.7 (27.7 – 43.6)
Stature, cm	170.6 (167.2 – 174.0)	166.6 (164.1 – 169.2)	176.5 (171.9 – 181.1)
Body Mass, kg	74.4 (66.7 – 82.1)	66.2 (62.3 – 70.2)	86.6 (73.1 – 100.1)
BMI, kg·m ⁻²	25.4 (23.5 – 27.3)	23.9 (22.4 – 25.3)	27.7 (24.1 – 31.2)
Body Fat %	28.8 (25.3 – 32.3)	31.6 (28.5 – 34.8)	24.5 (18.4 – 30.6)
Waist Circumference, cm	83.9 (76.53 – 91.2)	75.5 (70.6 – 80.5)	96.4 (85.5 – 107.3)
Hip Circumference, cm	102.6 (99.2 – 105.9)	101.2 (98.1 – 104.2)	104.7 (97.5 – 111.8)
W:H ratio	0.81 (0.76 -0.87)	0.75 (0.71 – 0.78)	0.92 (0.86 – 0.97)
$\dot{V}O_2$ peak, L·min ⁻¹	2.4 (2.2 – 2.6)	2.2 (2.0 – 2.3)	2.8 (2.6 – 3.0)
$\dot{V}O_2$ peak, mL·kg ⁻¹ ·min ⁻¹	32.9 (30.8 – 35.1)	32.9 (30.6 – 35.2)	33.0 (28.6 – 37.3)
Peak Power Output, W	153 (142 – 163)	139 (132 – 146)	174 (164 – 183)
HR _{MAX} , beats·min ⁻¹	185 (176 – 193)	188 (176 – 200)	180 (171 – 189)
PFO ^a , g·min ⁻¹	0.23 (0.19 – 0.27) (Range: 0.11 - 0.35)	0.24 (0.19 – 0.29) (Range: 0.14 - 0.35)	0.22 (0.15 – 0.28) (Range: 0.11 - 0.34)
FAT _{MAX} ^a , % $\dot{V}O_2$ peak	36 (33 – 40) (Range: 28 – 51)	39 (35 – 43) (Range: 33 - 51)	33 (29 – 37) (Range: 29 - 41)
FAT _{MAX} ^a , %PPO	29 (26 – 32)	31 (27 – 35)	25 (22 – 29)
FAT _{MAX} ^a , %HR _{MAX}	55 (51 – 58)	58 (53 – 62)	51 (46 – 55)

^a = *n* of 14, one participant excluded due to hyperventilation. BMI = Body Mass Index; W:H ratio = Waist-to-Hip ratio; $\dot{V}O_2$ peak = peak oxygen consumption; HR_{MAX} = Maximum Heart Rate; PFO = Peak Fat Oxidation; FAT_{MAX} = the exercise intensity that elicits PFO; Data presented as mean (\pm 95 % CI).

Table 2. Metabolic variables during the short continuous exercise sessions (SCE)

		Minute 3-4	Minute 5-6	Minute 7-8	Minute 9-10	RM ANOVA
$\dot{V}E$, L·min ⁻¹	SCE _{pre}	17.94 (16.26 – 19.62)	17.41 (16.15 – 18.66)	17.94 (16.39 – 19.48)	18.13 (16.41 – 19.85)	F (1,959, 27,428) = 0.821, <i>p</i> = 0.448
	SCE _{equal}	21.40 (19.16 – 23.63)	21.36 (18.97 – 23.74)	22.13 (19.93 – 24.32)	21.26 (18.93 – 23.58)	F (3, 36) = 1.42, <i>p</i> = 0.251
	SCE _{post}	28.17 (25.91 – 30.42)	28.00 (25.71 – 30.29)	28.37 (26.10 – 30.65)	28.43 (26.06 – 30.80)	F (3, 42) = 0.387, <i>p</i> = 0.763
RER	SCE _{pre}	0.86 (0.82 – 0.89)	0.85 (0.83 – 0.88)	0.86 (0.86 – 0.88)	0.86 (0.83 – 0.89)	F (3, 42) = 0.145, <i>p</i> = 0.932
	SCE _{equal}	0.85 (0.82 – 0.88)	0.87 (0.83 – 0.90)	0.87 (0.85 – 0.90)	0.86 (0.84 – 0.87)	F (3, 36) = 2.779, <i>p</i> = 0.055
	SCE _{post}	0.93* (0.91 – 0.96)	0.93** (0.91 – 0.95)	0.91 (0.90 – 0.93)	0.91 (0.89 – 0.92)	F (3, 42) = 6.904, <i>p</i> = 0.001

SCE_{pre} *n* = 15; SCE_{equal} *n* = 13; SCE_{post} *n* = 15; ** *p* ≤ 0.01 vs minute 9-10; **p* < 0.05 vs minute 9-10. Data presented as

mean (± 95 % CI).

Table 3. Surrogate validity of fat oxidation ($\text{g}\cdot\text{min}^{-1}$) within the short continuous exercise sessions (SCE) at the stage preceding and equal to FAT_{MAX} .

Fat Oxidation ($\text{g}\cdot\text{min}^{-1}$)		Minute 3-4 vs 5-6	Minute 3-4 vs 7-8	Minute 3-4 vs 9-10
SCE _{pre} ($n = 15$)	r	0.84 (0.58 – 0.95)	0.71 (0.30 – 0.89)	0.64 (0.18 – 0.87)
	ICC	0.85 (0.60 – 0.95)	0.68 (0.27 – 0.88)	0.63 (0.19 – 0.86)
	CV (%)	17 (10 – 22)	23 (17 – 27)	25 (13 – 32)
	TE ($\text{g}\cdot\text{min}^{-1}$)	0.03 (0.02 – 0.05)	0.04 (0.03 – 0.06)	0.04 (0.03 – 0.07)
SCE _{equal} ($n = 13$)	r	0.82 (0.48 – 0.94)	0.83 (0.52 – 0.95)	0.65 (0.15 – 0.88)
	ICC	0.77 (0.35 – 0.93)	0.72 (0.16 – 0.91)	0.61 (0.11 – 0.86)
	CV (%)	25 (10 – 34)	19 (12 – 23)	19 (13 – 23)
	TE ($\text{g}\cdot\text{min}^{-1}$)	0.03 (0.02 – 0.06)	0.03 (0.02 – 0.05)	0.04 (0.03 – 0.07)

r = Pearson correlation coefficient; ICC = Intra-Class Correlation; CV = Within-Subject Coefficient of Variation;

TE = Typical Error. Data presented as mean (\pm 95 % CI).

FIGURES




Figure 1a – 1h. Gas exchange kinetics ($\dot{V}O_2$ and $\dot{V}CO_2$) during the short continuous exercise bouts (SCE) at the stage before FAT_{MAX} [SCE_{pre} (a and b; $n = 15$)], equal to FAT_{MAX} [SCE_{equal} (c and d; $n = 13$)], after FAT_{MAX} [SCE_{post} (e and f; $n = 15$)] and graded test (GT) vs SCE minute 3-4 at FAT_{MAX} (g and h; $n = 12$). Fig. 1h displays Wilcoxon signed-rank test (due to the differences in $\dot{V}CO_2$ between the graded test (GT) vs SCE minute 3-4 at FAT_{MAX} were non-normally distributed). A solid line represents the group mean \pm 95 % CI and dashed lines denote individual data. Overall group means are signified by triangles, females are denoted by stars and males are indicated by circles.




Figure 2a – 2d. Determination of fat oxidation ($\text{g}\cdot\text{min}^{-1}$) during the short continuous exercise bouts (SCE) at the stage before FAT_{MAX} [SCE_{pre} (a; $n = 15$)], equal to FAT_{MAX} [$\text{SCE}_{\text{equal}}$ (b; $n = 13$)], after FAT_{MAX} [SCE_{post} (c; $n = 15$)] and graded test (GT) vs SCE minute 3-4 at FAT_{MAX} (d; $n = 12$). A solid line represents the group mean \pm 95 % CI and dashed lines denote individual data. Overall group means are signified by triangles, females are denoted by stars and males are indicated by circles. ** $p \leq 0.01$.





Figure 3a – 3d. Bland-Altman plots of the agreement between fat oxidation ($\text{g}\cdot\text{min}^{-1}$) in the short continuous exercise bout (SCE) at the stage equal to FAT_{MAX} (SCE_{equal}) minute 3-4 vs 5-6 (a; $n = 13$), 7-8 (b; $n = 13$) and 9-10 (c; $n = 13$) and graded test (GT) vs SCE minute 3-4 (d; $n = 12$) at FAT_{MAX} . A solid line represents bias and the dashed lines represent lower and upper 95 % limits of agreement. Females are denoted by stars and males are indicated by circles.

SUPPLEMENTARY MATERIAL S1**S1a. Detailed recruitment process**

S1b. Criteria for attainment of steady-state gas exchange

To assess whether steady-state gas exchange was achieved for each participant the following criteria was applied:

- a) If the difference in $\dot{V}O_2$ and $\dot{V}CO_2$ between SCE minute 3-4 and the respective time point (i.e. SCE minute 5-6, 7-8, 9-10 or Graded Test minute 3-4) was within $1.96 * \text{Standard Deviation}$ of the two concerned time-points.

And

- b) If the Coefficient of Variation (CV) for $\dot{V}O_2$ and $\dot{V}CO_2$ was $\leq 10 \%$ when comparing SCE minute 3-4 and each respective time point (i.e. SCE minute 5-6, 7-8, 9-10 or Graded Test minute 3-4). The CV was calculated via a slightly adapted version of Compher et al. (2006) guidelines on assessing whether steady-state conditions have been achieved to measure resting metabolic rate:

*(Standard Deviation of SCE minute 3-4 vs respective time point * (mean of SCE minute 3-4 and respective time point)*100.*

S1c. Sample sizes of statistical analysis tests

One participant hyperventilated in the short continuous exercise bout (SCE) at the stage equal to FAT_{MAX} (SCE_{equal}) at all time points (ID 13) and was excluded from all analyses involving data at the stage equal to FAT_{MAX} (i.e. SCE_{equal} and vs graded test at FAT_{MAX}).

One participant hyperventilated at all time points in the graded test (ID 19) and was excluded from all analyses involving the graded test (i.e. graded test vs SCE_{equal}).

In one participant (ID 15) the expired gas sample at minute 3-4 was delayed by 40 seconds and thus was excluded from all analyses involving minute 3-4 data at the stage equal to FAT_{MAX} (i.e. SCE_{equal} and vs graded test at FAT_{MAX} minute 3-4).

Repeated Measures ANOVA sample sizes:

- SCE at the stage before FAT_{MAX} (SCE_{pre}), $n = 15$
- SCE_{equal} , $n = 13$ (Exclude IDs 13 and 15)
- SCE at the stage after FAT_{MAX} (SCE_{post}), $n = 15$

Dependent Sample T-Tests or Wilcoxon Signed-Rank test sample sizes:

- Graded test at FAT_{MAX} minute 3-4 vs SCE_{equal} minute 3-4, $n = 12$ (Exclude IDs 13, 15 and 19)




Reliability and Intra-Individual Variability tests sample sizes:

- SCE_{pre} , $n = 15$
- SCE_{equal} , $n = 13$ (IDs 13 and 15 excluded)
- Graded test at FAT_{MAX} minute 3-4 vs SCE_{equal} minute 3-4, $n = 12$ (IDs 13, 15 and 19 excluded)

Supplementary Table S1. Metabolic variables in the graded test vs short continuous exercise session at stage equal to FAT_{MAX}

	Graded Test	SCE minute 3-4	<i>t</i> -Test
$\dot{V}E$, L·min ⁻¹	19.65 (17.23 – 22.07)	21.10 (18.75 – 23.45)	T (11) = -1.913, <i>p</i> = 0.082
RER	0.83 (0.80 – 0.86)	0.85 (0.81 – 0.88)	T (11) = -1.045, <i>p</i> = 0.319

Graded Test = minute 3-4; n = 12. Data presented as mean (± 95 % CI).



Supplementary Figure S1. Bland-Altman plots of the agreement between fat oxidation ($\text{g}\cdot\text{min}^{-1}$) in the short continuous exercise bout at the stage before FAT_{MAX} (SCE_{pre}) minute 3-4 vs minute 5-6 (a; $n = 15$), 7-8 (b; $n = 15$) and 9-10 (c; $n = 15$). A solid line represents bias and the dashed lines represent lower and upper 95 % limits of agreement. Females are denoted by stars and males are indicated by circles.