



PHD

Novel transition metal-catalysed syntheses of carboxylic acid derivatives

Owston, Nathan

Award date:
2008

Awarding institution:
University of Bath

[Link to publication](#)

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Appendix

Anand, N., N. A. Owston, et al. (2007). "Ruthenium-catalysed conversion of oxime ethers into nitriles." Tetrahedron Letters **48**(44): 7761-7763.

The conversion of oxime ethers into nitriles has been achieved under neutral conditions using $\text{Ru}(\text{CO})(\text{PPh}_3)_3\text{H}_2$ and the bidentate ligand Xantphos as the catalyst. DOI:10.1016/j.tetlet.2007.09.028

Owston, N. A., A. J. Parker, et al. (2007). "Highly efficient ruthenium-catalyzed oxime to amide rearrangement." Organic Letters **9**(18): 3599-3601.

A wide range of aldoximes has been converted into the corresponding amides using the ruthenium-based catalyst $\text{Ru}(\text{PPh}_3)_3(\text{CO})\text{H}_2/\text{dppe}/\text{TsOH}$. The amides are generated in high yield and selectivity, with catalyst loading as low as 0.04 mol %. DOI: 10.1021/ol701445n

Owston, N. A., A. J. Parker, et al. (2007). "Iridium-catalyzed conversion of alcohols into amides via oximes." Organic Letters **9**(1): 73-75.

The iridium catalyst $[\text{Ir}(\text{Cp}^*)\text{Cl}_2]_2$ is effective for the rearrangement of oximes to furnish amides. The reaction has been combined with catalytic transfer hydrogenation between an alcohol and alkene to allow the conversion of alcohols into amides in a one-pot process. DOI: 10.1021/ol062549u

Owston, N. A., A. J. Parker, et al. (2008). "Oxidation of primary alcohols to methyl esters by hydrogen transfer." Chemical Communications(5): 624-625.

The oxidation of alcohols in the presence of methanol has been achieved using a ruthenium catalyst with crotononitrile as the hydrogen acceptor. DOI: 10.1039/b717073d