



*Citation for published version:*

Chen, W, Dávila, J & Guerra, I 2016, 'Bubble tower solutions for a supercritical elliptic problem in  $\mathbb{R}^N$ ', *Annali della Scuola Normale Superiore di Pisa*, vol. 15, pp. 85-116. [https://doi.org/10.2422/2036-2145.201310\\_013](https://doi.org/10.2422/2036-2145.201310_013)

*DOI:*

[10.2422/2036-2145.201310\\_013](https://doi.org/10.2422/2036-2145.201310_013)

*Publication date:*

2016

*Document Version*

Peer reviewed version

[Link to publication](#)

This is the authors' accepted manuscript of a paper published in final form in Chen, W, Dávila, J & Guerra, I 2016, 'Bubble tower solutions for a supercritical elliptic problem in  $\mathbb{R}^N$ ', *Annali della Scuola Normale Superiore di Pisa*, vol. 15, pp. 85-116 and available online via [https://doi.org/10.2422/2036-2145.201310\\_013](https://doi.org/10.2422/2036-2145.201310_013)

## University of Bath

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# BUBBLE TOWER SOLUTIONS FOR SUPERCRITICAL ELLIPTIC PROBLEM IN $\mathbb{R}^N$

WENJING CHEN, JUAN DÁVILA, AND IGNACIO GUERRA

ABSTRACT. We consider the following problem

$$\begin{cases} -\Delta u + u = u^p + \lambda u^q, & u > 0 \quad \text{in } \mathbb{R}^N; \\ u(z) \rightarrow 0 \quad \text{as } |z| \rightarrow \infty, \end{cases}$$

where  $p = p^* + \varepsilon$ , with  $p^* = \frac{N+2}{N-2}$ ,  $1 < q < \frac{N+2}{N-2}$  if  $N \geq 4$ ,  $3 < q < 5$  if  $N = 3$ ,  $\lambda > 0$ , and  $\varepsilon$  is a positive parameter. We prove that for  $\varepsilon > 0$  small enough, it has a solution with the shape of a tower of bubbles.

**Keywords:** elliptic equation, non-uniqueness, bubble-tower solutions

## 1. INTRODUCTION

We are interested in the elliptic equation

$$(1.1) \quad \begin{cases} -\Delta u + u = u^p + \lambda u^q, & u > 0 \quad \text{in } \mathbb{R}^N, \\ u(x) \rightarrow 0 \quad \text{as } |x| \rightarrow \infty, \end{cases}$$

where  $N \geq 3$ ,  $\lambda > 0$  and  $1 < q < p$ . This problem arises in the study of standing waves of a nonlinear Schrödinger equation with two power type nonlinearities, see for example Tao, Visan and Zhang [28].

If  $p = q$ , equation (1.1) reduces to

$$(1.2) \quad \begin{cases} -\Delta u + u = u^p, & u > 0 \quad \text{in } \mathbb{R}^N, \\ u(x) \rightarrow 0 \quad \text{as } |x| \rightarrow \infty, \end{cases}$$

after a suitable scaling.

Thanks to the classical result of Gidas, Ni and Nirenberg [15], solutions of (1.1) and (1.2) are radially symmetric about some point, which we will assume is always the origin.

It is well known that problem (1.2) has a solution if and only if  $1 < p < \frac{N+2}{N-2}$ . Existence was proved by Berestycki and Lions [2], while non-existence follows from the Pohozaev identity [26]. Uniqueness also holds and was fully settled by Kwong [16], after a series of contributions [4, 17, 23, 24, 22, 21]. See also Felmer, Quaas, Tang and Yu [10] for further properties.

Concerning (1.1), the work of Berestycki and Lions [2] is still applicable if  $1 < q < p < \frac{N+2}{N-2}$ , and one obtains existence of a solution. If  $p, q \geq \frac{N+2}{N-2}$  there is no solution, again from the Pohozaev identity.

Recently, Dávila, del Pino and Guerra [5] proved that uniqueness does not hold in general for (1.1), if  $1 < q < p < \frac{N+2}{N-2}$ . More precisely if  $N = 3$ , the authors

obtained at least three solutions to problem (1.1) if  $1 < q < 3$ ,  $\lambda > 0$  is sufficiently large and fixed, and  $p < 5$  is close enough to 5.

Let us mention some contributions to the question of existence for (1.1) when one exponent is subcritical and other is critical or supercritical. If  $1 < q < p = \frac{N+2}{N-2}$  in (1.1), Alves, de Morais Filho and Souto [1] proved:

- when  $N \geq 4$ , there exists a nontrivial classical solution for all  $\lambda > 0$  and  $1 < q < \frac{N+2}{N-2}$ ;
- when  $N = 3$ , there exists a nontrivial classical solution for all  $\lambda > 0$  and  $3 < q < 5$ ;
- when  $N = 3$ , there exists a nontrivial classical solution for  $\lambda > 0$  large enough and  $1 < q \leq 3$ .

Moreover, Ferrero and Gazzola [11] proved that for  $q < \frac{N+2}{N-2} \leq p$ , there exists  $\bar{\lambda} > 0$ , such that if  $\lambda > \bar{\lambda}$ , then (1.1) has at least one solution, while for  $q < \frac{N+2}{N-2} < p$ , there exists  $0 < \underline{\lambda} < \bar{\lambda}$  such that if  $\lambda < \underline{\lambda}$ , then there is no solution.

In this paper, we are interested in multiplicity of solutions of (1.1), and for this we take an asymptotic approach, that is, we consider

$$(1.3) \quad \begin{cases} -\Delta u + u = u^p + \lambda u^q, & u > 0 \quad \text{in } \mathbb{R}^N, \\ u(z) \rightarrow 0 \quad \text{as } |z| \rightarrow \infty, \end{cases}$$

where  $p = p^* + \varepsilon$ , with  $p^* = \frac{N+2}{N-2}$ ,  $\lambda > 0$  and  $\varepsilon > 0$  are parameters, and  $q$  satisfies

$$(1.4) \quad 1 < q < \frac{N+2}{N-2} \quad \text{if } N \geq 4; \quad 3 < q < 5 \quad \text{if } N = 3.$$

Our result can be stated as follows.

**Theorem 1.1.** *Let  $\lambda > 0$  and let  $q$  satisfy (1.4). Given an integer  $k \geq 1$ , then there exists  $\varepsilon_0 > 0$  such that for any  $\varepsilon \in (0, \varepsilon_0)$ , there is a solution  $u_\varepsilon(z)$  of problem (1.3) of the form*

$$(1.5) \quad u_\varepsilon(z) = (N(N-2))^{\frac{N-2}{4}} \sum_{j=1}^k \frac{\varepsilon^{-[(j-1) + \frac{1}{p^*-q}]} (\Lambda_j^*)^{-\frac{N-2}{2}}}{\left(1 + \varepsilon^{-\frac{4}{N-2}[(j-1) + \frac{1}{p^*-q}]} (\Lambda_j^*)^{-2} |z|^2\right)^{\frac{N-2}{2}}} (1 + o(1)),$$

where the constants  $\Lambda_j^* > 0$ ,  $j = 1, 2, \dots, k$ , can be computed explicitly and depend on  $k, N, q$ .

The expansion (1.5) is valid if  $\frac{1}{C} \varepsilon^{\frac{2}{N-2}[(i-1) + \frac{1}{p^*-q}]} \leq |z| \leq C \varepsilon^{\frac{2}{N-2}[(i-1) + \frac{1}{p^*-q}]}$ , with some  $i \in \{1, 2, \dots, k\}$ , and  $o(1) \rightarrow 0$  uniformly as  $\varepsilon \rightarrow 0$  in this region.

The solutions described in this result behave like a superposition of “bubbles” of different blow-up orders centered at the origin, and hence have been called bubble-tower solutions. By bubbles we mean the functions

$$(1.6) \quad w_\mu(z) = \alpha_N \frac{\mu^{\frac{N-2}{2}}}{(\mu^2 + |z|^2)^{\frac{N-2}{2}}}, \quad \text{with } \alpha_N = (N(N-2))^{\frac{N-2}{4}},$$

where  $\mu > 0$ , which are the unique positive solutions (except translations) of

$$-\Delta w = w^{p^*} \quad \text{in } \mathbb{R}^N.$$

Based on numerical simulations we present bifurcation diagrams for solutions of (1.3) where  $q$  satisfies (1.4). In Figure 1 (left) we show the bifurcation diagram as




FIGURE 1. Left:  $u(0)$  vs.  $p$  for  $\lambda$  large and fixed. Right:  $u(0)$  vs.  $\lambda$  for  $p = p^* + \varepsilon$ ,  $\varepsilon > 0$  small and fixed.

a function of  $p$  for a fixed large  $\lambda$ , and in Figure 1 (right) we show the diagram as a function of  $\lambda$  for  $p = p^* + \varepsilon$ ,  $\varepsilon > 0$  small and fixed. In both diagrams we observe branches of solutions, with the upper part having unbounded solutions as  $\varepsilon \rightarrow 0$  or  $\lambda \rightarrow \infty$ . We believe that the solutions constructed in Theorem 1.1 are located on these upper branches, and are shown in the diagrams for the cases of 1 and 2 bubbles.

Bubble-tower solutions were found by del Pino, Dolbeault and Musso [6] for a slightly supercritical Brezis-Nirenberg problem in a ball, and after that have been studied intensively [3, 7, 8, 9, 13, 14, 18, 19, 20, 25]. In particular we mention the work of Campos [3] who considered the existence of bubble-tower solutions to a problem related to ours:

$$\begin{cases} -\Delta u = u^{p^* \pm \varepsilon} + u^q, & u > 0 \quad \text{in } \mathbb{R}^N; \\ u(z) \rightarrow 0 \quad \text{as } |z| \rightarrow \infty, \end{cases}$$

with  $\frac{N}{N-2} < q < p^* = \frac{N+2}{N-2}$ ,  $N \geq 3$ .

For the proof of Theorem 1.1, we consider a variation of the so-called Emden-Fowler transformation:

$$v(x) = \left( \frac{p^* - 1}{2} \right)^{\frac{2}{p^*-1}} r^{\frac{2}{p^*-1}} u(r),$$

with

$$r = |z| = e^{-\frac{p^*-1}{2}x}, \quad x \in (-\infty, +\infty).$$

Then finding a radial solution  $u(r)$  to (1.3) corresponds to solving the problem

$$(1.7) \quad \begin{cases} \mathcal{L}_0(v) = \alpha_\varepsilon e^{\varepsilon x} v^{p^* + \varepsilon} + \lambda \beta_N e^{-(p^*-q)x} v^q & \text{in } (-\infty, +\infty); \\ v(x) > 0 & \text{for } x \in (-\infty, +\infty); \\ v(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty, \end{cases}$$

where

$$(1.8) \quad \mathcal{L}_0(v) = -v'' + v + \left( \frac{2}{N-2} \right)^2 e^{-\frac{4}{N-2}x} v,$$

is the transformed operator associated to  $-\Delta + I$ , and  $\alpha_\varepsilon, \beta_N$  are constants, see (2.5).

Under the Emden-Fowler transformation the bubbles  $w_\mu$  take the form

$$(1.9) \quad W(x - \xi) = \left( \frac{4N}{N-2} \right)^{\frac{N-2}{4}} e^{-(x-\xi)} \left( 1 + e^{-\frac{4}{N-2}(x-\xi)} \right)^{-\frac{N-2}{2}}$$

with  $\mu = e^{-\frac{2}{N-2}\xi}$ , and solve

$$\begin{cases} W'' - W + W^{p^*} = 0, & \text{in } (-\infty, +\infty); \\ W'(0) = 0; \\ W(x) > 0, \quad W(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty. \end{cases}$$

In Section 2, we build an approximate solution to (1.7) as a sum of suitable projections of the transformed bubbles  $W$  centered at  $0 < \xi_1 < \dots < \xi_k$  with  $\xi_1 \rightarrow \infty$ . After the study of the linearized problem at the approximate solution in Section 3, and solvability of a nonlinear projected problem in Section 4, we perform a Lyapunov-Schmidt reduction procedure as in [12, 18, 3]. Then the problem becomes to find a critical point of some functional depending on  $0 < \xi_1 < \dots < \xi_k$ . This is done in Section 5 where Theorem 1.1 is proved.

From the technical point of view, one difficulty is due to the form of the linearized operator. As  $r \rightarrow \infty$  dominates  $-\Delta + I$  (or  $\mathcal{L}_0$  as  $x \rightarrow -\infty$  after the change of variables) while near the regions of concentration the important part of the linearization is  $\Delta + p^*w_\mu^{p^*-1}$ . This is taken into account in the norm we use for the solutions of linearized problem, and it is more naturally written for the functions after the Emden-Fowler transformation. This is different from many previous works, but is already contained in [5].

## 2. THE FIRST APPROXIMATE SOLUTION

In this section, we build the first approximate solution to (1.3). In order to do this, we introduce  $U_\mu$  as the unique solution of the following problem

$$(2.1) \quad \begin{cases} -\Delta U_\mu + U_\mu = w_\mu^{p^*} & \text{in } \mathbb{R}^N, \\ U_\mu(z) \rightarrow 0 & \text{as } |z| \rightarrow \infty, \end{cases}$$

where  $w_\mu$  are the bubbles (1.6). We write

$$U_\mu(z) = w_\mu(z) + R_\mu(z).$$

Then  $R_\mu(z)$  satisfies

$$-\Delta R_\mu(z) + R_\mu(z) = -w_\mu(z) \quad \text{in } \mathbb{R}^N, \quad R_\mu(z) \rightarrow 0 \quad \text{as } |z| \rightarrow \infty.$$

We have the following result, whose proof is postponed to the Appendix.

**Lemma 2.1.** *Assume  $0 < \mu \leq 1$ , we have*

- (a)  $0 < U_\mu(z) \leq w_\mu(z)$ , for  $z \in \mathbb{R}^N$ .
- (b) One has

$$U_\mu(z) \leq C\mu^{\frac{N-2}{2}}|z|^{-(N+2)}, \quad \text{for } |z| \geq R,$$

where  $R$  is a large positive number but fixed.

(c) Given any  $\mu > 0$  small, we have

$$(2.2) \quad |R_\mu(z)| \leq C \frac{\mu^{\frac{N-2}{2}}}{|z|^{N-2}} \quad \text{for } N \geq 3, \quad |z| \geq 1.$$

$$(2.3) \quad |R_\mu(z)| \leq C \begin{cases} \mu^{-\frac{N-6}{2}} & \text{for } N \geq 5; \\ \mu \log \frac{1}{\mu} & \text{for } N = 4; \\ \mu^{\frac{1}{2}} & \text{for } N = 3. \end{cases} \quad |z| \leq \frac{\mu}{2}.$$

$$(2.4) \quad |R_\mu(z)| \leq C \begin{cases} \mu^{-\frac{N-6}{2}} \frac{1}{(1+|\frac{z}{\mu}|^2)^{\frac{N-4}{2}}} & \text{for } N \geq 5; \\ \mu \log \frac{1}{|z|} & \text{for } N = 4; \\ \mu^{\frac{1}{2}} & \text{for } N = 3. \end{cases} \quad \frac{\mu}{2} \leq |z| \leq 1.$$

We define the following Emden-Fowler transformation

$$v(x) = \mathcal{T}(u(r)) = \left( \frac{p^* - 1}{2} \right)^{\frac{2}{p^* - 1}} r^{\frac{2}{p^* - 1}} u(r), \quad r = |z| = e^{-\frac{p^* - 1}{2}x}$$

with  $x \in (-\infty, +\infty)$ . Using this transformation, finding a radial solution  $u(r)$  to problem (1.3) corresponds to that of solving problem (1.7), where

$$(2.5) \quad \alpha_\varepsilon = \left( \frac{p^* - 1}{2} \right)^{-\frac{2\varepsilon}{p^* - 1}}, \quad \beta_N = \left( \frac{p^* - 1}{2} \right)^{\frac{2(p^* - q)}{p^* - 1}}.$$

Define

$$V_\xi(x) = \mathcal{T}(U_\mu)(r), \quad \text{with } r = e^{-\frac{p^* - 1}{2}x}, \quad \mu = e^{-\frac{2}{N-2}\xi}.$$

Then  $V_\xi(x)$  is the solution of the problem

$$\begin{cases} \mathcal{L}_0 V_\xi(x) = W(x - \xi)^{p^*} & \text{in } (-\infty, +\infty); \\ V_\xi(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty. \end{cases}$$

Note that  $\mathcal{L}_0$  is the transformed operator associated to  $-\Delta + Id$  and given in (1.8).

We write

$$V_\xi(x) = W(x - \xi) + R_\xi(x),$$

where  $W$  is given in (1.9) and  $R_\xi(x) = \mathcal{T}(R_\mu)(r)$ . By the Emden-Fowler transformation and as a consequence of Lemma 2.1, we have the following estimates.

**Lemma 2.2.** For  $\xi > 0$ , we have

- (a)  $0 < V_\xi(x) \leq W(x - \xi) = O(e^{-|x - \xi|})$ , for  $x \in \mathbb{R}$ .  
(b)

$$(2.6) \quad V_\xi(x) \leq C e^{\frac{N+6}{N-2}x} e^{-\xi}, \quad \text{for } -\infty < x \leq -\frac{N-2}{2} \log R,$$

for  $R > 0$  is a fixed large number as Lemma 2.1.

(c) For  $N \geq 3$ , there is a positive constant  $C$ , such that

$$|R_\xi(x)| \leq C \begin{cases} e^{-|x - \xi|} & \text{if } x \leq 0; \\ e^{-|x - \xi|} e^{-\frac{2}{N-2} \min\{x, \xi\}} & \text{if } x \geq 0. \end{cases}$$

Define

$$Z_\xi(x) := \partial_\xi V_\xi(x) = \partial_\xi W(x - \xi) + \partial_\xi R_\xi(x).$$

Note that  $\partial_\xi W(x - \xi) = O(e^{-|x-\xi|})$  and

$$\partial_\xi W(x - \xi) = -\frac{2}{N-2} \mu \mathcal{T}(\partial_\mu w_\mu(r)),$$

$$(2.7) \quad Z_\xi(x) = -\frac{2}{N-2} \mu \mathcal{T}(\tilde{Z}_\mu(r)) \quad \text{with} \quad \tilde{Z}_\mu(z) = \partial_\mu U_\mu(z),$$

$$(2.8) \quad \partial_\xi R_\xi(x) = -\frac{2}{N-2} \mu \mathcal{T}(\partial_\mu R_\mu(r)).$$

Then from (6.1), (2.8) and Lemma 2.2 (c), we have for  $N \geq 3$ ,

$$|\partial_\xi R_\xi(x)| \leq C \begin{cases} e^{-|x-\xi|} & \text{if } x \leq 0; \\ e^{-|x-\xi|} e^{-\frac{2}{N-2} \min\{x,\xi\}} & \text{if } x \geq 0. \end{cases}$$

Therefore

$$Z_\xi(x) = O(e^{-|x-\xi|}) \quad \text{for } \forall x \in \mathbb{R}.$$

Moreover, from (6.2) and (2.7), we find

$$|Z_\xi(x)| \leq C e^{\frac{N+6}{N-2}x} e^{-\xi}, \quad \text{for } -\infty < x \leq -\frac{N-2}{2} \log R,$$

for a fixed large  $R > 0$ .

Let  $\eta > 0$  be a small but fixed number. Given an integer number  $k$ , let  $\Lambda_j$ , for  $j = 1, \dots, k$ , be positive numbers and satisfy

$$(2.9) \quad \eta < \Lambda_j < \frac{1}{\eta}.$$

Set

$$(2.10) \quad \mu_1 = \varepsilon^{\frac{2}{(N+2)-(N-2)q}} \Lambda_1 \quad \text{and} \quad \mu_j = \varepsilon^{\frac{2}{N-2}(j-1) + \frac{2}{(N+2)-(N-2)q}} \Lambda_j$$

for  $j = 2, \dots, k$ . We observe that

$$\frac{\mu_{j+1}}{\mu_j} = \varepsilon^{\frac{2}{N-2}} \frac{\Lambda_{j+1}}{\Lambda_j}, \quad j = 1, \dots, k-1.$$

Define  $k$  points in  $\mathbb{R}$  as

$$\mu_j = e^{-\frac{2}{N-2}\xi_j}, \quad j = 1, \dots, k.$$

Then we have that

$$0 < \xi_1 < \xi_2 < \dots < \xi_k,$$

and

$$(2.11) \quad \begin{cases} \xi_1 = -\frac{1}{p^*-q} \log \varepsilon - \frac{N-2}{2} \log \Lambda_1, \\ \xi_j - \xi_{j-1} = -\log \varepsilon - \frac{N-2}{2} \log \frac{\Lambda_j}{\Lambda_{j-1}}, \quad j = 2, \dots, k. \end{cases}$$

Set

$$(2.12) \quad W_j = W(x - \xi_j), \quad R_j = R_{\xi_j}(x), \quad V_j = W_j + R_j, \quad V = \sum_{j=1}^k V_j.$$

We look for a solution of (1.3) of the form  $u = \sum_{j=1}^k U_{\mu_j} + \psi$  corresponds to find a solution of (1.7) of the form  $v = V + \phi$ , where  $V$  is given by (2.12) and  $\phi = \mathcal{T}(\psi)$  is a small term. Thus problem (1.7) becomes

$$(2.13) \quad \begin{cases} \mathcal{L}_\varepsilon(\phi) = N(\phi) + E & \text{in } (-\infty, +\infty); \\ \phi(x) > 0 & \text{for } x \in (-\infty, +\infty); \\ \phi(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty, \end{cases}$$

where

$$\begin{aligned} \mathcal{L}_\varepsilon(\phi) &= \mathcal{L}_0(\phi) - \alpha_\varepsilon(p^* + \varepsilon)e^{\varepsilon x}V^{p^*+\varepsilon-1}\phi - \lambda q\beta_N e^{-(p^*-q)x}V^{q-1}\phi, \\ N(\phi) &= \alpha_\varepsilon e^{\varepsilon x} \left[ (V + \phi)^{p^*+\varepsilon} - V^{p^*+\varepsilon} - (p^* + \varepsilon)V^{p^*+\varepsilon-1}\phi \right] \\ &\quad + \lambda\beta_N e^{-(p^*-q)x} \left[ (V + \phi)^q - V^q - qV^{q-1}\phi \right] \end{aligned}$$

and

$$\begin{aligned} E &= \alpha_\varepsilon e^{\varepsilon x}V^{p^*+\varepsilon} - \mathcal{L}_0(V) + \lambda\beta_N e^{-(p^*-q)x}V^q \\ &= \alpha_\varepsilon e^{\varepsilon x}V^{p^*+\varepsilon} - \sum_{j=1}^k W_j^{p^*} + \lambda\beta_N e^{-(p^*-q)x}V^q. \end{aligned}$$

where  $\mathcal{L}_0$  is defined by (1.8).

### 3. THE LINEAR PROBLEM

In order to solve problem (2.13), we consider first the following problem: given points  $\xi = (\xi_1, \dots, \xi_k)$ , finding a function  $\phi$  such that for certain constants  $c_1, c_2, \dots, c_k$ ,

$$(3.1) \quad \begin{cases} \mathcal{L}_\varepsilon(\phi) = N(\phi) + E + \sum_{j=1}^k c_j Z_j & \text{in } (-\infty, +\infty); \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0; \\ \int_{\mathbb{R}} Z_j \phi = 0, \quad \forall j = 1, \dots, k, \end{cases}$$

where  $Z_j(x) = Z_{\xi_j}(x) = \partial_{\xi_j} V_{\xi_j}(x)$  for  $j = 1, 2, \dots, k$ .

To solve (3.1), it is important to understand its linear part, thus we consider the following problem: given a function  $h$ , finding  $\phi$  such that

$$(3.2) \quad \begin{cases} \mathcal{L}_\varepsilon(\phi) = h + \sum_{j=1}^k c_j Z_j & \text{in } (-\infty, +\infty); \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0; \\ \int_{\mathbb{R}} Z_j \phi = 0, \quad \forall j = 1, \dots, k, \end{cases}$$

for certain constants  $c_j$ .

Now we analyze invertibility properties of the operator  $\mathcal{L}_\varepsilon$  under the orthogonality conditions. Let  $\sigma$  satisfy

$$(3.3) \quad 0 < \sigma < \min \left\{ q - 1, 1, \frac{(N + 2)(2q - 1)}{N + 6}, \frac{3q - p^*}{2} \right\}.$$

We define the real number  $M$  as follows

$$(3.4) \quad M = \begin{cases} 0, & \text{if } 1 \geq \frac{4}{N-2} + \sigma; \\ \max\{0, \gamma\}, & \text{if } 1 \leq \frac{4}{N-2} + \sigma, \end{cases}$$



where  $\gamma$  satisfies

$$\left(1 - \left(\frac{4}{N-2} + \sigma\right)^2\right) e^{-\frac{4}{N-2}\gamma} = -\frac{1}{2} \left(\frac{2}{N-2}\right)^2.$$

We define the following norms for functions  $\phi, h$  defined on  $\mathbb{R}$ ,

$$(3.5) \quad \|\phi\|_* = \sup_{x \leq -M} e^{-(\frac{4}{N-2} + \sigma)x} e^{\sigma\xi_1} |\phi(x)| + \sup_{x \in \mathbb{R}} \left( \sum_{j=1}^k e^{-\sigma|x-\xi_j|} \right)^{-1} |\phi(x)|,$$

and

$$\|h\|_{**} = \sup_{x \in \mathbb{R}} \left( \sum_{j=1}^k e^{-\sigma|x-\xi_j|} \right)^{-1} |h(x)|.$$

The choice of norm here is motivated by the presence of 2 regimes in the solution of the linearized problem. Near the concentration points  $\xi_j$  we have a right hand side of the form  $|h(x)| \leq C e^{-\sigma|x-\xi_j|}$  and near these points the dominant terms in the linear operator  $\mathcal{L}_\varepsilon$  are

$$-\phi'' + \phi - \alpha_\varepsilon(p^* + \varepsilon) e^{\varepsilon x} V^{p^* + \varepsilon - 1} \phi,$$

so we can expect the solution  $\phi$  to be controlled by  $|\phi(x)| \leq C e^{-\sigma|x-\xi_j|}$ . For  $x \leq 0$  the dominant part of the linear operator is

$$\left(\frac{2}{N-2}\right)^2 e^{-\frac{4}{N-2}x} \phi.$$

Since the right hand side is controlled by  $e^{-\sigma|x-\xi_1|}$ , we can control  $\phi$  using as supersolution  $e^{(\frac{4}{N-2} + \sigma)x} e^{-\sigma\xi_1}$ . Actually this will be a super solution for the whole linear operator for  $x \leq -M$ , where  $M$  is defined in (3.4).

The main result in this section is solvability of problem (3.2).

**Proposition 3.1.** *There exist positive numbers  $\varepsilon_0$ , and  $C > 0$  such that if the points  $0 < \xi_1 < \xi_2 < \dots < \xi_k$  satisfy (2.11), then for all  $0 < \varepsilon < \varepsilon_0$  and all functions  $h \in C(\mathbb{R}; \mathbb{R})$  with  $\|h\|_{**} < +\infty$ , problem (3.2) has a unique solution  $\phi =: T_\varepsilon(h)$  with  $\|\phi\|_* < +\infty$ . Moreover,*

$$(3.6) \quad \|\phi\|_* \leq C \|h\|_{**} \quad \text{and} \quad |c_j| \leq C \|h\|_{**}.$$

We first consider a simpler problem

$$(3.7) \quad \begin{cases} \mathcal{L}_0(\phi) - \alpha_\varepsilon(p^* + \varepsilon) e^{\varepsilon x} V^{p^* + \varepsilon - 1} \phi = h + \sum_{j=1}^k c_j Z_j & \text{in } (-\infty, +\infty); \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0; \\ \int_{\mathbb{R}} Z_j \phi = 0, \quad \forall j = 1, \dots, k, \end{cases}$$

for certain constants  $c_j$ , here  $\mathcal{L}_0$  is defined by (1.8).

**Lemma 3.2.** *Under the assumptions of Proposition 3.1, then for all  $0 < \varepsilon < \varepsilon_0$  and any  $h, \phi$  solution of (3.7), we have*

$$(3.8) \quad \|\phi\|_* \leq C \|h\|_{**},$$

and

$$(3.9) \quad |c_j| \leq C \|h\|_{**}.$$

*Proof.* To prove (3.8), by contradiction, we suppose that there exist sequences  $\phi_n$ ,  $h_n$ ,  $\varepsilon_n$  and  $c_j^n$  that satisfy (3.7), with

$$\|\phi_n\|_* = 1, \quad \|h_n\|_{**} \rightarrow 0, \quad \varepsilon_n \rightarrow 0.$$

We get a contradiction by the following steps.

*Step 1:*  $c_j^n \rightarrow 0$  as  $n \rightarrow +\infty$ .

Multiplying (3.7) by  $Z_i^n$  and integrating by parts twice, we get that

$$(3.10) \quad \sum_{j=1}^k c_j^n \int_{\mathbb{R}} Z_j^n Z_i^n = - \int_{\mathbb{R}} h_n Z_i^n + \int_{\mathbb{R}} \left[ \mathcal{L}_0(Z_i^n) - \alpha_{\varepsilon_n} (p^* + \varepsilon_n) e^{\varepsilon_n x} V^{p^* + \varepsilon_n - 1} Z_i^n \right] \phi_n.$$

Note that

$$\int_{\mathbb{R}} Z_j^n Z_i^n = C \delta_{ij} + o(1),$$

where  $\delta_{ij}$  is Kronecker's delta. Then (3.10) defines a linear system in the  $c_j^n$ 's which is almost diagonal as  $n \rightarrow \infty$ .

Since  $Z_i^n(x) = \partial_{\xi_i^n} V_{\xi_i^n}(x) = O(e^{-|x - \xi_i^n|})$ , we then have

$$(3.11) \quad \begin{aligned} \left| \int_{\mathbb{R}} h_n Z_i^n \right| &\leq C \|h_n\|_{**} \int_{\mathbb{R}} \left( \sum_{j=1}^k e^{-\sigma|x - \xi_j^n|} \right) e^{-|x - \xi_i^n|} dx \\ &\leq Ck \|h_n\|_{**} \int_{\mathbb{R}} e^{-|y|} dy \leq C \|h_n\|_{**}. \end{aligned}$$

Moreover,  $Z_i^n$  satisfy

$$\mathcal{L}_0(Z_i^n) = p^* W^{p^* - 1}(x - \xi_i^n) \partial_{\xi_i^n} W(x - \xi_i^n),$$

so we get

$$(3.12) \quad \left| \int_{\mathbb{R}} \left[ \mathcal{L}_0(Z_i^n) - \alpha_{\varepsilon_n} (p^* + \varepsilon_n) e^{\varepsilon_n x} V^{p^* + \varepsilon_n - 1} Z_i^n \right] \phi_n \right| = o(1) \|\phi_n\|_*.$$

From (3.10)-(3.12), we obtain

$$(3.13) \quad |c_j^n| \leq C \|h_n\|_{**} + o(1) \|\phi_n\|_*.$$

Thus  $\lim_{n \rightarrow \infty} c_j^n = 0$ .

*Step 2:* For any  $L > 0$ , any  $l \in \{1, 2, \dots, k\}$ , we have

$$(3.14) \quad \sup_{x \in [\xi_l^n - L, \xi_l^n + L]} |\phi_n(x)| \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Indeed, suppose not, we assume that there exist  $L > 0$  and some  $l \in \{1, 2, \dots, k\}$  such that

$$|\phi_n(x_{n,l})| \geq c > 0, \quad \text{for some } x_{n,l} \in [\xi_l^n - L, \xi_l^n + L].$$

By elliptic estimates, there is a subsequence of  $\phi_n$  converging uniformly on compact sets to a nontrivial bounded solution  $\tilde{\phi}$  of

$$\mathcal{L}_0(\tilde{\phi}) = p^* W^{p^* - 1}(x - \xi_l) \tilde{\phi},$$

where  $\xi_l = \lim_{n \rightarrow \infty} \xi_l^n$ . By nondegeneracy [27], it is well known that  $\tilde{\phi} = cZ_l$  for some constant  $c \neq 0$ . But taking the limit in the orthogonality condition  $\int_{\mathbb{R}} Z_l^n \phi_n = 0$ , we obtain  $\tilde{\phi} = 0$ , which is a contradiction. Thus (3.14) holds.

*Step 3: We prove that  $\|\phi_n\|_* \rightarrow 0$  as  $n \rightarrow \infty$ .*

**Claim:** For any  $L > 0$  and  $j \in \{1, 2, \dots, k\}$ , we have

$$(3.15) \quad \sup_{\mathbb{R} \setminus \cup_{j=1}^k [\xi_j^n - L, \xi_j^n + L]} \left( \sum_{j=1}^k e^{-\sigma|x - \xi_j^n|} \right)^{-1} |\phi_n(x)| \rightarrow 0,$$

and

$$(3.16) \quad \sup_{x \leq -M} e^{-(\frac{4}{N-2} + \sigma)x} e^{\sigma \xi_1^n} |\phi_n(x)| \rightarrow 0,$$

as  $n \rightarrow +\infty$ .

By the definition of  $\|\cdot\|_*$  in (3.5), using (3.14), (3.15) and (3.16), we get that  $\|\phi_n\|_* \rightarrow 0$  as  $n \rightarrow \infty$ .

Now we prove the above claim. We note that

$$h_n + \sum_{j=1}^k c_j^n Z_j^n \leq (C_0 \|h_n\|_{**} + o(\|\phi_n\|_*)) \sum_{j=1}^k e^{-\sigma|x - \xi_j^n|}, \quad \text{with } C_0 > 0.$$

For  $x \in \mathbb{R} \setminus \cup_{j=1}^k [\xi_j^n - L, \xi_j^n + L]$ , let us define

$$\begin{aligned} \tilde{\psi}_n(x) &= \left( C_0 \|h_n\|_{**} + e^{\sigma L} \sup_{\cup_{j=1}^k [\xi_j^n - L, \xi_j^n + L]} |\phi_n(x)| + o(\|\phi_n\|_*) \right) \sum_{j=1}^k e^{-\sigma|x - \xi_j^n|} \\ &\quad + \varrho \sum_{j=1}^k e^{-\bar{\sigma}|x - \xi_j^n|} \end{aligned}$$

with  $\varrho > 0$  small but fixed and  $0 < \bar{\sigma} < \sigma$ . Then by choosing suitable large  $L > 0$ , we get

$$\begin{aligned} \mathcal{L}_0(\tilde{\psi}_n(x)) - \alpha_{\varepsilon_n}(p^* + \varepsilon_n) e^{\varepsilon_n x} V^{p^* + \varepsilon_n - 1} \tilde{\psi}_n(x) \\ \geq \mathcal{L}_0(\phi_n(x)) - \alpha_{\varepsilon_n}(p^* + \varepsilon_n) e^{\varepsilon_n x} V^{p^* + \varepsilon_n - 1} \phi_n(x). \end{aligned}$$

On the other hand, we have that for any  $L > 0$  and  $j \in \{1, 2, \dots, k\}$ ,

$$\tilde{\psi}_n(\xi_j^n - L) \geq \phi_n(\xi_j^n - L) \quad \text{and} \quad \tilde{\psi}_n(\xi_j^n + L) \geq \phi_n(\xi_j^n + L).$$

Moreover, there exists  $R > 0$  large enough, such that

$$\tilde{\psi}_n(R) \geq \phi_n(R),$$

and

$$\tilde{\psi}_n(-R) \geq \phi_n(-R).$$

By the maximum principle, we get

$$\phi_n(x) \leq \tilde{\psi}_n(x) \quad \text{for } x \in [-R, R] \setminus \cup_{j=1}^k [\xi_j^n - L, \xi_j^n + L].$$

Similarly, we obtain  $\phi_n(x) \geq -\tilde{\psi}_n(x)$  for  $x \in [-R, R] \setminus \cup_{j=1}^k [\xi_j^n - L, \xi_j^n + L]$ . Thus

$$|\phi_n(x)| \leq \tilde{\psi}_n(x) \quad \text{for } x \in [-R, R] \setminus \cup_{j=1}^k [\xi_j^n - L, \xi_j^n + L].$$

Letting  $R \rightarrow +\infty$ , we get

$$|\phi_n(x)| \leq \tilde{\psi}_n(x) \quad \text{for } x \in \mathbb{R} \setminus \cup_{j=1}^k [\xi_j^n - L, \xi_j^n + L].$$

Letting  $\rho \rightarrow 0$ , for  $x \in \mathbb{R} \setminus \cup_{j=1}^k [\xi_j^n - L, \xi_j^n + L]$ , we have that

$$|\phi_n(x)| \leq \left( C_0 \|h_n\|_{**} + e^{\sigma L} \sup_{\cup_{j=1}^k [\xi_j^n - L, \xi_j^n + L]} |\phi_n(x)| + o(\|\phi_n\|_*) \right) \sum_{j=1}^k e^{-\sigma|x-\xi_j^n|}.$$

So (3.15) holds.

For  $x \leq -M$ , let  $\rho > 0$  small and  $C_1 > 0$  be chosen later, we define

$$\psi_n(x) = C_1 (C_0 \|h_n\|_{**} + o(\|\phi_n\|_*)) e^{(\frac{4}{N-2} + \sigma)x} e^{-\sigma\xi_1^n} + \rho e^{\frac{4}{N-2}x}.$$

By the maximum principle, we get

$$\phi_n(x) \leq \psi_n(x) \quad \text{for } x \in [-R, -M],$$

if  $R > 0$  is large enough. By a similar argument, we obtain  $\phi_n(x) \geq -\psi_n(x)$  for  $x \in [-R, -M]$ . Thus

$$|\phi_n(x)| \leq \psi_n(x) \quad \text{for } x \in [-R, -M].$$

Let  $R \rightarrow +\infty$ , we get

$$|\phi_n(x)| \leq \psi_n(x) \quad \text{for } x \in [-\infty, -M].$$

Let  $\rho \rightarrow 0$ , we have

$$|\phi_n(x)| \leq C_1 (C_0 \|h_n\|_{**} + o(\|\phi_n\|_*)) e^{(\frac{4}{N-2} + \sigma)x} e^{-\sigma\xi_1^n} \quad \text{for } x \in [-\infty, -M].$$

So we obtain that (3.16) holds.

Moreover, estimate (3.9) follows from (3.13) and (3.8).  $\square$

**Proof of Proposition 3.1.** From Lemma 3.2, for  $\phi$  and  $h$  satisfying (3.2), we then have

$$\|\phi\|_* \leq C \left( \|h\|_{**} + \|e^{-(p^*-q)x} V^{q-1} \phi\|_{**} \right),$$

and

$$|c_j| \leq C \left( \|h\|_{**} + \|e^{-(p^*-q)x} V^{q-1} \phi\|_{**} \right).$$

In order to establish (3.6), it is sufficient to show that

$$(3.17) \quad \|e^{-(p^*-q)x} V^{q-1} \phi\|_{**} \leq o(1) \|\phi\|_*.$$

Indeed,

$$(3.18) \quad \begin{aligned} \|e^{-(p^*-q)x} V^{q-1} \phi\|_{**} &\leq \sup_{x \leq -M} \left( \sum_{j=1}^k e^{-\sigma|x-\xi_j^n|} \right)^{-1} \left| e^{-(p^*-q)x} V^{q-1} \phi \right| \\ &\quad + \sup_{x \geq -M} \left( \sum_{j=1}^k e^{-\sigma|x-\xi_j^n|} \right)^{-1} \left| e^{-(p^*-q)x} V^{q-1} \phi \right| := Q_1 + Q_2. \end{aligned}$$

Now we estimate  $Q_1$  and  $Q_2$  respectively, we first have

$$(3.19) \quad \begin{aligned} Q_1 &\leq C \sup_{x \leq -M} e^{\sigma|x-\xi_1^n|} |\phi(x)| e^{-(p^*-q)x} V^{q-1} \\ &\leq C e^{-(q-1)\xi_1^n} \sup_{x \leq -M} e^{-(\frac{4}{N-2} + \sigma)x} e^{\sigma\xi_1^n} |\phi(x)|. \end{aligned}$$

For  $Q_2$ , if  $-M \leq x \leq \xi_1$ , then we have

$$\begin{aligned} e^{-(p^*-q)x} V^{q-1} &\leq \sum_{j=1}^k e^{-(p^*-q)x} e^{-(q-1)|x-\xi_j|} \leq C e^{(2q-p^*-1)x} e^{-(q-1)\xi_1} \\ &\leq C \max \left\{ e^{-(p^*-q)\xi_1}, e^{-(q-1)\xi_1} \right\}. \end{aligned}$$

If  $x \geq \xi_1$ , then we have

$$e^{-(p^*-q)x} V^{q-1} \leq \sum_{j=1}^k e^{-(p^*-q)x} e^{-(q-1)|x-\xi_j|} \leq C e^{-(p^*-q)x} \leq C e^{-(p^*-q)\xi_1}.$$

Thus we find

$$(3.20) \quad Q_2 \leq C \max \left\{ e^{-(p^*-q)\xi_1}, e^{-(q-1)\xi_1} \right\} \sup_{x \geq -M} \left( \sum_{j=1}^k e^{-\sigma|x-\xi_j|} \right)^{-1} |\phi(x)|.$$

From (3.18), (3.19) and (3.20), we get

$$\|e^{-(p^*-q)x} V^{q-1} \phi\|_{**} \leq C \max \left\{ e^{-(p^*-q)\xi_1}, e^{-(q-1)\xi_1} \right\} \|\phi\|_* = o(1) \|\phi\|_*.$$

So estimate (3.17) holds.

We now prove the existence and uniqueness of solution to (3.2). Consider the Hilbert space

$$H = \left\{ \phi \in H^1(\mathbb{R}) : \int_{\mathbb{R}} Z_j \phi = 0, \quad \forall j = 1, 2, \dots, k. \right\}$$

with inner product

$$\langle \phi, \psi \rangle = \int_{\mathbb{R}} (\phi' \psi' + \phi \psi) dx.$$

Then problem (3.7) is equivalent to find  $\phi \in H$  such that

$$(3.21) \quad \begin{aligned} \langle \phi, \psi \rangle &= \int_{\mathbb{R}} \left[ \alpha_\varepsilon (p^* + \varepsilon) V^{p^*+\varepsilon-1} \phi + \lambda q \beta_N e^{-(p^*-q)x} V^{q-1} \phi \right. \\ &\quad \left. + \left( \frac{2}{N-2} \right)^2 e^{-\frac{4}{N-2}x} \phi + h \right] \psi dx, \end{aligned}$$

for all  $\psi \in H$ . By the Riesz representation theorem, (3.21) is equivalent to solve

$$(3.22) \quad \phi = K(\phi) + \tilde{h}$$

with  $\tilde{h} \in H$  depending linearly on  $h$  and  $K : H \rightarrow H$  being a compact operator. Fredholm's alternative yields there is a unique solution to problem (3.22) for any  $h$  provided that

$$(3.23) \quad \phi = K(\phi)$$

has only the zero solution in  $H$ . (3.23) is equivalent to problem (3.2) with  $h = 0$ . If  $h = 0$ , estimate (3.6) implies that  $\phi = 0$ . This ends the proof of Proposition 3.1.

Now we study the differentiability of the operator  $T_\varepsilon$  with respect to  $\xi = (\xi_1, \dots, \xi_k)$ . Consider the Banach space

$$C_* = \{f \in C(\mathbb{R}) : \|f\|_{**} < \infty\}$$

endowed with the  $\|\cdot\|_{**}$  norm. The following result holds.

**Proposition 3.3.** *Under the assumptions of Proposition 3.1, the map  $\xi \mapsto T_\varepsilon$  is of class  $C^1$ . Moreover,*

$$\|D_\xi T_\varepsilon(h)\|_* \leq C\|h\|_{**}$$

uniformly on the vectors  $\xi$  which satisfy (2.11).

*Proof.* Fix  $h \in \mathcal{C}_*$  and let  $\phi = T_\varepsilon(h)$  for  $\varepsilon < \varepsilon_0$ . Let us recall that  $\phi$  satisfies

$$\begin{cases} \mathcal{L}_\varepsilon(\phi) = h + \sum_{j=1}^k c_j Z_j & \text{in } (-\infty, +\infty); \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0; \\ \int_{\mathbb{R}} Z_j \phi = 0, \quad \forall j = 1, \dots, k, \end{cases}$$

for certain constants  $c_j$ . Differentiating above equation with respect to  $\xi_l$ ,  $l \in \{1, \dots, k\}$ . Set  $Y = \partial_{\xi_l} \phi$  and  $d_j = \partial_{\xi_l} c_j$ , we have

$$\begin{cases} \mathcal{L}_\varepsilon(Y) = \bar{h} + \sum_{j=1}^k d_j Z_j & \text{in } (-\infty, +\infty); \\ \lim_{|x| \rightarrow \infty} Y(x) = 0; \\ \int_{\mathbb{R}} Y Z_j + \phi \partial_{\xi_l} Z_j = 0, \quad \forall j = 1, \dots, k, \end{cases}$$

where

$$\bar{h} = \alpha_\varepsilon(p^* + \varepsilon)(p^* + \varepsilon - 1)e^{\varepsilon x} V^{p^* + \varepsilon - 2} Z_l \phi + \lambda q(q-1)\beta_N e^{-(p^* - q)x} V^{q-2} Z_l \phi + c_l \partial_{\xi_l} Z_l.$$

Let  $\eta = Y - \sum_{i=1}^k b_i Z_i$ , where  $b_i \in \mathbb{R}$  is chosen such that

$$\int_{\mathbb{R}} \eta Z_j = 0,$$

that is,

$$(3.24) \quad \sum_{i=1}^k b_i \int_{\mathbb{R}} Z_i Z_j = \int_{\mathbb{R}} Y Z_j = \int_{\mathbb{R}} \partial_{\xi_l} \phi Z_j = - \int_{\mathbb{R}} \phi \partial_{\xi_l} Z_j.$$

This is an almost diagonal system, it has a unique solution and we have

$$(3.25) \quad |b_i| \leq C\|\phi\|_*.$$

Moreover,  $\eta$  satisfies

$$(3.26) \quad \begin{cases} \mathcal{L}_\varepsilon(\eta) = g + \sum_{j=1}^k d_j Z_j & \text{in } (-\infty, +\infty); \\ \lim_{|x| \rightarrow \infty} \eta(x) = 0; \\ \int_{\mathbb{R}} \eta Z_j = 0, \quad \forall j = 1, \dots, k, \end{cases}$$

with

$$g = \bar{h} - \sum_{i=1}^k b_i \mathcal{L}_\varepsilon(Z_i).$$

From Proposition 3.1, there is a unique solution  $\eta = T_\varepsilon(g)$  to (3.26) and

$$(3.27) \quad \|\eta\|_* \leq C\|g\|_{**}.$$

Moreover, we have

$$\begin{aligned}
\|g\|_{**} &\leq C\|e^{\varepsilon x}V^{p^*+\varepsilon-2}Z_l\phi\|_{**} + C\|e^{-(p^*-q)x}V^{q-2}Z_l\phi\|_{**} \\
&\quad + \|c_l\partial_{\xi_l}Z_l\|_{**} + \sum_{i=1}^k |b_i|\|\mathcal{L}_\varepsilon(Z_i)\|_{**} \\
(3.28) \quad &\leq C(\|\phi\|_* + |c_l| + |b_i|) \leq C\|h\|_{**},
\end{aligned}$$

because  $|b_i| \leq C\|\phi\|_*$ ,  $\|\phi\|_* \leq C\|h\|_{**}$  and  $|c_l| \leq C\|h\|_{**}$ .

By (3.25), (3.27), (3.28) and  $\|Z_i\|_* \leq C$ , we obtain that

$$\|\partial_{\xi_l}\phi\|_* \leq \|\eta\|_* + \sum_{i=1}^k |b_i|\|Z_i\|_* \leq C\|h\|_{**}.$$

Besides  $\partial_{\xi_l}\phi$  depends continuously on  $\xi$  in the considered region for this norm.  $\square$

#### 4. NONLINEAR PROBLEM

In this section, our purpose is to study nonlinear problem. We first have the validity of the following result.

**Lemma 4.1.** *We have*

$$(4.1) \quad \|N(\phi)\|_{**} \leq C \left( \|\phi\|_*^{\min\{p^*, 2\}} + \|\phi\|_*^{\min\{q, 2\}} \right),$$

and

$$(4.2) \quad \|\partial_\phi N(\phi)\|_{**} \leq C \left( \|\phi\|_*^{\min\{p^*-1, 1\}} + \|\phi\|_*^{\min\{q-1, 1\}} \right),$$

for  $\|\phi\|_* \leq 1$ .

*Proof.* By the fundamental theorem of calculus and the definition of  $\|\cdot\|_{**}$ , we have

$$\begin{aligned}
&\|N(\phi)\|_{**} \\
&\leq \alpha_\varepsilon(p^* + \varepsilon) \sup_{x \in \mathbb{R}} \left( \sum_{j=1}^k e^{-\sigma|x-\xi_j|} \right)^{-1} e^{\varepsilon x} \left| \int_0^1 [(V + t\phi)^{p^*+\varepsilon-1} - V^{p^*+\varepsilon-1}] \phi \, dt \right| \\
&\quad + \lambda q \beta_N \sup_{x \in \mathbb{R}} \left( \sum_{j=1}^k e^{-\sigma|x-\xi_j|} \right)^{-1} e^{-(p^*-q)x} \left| \int_0^1 [(V + t\phi)^{q-1} - V^{q-1}] \phi \, dt \right| \\
&:= N_1 + N_2.
\end{aligned}$$

Using

$$\|a + b\|^q - \|a\|^q \leq C \begin{cases} |a|^{q-1}|b| + |b|^q & \text{if } q \geq 1; \\ \min\{|a|^{q-1}|b|, |b|^q\} & \text{if } 0 < q < 1, \end{cases}$$

if  $p^* \geq 2$  and for  $\|\phi\|_* \leq 1$ , we have

$$\begin{aligned}
N_1 &\leq C \sup_{x \in \mathbb{R}} \left( \sum_{j=1}^k e^{-\sigma|x-\xi_j|} \right)^{-1} e^{\varepsilon x} V^{p^*+\varepsilon-2} |\phi|^2 + C \sup_{x \in \mathbb{R}} \left( \sum_{j=1}^k e^{-\sigma|x-\xi_j|} \right)^{-1} e^{\varepsilon x} |\phi|^{p^*+\varepsilon} \\
&\leq C\|\phi\|_*^2 + C\|\phi\|_*^{p^*+\varepsilon} \leq C\|\phi\|_*^2.
\end{aligned}$$

Similarly, if  $1 < p^* < 2$ , we find that  $N_1 \leq C\|\phi\|_*^{p^*}$ . Thus we get

$$N_1 \leq C\|\phi\|_*^{\min\{p^*, 2\}}.$$

Moreover, by similar computations as  $N_1$ , we can conclude that

$$N_2 \leq C\|\phi\|_*^{\min\{q, 2\}}.$$

Thus we get (4.1).

We differentiate  $N(\phi)$  with respect to  $\phi$ , we have

$$\partial_\phi N(\phi) = \alpha_\varepsilon(p^* + \varepsilon)e^{\varepsilon x} \left[ (V + \phi)^{p^* + \varepsilon - 1} - V^{p^* + \varepsilon - 1} \right] + \lambda\beta_N q e^{-(p^* - q)x} \left[ (V + \phi)^{q-1} - V^{q-1} \right].$$

By a similar argument as  $\|N(\phi)\|_{**}$ , (4.2) holds.  $\square$

**Lemma 4.2.** *Let  $\sigma > 0$  satisfy (3.3) and  $0 < \xi_1 < \xi_2 < \dots < \xi_k$  satisfy (2.11). If  $q$  satisfies (1.4), then there exist  $\tau \in (\frac{1}{2}, 1)$  and a constant  $C > 0$ , such that*

$$\|E\|_{**} \leq C\varepsilon^\tau, \quad \|\partial_\xi E\|_{**} \leq C\varepsilon^\tau.$$

*Proof.* We have

$$\begin{aligned} E &= \alpha_\varepsilon e^{\varepsilon x} \left( V^{p^* + \varepsilon} - V^{p^*} \right) + (\alpha_\varepsilon e^{\varepsilon x} - 1)V^{p^*} + \left( V^{p^*} - \left( \sum_{j=1}^k W_j \right)^{p^*} \right) \\ &\quad + \left( \left( \sum_{j=1}^k W_j \right)^{p^*} - \sum_{j=1}^k W_j^{p^*} \right) + \lambda\beta_N e^{-(p^* - q)x} V^q \\ (4.3) \quad &:= E_1 + E_2 + E_3 + E_4 + E_5. \end{aligned}$$

*Estimate of  $E_1$ :*

$$|E_1| = \left| \varepsilon \alpha_\varepsilon e^{\varepsilon x} \int_0^1 V^{p^* + t\varepsilon} \log V dt \right| \leq C\varepsilon \sum_{j=1}^k e^{-\sigma|x - \xi_j|}.$$

*Estimate of  $E_2$ :* By the Taylor expansion, we have

$$\begin{aligned} |E_2| &= \left| \left( \left( \frac{p^* - 1}{2} \right)^{-\frac{2\varepsilon}{p^* - 1}} e^{\varepsilon x} - 1 \right) V^{p^*} \right| \\ &= \left( \varepsilon x \int_0^1 e^{t\varepsilon x} dt + O(\varepsilon^2) e^{\varepsilon x} \right) V^{p^*} \leq C\varepsilon |\log \varepsilon| \sum_{j=1}^k e^{-\sigma|x - \xi_j|}. \end{aligned}$$

*Estimate of  $E_3$ :* Since

$$|E_3| = \left| V^{p^*} - \left( \sum_{j=1}^k W_j \right)^{p^*} \right| \leq C V^{p^* - 1} \sum_{j=1}^k |R_{\xi_j}(x)|.$$

Thanks to Lemma 2.2, for  $x \leq 0$ , we have

$$|E_3| \leq C V^{p^* - 1} \sum_{j=1}^k e^{-|x - \xi_j|} \leq C V^{p^* - 1} e^{-\xi_1} \leq C \varepsilon^{\frac{1}{p^* - q}} \sum_{j=1}^k e^{-\sigma|x - \xi_j|}.$$



For  $0 \leq x \leq \xi_1$ ,

$$\begin{aligned} |E_3| &\leq CV^{p^*-1} \sum_{j=1}^k e^{-|x-\xi_j|} e^{-\frac{2}{N-2} \min\{x, \xi_j\}} \\ &\leq C \sum_{j=1}^k e^{-\sigma|x-\xi_j|} \begin{cases} \varepsilon^{\frac{2}{N+2-(N-2)q}} & \text{if } N \geq 4; \\ \varepsilon^{\frac{1}{5-q}} & \text{if } N = 3. \end{cases} \end{aligned}$$

If  $x \geq \xi_1$ , for  $0 < \sigma < p^* - 1$ , we have

$$\begin{aligned} |E_3| &\leq CV^{p^*-1} \sum_{j=1}^k e^{-|x-\xi_j|} e^{-\frac{2}{N-2} \min\{x, \xi_j\}} \\ &\leq CV^{p^*-1} e^{-\frac{2}{N-2} \xi_1} \leq C \varepsilon^{\frac{2}{N+2-(N-2)q}} \sum_{j=1}^k e^{-\sigma|x-\xi_j|}. \end{aligned}$$

Therefore we get for  $x \in \mathbb{R}$ ,

$$|E_3| \leq C \sum_{j=1}^k e^{-\sigma|x-\xi_j|} \begin{cases} \varepsilon^{\frac{2}{N+2-(N-2)q}} & \text{if } N \geq 4; \\ \varepsilon^{\frac{1}{5-q}} & \text{if } N = 3. \end{cases}$$

*Estimate of  $E_4$ :* If  $-\infty < x \leq \frac{\xi_1 + \xi_2}{2}$ , we have

$$\begin{aligned} |E_4| &\leq \left| \left( \sum_{j=1}^k W(x - \xi_j) \right)^{p^*} - W(x - \xi_1)^{p^*} \right| + \left| \sum_{j=2}^k W(x - \xi_j)^{p^*} \right| \\ &\leq p^* \left( \sum_{j=1}^k W(x - \xi_j) \right)^{p^*-1} \sum_{j=2}^k W(x - \xi_j) + \sum_{j=2}^k W(x - \xi_j)^{p^*} \\ &= p^* \left( \sum_{j=1}^k W(x - \xi_j) \right)^{p^*-1-\theta} \left( \sum_{j=1}^k W(x - \xi_j) \right)^{\theta} \sum_{j=2}^k W(x - \xi_j) + \sum_{j=2}^k W(x - \xi_j)^{p^*} \end{aligned}$$

with a positive number  $\theta$ , satisfying  $0 < \theta < p^* - 1 - \sigma$ . Note that

$$\left( \sum_{j=1}^k W(x - \xi_j) \right)^{\theta} \sum_{j=2}^k W(x - \xi_j) \leq C \varepsilon^{\frac{1+\theta}{2}}.$$

Moreover,

$$\sum_{j=2}^k W(x - \xi_j)^{p^*} \leq C \varepsilon^{\frac{p^*-\sigma}{2}} \sum_{j=1}^k e^{-\sigma|x-\xi_j|}.$$

Thus

$$|E_4| \leq C \varepsilon^{\frac{1+\theta}{2}} \sum_{j=1}^k e^{-\sigma|x-\xi_j|}, \quad \text{for } -\infty < x \leq \frac{\xi_1 + \xi_2}{2},$$

with  $0 < \theta < p^* - 1 - \sigma$ . Similarly, for  $\frac{\xi_{l-1} + \xi_l}{2} \leq x \leq \frac{\xi_l + \xi_{l+1}}{2}$  with  $l = 2, \dots, k-1$ , and  $x \geq \frac{\xi_{k-1} + \xi_k}{2}$ , we get

$$|E_4| \leq C\varepsilon^{\frac{1+\theta}{2}} \sum_{j=1}^k e^{-\sigma|x-\xi_j|}.$$

Therefore for  $x \in \mathbb{R}$ , we have

$$|E_4| \leq C\varepsilon^{\frac{1+\theta}{2}} \sum_{j=1}^k e^{-\sigma|x-\xi_j|}, \quad \text{where } 0 < \theta < p^* - 1 - \sigma.$$

The estimate of  $E_5$  is similar as the previous ones and we get

$$|E_5| \leq C \max\{\varepsilon, \varepsilon^{\frac{q-\sigma}{p^*-q}}\} \sum_{j=1}^k e^{-\sigma|x-\xi_j|}.$$

From (4.3) and the previous estimates, for  $0 < \theta < p^* - 1 - \sigma$  with  $\sigma$  satisfying (3.3), we have

$$\|E\|_{**} \leq C \begin{cases} \max\left\{\varepsilon|\log \varepsilon|, \varepsilon^{\frac{2}{N+2-(N-2)q}}, \varepsilon^{\frac{1+\theta}{2}}, \varepsilon^{\frac{q-\sigma}{p^*-q}}\right\} & \text{if } N \geq 4; \\ \max\left\{\varepsilon|\log \varepsilon|, \varepsilon^{\frac{1}{5-q}}, \varepsilon^{\frac{1+\theta}{2}}, \varepsilon^{\frac{q-\sigma}{p^*-q}}\right\} & \text{if } N = 3. \end{cases}$$

Therefore if  $q$  satisfies (1.4), we find that there exists  $\tau \in (\frac{1}{2}, 1)$  such that

$$\|E\|_{**} \leq C\varepsilon^\tau.$$

Differentiating  $E$  with respect to  $\xi_i$  ( $i = 1, 2, \dots, k$ ), we have

$$\begin{aligned} \partial_{\xi_i} E &= \alpha_\varepsilon(p^* + \varepsilon)e^{\varepsilon x} V p^* + \varepsilon^{-1} \partial_{\xi_i} V - p^* \sum_{j=1}^k W(x - \xi_j)^{p^*-1} \partial_{\xi_i} W(x - \xi_j) \\ &\quad + \lambda \beta_N q e^{-(p^*-q)x} V q^{-1} \partial_{\xi_i} V. \end{aligned}$$

The proof of estimate for  $\|\partial_{\xi_i} E\|_{**}$  is similar to that of  $\|E\|_{**}$ .  $\square$

**Proposition 4.3.** *Assume that  $0 < \xi_1 < \xi_2 < \dots < \xi_k$  satisfy (2.11). Then there exists  $C > 0$  such that for  $\varepsilon > 0$  small enough, there exists a unique solution  $\phi = \phi(\xi)$  to problem (3.1) with*

$$\|\phi\|_* \leq C\varepsilon^\tau,$$

for some  $\tau \in (\frac{1}{2}, 1)$  satisfying Lemma 4.2. Moreover, the map  $\xi \mapsto \phi(\xi)$  is of class  $C^1$  for the  $\|\cdot\|_*$  norm, and

$$\|\partial_{\xi_i} \phi\|_* \leq C\varepsilon^\tau.$$

*Proof.* Problem (3.1) is equivalent to solve a fixed point problem

$$\phi = T_\varepsilon(N(\phi) + E) := A_\varepsilon(\phi).$$

We will show that the operator  $A_\varepsilon$  is a contraction map in a proper region. Set

$$\mathcal{F}_\gamma = \{\phi \in C(\mathbb{R}) : \|\phi\|_* \leq \gamma\varepsilon^\tau\},$$

where  $\gamma > 0$  will be chosen later.

For  $\phi \in \mathcal{F}_\gamma$ , by Lemmas 4.1 and 4.2, we get

$$\|A_\varepsilon(\phi)\|_* = \|T_\varepsilon(N(\phi) + E)\|_* \leq C\|N(\phi)\|_{**} + \|E\|_{**}$$

$$\leq C \left( \gamma^{\min\{p^*, 2\}} \varepsilon^{\min\{p^* - 1, 1\}\tau} + \gamma^{\min\{q, 2\}} \varepsilon^{\min\{q - 1, 1\}\tau} + 1 \right) \varepsilon^\tau.$$

Then we have  $A_\varepsilon(\phi) \in \mathcal{F}_\gamma$  for  $\phi \in \mathcal{F}_\gamma$  by choosing  $\gamma$  large enough but fixed.

Moreover, for  $\phi_1, \phi_2 \in \mathcal{F}_\gamma$ , by writing

$$N(\phi_1) - N(\phi_2) = \int_0^1 N'(\phi_2 + t(\phi_1 - \phi_2)) dt (\phi_1 - \phi_2).$$

By Proposition 3.1 and using (4.2), we find

$$\begin{aligned} & \|A_\varepsilon(\phi_1) - A_\varepsilon(\phi_2)\|_* \leq C \|N(\phi_1) - N(\phi_2)\|_{**} \\ & \leq C \left( \left( \max_{i=1,2} \|\phi_i\|_* \right)^{\min\{p^* - 1, 1\}} + \left( \max_{i=1,2} \|\phi_i\|_* \right)^{\min\{q - 1, 1\}} \right) \|\phi_1 - \phi_2\|_* \\ & \leq C \varepsilon^\kappa \|\phi_1 - \phi_2\|_* \end{aligned}$$

with some  $\kappa > 0$ . This implies that  $A_\varepsilon$  is a contraction map from  $\mathcal{F}_\gamma$  to  $\mathcal{F}_\gamma$ . Thus  $A_\varepsilon$  has a unique fixed point in  $\mathcal{F}_\gamma$ .

Now we consider the differentiability of  $\xi \mapsto \phi(\xi)$ . We write

$$B(\xi, \phi) := \phi - T_\varepsilon(N(\phi) + E).$$

First we observe that  $B(\xi, \phi) = 0$ . Moreover,

$$\partial_\phi B(\xi, \phi)[\theta] = \theta - T_\varepsilon(\theta(\partial_\phi(N(\phi)))) \equiv \theta + M(\theta),$$

where

$$M(\theta) = -T_\varepsilon(\theta(\partial_\phi(N(\phi)))).$$

By a direct calculation, we get

$$\|M(\theta)\|_* \leq C \|\theta(\partial_\phi(N(\phi)))\|_{**} \leq C \varepsilon^\kappa \|\theta\|_*.$$

So for  $\varepsilon > 0$  small enough, the operator  $\partial_\phi B(\xi, \phi)$  is invertible with uniformly bounded inverse in  $\|\cdot\|_*$ . It also depends continuously on its parameters. Let us differentiate with respect to  $\xi$ , we have

$$\partial_\xi B(\xi, \phi) = -(\partial_\xi T_\varepsilon)(N(\phi) + E) - T_\varepsilon((\partial_\xi N)(\xi, \phi) + \partial_\xi E),$$

where all these expressions depend continuously on their parameters. The implicit function theorem yields that  $\phi(\xi)$  is of class  $C^1$  and

$$\partial_\xi \phi = -(\partial_\phi B(\xi, \phi))^{-1} [\partial_\xi B(\xi, \phi)]$$

so that

$$\|\partial_\xi \phi\|_* \leq C (\|N(\phi)\|_{**} + \|E\|_{**} + \|(\partial_\xi N)(\xi, \phi)\|_{**} + \|\partial_\xi E\|_{**}) \leq C \varepsilon^\tau.$$

□

## 5. THE FINITE-DIMENSIONAL VARIATIONAL REDUCTION

According to the results of the previous section, our problem has been reduced to find points  $\xi = (\xi_1, \xi_2, \dots, \xi_k)$ , such that

$$(5.1) \quad c_j(\xi) = 0 \quad \text{for all } j = 1, \dots, k.$$

If (5.1) holds, then  $v = V + \phi$  is a solution to (1.7), and  $u = \sum_{j=1}^k U_{\mu_j} + \psi$  is the solution to problem (1.3) with  $\psi = \mathcal{T}^{-1}(\phi)$ .

Define the function  $\mathcal{I}_\varepsilon : (\mathbb{R}^+)^k \rightarrow \mathbb{R}$  as

$$\mathcal{I}_\varepsilon(\xi) := I_\varepsilon(V + \phi),$$

where  $V$  is defined by (2.12) and  $I_\varepsilon$  is the energy functional of (1.7) defined by

$$\begin{aligned} I_\varepsilon(v) &= \frac{1}{2} \int_{-\infty}^{+\infty} (|v'(x)|^2 + |v|^2) dx + \frac{1}{2} \left( \frac{2}{N-2} \right)^2 \int_{-\infty}^{+\infty} e^{-\frac{4}{N-2}x} v^2 dx \\ &\quad - \frac{1}{p^* + \varepsilon + 1} \alpha_\varepsilon \int_{-\infty}^{+\infty} e^{\varepsilon x} |v|^{p^* + \varepsilon + 1} dx - \frac{1}{q+1} \lambda \beta_N \int_{-\infty}^{+\infty} e^{-(p^* - q)x} |v|^{q+1} dx. \end{aligned}$$

We have the following fact.

**Lemma 5.1.** *The function  $V + \phi$  is a solution to (1.7) if and only if  $\xi = (\xi_1, \dots, \xi_k)$  is a critical point of  $\mathcal{I}_\varepsilon(\xi)$ , where  $\phi = \phi(\xi)$  is given by Proposition 4.3.*

*Proof.* For  $s \in \{1, 2, \dots, k\}$ , we have

$$\begin{aligned} \partial_{\xi_s} \mathcal{I}_\varepsilon(\xi) &= \partial_{\xi_s} (I_\varepsilon(V + \phi)) = DI_\varepsilon(V + \phi) [\partial_{\xi_s} V + \partial_{\xi_s} \phi] \\ &= \sum_{j=1}^k c_j \int_{\mathbb{R}} Z_j [\partial_{\xi_s} V + \partial_{\xi_s} \phi] = \sum_{j=1}^k c_j \left( \int_{\mathbb{R}} Z_j Z_s dx + o(1) \right), \end{aligned}$$

where  $o(1) \rightarrow 0$  as  $\varepsilon \rightarrow 0$  uniformly for the norm  $\|\cdot\|_*$ . This implies that the above relations define an almost diagonal homogeneous linear equation system for the  $c_j$ . Thus  $\xi$  is the critical point of  $I_\varepsilon$  if and only if  $c_j = 0$  for all  $j = 1, 2, \dots, k$ .  $\square$

**Lemma 5.2.** *The following expansion holds*

$$\mathcal{I}_\varepsilon(\xi) = I_\varepsilon(V) + o(\varepsilon)$$

as  $\varepsilon \rightarrow 0$ , where  $o(\varepsilon)$  is uniform in the  $C^1$ -sense on the vectors  $\xi$  satisfying (2.11).

*Proof.* By the fact that  $DI_\varepsilon(V + \phi)[\phi] = 0$  and using the Taylor expansion, we have

$$\begin{aligned} \mathcal{I}_\varepsilon(\xi) - I_\varepsilon(V) &= I_\varepsilon(V + \phi) - I_\varepsilon(V) = \int_0^1 D^2 I_\varepsilon(V + t\phi) [\phi^2] t dt \\ &= \int_0^1 t dt \int_{-\infty}^{+\infty} (N(\phi) + E) \phi dx \\ &\quad + (p^* + \varepsilon) \alpha_\varepsilon \int_0^1 t dt \int_{-\infty}^{+\infty} e^{\varepsilon x} [V^{p^* + \varepsilon - 1} - (V + t\phi)^{p^* + \varepsilon - 1}] \phi^2 dx \\ &\quad + \lambda \beta_N q \int_0^1 t dt \int_{-\infty}^{+\infty} e^{-(p^* - q)x} [V^{q-1} - (V + t\phi)^{q-1}] \phi^2 dx. \end{aligned}$$

Since  $\|\phi\|_* \leq C\varepsilon^\tau$  and  $\|E\|_{**} \leq C\varepsilon^\tau$  with  $\tau > \frac{1}{2}$ , we get

$$\mathcal{I}_\varepsilon(\xi) - I_\varepsilon(V) = O(\varepsilon^{2\tau}) = o(\varepsilon)$$

uniformly on the points  $\xi$  which satisfy (2.11).

Moreover, differentiating with respect to  $\xi_s$ , we have

$$\begin{aligned} \partial_{\xi_s} (\mathcal{I}_\varepsilon(\xi) - I_\varepsilon(V)) &= \int_0^1 \int_{-\infty}^{+\infty} \partial_{\xi_s} [(N(\phi) + E) \phi] t dx dt \\ &\quad + \alpha_\varepsilon (p^* + \varepsilon) \int_0^1 t dt \int_{-\infty}^{+\infty} e^{\varepsilon x} \partial_{\xi_s} \left( [V^{p^* + \varepsilon - 1} - (V + t\phi)^{p^* + \varepsilon - 1}] \phi^2 \right) dx \\ &\quad + \lambda \beta_N q \int_0^1 t dt \int_{-\infty}^{+\infty} e^{-(p^* - q)x} \partial_{\xi_s} \left( [V^{q-1} - (V + t\phi)^{q-1}] \phi^2 \right) dx. \end{aligned}$$

By the fact that  $\|\partial_\xi \phi\|_* \leq C\varepsilon^\tau$  and  $\|\partial_\xi E\|_{**} \leq C\varepsilon^\tau$  with  $\tau > \frac{1}{2}$ , we deduce that

$$\partial_{\xi_s} (\mathcal{I}_\varepsilon(\xi) - I_\varepsilon(V)) = O(\varepsilon^{2\tau}) = o(\varepsilon).$$

□

Now we consider the energy functional of problem (1.3), which is defined by

$$J(u) = \frac{1}{2} \int_{\mathbb{R}^N} (|\nabla u|^2 + u^2) - \frac{1}{p^* + 1 + \varepsilon} \int_{\mathbb{R}^N} |u|^{p^* + 1 + \varepsilon} - \frac{\lambda}{q + 1} \int_{\mathbb{R}^N} |u|^{q+1}.$$

By a direct calculation, we have that

$$(5.2) \quad I_\varepsilon(V) = \left( \frac{2}{N-1} \right)^{N-1} \frac{1}{\omega_{N-1}} J(U),$$

where  $V$  is defined by (2.12),  $\omega_{N-1}$  is the volume of the unit sphere in  $\mathbb{R}^N$  and  $U(z) = \sum_{j=1}^k U_{\mu_j}(z)$  with  $U_{\mu_j}$  satisfying problem (2.1).

We give the following expansion of  $J(U)$ , whose proof is in the Appendix.

**Lemma 5.3.** *Assume that (2.9) and (2.10) hold, then we have the following expansion:*

$$(5.3) \quad J(U) = a_1 + a_2\varepsilon - \varphi(\Lambda_1, \dots, \Lambda_k)\varepsilon + a_3\varepsilon \log \varepsilon + o(\varepsilon),$$

where

$$(5.4) \quad \varphi(\Lambda_1, \dots, \Lambda_k) = a_4 \Lambda_1^{\frac{N+2-(N-2)q}{2}} - a_5 \sum_{i=1}^k \log \Lambda_i + a_6 \sum_{l=1}^{k-1} \left( \frac{\Lambda_{l+1}}{\Lambda_l} \right)^{\frac{N-2}{2}},$$

and as  $\varepsilon \rightarrow 0$ ,  $o(\varepsilon)$  is uniform in the  $C^1$ -sense on the  $\Lambda_i$ 's satisfying (2.9), and

$$\begin{aligned} a_1 &= \frac{k}{N} \alpha_N^{p^*+1} \int_{\mathbb{R}^N} \frac{1}{(1+|z|^2)^N} dz, \\ a_2 &= \frac{k}{(p^*+1)^2} \alpha_N^{p^*+1} \int_{\mathbb{R}^N} \frac{1}{(1+|z|^2)^N} dz \\ &\quad - \frac{k}{p^*+1} \alpha_N^{p^*+1} \int_{\mathbb{R}^N} \frac{1}{(1+|z|^2)^N} \log \frac{\alpha_N}{(1+|z|^2)^{\frac{N-2}{2}}} dz, \\ a_3 &= \frac{(N-2)^2}{4N} \left( \alpha_N^{p^*+1} \int_{\mathbb{R}^N} \frac{1}{(1+|z|^2)^N} dz \right) \\ &\quad \times \sum_{i=1}^k \left( \frac{2(i-1)}{N-2} + \frac{2}{N+2-(N-2)q} \right), \\ a_4 &= \frac{\lambda}{q+1} \int_{\mathbb{R}^N} \frac{1}{(1+|z|^2)^{\frac{(N-2)(q+1)}{2}}} dz, \\ a_5 &= \frac{(N-2)^2}{4N} \left( \alpha_N^{p^*+1} \int_{\mathbb{R}^N} \frac{1}{(1+|z|^2)^N} dz \right), \\ a_6 &= \alpha_N^{p^*+1} \int_{\mathbb{R}^N} \frac{1}{(1+|z|^2)^{\frac{N+2}{2}}} \frac{1}{|z|^{N-2}} dz. \end{aligned}$$

Now we are ready to prove our main result.

**Proof of Theorem 1.1.** Thanks to Lemma 5.1, we know that

$$u = \sum_{j=1}^k U_{\mu_j} + \psi \quad \text{with } \psi = \mathcal{T}^{-1}(\phi)$$

is a solution to problem (1.3) if and only if  $\xi$  is a critical point of  $\mathcal{I}_\varepsilon(\xi)$ , where the existence of  $\phi$  is guaranteed by Proposition 4.3.

Finding a critical point of  $\mathcal{I}_\varepsilon(\xi)$  is equivalent to find that of  $\tilde{\mathcal{I}}_\varepsilon(\xi)$ , which is defined as

$$\tilde{\mathcal{I}}_\varepsilon(\xi) = - \left( \frac{N-1}{2} \right)^{N-1} \frac{\omega_{N-1}}{\varepsilon} \mathcal{I}_\varepsilon(\xi) + \frac{a_1}{\varepsilon} + a_2 + a_3 \log \varepsilon.$$

On the other hand, from Lemmas 5.2 and 5.3, using (5.2), we have

$$\begin{aligned} \mathcal{I}_\varepsilon(\xi) &= I_\varepsilon(V) + o(\varepsilon) = \left( \frac{2}{N-1} \right)^{N-1} \frac{1}{\omega_{N-1}} J(U) + o(\varepsilon) \\ &= \left( \frac{2}{N-1} \right)^{N-1} \frac{1}{\omega_{N-1}} [a_1 + a_2 \varepsilon - \varphi(\Lambda_1, \dots, \Lambda_k) \varepsilon + a_3 \varepsilon \log \varepsilon] + o(\varepsilon), \end{aligned}$$

as  $\varepsilon \rightarrow 0$ , where  $\varphi(\Lambda)$  is defined by (5.4) and  $o(\varepsilon)$  is uniform in the  $C^1$ -sense. Then we have

$$(5.5) \quad \tilde{\mathcal{I}}_\varepsilon(\xi) = \varphi(\Lambda) + o(1),$$

where  $o(1)$  is uniform in the  $C^1$ -sense as  $\varepsilon \rightarrow 0$ .

We set  $s_1 = \Lambda_1$ ,  $s_j = \frac{\Lambda_j}{\Lambda_{j-1}}$ , then we can write  $\varphi(\Lambda_1, \dots, \Lambda_k)$  as

$$\begin{aligned} \varphi(s_1, \dots, s_k) &= a_4 s_1^{\frac{N+2-(N-2)q}{2}} - a_5 k \log s_1 - \sum_{j=2}^k \left[ a_5 (k-j+1) \log s_j - a_6 s_j^{\frac{N-2}{2}} \right] \\ &:= \tilde{\varphi}_1 - \sum_{j=2}^k \tilde{\varphi}_j, \end{aligned}$$

with

$$\tilde{\varphi}_1 = a_4 s_1^{\frac{N+2-(N-2)q}{2}} - a_5 k \log s_1$$

and

$$\tilde{\varphi}_j = a_5 (k-j+1) \log s_j - a_6 s_j^{\frac{N-2}{2}}, \quad j = 2, \dots, k.$$

We note that

$$(5.6) \quad \bar{s}_1 = \left( \frac{2a_5 k}{a_4 (N+2-(N-2)q)} \right)^{\frac{2}{N+2-(N-2)q}}$$

is the critical point of  $\tilde{\varphi}_1$ , and

$$(5.7) \quad \bar{s}_j = \left( \frac{2a_5 (k-j+1)}{(N-2)a_6} \right)^{\frac{2}{N-2}}, \quad j = 2, \dots, k,$$

is the critical point of  $\tilde{\varphi}_j$ . Moreover

$$\tilde{\varphi}_1''(\bar{s}_1) < 0, \quad \tilde{\varphi}_j''(\bar{s}_j) < 0, \quad j = 2, \dots, k.$$

So  $(\bar{s}_1, \bar{s}_2, \dots, \bar{s}_k)$  is a nondegenerate critical point of  $\varphi(s_1, \dots, s_k)$ . Thus

$$\Lambda^* := (\bar{s}_1, \bar{s}_2 \bar{s}_1, \bar{s}_3 \bar{s}_2 \bar{s}_1, \dots, \bar{s}_k \times \dots \times \bar{s}_2 \bar{s}_1)$$

is a nondegenerate critical point of  $\varphi(\Lambda)$ . It follows that the local degree  $\deg(\nabla\varphi(\Lambda), \mathcal{O}, 0)$  is well defined and is nonzero, here  $\mathcal{O}$  is an arbitrarily small neighborhood of  $\Lambda^*$ . Hence from (5.5), for  $\varepsilon > 0$  small enough, we have that  $\deg(\nabla_\xi \tilde{\mathcal{I}}_\varepsilon(\xi), \bar{\mathcal{O}}, 0) \neq 0$ , where  $\bar{\mathcal{O}}$  is a small neighborhood of  $\xi^* = (\xi_1^*, \dots, \xi_k^*)$  and

$$\xi_j^* = \left[ (j-1) + \frac{1}{p^* - q} \right] \log \frac{1}{\varepsilon} - \frac{N-2}{2} \log(\bar{s}_j \bar{s}_{j-1} \cdots \bar{s}_1), \text{ for } \forall j = 1, \dots, k.$$

So  $\xi^*$  is a critical point of  $\tilde{\mathcal{I}}_\varepsilon(\xi)$ , which implies there is a critical point of  $\mathcal{I}_\varepsilon$ .

Furthermore, if for some  $i$ ,  $|x - \xi_i| \leq C_0$  with some  $C_0 > 0$ , then we have  $|\phi| = o(W(x - \xi_i))$ . Thus  $\psi(|z|) = \mathcal{T}^{-1}(\phi(x)) = o(w_{\mu_i})$  for  $\frac{1}{\mathcal{O}}\mu_i \leq |z| \leq C\mu_i$ . Moreover, from (c) of Lemma 2.1, we get that  $R_{\mu_i} = o(w_{\mu_i})$  for  $\frac{1}{\mathcal{O}}\mu_i \leq |z| \leq C\mu_i$ . Therefore we obtain (1.5) holds with

$$\Lambda_j^* = \bar{s}_j \bar{s}_{j-1} \cdots \bar{s}_1, \quad j = 1, \dots, k,$$

where  $\bar{s}_j$  are given by (5.6) and (5.7). This finishes the proof.  $\square$

## 6. APPENDIX

**6.1. Proof of Lemma 2.1.** In order to prove Lemma 2.1, we introduce the Green function. For a fixed  $z \in \mathbb{R}^N$ , let  $G(z, y)$  be the Green function of  $-\Delta + I$ , which satisfies

$$\begin{aligned} -\Delta G(z, y) + G(z, y) &= \delta_z(y) \quad \text{in } \mathbb{R}^N, \\ G(z, y) &\rightarrow 0 \quad |y| \rightarrow \infty. \end{aligned}$$

We have the following result.

**Lemma 6.1.** *We have*

$$|G(z, y)| \leq \frac{C}{|y - z|^{N-2}} \quad \text{for } 0 < |y - z| \leq 1,$$

and

$$|G(z, y)| \leq C|y - z|^{\frac{1-N}{2}} e^{-|y-z|} \quad \text{for } |y - z| \geq 1.$$

*Proof.* By radial symmetry, we can write  $G(z, y) = G(r)$  with  $r = |y - z|$ . Since  $G(r)$  is singular at zero and tends to zero at infinity, we can verify that  $G$  is given by

$$G(r) = \frac{N-2}{(2\pi)^{\frac{N}{2}} \Gamma(\frac{N}{2})^2} r^{\frac{2-N}{2}} K_{\frac{N-2}{2}}(r),$$

where  $K_{\frac{N-2}{2}}(r)$  is a Modified Bessel Function of the Second Kind, see [15]. For  $N = 3$ , the function  $G$  has the explicit form  $G(r) = \frac{e^{-r}}{4\pi r}$ . In general, we have that  $K_{\frac{N-2}{2}}(r) \sim \frac{\Gamma(\frac{N-2}{2})}{2} (\frac{2}{r})^{\frac{N-2}{2}}$  for  $r$  close to 0, and  $K_{\frac{N-2}{2}}(r) \sim \sqrt{\frac{\pi}{2r}} e^{-r}$  for  $r$  large. Using these estimates, we obtain the result.  $\square$

*Proof of Lemma 2.1.* (a) It is a direct consequence of the maximum principle.

(b) Define the barrier function  $Q(z) = \mu^{\frac{N-2}{2}} |z|^{-(N+2)}$ . It satisfies  $-\Delta Q(z) + Q(z) \geq c\mu^{\frac{N-2}{2}} |z|^{-(N+2)}$  for all  $|z| \geq R$  with  $R > 0$  a large constant, here  $c$  is positive constant. Since  $Q(z) = \mu^{\frac{N-2}{2}} R^{-(N+2)}$  for  $|z| = R$  and  $U_\mu(z) \leq w_\mu(z) \leq \alpha_N \mu^{\frac{N-2}{2}} |z|^{-(N-2)}$  for all  $|z| \geq 0$ . Set  $\varphi(z) = AQ(z) - U_\mu(z)$  for some constant  $A > 0$ , we then have  $-\Delta\varphi(z) + \varphi(z) \geq 0$  for  $|z| \geq R$ , and  $\varphi(z) \geq 0$  for  $|z| = R$  by

choosing suitable constant  $A$ . By the maximum principle we get  $U_\mu(z) \leq AQ(z) = A\mu^{\frac{N-2}{2}}|z|^{-(N+2)}$  for  $|z| \geq R$ .

(c) Using the representation

$$R_\mu(z) = \int_{\mathbb{R}^N} G(y-z)w_\mu(y)dy$$

and standard convolution estimates we can obtain the stated bounds for  $R_\mu$ .  $\square$

Set

$$\tilde{Z}_\mu(z) = \partial_\mu U_\mu(z), \quad \bar{Z}_\mu(z) = \partial_\mu w_\mu(z),$$

then  $\tilde{Z}_\mu(z)$  satisfies

$$\begin{cases} -\Delta \tilde{Z}_\mu + \tilde{Z}_\mu = \frac{N+2}{N-2} w_\mu^{\frac{4}{N-2}} \bar{Z}_\mu & \text{in } \mathbb{R}^N, \\ \tilde{Z}_\mu(z) \rightarrow 0 & \text{as } |z| \rightarrow \infty. \end{cases}$$

We can write

$$\tilde{Z}_\mu(z) = \bar{Z}_\mu(z) + \partial_\mu R_\mu(z),$$

then  $\partial_\mu R_\mu(z)$  satisfies

$$\begin{cases} -\Delta(\partial_\mu R_\mu(z)) + \partial_\mu R_\mu(z) = -\partial_\mu w_\mu(z) & \text{in } \mathbb{R}^N, \\ \partial_\mu R_\mu(z) \rightarrow 0 & \text{as } |z| \rightarrow \infty. \end{cases}$$

We observe that  $|\partial_\mu w_\mu(z)| \leq C\mu^{-1}w_\mu$ , then we have

**Corollary 6.2.** *One has*

$$(6.1) \quad |\partial_\mu R_\mu(z)| \leq C\mu^{-1}|R_\mu(z)| \quad \text{for } \forall z \in \mathbb{R}^N.$$

Moreover, by the maximum principle, we have that

$$(6.2) \quad |\tilde{Z}_\mu(z)| \leq C\mu^{\frac{N-4}{2}}|z|^{-(N+2)} \quad \text{for } |z| \geq R,$$

where  $R$  is a large positive number but fixed in Lemma 2.1.

## 6.2. Expansion of energy.

*Proof of Lemma 5.3.* The proof is very similar to the one in [20]. The difference is that we have more terms in the energy and the initial approximation is also somewhat different. We have

$$\begin{aligned} J(U) &= \left[ \frac{1}{2} \int_{\mathbb{R}^N} (|\nabla U|^2 + U^2) - \frac{1}{p^*+1} \int_{\mathbb{R}^N} U^{p^*+1} \right] \\ &\quad + \left[ \frac{1}{p^*+1} \int_{\mathbb{R}^N} U^{p^*+1} - \frac{1}{p^*+1+\varepsilon} \int_{\mathbb{R}^N} U^{p^*+1+\varepsilon} \right] - \frac{\lambda}{q+1} \int_{\mathbb{R}^N} U^{q+1} \\ (6.3) \quad &:= J_1 + J_2 + J_3, \end{aligned}$$

where  $U = \sum_{j=1}^k U_{\mu_j}$  with  $U_{\mu_j} = w_{\mu_j} + R_{\mu_j}$ .

As in [20] but using the estimates of  $R_\mu$  in Lemma 2.1 we can get

$$\begin{aligned} J_1 &= \frac{k}{N} \alpha_N^{p^*+1} \int_{\mathbb{R}^N} \frac{1}{(1+|z|^2)^N} dz \\ (6.4) \quad &- \varepsilon \sum_{l=1}^{k-1} \left( \frac{\Lambda_{l+1}}{\Lambda_l} \right)^{\frac{N-2}{2}} \alpha_N^{p^*+1} \int_{\mathbb{R}^N} \frac{1}{(1+|z|^2)^{\frac{N+2}{2}}} \frac{1}{|z|^{N-2}} dz + o(\varepsilon). \end{aligned}$$



Also as in [20] we obtain

$$\begin{aligned}
J_2 &= \varepsilon \frac{k}{(p^* + 1)^2} \alpha_N^{p^* + 1} \int_{\mathbb{R}^N} \frac{1}{(1 + |z|^2)^N} dz \\
&\quad - \varepsilon \frac{k}{p^* + 1} \alpha_N^{p^* + 1} \int_{\mathbb{R}^N} \frac{1}{(1 + |z|^2)^N} \log \frac{\alpha_N}{(1 + |z|^2)^{\frac{N-2}{2}}} dz \\
&\quad + \varepsilon \frac{(N-2)^2}{4N} \left( \alpha_N^{p^* + 1} \int_{\mathbb{R}^N} \frac{1}{(1 + |z|^2)^N} dz \right) \sum_{i=1}^k \log \Lambda_i \\
&\quad + \frac{(N-2)^2}{4N} \left( \alpha_N^{p^* + 1} \int_{\mathbb{R}^N} \frac{1}{(1 + |z|^2)^N} dz \right) \\
(6.5) \quad &\quad \times \sum_{i=1}^k \left( \frac{2(i-1)}{N-2} + \frac{2}{N+2-(N-2)q} \right) \varepsilon \log \varepsilon + o(\varepsilon).
\end{aligned}$$

We will do with detail the estimate of the term  $J_3$ .

Given  $\delta > 0$  small but fixed. Let  $\mu_1, \dots, \mu_k$  be given by (2.10), set  $\mu_0 = \frac{\delta^2}{\mu_1}$  and  $\mu_{k+1} = 0$ . Define the following annulus

$$A_i := B(0, \sqrt{\mu_i \mu_{i-1}}) \setminus B(0, \sqrt{\mu_i \mu_{i+1}}), \quad \text{for } i = 1, \dots, k.$$

We observe that  $B(0, \delta) = \bigcup_{i=1}^k A_i$ . On each  $A_i$ , the leading term in  $\sum_{j=1}^k U_{\mu_j}$  is  $U_{\mu_i}$ .

Then we have

$$\begin{aligned}
-(q+1)J_3 &= \lambda \sum_{l=1}^k \int_{A_l} \left[ \left( U_{\mu_l} + \sum_{j=1, j \neq l}^k U_{\mu_j} \right)^{q+1} - U_{\mu_l}^{q+1} - (q+1) U_{\mu_l}^q \sum_{j=1, j \neq l}^k U_{\mu_j} \right] \\
&\quad + \lambda \sum_{l=1}^k \int_{A_l} U_{\mu_l}^{q+1} + \lambda(q+1) \sum_{l=1}^k \int_{A_l} \sum_{j=1, j \neq l}^k U_{\mu_l}^q U_{\mu_j} + \lambda \int_{\mathbb{R}^N \setminus B(0, \delta)} \left( \sum_{j=1}^k U_{\mu_j} \right)^{q+1} \\
&:= J_{3,1} + J_{3,2} + J_{3,3} + J_{3,4}.
\end{aligned}$$

By the mean value theorem, for some  $t \in [0, 1]$ , we have

$$\begin{aligned}
J_{3,1} &= \lambda \frac{q(q+1)}{2} \sum_{l=1}^k \int_{A_l} \left( U_{\mu_l} + t \sum_{j=1, j \neq l}^k U_{\mu_j} \right)^{q-1} \left( \sum_{j=1, j \neq l}^k U_{\mu_j} \right)^2 \\
&\leq C\lambda \sum_{j,l=1, j \neq l}^k \int_{A_l} w_{\mu_l}^{q-1} w_{\mu_j}^2 + C\lambda \sum_{i,j,l=1, i, j \neq l}^k \int_{A_l} w_{\mu_i}^{q-1} w_{\mu_j}^2.
\end{aligned}$$

Since

$$\begin{aligned}
\sum_{j,l=1, j \neq l}^k \int_{A_l} w_{\mu_l}^{q-1} w_{\mu_j}^2 &= \sum_{j,l=1, j \neq l}^k \int_{A_l} (w_{\mu_l}^{q-1} w_{\mu_j}^{\frac{q-1}{q}}) w_{\mu_j}^{\frac{q+1}{q}} \\
(6.6) \quad &\leq \sum_{j,l=1, j \neq l}^k \left( \int_{A_l} w_{\mu_l}^q w_{\mu_j} \right)^{\frac{q-1}{q}} \left( \int_{A_l} w_{\mu_j}^{q+1} \right)^{\frac{1}{q}},
\end{aligned}$$

and

$$(6.7) \quad \sum_{i,j,l=1, i,j \neq l}^k \int_{A_l} w_{\mu_i}^{q-1} w_{\mu_j}^2 \leq \sum_{i,j,l=1, i,j \neq l}^k \left( \int_{A_l} w_{\mu_i}^{q+1} \right)^{\frac{q-1}{q+1}} \left( \int_{A_l} w_{\mu_j}^{q+1} \right)^{\frac{2}{q+1}}.$$

If  $j > l$ , then

$$\begin{aligned} \int_{A_l} w_{\mu_i}^q w_{\mu_j} dz &= \alpha_N^{q+1} \int_{\sqrt{\mu_l \mu_{l+1}} \leq |z| \leq \sqrt{\mu_l \mu_{l-1}}} \frac{\mu_l^{\frac{N-2}{2}q}}{(\mu_l^2 + |z|^2)^{\frac{N-2}{2}q}} \frac{\mu_j^{\frac{N-2}{2}}}{(\mu_j^2 + |z|^2)^{\frac{N-2}{2}}} dz \\ &= \left( \frac{\mu_j}{\mu_l} \right)^{\frac{N-2}{2}} \mu_l^{-\frac{N-2}{2}q + \frac{N+2}{2}} \left[ \alpha_N^{q+1} \int_{\mathbb{R}^N} \frac{1}{(1 + |z|^2)^{\frac{N-2}{2}q}} \frac{1}{|z|^{N-2}} dz + o(1) \right]. \end{aligned} \quad (6.8)$$

If  $j < l$ , then

$$\begin{aligned} \int_{A_l} w_{\mu_i}^q w_{\mu_j} dx &= \alpha_N^{q+1} \int_{\sqrt{\mu_l \mu_{l+1}} \leq |z| \leq \sqrt{\mu_l \mu_{l-1}}} \frac{\mu_l^{\frac{N-2}{2}q}}{(\mu_l^2 + |z|^2)^{\frac{N-2}{2}q}} \frac{\mu_j^{\frac{N-2}{2}}}{(\mu_j^2 + |z|^2)^{\frac{N-2}{2}}} dz \\ &= \left( \frac{\mu_l}{\mu_j} \right)^{\frac{N-2}{2}} \mu_l^{-\frac{N-2}{2}q + \frac{N+2}{2}} \alpha_N^{q+1} \int_{\sqrt{\frac{\mu_{l+1}}{\mu_l}} \leq |z| \leq \sqrt{\frac{\mu_{l-1}}{\mu_l}}} \frac{1}{(1 + |z|^2)^{\frac{N-2}{2}q}} \frac{1}{(1 + (\frac{\mu_l}{\mu_j})^2 |z|^2)^{\frac{N-2}{2}}} dz \\ &\leq \left( \frac{\mu_l}{\mu_j} \right)^{\frac{N-2}{2}} \mu_l^{-\frac{N-2}{2}q + \frac{N+2}{2}} \alpha_N^{q+1} \int_{\sqrt{\frac{\mu_{l+1}}{\mu_l}} \leq |z| \leq \sqrt{\frac{\mu_{l-1}}{\mu_l}}} \frac{1}{(1 + |z|^2)^{\frac{N-2}{2}q}} dz. \end{aligned} \quad (6.9)$$

For  $i \neq l$ , we have

$$(6.10) \quad \int_{A_l} w_{\mu_i}^{q+1} \leq C \mu_i^{-\frac{N-2}{2}q + \frac{N+2}{2}} \begin{cases} \left( \frac{\mu_l}{\mu_i} \right)^{\frac{N}{2}} & \text{if } i \leq l-1 < l; \\ \left( \frac{\mu_i^2}{\mu_l \mu_{l-1}} \right)^{\frac{N-2}{2}q-1} & \text{if } i \geq l+1 > l. \end{cases}$$

From (6.6)-(6.10), (1.4) and (2.10), we get  $J_{3,1} = o(\varepsilon)$ .

Moreover,

$$\begin{aligned} J_{3,2} &= \lambda \sum_{l=1}^k \int_{A_l} w_{\mu_l}^{q+1} + \lambda \sum_{l=1}^k \int_{A_l} (U_{\mu_l}^{q+1} - w_{\mu_l}^{q+1}) \\ &= \varepsilon \Lambda_1^{\frac{N+2-(N-2)q}{2}} \lambda \int_{\mathbb{R}^N} \frac{1}{(1 + |z|^2)^{\frac{(N-2)(q+1)}{2}}} dz + o(\varepsilon). \end{aligned}$$

From (6.8) and (6.9), we have

$$J_{3,3} \leq C \lambda \sum_{l=1}^k \int_{A_l} \sum_{j=1, j \neq l}^k U_{\mu_l}^q U_{\mu_j} \leq C \lambda \sum_{l=1}^k \int_{A_l} \sum_{j=1, j \neq l}^k w_{\mu_l}^q w_{\mu_j} = o(\varepsilon).$$

Finally,

$$J_{3,4} = \lambda \int_{\mathbb{R}^N \setminus B(0,\delta)} \left( \sum_{j=1}^k U_{\mu_j} \right)^{q+1} \leq C \sum_{j=1}^k \int_{\mathbb{R}^N \setminus B(0,\delta)} w_{\mu_j}^{q+1} dz = o(\varepsilon).$$

Thus we get

$$(6.11) \quad J_3 = -\varepsilon \Lambda_1^{\frac{N+2-(N-2)q}{2}} \frac{\lambda}{q+1} \int_{\mathbb{R}^N} \frac{1}{(1+|z|^2)^{\frac{(N-2)(q+1)}{2}}} dz + o(\varepsilon).$$

From (6.3), (6.4), (6.5) and (6.11), we obtain (5.3).  $\square$

**Acknowledgments.** W. Chen was supported by a doctoral grant of CONICYT Chile. J. Dávila was supported by Fondecyt 1130360 and Fondo Basal CMM. I. Guerra was supported by Fondecyt 1130790.

#### REFERENCES

- [1] C. O. Alves, D. C. de Moraes Filho & M. A. S. Souto, *Radially Symmetric Solutions for a Class of Critical Exponent Elliptic Problems in  $\mathbb{R}^N$* . Electron. J. Differential Equations 1996, No. 07.
- [2] H. Beresticki, P. L. Lions, *Nonlinear scalar field equations. I. Existence of a ground state*. Arch. Rational Mech. Anal. 82 (1983), no. 4, 313-345.
- [3] J. Campos, *Bubble-tower phenomena in a semilinear elliptic equation with mixed Sobolev growth*. Nonlinear Anal. 68 (2008), no. 5, 1382-1397.
- [4] C.V. Coffman, *On the positive solutions of boundary-value problems for a class of nonlinear differential equations*. J. Differential Equations 3 (1967), 92-111.
- [5] J. Dávila, M. Del Pino, I. Guerra, *Non-uniqueness of positive ground states of nonlinear schrödinger equations*. Proc. London Math. Soc. (3) 106 (2013) 318-344.
- [6] M. del Pino, J. Dolbeault, M. Musso, *"Bubble-tower" radial solutions in the slightly supercritical Brezis-Nirenberg problem*. J. Differential Equations 193 (2003), no. 2, 280-306.
- [7] M. del Pino, J. Dolbeault, M. Musso, *The Brezis-Nirenberg problem near criticality in dimension 3*. J. Math. Pures Appl. (9) 83 (2004), no. 12, 1405-1456.
- [8] M. del Pino, I. Guerra, *Ground states of a prescribed mean curvature equation*. J. Differential Equations 241 (2007), no. 1, 112-129.
- [9] M. del Pino, M. Musso, A. Pistoia, *Super-critical boundary bubbling in a semilinear Neumann problem*. Ann. Inst. H. Poincaré Anal. Non Linéaire 22 (2005), no. 1, 45-82.
- [10] P.L. Felmer, A. Quaas, M. Tang, J. Yu, *Monotonicity properties for ground states of the scalar field equation*. Ann. Inst. H. Poincaré Anal. Non Linéaire 25 (2008), no. 1, 105-119.
- [11] A. Ferrero, F. Gazzola, *On subcriticality assumptions for the existence of ground states of quasilinear elliptic equations*. Adv. Differential Equations 8 (2003), 1081-1106.
- [12] A. Floer, A. Weinstein, *Nonspreading wave packets for the cubic Schrödinger equation with bounded potential*. J. Funct. Anal. 69(3)(1986), 397-408.
- [13] Y. Ge, R. Jing, F. Pacard, *Bubble towers for supercritical semilinear elliptic equations*. J. Funct. Anal. 221 (2005), no. 2, 251-302.
- [14] Y. Ge, M. Musso, A. Pistoia, *Sign changing tower of bubbles for an elliptic problem at the critical exponent in pierced non-symmetric domains*, Comm. Partial Differential Equations 35 (2010), no. 8, 1419-1457.
- [15] B. Gidas, W.M. Ni, L. Nirenberg, *Symmetry of positive solutions of nonlinear elliptic equations in  $\mathbb{R}^N$* . Mathematical analysis and applications, Part A, Advances in Math. Suppl. Studies 7A, Academic Press, New York-London, 1981, 369-402.
- [16] M.K. Kwong, *Uniqueness of positive solutions of  $\Delta u - u + u^p = 0$  in  $\mathbf{R}^n$* . Arch. Rational Mech. Anal. 105 (1989), no. 3, 243-266.
- [17] K. McLeod, J. Serrin, *Uniqueness of positive radial solutions of  $\Delta u + f(u) = 0$  in  $\mathbf{R}^n$* . Arch. Rational Mech. Anal. 99 (1987), no. 2, 115-145.
- [18] A. M. Micheletti, M. Musso, A. Pistoia, *Super-position of spikes for a slightly super-critical elliptic equation in  $\mathbb{R}^N$* . Discrete Contin. Dyn. Syst. 12 (2005), no. 4, 747-760.

- [19] M. Musso, A. Pistoia, *Sign changing solutions to a nonlinear elliptic problem involving the critical Sobolev exponent in pierced domains*. J. Math. Pures Appl. (9) 86 (2006), no. 6, 510-528.
- [20] M. Musso, A. Pistoia, *Tower of bubbles for almost critical problems in general domains*. J. Math. Pures Appl. (9) 93 (2010), no. 1, 1-40.
- [21] W.-M. Ni, *Uniqueness of solutions of nonlinear Dirichlet problems*. J. Differential Equations 50 (1983), no. 2, 289-304.
- [22] W.-M. Ni, R.D. Nussbaum, *Uniqueness and nonuniqueness for positive radial solutions of  $\Delta u + f(u, r) = 0$* . Comm. Pure Appl. Math. 38 (1985), no. 1, 67-108.
- [23] L.A. Peletier, J. Serrin, *Uniqueness of positive solutions of semilinear equations in  $\mathbf{R}^n$* . Arch. Rational Mech. Anal. 81 (1983), no. 2, 181-197.
- [24] L.A. Peletier, J. Serrin, *Uniqueness of nonnegative solutions of semilinear equations in  $\mathbf{R}^n$* . J. Differential Equations 61 (1986), no. 3, 380-397.
- [25] A. Pistoia, T. Weth, *Sign changing bubble tower solutions in a slightly subcritical semilinear Dirichlet problem*. Ann. Inst. H. Poincaré Anal. Non Linéaire 24 (2007), no. 2, 325-340.
- [26] S. I. Pohožaev, *Eigenfunctions of the equation  $\Delta u + \lambda f(u) = 0$* , Sov. Math. Doklady 5(1965), 1408-1411.
- [27] O. Rey, *The role of the Greens function in a nonlinear elliptic equation involving the critical Sobolev exponent*, J. Funct. Anal. 89 (1990) 1-52.
- [28] T. Tao, M. Visan, X. Zhang, *The nonlinear Schrödinger equation with combined power-type nonlinearities*. Comm. Partial Differential Equations 32 (2007), no. 7-9, 1281-1343.
- [29] J. Wei, S. Yan, *Infinitely many solutions for the prescribed scalar curvature problem on  $S^N$* . J. Funct. Anal. 258 (2010), no. 9, 3048-3081.

W. CHEN—DEPARTAMENTO DE INGENIERÍA MATEMÁTICA AND CMM, UNIVERSIDAD DE CHILE,  
CASILLA 170 CORREO 3, SANTIAGO, CHILE.  
*E-mail address:* wjchen1102@gmail.com

J. DÁVILA—DEPARTAMENTO DE INGENIERÍA MATEMÁTICA AND CMM, UNIVERSIDAD DE CHILE,  
CASILLA 170 CORREO 3, SANTIAGO, CHILE.  
*E-mail address:* jdavila@dim.uchile.cl

I. GUERRA—DEPARTAMENTO DE MATEMÁTICA Y CIENCIA DE LA COMPUTACION, UNIVERSIDAD  
DE SANTIAGO DE CHILE, 9170125, SANTIAGO, CHILE.  
*E-mail address:* ignacio.guerra@usach.cl