



Citation for published version:

Cammarota, V & Morters, P 2012, 'On the most visited sites of planar Brownian motion', *Electronic Communications in Probability*, vol. 17, 15, pp. 1-9. <https://doi.org/10.1214/ECP.v17-1809>

DOI:

[10.1214/ECP.v17-1809](https://doi.org/10.1214/ECP.v17-1809)

Publication date:

2012

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

On the most visited sites of planar Brownian motion

VALENTINA CAMMAROTA* and PETER MÖRTERS†

Abstract: Let $(B_t: t \geq 0)$ be a planar Brownian motion and define gauge functions $\phi_\alpha(s) = \log(1/s)^{-\alpha}$ for $\alpha > 0$. If $\alpha < 1$ we show that almost surely there exists a point x in the plane such that $\mathcal{H}^{\phi_\alpha}(\{t \geq 0: B_t = x\}) > 0$, but if $\alpha > 1$ almost surely $\mathcal{H}^{\phi_\alpha}(\{t \geq 0: B_t = x\}) = 0$ simultaneously for all $x \in \mathbb{R}^2$. This resolves a longstanding open problem posed by S. J. Taylor in 1986.

MSc classification (2010): 60J65.

Keywords: Brownian motion, Hausdorff dimension, Hausdorff gauge, exact Hausdorff measure, local time, point of infinite multiplicity, random fractal, uniform dimension estimates.

1. INTRODUCTION AND STATEMENT OF MAIN RESULTS

Let $(B_t: t \geq 0)$ be a standard planar Brownian motion. Dvoretzky, Erdős and Kakutani (1958) first showed that, almost surely, there exist points x in the plane such that $\{t \geq 0: B_t = x\}$, the set of times where the Brownian path visits x , is uncountably infinite. Modern proofs of this fact are given in Le Gall (1987) and Mörters and Peres (2010). The result naturally raises the question: How large can the sets $\{t \geq 0: B_t = x\}$ be? Kaufman's famous dimension-doubling theorem implies that, almost surely, for all points x in the plane this set has Hausdorff dimension zero.

S. James Taylor in his influential survey paper 'The measure theory of random fractals' of 1986 raises the problem in terms of Hausdorff gauge functions. Letting $\phi: (0, \varepsilon) \rightarrow [0, 1]$ be a right continuous, increasing function with $\phi(0+) = 0$, we denote

$$\mathcal{H}^\phi(E) = \liminf_{\delta \downarrow 0} \left\{ \sum_{i=1}^{\infty} \phi(\text{diam}(E_i)): E \subseteq \bigcup_{i=1}^{\infty} E_i, 0 < \text{diam}(E_i) < \delta \right\},$$

the ϕ -Hausdorff measure of the set $E \subset \mathbb{R}$. Problem 5 in Taylor (1986) is the following question:

Which gauge functions ϕ are such that, almost surely, we have $\mathcal{H}^\phi(\{t \geq 0: B_t = x\}) = 0$ for all $x \in \mathbb{R}^2$?

In the paragraph following this question Taylor focuses on the gauge functions of the form $\phi_\alpha(s) = \log(1/s)^{-\alpha}$ for $\alpha > 0$, moving to a weaker but still challenging form of the problem. He notes that the results of Perkins and Taylor (1987) imply that these functions satisfy the above condition for $\alpha > 2$ and states that

It is not much more than a guess, but my hunch is that ϕ_α satisfies the condition for $\alpha > 1$, but not for $0 < \alpha < 1$.

*Department of Statistics, "Sapienza" University of Rome, P. le Aldo Moro, 5 – 00185 Rome, Italy.
E-mail: valentina.cammarota@uniroma1.it.

†Department of Mathematical Sciences, University of Bath, Bath BA2 7AY, England.
E-mail: maspm@bath.ac.uk.

The weaker form of the problem is reiterated in Perkins and Taylor (1987) as Problem 3.11. More than fifteen years later Xiao (2004) notes that the problem is still open even in the weaker form. The problem has recently been reiterated as Problem 5 in the open problems section of Mörters and Peres (2010). It is the aim of the present paper to solve the weaker form of the problem and to confirm Taylor's conjecture.

Although the solution involves both a new lower and upper bound, only the lower bound involves substantial work. It relies on the construction of an intersection local time for points of infinite multiplicity, which is due to Bass, Burdzy and Khoshnevisan (1994). Denoting by $\mathcal{B}(x, \varepsilon) \subset \mathbb{R}^2$ the open disc with centre x and radius ε and by \mathcal{B} the open disc $\mathcal{B}(0, 1)$ we let N_ε^x be the number of times the Brownian path $(B_t: t \geq 0)$ travels from the point $x \in \mathcal{B}$ to the circle $\partial\mathcal{B}(x, \varepsilon)$ before it leaves \mathcal{B} for the first time. Naturally for most points x we have $N_\varepsilon^x = 0$, but some points satisfy

$$\lim_{\varepsilon \downarrow 0} \frac{N_\varepsilon^x}{\log(1/\varepsilon)} = a, \quad (1.1)$$

for some $a > 0$. If (1.1) holds we say that Brownian motion 'spends a units of local time at x '. Note that the points x are by definition points of infinite multiplicity. Bass, Burdzy and Khoshnevisan (1994) in their Theorem 1.1 show that, for $0 < a < \frac{1}{2}$ there exists a (random) measure β_a on the plane such that (1.1) holds for β_a -almost every x . Moreover the measure is nonzero and has carrying Hausdorff dimension $2 - a$ almost surely.

Our main result shows that the points selected according to β_a also have a large inverse image under the Brownian motion, thus providing a lower bound for Taylor's Problem 5.

Theorem 1.1. *Let $0 < \alpha < \frac{1}{2}$ and take β_a to be the measure on points with local time a . Define the gauge function*

$$\varphi(\varepsilon) := \frac{\log \log \log(1/\varepsilon)}{\log(1/\varepsilon)}.$$

Then, almost surely,

$$\mathcal{H}^\varphi(\{t \geq 0: B_t = x\}) \geq a \sqrt{\frac{\pi}{2}} > 0,$$

for β_a -almost every x .

The proof of Theorem 1.1 is given in Section 2. To solve the weaker form of Taylor's Problem 5 we also need an upper bound confirming that the bound above has the right power of the logarithm.

Theorem 1.2. *For every gauge function ϕ with $\phi(\varepsilon) \log(1/\varepsilon) \rightarrow 0$, almost surely,*

$$\mathcal{H}^\phi(\{t \geq 0: B_t = x\}) = 0 \quad \text{for every } x \in \mathbb{R}^2.$$

The proof of Theorem 1.2, which uses a simple first moment estimate, is given in Section 3. Combining Theorems 1.1 and 1.2 confirms Taylor's conjecture on the weaker form of Problem 5 as stated in the abstract. The strong form of the problem remains open at this point, see Section 4 for a new conjecture and some further remarks on this.

2. PROOF OF THE LOWER BOUND

Let \mathbb{P}^z and \mathbb{E}^z be the distribution and the corresponding expectation of planar Brownian motion started at $z \in \mathbb{R}^2$, and let $\tau(A)$ be the first hitting time of a Borel set $A \subset \mathcal{B}$. In

particular we let $\tau = \tau(\partial\mathcal{B})$ be the first hitting time of the unit disc and denote by $p_t^{\mathcal{B}}(\cdot, \cdot)$ the transition (sub-)density for the Brownian motion killed at τ .

Let $B = (B_t: 0 \leq t \leq \tau)$ be the Brownian motion started at the origin and killed upon leaving \mathcal{B} . The idea of the proof of Theorem 1.1 is to use the representation of B as seen from a typical point chosen accordingly to β_a , which is given in Section 5 of Bass et al. (1994). The main ingredient of this representation is the construction of the process $(Z_a^x(t): 0 \leq t \leq \tau)$ which we now recall.

Fix some $x \in \mathcal{B}$ and let h be a strictly positive harmonic function in $\mathcal{B} \setminus \{x\}$ with zero boundary values on $\partial\mathcal{B}$ and a pole at x such that

$$\lim_{z \rightarrow x} \frac{h(z)}{|\log |z - x||} = 1. \quad (2.1)$$

The h -transform of B is a Markov process in $\mathcal{B} \setminus \{x\}$ with transition density p_t^h given by

$$p_t^h(z, y) = \frac{h(y)}{h(z)} p_t^{\mathcal{B}}(z, y). \quad (2.2)$$

The distribution of this process started in $y \in \mathcal{B} \setminus \{x\}$ is denoted \mathbb{P}_h^y . Let $C_*[0, \infty)$ be the set of all paths $e: [0, \infty) \rightarrow \mathcal{B} \cup \{\Delta\}$ which are continuous on some interval $[0, \sigma)$ and then jump to the isolated coffin state Δ . The canonical process on $C_*[0, \infty)$ will be denoted by X , i.e. $X_t(e) = e_t$ for all $e \in C_*[0, \infty)$ and $t \geq 0$. There is an, up to a constant factor unique, positive and σ -finite measure H^x on $C_*[0, \infty)$ such that

- $\lim_{t \downarrow 0} X_t = x$, H^x -almost surely,
- H^x is strong Markov for the transition densities $p_t^h(\cdot, \cdot)$,
- $H^x(\tau(A) < \infty) < \infty$, for any compact set $A \subset \mathcal{B} \setminus \{x\}$.

The excursion law H^x will be normalised so that

$$\lim_{\varepsilon \downarrow 0} \frac{H^x(\tau(\partial\mathcal{B}(x, \varepsilon)) < \infty)}{|\log \varepsilon|} = a, \quad (2.3)$$

see Lemma 5.1 in Bass et al. (1994) for existence of this normalisation, and Burdzy (1987) for background on excursion laws.

Let Leb denote the Lebesgue measure on $[0, \infty)$ and Y be a Poisson point process on $[0, \infty) \times C_*[0, \infty)$ with mean measure $\text{Leb} \otimes H^x$. Bass et al. (1994), Lemma 5.1, show that, for every $t < \infty$,

$$\sum_{\substack{(s, e_s) \in Y \\ s < t}} \sigma(e_s) < \infty, \quad (2.4)$$

almost surely, where $\sigma(e_s)$ is the lifetime of the excursion e_s .

The trajectories of Z_a^x are assembled from three parts,

$$Z_a^x(t) = \begin{cases} Z_1(t) & \text{if } 0 \leq t \leq t_1, \\ Z_2(t - t_1) & \text{if } t_1 \leq t \leq t_1 + t_2, \\ Z_3(t - t_1 - t_2) & \text{if } t_1 + t_2 \leq t \leq \tau = t_1 + t_2 + t_3. \end{cases}$$

(i) $(Z_1(t): 0 \leq t \leq t_1)$ is an h -process in $\mathcal{B} \setminus \{x\}$ which starts from 0 and is stopped when it approaches x at time t_1 .

(ii) For $u > 0$ let

$$T(u) = \sup \left\{ t: \sum_{\substack{(s, e_s) \in Y \\ s < t}} \sigma(e_s) \leq u \right\}.$$

By (2.4), $T(u)$ is well defined for all $u < \infty$ almost surely. Note that, almost surely, $T(u) < \infty$ for each u and, for almost all u , there is a point $(s, e_s) \in Y$ such that $s = T(u)$. For such u we define

$$Z_2(u) = e_{T(u)} \left(u - \sum_{\substack{(s, e_s) \in Y \\ s < T(u)}} \sigma(e_s) \right)$$

and for the remaining u we define $Z_2(u) = x$. Let $t_2 = \sum_{s < 1} \sigma(e_s)$ and observe that this defines a continuous process $(Z_2(t): 0 \leq t \leq t_2)$.

(iii) $(Z_3(t): 0 \leq t \leq t_3)$ is a Brownian motion starting from x and killed at the first exit from the unit ball \mathcal{B} .

We assume that Z_1 , Z_2 and Z_3 are independent. The distribution of the process $(Z_a^x(t): 0 \leq t \leq \tau)$ will be denoted by \mathbb{Q}_a^x .

The process Z_a^x may be thought of as a Brownian motion conditioned to spend a units of local time at x . It is possible to interpret \mathbb{Q}_a^x as the distribution of B conditioned by the event that x is in the support of β_a . This follows from the ‘Palm measure’ decomposition of β_a stated below and proved in Bass et al. (1994), Theorem 5.2.

Lemma 2.1. *For every $a \in (0, \frac{1}{2})$ and every nonnegative measurable function f on $\mathbb{R}^2 \times C_*[0, \infty)$ we have*

$$\mathbb{E}^0 \int f(y, B) \beta_a(dy) = \int_{\mathcal{B}} \int \mathbf{1}_{f(y, B)} |1 - y^2|^a |\log |y|| \mathbb{Q}_a^y(dB) dy.$$

Let L_a^x be the right-continuous generalised inverse of the sum of the excursion lengths of Z_a^x from x ,

$$L_a^x(t) := \sup \left\{ \theta: \sum_{\substack{(s, e_s) \in Y \\ s < \theta}} \sigma(e_s) \leq t \right\}.$$

L_a^x is the local time (in the sense of excursion theory) of Z_a^x at x . The following proposition is the main ingredient of the proof of the lower bound.

Proposition 2.2. *For every $a \in (0, \frac{1}{2})$ and $x \in \mathbb{R}^2$, we have*

$$\limsup_{\varepsilon \downarrow 0} \frac{L_a^x(t_1 + \varepsilon)}{\varphi(\varepsilon)} = a^{-1} \sqrt{\frac{2}{\pi}} < \infty, \quad \mathbb{Q}_a^x\text{-almost surely}, \quad (2.5)$$

where $\varphi(\varepsilon) := \frac{\log \log \log(1/\varepsilon)}{\log(1/\varepsilon)}$.

Before proving Proposition 2.2, we show how this implies Theorem 1.1.

First, the natural link between the Hausdorff measure $\mathcal{H}^\varphi(\{t \geq 0: B_t = x\})$ and the law of the iterated logarithm in (2.5) is the Rogers-Taylor theorem stated below, see Mörters and Peres (2010), Proposition 6.44, for a proof.

Lemma 2.3. *Let μ be a Borel measure on \mathbb{R} and let ϕ be a Hausdorff gauge function. If $\Lambda \subset \mathbb{R}$ is a closed set and*

$$A := \left\{ t \in \Lambda : \limsup_{\varepsilon \downarrow 0} \frac{\mu[t, t + \varepsilon]}{\phi(\varepsilon)} < \alpha \right\},$$

then $\mathcal{H}^\phi(A) \geq \frac{\mu(A)}{\alpha}$.

In our case we consider the closed set $\Lambda_x := \{t \geq 0: Z_a^x(t) = x\}$, the (probability) measure ℓ_a^x given by $\ell_a^x[t, t + \varepsilon] := L_a^x(t + \varepsilon) - L_a^x(t)$, the gauge function φ and the sets

$$A_\alpha := \left\{ t \in \Lambda_x : \limsup_{\varepsilon \downarrow 0} \frac{\ell_a^x[t, t + \varepsilon]}{\varphi(\varepsilon)} < \alpha \right\}.$$

Then, by Lemma 2.3, we have $\mathcal{H}^\varphi(\Lambda_x) \geq \mathcal{H}^\varphi(A_\alpha) \geq \frac{1}{\alpha} \ell_a^x(A_\alpha)$, and we now show that, \mathbb{Q}_a^x -almost surely, $\ell_a^x(A_\alpha) = 1$ for suitable $\alpha > 0$.

For $0 \leq t \leq 1$ let $T_t := \inf\{s > 0: L_a^x(s) \geq t\} < \infty$. By construction the process $(Z_a^x(T_t + s): 0 \leq s \leq T_1 - T_t)$ has the same law as $(Z_a^x(t_1 + s): 0 \leq s \leq T_{1-t})$. Hence it follows from Proposition 2.2 that, for all $t \in (0, 1)$,

$$\limsup_{\varepsilon \downarrow 0} \frac{\ell_a^x[T_t, T_t + \varepsilon]}{\varphi(\varepsilon)} = a^{-1} \sqrt{\frac{2}{\pi}}, \quad \mathbb{Q}_a^x\text{-almost surely.}$$

By applying Fubini's theorem we get

$$\text{Leb} \left\{ t \in (0, 1): \limsup_{\varepsilon \downarrow 0} \frac{\ell_a^x[T_t, T_t + \varepsilon]}{\varphi(\varepsilon)} = a^{-1} \sqrt{\frac{2}{\pi}} \right\} = 1, \quad \mathbb{Q}_a^x\text{-almost surely,}$$

that is

$$\ell_a^x \left\{ s \in \Lambda_x : \limsup_{\varepsilon \downarrow 0} \frac{\ell_a^x[s, s + \varepsilon]}{\varphi(\varepsilon)} = a^{-1} \sqrt{\frac{2}{\pi}} \right\} = 1, \quad \mathbb{Q}_a^x\text{-almost surely.}$$

Hence $\ell_a^x(A_\alpha) = 1$ for all $\alpha > a^{-1} \sqrt{2/\pi}$, and so $\mathcal{H}^\varphi(\Lambda_x) \geq a \sqrt{\pi/2} > 0$, \mathbb{Q}_a^x -almost surely.

To complete the proof of the lower bound we apply Lemma 2.1 to the nonnegative, measurable function $f(x, B) = \mathbf{1}_{\{(x, B): \mathcal{H}^\varphi(\Lambda_x) < a \sqrt{\pi/2}\}}$. We have seen that

$$\int_{\mathcal{B}} \int f(x, B) |1 - x^2|^a |\log |x|| \mathbb{Q}_a^x(dB) dx = 0,$$

and this implies that

$$\mathbb{E}^0 \int f(x, B) \beta_a(dx) = 0,$$

that is $\mathcal{H}^\varphi(\Lambda_x) \geq a \sqrt{\pi/2}$ for β_a -almost every x , \mathbb{P}^0 -almost surely, as required to prove Theorem 1.1 subject to the proof of Proposition 2.2.

We now prove Proposition 2.2. Bertoin and Caballero (1995), making use of the work of Fristedt and Pruitt (1971), have shown the following result, see their Theorem 1.

Lemma 2.4. *Let ξ be a subordinator, Φ its Laplace exponent and S its right-continuous generalised inverse. If Φ is slowly varying at infinity and $\Phi(\infty) = \infty$, then*

$$\limsup_{t \downarrow 0} \frac{S(t)\Phi(t^{-1} \log |\log \Phi(1/t)|)}{\log |\log \Phi(1/t)|} = 1, \quad \text{almost surely.}$$

In our case $\xi_t = \sum_{s < t} \sigma(e_s)$ and for the Laplace exponent Φ of the right-continuous generalised inverse of the local time we have

$$\begin{aligned} \exp\{-t\Phi(\lambda)\} &= \mathbb{E}\left\{\exp\left\{-\lambda \sum_{\substack{(s, e_s) \in Y \\ s < t}} \sigma(e_s)\right\}\right\} \\ &= \exp\left\{-t \int dH^x(e)(1 - e^{-\lambda\sigma(e)})\right\}, \end{aligned}$$

where we used (3.29) in Kingman (1993). Since

$$\Phi(\lambda) = \int dH^x(e)(1 - e^{-\lambda\sigma(e)}) = \int_0^1 dt H^x(\sigma(e) > \frac{-\log t}{\lambda}), \quad (2.6)$$

to get the asymptotics of $\Phi(\lambda)$ as $\lambda \uparrow \infty$ we first look at $H^x(\sigma(e) > \theta)$ as $\theta \downarrow 0$.

Lemma 2.5. *We have*

$$H^x(\sigma(e) > \theta) = a \int_{\mathcal{B} \setminus \{x\}} h(\xi) p_\theta^{\mathcal{B}}(x, \xi) d\xi.$$

Proof. We observe that

$$\begin{aligned} H^x(\sigma(e) > \theta) &= \int_{\mathcal{B} \setminus \{x\}} H^x(X_\theta \in d\xi) \\ &= \int_{\mathcal{B} \setminus \{x\}} \lim_{\epsilon \downarrow 0} H^x(X_{\zeta(\epsilon) + \theta} \in d\xi; \zeta(\epsilon) + \theta < \sigma), \end{aligned}$$

where $\zeta(\epsilon)$ is the first hitting time of $\partial\mathcal{B}(x, \epsilon)$. By the strong Markov property of the excursion measure,

$$\begin{aligned} H^x(\sigma(e) > \theta) &= \int_{\mathcal{B} \setminus \{x\}} \lim_{\epsilon \downarrow 0} H^x(\mathbb{P}_h^{X_{\zeta(\epsilon)}}(X_\theta \in d\xi; \theta < \sigma); \zeta(\epsilon) < \sigma) \\ &\leq \int_{\mathcal{B} \setminus \{x\}} \lim_{\epsilon \downarrow 0} \sup_{z \in \partial\mathcal{B}(x, \epsilon)} \mathbb{P}_h^z(X_\theta \in d\xi) H^x(\zeta(\epsilon) < \sigma) \\ &= \int_{\mathcal{B} \setminus \{x\}} \lim_{\epsilon \downarrow 0} \sup_{z \in \partial\mathcal{B}(x, \epsilon)} \frac{h(\xi)}{h(z)} p_\theta^{\mathcal{B}}(z, \xi) H^x(\zeta(\epsilon) < \sigma) d\xi, \end{aligned}$$

and in view of (2.1) and (2.3), we have

$$\begin{aligned} H^x(\sigma(e) > \theta) &\leq \int_{\mathcal{B} \setminus \{x\}} h(\xi) \lim_{\epsilon \downarrow 0} \sup_{z \in \partial\mathcal{B}(x, \epsilon)} p_\theta^{\mathcal{B}}(z, \xi) \frac{H^x(\zeta(\epsilon) < \sigma)}{|\log \epsilon|} d\xi \\ &= a \int_{\mathcal{B} \setminus \{x\}} h(\xi) p_\theta^{\mathcal{B}}(x, \xi) d\xi. \end{aligned}$$

Analogously, we get

$$\begin{aligned} H^x(\sigma(e) > \theta) &\geq \int_{\mathcal{B} \setminus \{x\}} \lim_{\epsilon \downarrow 0} \inf_{z \in \partial\mathcal{B}(x, \epsilon)} \mathbb{P}_h^z(X_\theta \in d\xi) H^x(\zeta(\epsilon) < \sigma) \\ &= a \int_{\mathcal{B} \setminus \{x\}} h(\xi) p_\theta^{\mathcal{B}}(x, \xi) d\xi. \end{aligned} \quad \square$$

The following lemma, see Port and Stone (1978), Proposition 4.1, will be used to replace $p_\theta^{\mathcal{B}}$ by the Brownian transition kernel in the main part of the integral.

Lemma 2.6. *Let U be an open set and $x \in U$. Then there is a $\delta_0 > 0$ such that $\lim_{t \downarrow 0} p_t^U(\zeta, \xi)/p_t(\zeta, \xi) = 1$ uniformly for all $\zeta, \xi \in \mathcal{B}(x, \delta_0)$.*

We obtain the following asymptotics for $H^x(\sigma(e) > \theta)$ as $\theta \downarrow 0$.

Lemma 2.7.

$$H^x(\sigma(e) > \theta) \sim a \sqrt{\frac{\pi}{2}} \log(1/\theta) \text{ as } \theta \downarrow 0.$$

Proof. For any small $\delta > 0$ we have

$$H^x(\sigma(e) > \theta) = a \int_{\mathcal{B} \setminus \{x\}} h(\xi) p_\theta^{\mathcal{B}}(x, \xi) d\xi \leq a \int_{\mathcal{B}(x, \delta) \setminus \{x\}} h(\xi) p_\theta^{\mathcal{B}}(x, \xi) d\xi + \text{const},$$

since the integral over the complement of $\mathcal{B}(x, \delta)$ is bounded. In view of (2.1) and Lemma 2.6, for every $\varepsilon > 0$, we find a small $\delta > 0$ and $d = d(\delta) > 0$ such that, for $\theta < d$,

$$\begin{aligned} H^x(\sigma(e) > \theta) &\leq (1 + \varepsilon) a \int_{\mathcal{B}(x, \delta) \setminus \{x\}} |\log|x - \xi|| p_\theta(x, \xi) d\xi + \text{const} \\ &= (1 + \varepsilon) a \sqrt{2\pi} \frac{1}{\theta} \int_0^\delta (-\log r) \exp\{-r^2/2\theta\} r dr + \text{const}, \end{aligned}$$

and, changing variables $s^2 = r^2/\theta$, we get

$$\begin{aligned} H^x(\sigma(e) > \theta) &\leq (1 + \varepsilon) a \sqrt{2\pi} \left[\int_0^{\delta/\sqrt{\theta}} (-\log s) \exp\{-s^2/2\} s ds \right. \\ &\quad \left. - \log \sqrt{\theta} \int_0^{\delta/\sqrt{\theta}} \exp\{-s^2/2\} s ds \right] + \text{const} \\ &\leq -(1 + \varepsilon) a \sqrt{2\pi} \log \sqrt{\theta} + \text{const}. \end{aligned}$$

Similarly, for the lower bound we have

$$H^x(\sigma(e) > \theta) \geq -(1 - \varepsilon) a \sqrt{2\pi} (1 - e^{-\frac{\delta^2}{2\theta}}) \log \sqrt{\theta} + \text{const},$$

which completes the proof. \square

We now derive the asymptotics of $\Phi(\lambda)$ as $\lambda \uparrow \infty$ from (2.6). For every fixed $d > 0$ we have

$$\int_{e^{-\lambda d}}^1 dt H^x(\sigma(e) > \frac{-\log t}{\lambda}) \leq \Phi(\lambda) \leq \text{const} + \int_{e^{-\lambda d}}^1 dt H^x(\sigma(e) > \frac{-\log t}{\lambda}),$$

where the constant depends on d but not on θ . Hence, for every $\varepsilon > 0$ there exists a small $d > 0$ such that

$$-(1 - \varepsilon) a \sqrt{\frac{\pi}{2}} \int_{e^{-\lambda d}}^1 \log\left(-\frac{\log t}{\lambda}\right) dt \leq \Phi(\lambda) \leq \text{const} - (1 + \varepsilon) a \sqrt{\frac{\pi}{2}} \int_{e^{-\lambda d}}^1 \log\left(-\frac{\log t}{\lambda}\right) dt,$$

and, as $\log \log(1/t)$ is integrable over $(0, 1)$, we get

$$\Phi(\lambda) \sim a \sqrt{\frac{\pi}{2}} \log \lambda \text{ as } \lambda \uparrow \infty,$$

which implies that Φ is slowly varying at infinity, and satisfies $\Phi(\infty) = \infty$.

We can now use Lemma 2.4 and get (2.5) for the gauge function

$$\varphi(\varepsilon) = a \sqrt{\frac{\pi}{2}} \frac{\log |\log \Phi(1/\varepsilon)|}{\Phi(\varepsilon^{-1} \log |\log \Phi(1/\varepsilon)|)} \sim \frac{\log \log \log(1/\varepsilon)}{\log(1/\varepsilon)}.$$

3. PROOF OF THE UPPER BOUND

Fix a ball $\mathcal{B}(0, R)$, $R > 1$, and stop Brownian motion at the first exit time τ from this ball. Given a cube Q of side length r inside this ball, we define recursively

$$\begin{aligned} \tau_1^Q &= \inf \{t \geq 0 : B_t \in Q\}, \\ \tau_{k+1}^Q &= \inf \{t \geq \tau_k^Q + r : B_t \in Q\}, \end{aligned}$$

with the usual convention that $\inf \emptyset = \infty$.

Lemma 3.1. *There exists $0 < \theta < 1$ such that for any $z \in \mathcal{B}(0, R)$ and $0 < r < \frac{1}{2}$,*

$$\mathbb{P}^z(\tau_k^Q < \tau) \leq \theta^k.$$

Proof. It suffices to bound $\mathbb{P}^z(\tau_{k+1}^Q \geq \tau \mid \tau_k^Q < \tau)$ from below by

$$\mathbb{P}^z(\tau_{k+1}^Q \geq \tau \mid |B_{\tau_k^Q + r} - x| > 2\sqrt{r}, \tau_k^Q < \tau) \mathbb{P}^z(|B_{\tau_k^Q + r} - x| > 2\sqrt{r} \mid \tau_k^Q < \tau),$$

where x is the centre of Q . The second factor can be bounded from below by a constant not depending on r . The first factor is bounded from below by the probability that planar Brownian motion started at any point in $\partial\mathcal{B}(0, 2\sqrt{r})$ hits $\partial\mathcal{B}(0, 2R)$ before $\partial\mathcal{B}(0, r)$. This probability is given by

$$\frac{\log 2\sqrt{r} - \log r}{\log 2R - \log r} \geq \frac{-\frac{1}{2} \log r}{-\log r + \log 2R},$$

which is bounded from zero by a positive constant independent of r . \square

Lemma 3.2. *Let \mathfrak{C}_m be the set of dyadic cubes of side length 2^{-m} inside a fixed unit cube $U \subset \mathcal{B}(0, R)$. Almost surely there exists a (random) integer C so that for all $m \geq 1$ and cubes $Q \in \mathfrak{C}_m$ and $r = 2^{-m}$ we have $\tau_{mC}^Q > \tau$.*

Proof. From Lemma 3.1 we get that, for any positive integer c ,

$$\sum_{m=1}^{\infty} \sum_{Q \in \mathfrak{C}_m} \mathbb{P}(\tau_{cm}^Q < \tau) \leq \sum_{m=1}^{\infty} 2^{2m} \theta^{cm}.$$

Now choose c so large that $4\theta^c < 1$. Then, by the Borel-Cantelli lemma, for all but finitely many m we have $\tau_{cm}^Q \geq \tau$ for all $Q \in \mathfrak{C}_m$. Finally, we can choose a random $C > c$ to handle the finitely many exceptional cubes. \square

To complete the proof we note that, on the event in the lemma, for a given m we can cover any set $\{0 < t < \tau : B_t \in Q\}$, $Q \in \mathfrak{C}_m$, with no more than Cm intervals of length $r = 2^{-m}$. This implies that, for any $z \in U$,

$$\mathcal{H}^\phi(\{0 < t < \tau : B_t = z\}) \leq \lim_{m \rightarrow \infty} Cm\phi(2^{-m}) = 0,$$

under the assumption on ϕ . Theorem 1.2 follows as $U \subset \mathcal{B}(0, R)$ and $R > 1$ were arbitrary.

4. OUTLOOK

As mentioned in the introduction, the strong form of Taylor's Problem 5 remains unresolved in this paper. Our work however allows us to make a conjecture. We believe that our lower bound is sharp, or in other words that for every gauge function ϕ with $\phi(\varepsilon)/\varphi(\varepsilon) \rightarrow 0$ where φ is as in Theorem 1.2, almost surely,

$$\mathcal{H}^\phi(\{t \geq 0: B_t = x\}) = 0 \quad \text{for every } x \in \mathbb{R}^2.$$

The reason for this belief is that our upper bound uses coverings of the level sets by intervals of fixed size. If we were able to adapt the size of the covering intervals to the fluctuations of the local time, one would expect a gain of order $\log \log \log(1/\varepsilon)$, similar to our lower bound. However, the technical difficulties in carrying out such an estimate appear to be considerable and we therefore defer verification of our conjecture to future work.

BIBLIOGRAPHY

- [BBK94] BASS, R.F., K. BURDZY and D. KHOSHNEVISAN Intersection local times for points of infinite multiplicity. *Ann. Probab.* 22:566–625 (1994).
- [BC95] BERTOIN, J. and M.E. CABALLERO On the rate of growth of subordinators with slowly varying Laplace exponent. In: *Séminaire de Probabilités XXIX*, Springer Lecture Notes in Mathematics Vol 1613, pp 125–132 (1995).
- [B87] BURDZY, K. *Multidimensional Brownian excursions and potential theory*. Longman, New York, 1987.
- [DEK58] DVORETZKY, A., P. ERDŐS and S. KAKUTANI Points of multiplicity \mathfrak{c} of planar Brownian motion. *Bull. Res. Council Israel* 7 (1958) 157–180.
- [FP71] FRISTEDT, B.E. and W.E. PRUITT Lower functions for increasing random walks and subordinators. *Z. Wahrscheinlichkeitstheorie* 18 (1971) 167–182.
- [K93] KINGMAN, J.F.C. *Poisson Processes*. Oxford University Press, Oxford, 1993.
- [LG87] LE GALL, J.-F. Le comportement du mouvement brownien entre les deux instants où il passe par un point double. *J. Funct. Anal.* 71 (1987) 246–262.
- [MP10] MÖRTERS, P. and Y. PERES *Brownian motion*. Cambridge University Press, Cambridge, 2010.
- [PS78] PORT, S. C. and C. J. STONE *Brownian motion and classical potential theory*. Academic Press, New York, 1978.
- [Ta86] TAYLOR, S.J. The measure theory of random fractals. *Math. Proc. Cambridge Phil. Soc.* 100 (1986) 383–406.
- [PT87] TAYLOR, S.J. and E. A. PERKINS Uniform measure results for the image of subsets under Brownian motion. *Probab. Theory Rel. Fields* 76 (1987) 257–289.
- [Xi04] XIAO, Y. Random fractals and Markov processes. In: *Fractal Geometry and Applications: A Jubilee of Benoît Mandelbrot*. Proc. Symposia Pure Math. 72, 261–338, American Mathematical Society, Providence, 2004.