



Citation for published version:

Zatoński, W & Zatoński, M 2015, 'Cytisine versus nicotine for smoking cessation', *New England Journal of Medicine*, vol. 372, no. 11, 1072. <https://doi.org/10.1056/NEJMc1500342>

DOI:

[10.1056/NEJMc1500342](https://doi.org/10.1056/NEJMc1500342)

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link to publication](#)

(C) The New England Journal of Medicine, 2015.

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

2. Jaiswal S, Fontanillas P, Flannick J, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med* 2014;371:2488-98.

3. Lawrence MS, Stojanov P, Mermel CH, et al. Discovery and saturation analysis of cancer genes across 21 tumour types. *Nature* 2014;505:495-501.

4. Shlush LI, Zandi S, Mitchell A, et al. Identification of pre-leukaemic haematopoietic stem cells in acute leukaemia. *Nature* 2014;506:328-33.

DOI: 10.1056/NEJMc1500684

Cytisine versus Nicotine for Smoking Cessation

TO THE EDITOR: Walker et al. (Dec. 18 issue)¹ describe the use of cytisine, as compared with nicotine, for smoking cessation. Although cytisine is perceived as a novelty in the West, it has been used in daily medical practice in Poland for more than 50 years.² In 1976, the first English-language study on the use of cytisine in Poland was published, based on the observations of 1968 patients.³ Cytisine is currently undergoing a renaissance in Poland. Two Polish studies — a cohort study conducted in 2003 through 2005⁴ and a randomized, controlled trial conducted in 2006 through 2010⁵ — showed the safety and effec-

tiveness of the drug. In 2012, a Polish generic drug company broke the monopoly by introducing a new cytisine product, Desmoxan. Cytisine is very inexpensive in Poland, costing under \$20 for full treatment. Its consumption has grown by factor of 10 since 2000 (Fig. 1). In 2013 alone, after cytisine became available over the counter and was heavily advertised, sales of the drug doubled. The use of cytisine is widespread, and it is considered to be safe by medical professionals and consumers. It seems that the use of cytisine is a major factor behind the decline in smoking in Poland — in the past 2 years, 500,000 smokers have quit.⁶ It is certainly desirable for this success to be replicated elsewhere.




Figure 1. Total Units of Cytisine Purchased as Smoking-Cessation Therapy in Poland, 2000–2013.

Each unit (or package) of cytisine is one full treatment for smoking cessation. The solid black line indicates the trend in sales between 2000 and 2013.

Witold Zatoński, M.D., Sc.D.

Maria Skłodowska-Curie Memorial Cancer Center
Warsaw, Poland
wazatonski@gmail.com

Mateusz Zatoński, M.A., M.Sc.

London School of Hygiene and Tropical Medicine
London, United Kingdom.

No potential conflict of interest relevant to this letter was reported.

- Walker N, Howe C, Glover M, et al. Cytisine versus nicotine for smoking cessation. *N Engl J Med* 2014;371:2353-62.
- Tutka P, Zatoński W. Cytisine for the treatment of nicotine addiction: from a molecule to therapeutic efficacy. *Pharmacol Rep* 2006;58:777-98.
- Granatowicz J. Smoking cessation through the use of cytisine and other chemotherapy. *World Smoking Health* 1976;1:8-11.
- Zatoński W, Cedzynska M, Tutka P, West R. An uncontrolled trial of cytisine (Tabex) for smoking cessation. *Tob Control* 2006;15:481-4.
- West R, Zatoński W, Cedzynska M, et al. Placebo-controlled trial of cytisine for smoking cessation. *N Engl J Med* 2011;365:1193-200.
- Zatonski W, Sulkowska U, Przewozniak K, Zatonski M. Malignant cancer epidemiology in Poland. In: Potrykowska A, Strzelecki Z, Szymborski J, Witkowski J, eds. *Cancer incidence and mortality versus the demographic situation in Poland*. Warsaw, Poland: Governmental Population Council, 2014:30-49. (In Polish.)

DOI: 10.1056/NEJMc1500342