



Citation for published version:

Baldasso, R & Stauffer, A 2022, 'Local survival of spread of infection among biased random walks', *Electronic Journal of Probability*, vol. 27, 135, pp. 1-28. <https://doi.org/10.1214/22-EJP861>

DOI:

[10.1214/22-EJP861](https://doi.org/10.1214/22-EJP861)

Publication date:

2022

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Local survival of spread of infection among biased random walks

Rangel Baldasso* Alexandre Stauffer†

October 7, 2020

Abstract

We study infection spread among biased random walks on \mathbb{Z}^d . The random walks move independently and an infected particle is placed at the origin at time zero. Infection spreads instantaneously when particles share the same site and there is no recovery. If the initial density of particles is small enough, the infected cloud travels in the direction of the bias of the random walks, implying that the infection does not survive locally. When the density is large, the infection spreads to the whole \mathbb{Z}^d . The proofs rely on two different techniques. For the small density case, we use a description of the infected cloud through genealogical paths, while the large density case relies on a renormalization scheme.

1 Introduction

We consider here an infection process that evolves on the d -dimensional integer lattice, where each individual performs a biased nearest-neighbor random walk. Let $p(\cdot)$ be a nearest-neighbor probability distribution on \mathbb{Z}^d . We assume without loss of generality, that, for each $i \in [d]$, $0 < p(e_i) \leq p(-e_i) < 1$, where $\{e_i\}_{i=1}^d$ is the canonical basis of \mathbb{Z}^d , and write $\vec{v} = \sum_{x \sim 0} p(x)\vec{x}$ for the d -dimensional drift of the distribution $p(\cdot)$.

We consider a ‘‘Poissonian cloud’’ of independent continuous-time random walks with jump distribution $p(\cdot)$. More formally, at time zero, each site $x \in \mathbb{Z}^d$ receives an i.i.d. number of particles $\eta_0(x) \sim \text{Poisson}(\rho)$, where $\rho > 0$ is a given parameter which we call density. Then, each particle evolves as an independent continuous-time random walk whose increments have distribution $p(\cdot)$. Denote by $\eta_t(x)$ the number of particles at position x at time t .

We now define the infection process we will consider. Particles will be of two types: either healthy or infected. At time zero, we add an additional particle at the origin that is declared infected. A given particle is infected at time t if there exists some time $s \in [0, t]$

*Email: r.baldasso@math.leidenuniv.nl; Mathematical Institute, Leiden University, P.O. Box 9512, 2300 RA Leiden, The Netherlands.

†Email: astauffer@mat.uniroma3.it; Università Roma Tre, Dip. di Matematica e Fisica, Largo S. Murialdo 1, 00146, Rome, Italy; University of Bath, Dept of Mathematical Sciences, BA2 7AY Bath, UK, supported by EPSRC Fellowship EP/N004566/1.

such that it shared a site with a previously infected particle. Alternatively, one might say that a healthy particle becomes immediately infected if it shares a site with an already infected particle.

Let $\xi_t(x)$ denote the number of infected particles at $x \in \mathbb{Z}^d$ at time t . Notice that $\xi_t(x) \in \{0, \eta_t(x)\}$, i.e., at any given time t and site $x \in \mathbb{Z}^d$, either all particles are healthy or all are infected.

Our interest lies in understanding the behavior of the process $\xi = (\xi_t)_{t \geq 0}$. We will focus on directions where the drift is negative, more precisely, we assume that $p(e_1) < p(-e_1)$ and examine the projection of the infected cloud in this direction¹. There are two forces here that are in opposition to each other: while each individual particle has a drift away from the origin, in order for the infection to travel towards the same direction, it is necessary that each site is visited only finitely many times by infected particles, which puts a strain on the system. The strength of this opposing force, though, is controlled by the density parameter ρ . This hints at the existence of different behaviors for the model, as the density parameter ρ changes.

We first consider the case when the density is small. Here, the time it takes for each infected location to be emptied is not enough to sustain the infection process locally, and the infection cloud travels towards \vec{v} . To precisely detect this phenomena, consider the quantity

$$r_t = \sup\{\langle x, e_1 \rangle : \xi_t(x) > 0\}, \quad (1.1)$$

the maximum displacement “towards the right” in the e_1 direction.

Theorem 1.1. *Assume that $p(e_1) < p(-e_1)$ and set $v_1 = p(e_1) - p(-e_1)$. For any $\delta > 0$, there exists a positive density $\rho_- > 0$ such that, for all $\rho \in (0, \rho_-)$, there exists a positive constant c_1 such that, for all $t \geq 0$,*

$$\mathbb{P}_\rho[r_t \geq (v_1 + \delta)t] \leq c_1 e^{-\frac{t}{c_1}}. \quad (1.2)$$

As a corollary of the theorem above, we conclude that, provided the density ρ is sufficiently small, every finite subset is eventually free of the infection, since it travels towards negative directions with positive speed.

Corollary 1.2. *If $p(e_1) < p(-e_1)$ and ρ is small enough, then each fixed site is visited finitely many times by infected particles, almost surely.*

When the density is large, the picture is different from Corollary 1.2. Even though each particle travels towards the direction \vec{v} , the time it takes for all particles in a given site to move allows the infection to locally survive and actually spread in the opposite direction. Our next theorem states that, provided the density is large enough, r_t actually grows with positive speed with large probability.

Theorem 1.3. *There exist a positive constant $\Delta > 0$ and density $\rho_+ > 0$ such that, for all $\rho > \rho_+$, there exist positive constants c_2, c_3 , and $t_0 = t_0(\rho)$ such that*

$$\mathbb{P}_\rho[r_t \leq c_3 t] \leq e^{-c_2(\log t)^{1+\Delta}}, \quad \text{for all } t \geq t_0. \quad (1.3)$$

¹This is not a restrictive assumption. In fact all our results remain valid if one assumes that $v_i = p(e_i) - p(-e_i) \neq 0$, for some $i \in [d]$. In order to simplify the statements, we assume without loss of generality that $v_1 < 0$. This can be achieved, for example, by permuting the coordinates.

According to Theorems 1.1 and 1.3, we have $r_t \leq -ct$, if $\rho < \rho_-$, and $r_t \geq \tilde{c}t$, for $\rho > \rho_+$. An interesting open problem would be to prove whether there is a critical point ρ_c such that, for $\rho < \rho_c$, the infection travels to the left, and, for $\rho > \rho_c$, the infection travels to the right.

Although the heuristics for the theorems above can be easily understood, turning them into proofs is not straightforward. Verifying such statements for similar models, such as in Kesten and Sidoravicius [10], relies on developing intricate renormalization schemes. In Baldasso and Teixeira [4], the authors consider another model, where particles evolve as a one-dimensional zero-range process without drift and also rely on multiscale renormalization to derive their results.

Proof overview. The proof of Theorem 1.1 follows by path-counting arguments. This proof has two distinct parts that rely on the fact that every site that is infected at time t can be reached by a concatenation of trajectories of random walks, with the first one starting at the origin. We will call such a concatenation of random walk trajectories as a *genealogical path*, but defer the formal definition to Section 2. The overall goal of the proof is to show that, for all times t , there exists no genealogical path that could bring the infection far from $t\vec{v}$.

The first part of the proof is similar to the analogous statement from Kesten and Sidoravicius [10], that consider the case when the random walks are balanced. In this part, we simply prove that, with sufficiently high probability, there exists no genealogical path starting at the origin that performs too many jumps. This will allow us to reduce the number of genealogical paths we need to consider in the rest of the proof.

The second part requires a more refined analysis, as we need to account for the contribution that each particle in a genealogical path could have to “move” the infection away from the drift. For this, we first verify that each particle that is followed by a genealogical path gives a contribution towards this displacement away from the drift that has good concentration. From this, we infer that, in order for a genealogical path to move too much away from the drift, it needs to follow too many different particles.

Theorem 1.3 has a much more intricate proof. Here we rely on multiscale renormalization by considering events where the infection does not travel fast enough in the positive direction. We prove that, provided the density is large enough, the probability of such events is small, by relating events of different scales. A central piece of the proof that allows us to establish such relation is the decoupling for biased random walks.

Decoupling. A decoupling is an estimate on the correlation decay of functions of the space-time configurations with supports that are sufficiently far away. They are powerful tools that replace the use of mixing properties for models that lack such estimates. We here prove a decoupling for the particle system composed of independent continuous-time biased random walks.

We regard the collection of space-time configurations as a subset of $\mathbb{N}_0^{\mathbb{Z}^d \times \mathbb{R}}$, and consider monotone functions of such configurations, which we assume are defined in this larger space.

We say that a function $f : \mathbb{N}_0^{\mathbb{Z}^d \times \mathbb{R}} \rightarrow \mathbb{R}$ has support on the space-time box $B \subset \mathbb{Z}^d \times \mathbb{R}$ if

$$\eta_t(x) = \bar{\eta}_t(x), \text{ for all } (x, t) \in B \text{ implies } f(\eta) = f(\bar{\eta}). \quad (1.4)$$

The function f is said decreasing if $\eta \preceq \bar{\eta}$ implies $f(\eta) \geq f(\bar{\eta})$.

Given two space-time supports B_1 and B_2 , their time distance is the quantity

$$d_V(B_1, B_2) = \inf \left\{ |t - s| : \begin{array}{l} \text{there exist } x, y \in \mathbb{Z}^d \text{ such that} \\ (x, t) \in B_1 \text{ and } (y, s) \in B_2 \end{array} \right\}. \quad (1.5)$$

We will prove a correlation estimate for decreasing functions of the space time with bounded supports that have sufficiently large time distances.

Theorem 1.4. *There exist positive constants c_4 and c_5 such that the following holds. Let B_1 and B_2 denote two space-time cubes of side-length $n > 0$ satisfying*

$$d_T = d_T(B_1, B_2) \geq c_5, \quad (1.6)$$

and assume, without loss of generality that the time coordinates in the box B_1 are smaller than the ones in B_2 . For any two decreasing functions of the space-time configurations $f_1, f_2 : \mathbb{N}_0^{\mathbb{Z}^d \times \mathbb{R}} \rightarrow [0, 1]$ with respective supports in B_1 and B_2 , we have, for any $\rho \geq 1$,

$$\mathbb{E}_{\rho^*}[f_1 f_2] \leq \mathbb{E}_{\rho^*}[f_1] \mathbb{E}_{\rho}[f_2] + \rho^{2d+2} (n + d_T)^{d+1} e^{-c_4 d_T^\delta}, \quad (1.7)$$

where $\rho^* = \rho \left(1 + d_T^{\frac{d}{4\sqrt{d+2}}} \right)$ and $\delta = \min \left\{ \frac{1}{2}, \frac{d}{2\sqrt{d+2}} \right\}$.

Remark 1.5. Notice that the estimate above is not a correlation estimate, since it relates expectations for different density parameters ρ and ρ^* . A natural question is whether it is possible to obtain such estimates without the use of the so-called sprinkling in the density. This is not the case, as verified in [8] for a very similar model, composed of discrete-time balanced random walks on \mathbb{Z} . In this case, they exhibit an example where correlations decay as $d_T^{-\frac{1}{2}}$ (see Equation (2.11) from [8]).

Proof overview of the decoupling. The proof of Theorem 1.4 relies on a construction of a coupling between two systems η and η^* with respective densities ρ and ρ^* such that, for a given subset of $H \in \mathbb{Z}^d$, we have $\eta_t(x) \leq \eta_t^*(x)$, for all $x \in H$ with large probability, provided t is large enough. With this coupling in hands, Theorem 1.4 follows easily.

The construction of the coupling is more intricate. Its nature resembles that of Bal-dasso and Teixeira [3, 4]. We start with two independent collections of particles η_0 and η_0^* . Inside a large set containing H , we match particles of η_0 to particles of η_0^* . We then let the random walks evolve. Whenever a pair of matched particles meet, they evolve together. Standard heat kernel estimates provide the bounds on the probability of this happening before some given time s . As those bounds are not strong enough for the estimates we need, we do a rematching of the particles a polynomial number of times to boost the aforementioned estimates to yield our desired stretched exponential bounds. We remark that this coupling is more robust than that of [7, 14, 15], since it does not require such a refined control of heat-kernel estimates

Related works. There are many different works that treat models for infection spread. Perhaps the most similar to ours is considered by Kesten and Sidoravicius [10].

In their case, particles evolve as continuous-time *unbiased* simple random walks and all particles placed initially at the origin begin infected. Once again, infection spreads through contact and there is no recovery. They consider the set $V(t)$ of sites visited by an infected particle up to time t and prove that there exist positive constants C_1 and C_2 such that, with large probability, $B(C_1t) \subset V(t) \subset B(C_2t)$, where $B(k) = [-k, k]^d$. In [12], Kesten and Sidoravicius strengthen the results of [10] and conclude that the set $V(t)$ satisfies a shape theorem, while in [11] they studied the case with recovery.

The proof in [10] shares some similarities with ours. The upper bound is also obtained through path-counting arguments while the lower bound revolves around the construction of a delicate renormalization structure. As we mentioned in the proof overview of our Theorems 1.1 and 1.3, the first part of our proof follows the path-counting argument for the upper bound in [10], but we need to proceed one step further to control that the infection does not move too much away from the bias of the random walks. With regard to the lower bound, we focus this discussion on the one-dimensional case to highlight the main differences between our proof and that in [10]. First, [10] observes that the infection front (say, the rightmost infected particle) behaves as a symmetric random walk when there is only one infected particle at the front, whereas the front has a drift to the right when there is more than one particle. This implies that, in order to prove that the infection grows linearly, it suffices to prove that the infection front has at least two particles a positive fraction of time, which they obtain via a renormalization scheme. In our case, where particles have a drift to the left, this strategy fails precisely because of two reasons. First, having two particles at the front may not be enough to overcome the drift to the left of the random walks, so one needs a sufficiently large number of particles. Second, even if two particles were enough to overcome the drift, just having a positive density of times with two particles at the front may not be enough to compensate for the drift to the left that the front undertake when it has just one particle. Our strategy is then to develop a multiscale renormalization scheme different from that of [10], with a target of controlling instances where the infection does not travel with a minimal positive speed to the right, and prove that events of this form have very small probability.

Regarding other works, Gracar and Stauffer [7, 6] analyzed a more general situation where the random walks move on top of the random conductance model. They prove the existence of a percolation structure (which they call Lipschitz surface) and use this to conclude that the infection spreads with positive speed for $d \geq 2$. A less structured percolating argument was obtained by Stauffer [19] in continuous space, where particles move as independent Brownian motions.

A simpler model that can be viewed as an infection process is the so-called frog model. Here, infected particles perform discrete-time simple random walks, while healthy particles do not move until an infected particle jumps onto their position. A thorough discussion about this model can be found in the survey paper by Popov [16]. We just remark that, under some minor conditions on the initial location of the particles, Alves, Machado, and Popov [1], and, independently, Ramírez and Sidoravicius [17], prove a shape theorem similar to the one in [12]. This was further strengthened by Alves, Machado, Popov and Ravishankar [2].

Let us now briefly review models where particles do not move independently. Baldasso and Teixeira [4] consider particles that move according to a one-dimensional zero-range process. Under mild conditions that guarantee the existence of invariant measures for the

process, they provide lower and upper bounds for the speed with which the front of the infection grows. Jara, Moreno, and Ramírez [9] consider an infection evolving on top of one-dimensional exclusion process and rely on regeneration arguments to prove a law of large numbers and central limit theorem for the infection front.

Regarding decoupling estimates (as in our Theorem 1.4), sprinkling ideas were first introduced in the context of random interacements by Sznitman [20] and in the context of independent Brownian motions by Sinclair and Stauffer [18] (see also [14]). These types of inequalities were used to study several conservative particle systems. Peres, Sinclair, Sousi, and Stauffer [14], Benjamini and Stauffer [5], and Stauffer [19] considered independent Brownian motions. Hilário, den Hollander, Sidoravicius, dos Santos, and Teixeira [8] treated discrete-time balanced random walks and built on the strategy from Popov and Teixeira [15] to provide decouplings for this system. The random conductance model was considered in [7], while Baldasso and Teixeira developed a decoupling inequality for the one-dimensional zero-range process [4] and the one-dimensional simple exclusion process [3].

2 Basic definitions

Let us now precisely construct the particle systems and infection process we consider.

Recall that $p(\cdot)$ denotes a nearest-neighbor probability distribution on \mathbb{Z}^d such that, for each $i \in [d]$,

$$0 < p(e_i) \leq p(-e_i) < 1, \quad (2.1)$$

where $\{e_i\}_{i=1}^d$ is the canonical basis of \mathbb{Z}^d . The vector $\vec{v} = \sum_{x \sim 0} p(x)\vec{x}$ is the d -dimensional drift of the distribution $p(\cdot)$. Due to (2.1), every coordinate of \vec{v} is non-positive. Furthermore, we assume that $p(e_1) < p(-e_1)$, so that v_1 , the first coordinate of \vec{v} , is negative.

For each $x \in \mathbb{Z}^d$ and $n \in \mathbb{N}$, let $S^{x,n} = (S_t^{x,n})_{t \geq 0}$ denote an independent copy of a continuous-time random walk with transition probability $p(\cdot)$ and $S_0^{x,n} = x$, for all $n \in \mathbb{N}$. Denote this collection by \mathcal{S} .

Given a non-negative parameter $\rho \geq 0$, consider, for each $x \in \mathbb{Z}^d$, an independent random variable $\eta_0(x)$ with distribution $\text{Poisson}(\rho)$. For each positive time $t > 0$, let

$$\eta_t(x) = \sum_{y \in \mathbb{Z}^d} \sum_{n=1}^{\infty} \mathbf{1}_{\{S_t^{y,n}=x\}} \mathbf{1}_{\{n \leq \eta_0(y)\}} \quad (2.2)$$

denote the number of particles at position x at time t . We write \mathbb{P}_ρ for the distribution of the process $\eta = (\eta_t)_{t \geq 0}$.

The product measure with marginals $\text{Poisson}(\rho)$ is invariant for the process, and we call the quantity ρ the density of the system. Besides, if $\rho' \leq \rho$, it is possible to define a order preserving coupling between two processes $\eta^{\rho'}$ and η^ρ with respective densities ρ' and ρ such that

$$\eta_t^{\rho'}(x) \leq \eta_t^\rho(x), \quad (2.3)$$

for all $t \geq 0$ and $x \in \mathbb{Z}^d$: one simply uses the same collection of walks \mathcal{S} to evolve both processes and consider the initial conditions $\eta_0^{\rho'}$ and η_0^ρ that satisfy $\eta_0^{\rho'}(x) \leq \eta_0^\rho(x)$, for all $x \in \mathbb{Z}^d$.




Figure 1: An example of a genealogical infected path. Different colors stand for different particles followed.

We now proceed to define the infection process $(\xi_t)_{t \geq 0}$. At any given time $t \geq 0$, $\xi_t(x)$ denotes the number of infected particles at $x \in \mathbb{Z}^d$. At time zero, only particles at the origin are infected, which means that $\xi_0(x) = \eta_0(0)\mathbf{1}_{\{x=0\}}$ (recall we add an infected particle at the origin at time zero). As for the evolution, each time an infected particle jumps to a site with healthy particles, all particles at that given site become infected. This in particular implies that $\xi_t(x) \in \{0, \eta_t(x)\}$, for all $t \geq 0$ and $x \in \mathbb{Z}^d$.

2.1 Genealogical infected paths

We can see the infection mechanism through genealogical paths. In order to define these paths, we need to consider paths that particles in the process follow. If X denotes a particle in the system, we denote by $(X(s))_{s \geq 0}$ the path that it performs.

We say that $\gamma : [0, T] \rightarrow \mathbb{Z}^d$ is a genealogical infected path up to time T (GIP(T)) if $\gamma(0) = 0$ and there exist a sequence of times $0 = t_0 < t_1 < \dots < t_n \leq T$ and a sequence of particles X_1, \dots, X_{n+1} such that $X_1(0) = 0$, X_k gets infected at time t_{k-1} , and, for all $t \in [t_{k-1}, t_k]$ we have $\gamma(t) = X_k(t)$, for all $k \in [n]$; for convenience of notation, we assume that $t_{n+1} = T$. Of course, $\xi_T(x) > 0$ if and only if there exists a GIP(T) γ with $\gamma(T) = x$. See Figure 1 for a representation of a GIP(T).

We will identify a GIP(T) by the following: a quantity k of jumps of γ , the number n of particles it follows, a vector (k_1, \dots, k_n) with non-negative entries that counts the number of jumps each particle performs before it stops being followed, and a vector with non-zero integer entries (j_1, \dots, j_n) that identifies which is the next particle to be followed. If $j_i > 0$, then when X_i makes its last jump we take X_{i+1} to be the j_i -th healthy particle that was present at the site where X_i jumped to. Whenever $j_i < 0$, we wait X_i to perform all its k_i jumps, and only at this moment wait $|j_i|$ healthy particles to jump on the site where X_i is, taking the $|j_i|$ -th such particle as X_{i+1} .

This identification has some particularities we need to address. First, observe that, if $k_i = 0$, we demand that $j_i < 0$, i.e., after following a particle that does not jump, we need to follow a particle that is not yet in the site we are considering. Moreover, we also have $\sum_{i=1}^n k_n = k$. Finally, it is not always the case that all possible choices for these quantifiers will yield a GIP, but all GIP can be identified by some choice of such values.

3 The small density case: proof of Theorem 1.1

We now consider the case when the density is very small. Using a first moment computation, we will prove that there exists a small density $\rho > 0$ such that, with large probability, the infection travels towards the negative direction in the first coordinate axis.

We will control how infection spreads by using genealogical paths. The first proposition we prove states that it is unlikely to exist a GIP that jumps many times. This is a general statement that does not require the density $\rho > 0$ to be small and does not depend on the probability distribution $p(\cdot)$. These quantities will be important when we consider finer properties of the model.

Proposition 3.1. *For any $\rho > 0$, there exists a positive constant $c_6 = c_6(\rho)$, which might be taken to be monotone non-decreasing as a function of ρ , such that, for all $T \geq 0$,*

$$\mathbb{P}_\rho \left[\begin{array}{l} \text{there exists a GIP}(T) \text{ that} \\ \text{jumps more than } c_6 T \text{ times} \end{array} \right] \leq e^{-c_6 T + 1}. \quad (3.1)$$

As a byproduct of the proposition above, we immediately obtain the following result.

Proposition 3.2. *For all densities $\rho > 0$, there exists $c_7 > 0$ and $\alpha > 0$ such that, for all $t \geq 0$,*

$$\mathbb{P}_\rho \left[\|x\| > \alpha t, \text{ for some } x \in \mathbb{Z}^d \text{ such that } \xi_t(x) > 0 \right] \leq c_7 e^{-\frac{t}{c_7}}. \quad (3.2)$$

Proof of Proposition 3.1. The proof of this statement relies on a first moment calculation. We will bound the expectation of the number of GIPs that jump more than $c_6 T$ times before time T .

The discussion gets simplified when we use the identification of GIP introduced in Subsection 2.1. A GIP is identified by the number of jumps k , the number of particles it follows n , a vector (k_1, \dots, k_n) with non-negative entries that counts the number of jumps each particle performs and a vector with non-zero integer entries (j_1, \dots, j_n) that identifies which is the next particle to be followed.

If $k_i > 0$, then the time it takes to follow particle X_i until its last jump is equal in distribution to the sum of k_i independent Exponential(1) random variables. We will call these exponential times \mathcal{T}_i . Besides, if $j_i < 0$, we gain a time contribution that comes from the fact we need to wait until j_i healthy particles jump into the site. This can be dominated by a collection of $|j_i|$ random variables with distribution Exponential($\rho + 1$), where each one of them counts the time until either one particle jumps into the current site, or the particle we are following jumps to another site. These waiting times will be denoted by $\mathcal{W}_{i,\ell}$. In the last case, when the particle we are following jumps before all the $|j_i|$ new particles arrived, we disregard the path, and control this by considering additional random variables $G_i \sim \text{Geometric}\left(\frac{1}{\rho+1}\right)$ that will control whether all the $|j_i|$ new particles were able to arrive into the correct site before the distinguished particle jumped.

Let \mathcal{G}_T denote the number of GIPs that jump more than $c_6 T$ times before time T , where c_6 is a constant that will be chosen later. The discussion above allows us to bound

$$\mathbb{E}_\rho[\mathcal{G}_T] \leq \mathbb{E}_\rho \left[\sum_{k=c_6 T}^{\infty} \sum_{n=1}^{\infty} \sum_{\substack{(k_1, \dots, k_n) \\ \sum_{i=1}^n k_i = k}} \sum_{(j_1, \dots, j_n)} \mathbf{1}_{\left\{ \sum_{i=1}^k \mathcal{T}_i + \sum_{i: j_i < 0} \sum_{\ell=1}^{|j_i|} \mathcal{W}_{i, \ell} < T \right\}} \right. \\ \left. \prod_{i=1}^n \left(\mathbf{1}_{\{\mathcal{N}_i > j_i\}} \mathbf{1}_{\{j_i \geq 1\}} + \mathbf{1}_{\{G_i > |j_i|\}} \mathbf{1}_{\{j_i \leq -1\}} \right) \right], \quad (3.3)$$

where \mathcal{N}_i counts the number of particles present at the site onto which X_i makes its last jump.

Let us now estimate the expectation above. Notice first that the possible choices for the partitions (k_1, \dots, k_n) such that $\sum_{i=1}^n k_i = k$ is bounded by

$$\binom{n+k-1}{n-1} \leq 2^{n+k}, \quad (3.4)$$

Second, write $J = \sum_{i: j_i < 0} |j_i|$ and notice that the random variables $\sum_{i=1}^k \mathcal{T}_i + \sum_{i: j_i < 0} \sum_{\ell=1}^{|j_i|} \mathcal{W}_{i, \ell}$ stochastically dominate a sum of $k+J$ i.i.d. $\text{Exponential}(1+\rho)$ random variables. This allows us to bound

$$\mathbb{P}_\rho \left[\sum_{i=1}^k \mathcal{T}_i + \sum_{i: j_i < 0} \sum_{\ell=1}^{|j_i|} \mathcal{W}_{i, \ell} \leq T \right] \leq \mathbb{P}[X \geq k+J], \quad (3.5)$$

where $X \sim \text{Poisson}((1+\rho)T)$. Notice also that the number of walks we follow n is clearly upper bounded by $k+J$, since, whenever $i \in [n]$ is such that $k_i = 0$, we have $j_i < 0$.

To bound the expectation, we first divide the sum according to which subset of indices $A \subset [n]$ is such that $j_i < 0$ for $i \in A$. Notice that, for a fixed choice of set A , standard concentration bounds (see Appendix A and, in particular, Remark A.2) yields

$$\mathbb{E}_\rho \left[\prod_{i=1}^n \left(\mathbf{1}_{\{\mathcal{N}_i > j_i\}} \mathbf{1}_{\{j_i \geq 1\}} + \mathbf{1}_{\{G_i > |j_i|\}} \mathbf{1}_{\{j_i \leq -1\}} \right) \right] \leq \left(\frac{\rho}{1+\rho} \right)^J \prod_{i \notin A} e^{-c_{16} j_i \mathbf{1}_{\{j_i \geq 10\rho\}}}, \quad (3.6)$$

where we bound the first term above by one if $j_i \leq 10\rho$ and use Chernoff's bound for the Poisson random variables \mathcal{N}_i if j_i is large.

From the discussion above we obtain the bound

$$\mathbb{E}_\rho[\mathcal{G}_T] \leq \sum_{k=c_6 T}^{\infty} \sum_{n=1}^{\infty} \sum_{A \subset [n]} \sum_{(j_1, \dots, j_n)} 2^{n+k} \mathbb{P}[X \geq k+J] \left(\frac{\rho}{1+\rho} \right)^J \mathbf{1}_{\{n \leq k+J\}} \prod_{i \notin A} e^{-c_{16} j_i \mathbf{1}_{\{j_i \geq 10\rho\}}} \\ \leq \sum_{k=c_6 T}^{\infty} \sum_{n=1}^{\infty} \sum_{A \subset [n]} \sum_{j_i: i \in A} 2^{n+k} \mathbb{P}[X \geq k+J] \left(\frac{\rho}{1+\rho} \right)^J (c\rho)^k \mathbf{1}_{\{n \leq k+J\}}, \quad (3.7)$$

where in the last inequality we summed out the terms j_i with $i \notin A$ using the fact that the amount of such indices is at most k . We now use the estimates from Proposition A.1. By choosing $c_6 \geq 2(\rho + 1)$, we obtain

$$\mathbb{E}_\rho[\mathcal{G}_T] \leq \sum_{k=c_6 T}^{\infty} \sum_{n=1}^{\infty} \sum_{A \subset [n]} \sum_{j_i: i \in A} 2^{n+k} \exp \left\{ -c_{16}(k+J) \log \frac{k+J}{(1+\rho)T} \right\} (c\rho)^{k+J} \mathbf{1}_{\{n \leq k+J\}}. \quad (3.8)$$

It remains to bound the summations above. We begin by observing that the number of choices of indices j_i with $i \in A$ such that $J = \sum_{i \in A} |j_i|$ is given is bounded by $\binom{J-1}{|A|-1}$. This allows us to bound the quantity above by

$$\sum_{k=c_6 T}^{\infty} \sum_{n=1}^{\infty} \sum_{A \subset [n]} \sum_{J=|A|}^{\infty} \binom{J-1}{|A|-1} 2^{n+k} \exp \left\{ -c_{16}(k+J) \log \frac{k+J}{(1+\rho)T} \right\} (c\rho)^{k+J} \mathbf{1}_{\{n \leq k+J\}}. \quad (3.9)$$

Second, we bound the number of choices for A according to its size. This yields the bound

$$\sum_{k=c_6 T}^{\infty} \sum_{n=1}^{\infty} \sum_{\ell=0}^n \sum_{J=\ell}^{\infty} \binom{n}{\ell} \binom{J}{\ell} 2^{n+k} \exp \left\{ -c_{16}(k+J) \log \frac{k+J}{(1+\rho)T} \right\} (c\rho)^{k+J} \mathbf{1}_{\{n \leq k+J\}}. \quad (3.10)$$

We now change the order of summations, and conclude that the r.h.s. of (3.8) is bounded by

$$\begin{aligned} & \sum_{k=c_6 T}^{\infty} \sum_{J=1}^{\infty} \sum_{\ell=0}^J \sum_{n=\ell}^{k+J} \binom{n}{\ell} \binom{J}{\ell} 2^{n+k} \exp \left\{ -c_{16}(k+J) \log \frac{k+J}{(1+\rho)T} \right\} (c\rho)^{k+J} \\ & \leq \sum_{k=c_6 T}^{\infty} \sum_{J=1}^{\infty} (k+J) 8^{(k+J)} \exp \left\{ -c_{16}(k+J) \log \frac{k+J}{(1+\rho)T} \right\} (c\rho)^{k+J} \\ & = \sum_{k=c_6 T}^{\infty} \sum_{J=1}^{\infty} (k+J) \exp \left\{ -c_{16}(k+J) \log \frac{k+J}{(1+\rho)T} \right\} (8c\rho)^{k+J}. \end{aligned} \quad (3.11)$$

To conclude, we set $L = \max\{k, J\} \leq k+J \leq 2L$ in the equation above to obtain that the r.h.s. of (3.8) is bounded by

$$\begin{aligned} & \sum_{L=c_6 T}^{\infty} 2L^3 \exp \left\{ -c_{16}L \log \frac{L}{(1+\rho)T} \right\} (8c\rho)^{2L} \\ & \leq \sum_{L=c_6 T}^{\infty} \exp \left\{ -L \left(c_{16} \log \frac{c_6}{(1+\rho)} - \log(64c^2\rho^2e^6) \right) \right\} \\ & = \sum_{L=c_6 T}^{\infty} \exp \left\{ -L \log \left(\left(\frac{c_6}{1+\rho} \right)^{c_{16}} \frac{1}{64c^2\rho^2e^6} \right) \right\} \\ & \leq e^{-c_6 T + 1}, \end{aligned} \quad (3.12)$$

if $c_6 = c_6(\rho) = \left(3(64e^6c^2\rho^2)^{\frac{1}{c_{16}}} + 2 \right) (1+\rho) > 0$.




Figure 2: The random variable \mathcal{R}_ϵ in dimension one. Notice that the random walk is completely contained in the gray area.

Combining the bound above with (3.8), we obtain that

$$\mathbb{E}_\rho[\mathcal{G}_T] \leq e^{-c_6 T+1}, \quad (3.13)$$

if $c_6 = c_6(\rho) > 0$ is large enough. To conclude the proof, simply apply Markov's inequality to obtain

$$\mathbb{P}_\rho[\mathcal{G}_T \geq 1] \leq \mathbb{E}_\rho[\mathcal{G}_T] \leq e^{-c_6 T+1}. \quad (3.14)$$

□

We now proceed to conclude the proof of Theorem 1.1. In view of Proposition 3.1, we may restrict ourselves to paths that do not jump many times until time T . Before presenting the proof, we provide some basic facts about biased random walks that will be used in the proof, since GIPs are constructed by concatenating such objects.

Lemma 3.3. *Let $(X_t)_{t \geq 0}$ be a random walk starting from the origin, and let \vec{v} denote its drift. For any $\epsilon > 0$, there exist a positive random variable \mathcal{R}_ϵ and a positive constant $c_8 = c_8(p(\cdot), d, \epsilon) > 0$ such that, almost surely*

$$\|X_t - t\vec{v}\| \leq \max\{\mathcal{R}_\epsilon, \epsilon t\}, \quad \text{for all } t \geq 0, \quad (3.15)$$

and, for every $u \geq 0$,

$$\mathbb{P}[\mathcal{R}_\epsilon \geq u] \leq c_8^{-1} e^{-c_8 u}. \quad (3.16)$$

In particular, \mathcal{R}_ϵ is stochastically dominated by $\frac{1}{c_8} Y + \frac{\log c_8}{c_8}$, where $Y \sim \text{Exponential}(1)$.

Proof. Begin by using Lemma A.3 to bound, for $T \geq \frac{2}{\epsilon}$,

$$\begin{aligned} & \mathbb{P}[\|X_t - t\vec{v}\| \geq \epsilon t, \text{ for some } t \geq T] \\ & \leq \sum_{k=1}^{\infty} \mathbb{P}[\|X_t - t\vec{v}\| \geq \epsilon kT, \text{ for some } t \in [kT, (k+1)T]] \\ & \leq \sum_{k=1}^{\infty} \mathbb{P}\left[\|X_t - t\vec{v}\| \geq \frac{\epsilon}{2}(k+1)T, \text{ for some } t \in [kT, (k+1)T]\right] \\ & \leq \sum_{k=1}^{\infty} c_{17}^{-1} e^{-c_{17}(k+1)T} \leq c^{-1} e^{-cT}, \end{aligned} \quad (3.17)$$

for some suitable positive constant $c > 0$.

Due to Borel-Cantelli Lemma, the random variable

$$\bar{\mathcal{R}}_\epsilon = \inf \left\{ u \in \left[\frac{2}{\epsilon}, \infty \right) \cap \mathbb{N} : \|X_t - t\bar{v}\| \leq \epsilon t, \text{ for all } t \geq u \right\} \quad (3.18)$$

is almost surely finite. Define now

$$\mathcal{R}_\epsilon = \max \left\{ \bar{\mathcal{R}}_\epsilon, \sup \{ \|X_t - t\bar{v}\| : t \in [0, \bar{\mathcal{R}}_\epsilon] \} \right\}, \quad (3.19)$$

so that (3.15) clearly holds. Finally, observe that, as an immediate consequence of (3.17) and Lemma A.3, one obtains that, for all $u \geq 0$,

$$\begin{aligned} \mathbb{P}[\mathcal{R}_\epsilon \geq u] &\leq \mathbb{P}[\bar{\mathcal{R}}_\epsilon \geq u] + \mathbb{P}[\sup \{ \|X_t - t\bar{v}\| : t \in [0, \bar{\mathcal{R}}_\epsilon] \} \geq u, \bar{\mathcal{R}}_\epsilon \leq u] \\ &\leq c^{-1}e^{-cu} + \mathbb{P}[\sup \{ \|X_t - t\bar{v}\| : t \in [0, u] \} \geq u] \\ &\leq c_8^{-1}e^{-c_8u}, \end{aligned} \quad (3.20)$$

concluding the proof. \square

We are now in position to prove Theorem 1.1. The proof is also based on the first moment method by bounding the expected number of GIPs that do not behave as expected. In view of Proposition 3.1, we may consider only GIPs that do not jump many times.

Proof of Theorem 1.1. By possibly increasing the value of c_1 , we can assume $T \geq 1$. Furthermore, by monotonicity, we may assume that $\delta < \frac{1}{2}|v_1|$ and denote by \mathcal{H}_T the number of GIPs that jump at most c_6T times and are such that their endpoint x satisfy $\langle x - T\bar{v}, e_1 \rangle \geq \delta T$, where c_6 is the constant from Proposition 3.1.

We first apply Lemma 3.3. Using the fact that $\delta < |v_1|$ and that $v_1 < 0$, the maximum displacement of a given particle towards the positive direction in the axis e_1 can be bounded by

$$\begin{aligned} \sup_t \langle X_t, e_1 \rangle &= \sup_t \{ \langle X_t - \bar{v}t, e_1 \rangle + v_1t \} \\ &\leq \max \left\{ \sup_t \{ v_1t + \delta t \}, \sup_t \{ v_1t + \mathcal{R}_\delta \} \right\} \\ &\leq \mathcal{R}_\delta, \end{aligned} \quad (3.21)$$

since $v_1 < 0$ and $v_1 + \delta < 0$. In particular, the maximum displacement of a given GIP can be bounded by a sum of i.i.d. random variables with distribution \mathcal{R}_δ , one for each followed particle in the path. Notice that independence of the random variables comes from the fact that we only follow newly infected particles. In particular, if the path follows at most αT particles, its maximum displacement is bounded by a sum of αT i.i.d. random variables $(\mathcal{R}_\delta^i)_{i=1}^{\alpha T}$ with distribution \mathcal{R}_δ . This implies that the probability that a given GIP has displacement to the right bigger than δT is bounded by

$$\mathbb{P} \left[\sum_{j=1}^{\alpha T} \mathcal{R}_\delta^j \geq \delta T \right] \leq \mathbb{P} \left[\sum_{j=1}^{\alpha T} Y_j \geq Tc_8\delta + \alpha T \log c_8 \right] = \mathbb{P}[Z \leq \alpha T], \quad (3.22)$$

where Y_j are i.i.d. Exponential(1) random variables and $Z \sim \text{Poisson}(Tc_8\delta + \alpha T \log c_8)$. Choose now α small enough such that $c_8\delta + \alpha \log c_8 > \alpha$ and observe that there exists a positive constant $c_9 = c_9(\alpha, \delta, c_8)$ such that

$$\mathbb{P} \left[\sum_{j=1}^{\alpha T} \mathcal{R}_\delta^j \geq \delta T \right] \leq \mathbb{P} [Z \leq \alpha T] \leq e^{-c_9 T}. \quad (3.23)$$

We are now in position to bound the expectation of \mathcal{H}_T . Reasoning similarly as in (3.3) and using the same notation, we can obtain the bound

$$\mathbb{E}_\rho[\mathcal{H}_T] \leq \mathbb{E}_\rho \left[\sum_{n=1}^{\infty} \sum_{(k_i)_{i=1}^n: \sum_{i=1}^n k_i \leq c_6 T} \sum_{(j_1, \dots, j_n)} \mathbf{1}_{\{\sum_{i=1}^n \mathcal{R}_\delta^i \geq \delta T\}} \prod_{i=1}^n (\mathbf{1}_{\{\mathcal{N}_i > j_i\}} \mathbf{1}_{\{j_i \geq 1\}} + \mathbf{1}_{\{G_i > |j_i|\}} \mathbf{1}_{\{j_i \leq -1\}}) \right], \quad (3.24)$$

where n denotes the number of particles followed in a given path, the vector $(k_i)_{i=1}^n$ counts how many jumps each of the particles performs and $(j_i)_{i=1}^n$ controls transitions between particles.

Proceeding as in the proof of Proposition 3.1 and using the bound in (3.23), we obtain

$$\mathbb{E}_\rho[\mathcal{H}_T] \leq \sum_{n=1}^{\infty} \sum_{(k_i)_{i=1}^n: \sum_{i=1}^n k_i \leq c_6 T} (\mathbf{1}_{\{n \geq \alpha T\}} + \mathbf{1}_{\{n \leq \alpha T\}} e^{-c_9 T}) (c\rho)^n. \quad (3.25)$$

To estimate the number of vectors $(k_i)_{i=1}^n$ such that $\sum_{i=1}^n k_i \leq c_6 T$, we bound this quantity by the number of vectors $(k_i)_{i=1}^{n+1}$ such that $\sum_{i=1}^{n+1} k_i = c_6 T$ and apply the bound in (3.4). We now combine this bound with (3.25) to obtain, for $\alpha \leq c_6$,

$$\begin{aligned} \mathbb{E}_\rho[\mathcal{H}_T] &\leq \sum_{n=1}^{\infty} (\mathbf{1}_{\{n \geq \alpha T\}} + \mathbf{1}_{\{n \leq \alpha T\}} e^{-c_9 T}) (c\rho)^n \binom{\lfloor c_6 T \rfloor + n}{n} \\ &\leq 2^{c_6 T} \sum_{n \geq \alpha T} (2c\rho)^n + e^{-c_9 T} \sum_{n=1}^{\alpha T} \left(\frac{2c\rho c_6 T e}{n} \right)^n \\ &\leq 2^{c_6 T} \frac{(2c\rho)^{\alpha T}}{1 - 2c\rho} + \alpha T e^{-c_9 T} e^{2c\rho c_6 T}, \end{aligned} \quad (3.26)$$

where the bound in the last summation is obtained by maximizing the expression $\left(\frac{2c\rho c_6 T e}{n} \right)^n$ in n .

From (3.26), one easily obtains that, provided ρ is small enough, there exists a positive constant c_1 such that

$$\mathbb{E}_\rho[\mathcal{H}_T] \leq c_1^{-1} e^{-c_1 T}, \quad (3.27)$$

for all $T \geq 1$. Markov's inequality concludes the proof. \square

4 Decoupling

This section contains the proof of Theorem 1.4, the main step towards the proof of Theorem 1.3. We first prove this theorem with the aid of an auxiliary proposition, whose proof can be found in Subsection 4.1.

In the following, we say that a process $\eta = (\eta_t)_{t \geq 0}$ has density $\rho > 0$ if the initial distribution of the process is a product of i.i.d. $\text{Poisson}(\rho)$ random variables.

The proof of the decoupling relies on the construction of a coupling between two processes with different densities such that the process with higher density dominates the less dense one inside a box with large probability.

Proposition 4.1. *There exists a positive constant $c_{10} = c_{10}(d) > 0$ such that the following holds. For any time $T \geq c_{10}$, density $\rho \geq 1$, and box $H = [1, n]^d \subset \mathbb{Z}^d$, there exists a coupling \mathbb{P} between two processes $\eta = (\eta_s)_{s \geq 0}$ and $\eta^* = (\eta_s^*)_{s \geq 0}$ such that*

1. η has density ρ and η^* has density $\rho^* = \rho(1 + T^{-\frac{d}{4\sqrt{d+2}}})$;

2. η is independent of η_0^* ;

- 3.

$$\mathbb{P}[\eta_T(x) > \eta_T^*(x), \text{ for some } x \in H] \leq \rho^{d+1}(n+T)^{d+1}e^{-c_{11}T^\delta}, \quad (4.1)$$

where $c_{11} = c_{11}(d) > 0$ and $\delta = \min\left\{\frac{1}{2}, \frac{d}{2\sqrt{d+2}}\right\}$.

We defer the proof of Proposition 4.1 to Subsection 4.1. For now, we use this proposition to conclude the proof of Theorem 1.4.

Proof of Theorem 1.4. Without loss of generality, assume that

$$B_1 = [1, n]^d \times [-n, 0] \text{ and } B_2 = [1, n]^d \times [T, T+n]. \quad (4.2)$$

In particular, $d_T = T$.

Set $H = [-2n - \rho T, 3n + \rho T]^d$ and consider the event

$$A = \left\{ \begin{array}{l} \text{some particle that is outside } H \text{ at time } T \\ \text{has a trajectory that intersects } B_2 \end{array} \right\}. \quad (4.3)$$

We now use the coupling from Proposition 4.1 for H (here we need to choose $d_T = T$ large enough). Denote by \mathbb{P} the probability measure of the coupling and by \mathbb{E} the expectation with respect to \mathbb{P} . We obtain two processes $\eta = (\eta_s)_{s \geq 0}$ and $\eta^* = (\eta_s^*)_{s \geq 0}$ with η independent of η_0^* such that

$$\begin{aligned} \mathbb{P}[\eta_T(x) > \eta_T^*(x), \text{ for some } x \in H] &\leq \rho^{d+1}(n + (1 + \rho)T)^{d+1} \exp\{-c_{11}T^\delta\} \\ &\leq \rho^{2d+2}(n+T)^{d+1}e^{-c_{11}T^\delta}, \end{aligned} \quad (4.4)$$

if T is taken large enough.

Observe now that, whenever $\eta_T \preceq_H \eta_T^*$ (we write $\eta \preceq_H \xi$ if $\eta(x) \leq \xi(x)$, for all $x \in H$) and $\eta \notin A$, we have $f_2(\eta^*) \leq f_2(\eta)$. This allows us to estimate

$$\begin{aligned}
\mathbb{E}_{\rho^*}[f_1(\eta^*)f_2(\eta^*)] &= \mathbb{E}_{\rho^*}[f_1(\eta^*)\mathbb{E}[f_2(\eta^*)|(\eta_s^*)_{s \leq 0}]] \\
&= \mathbb{E}_{\rho^*}[f_1(\eta^*)\mathbb{E}[f_2(\eta^*)|\eta_0^*]] \\
&\leq \mathbb{E}\left[f_1(\eta^*)\mathbb{E}[f_2(\eta^*)\mathbf{1}_{\{\eta \notin A\}}\mathbf{1}_{\{\eta_T \preceq_H \eta_T^*\}}|\eta_0^*]\right] + \mathbb{P}[\eta \in A] \\
&\quad + \mathbb{P}[\eta_T(x) > \eta_T^*(x), \text{ for some } x \in H] \\
&\leq \mathbb{E}[f_1(\eta^*)\mathbb{E}[f_2(\eta)|\eta_0^*]] + \mathbb{P}[\eta \in A] + \rho^{2d+2}(n+T)^{d+1}e^{-c_{11}T^\delta} \\
&\leq \mathbb{E}_{\rho^*}[f_1]\mathbb{E}_{\rho}[f_2] + \mathbb{P}[\eta \in A] + \rho^{2d+2}(n+T)^{d+1}e^{-c_{11}T^\delta}.
\end{aligned} \tag{4.5}$$

It remains to estimate the probability of the event A . We split this probability according to the position of the particles at time T . For $k \geq 1$, let

$$H(k) = \{x \in \mathbb{Z}^d : d_\infty(x, H) = k\}, \tag{4.6}$$

and notice that there exists a positive constant $c = c(d)$ such that

$$|H(k)| \leq c(n + \rho T + k)^{d-1}. \tag{4.7}$$

Besides, in order for a particle that is at $H(k)$ at time T to reach B_2 , it needs to perform at least $2n + k + \rho T$ steps. Consider the event

$$A(k) = \left\{ \begin{array}{l} \text{some particle that is in } H(k) \text{ at time } T \\ \text{has a trajectory that intersects } B_2 \end{array} \right\}. \tag{4.8}$$

We will bound the probability of $A(k)$ by considering the number of particles that are inside $H(k)$ at time T . We obtain

$$\begin{aligned}
\mathbb{P}_\rho[\eta \in A(k)] &\leq \mathbb{P}_\rho \left[\sum_{x \in H(k)} \eta_T(x) \geq 2c\rho(n + \rho T + k)^{d-1} + T + k \right] \\
&\quad + [2c\rho(n + \rho T + k)^{d-1} + T + k] \mathbb{P}[\text{Poisson}(n) \geq 2n + k + \rho T] \\
&\leq e^{-T-k} + [2c\rho(n + \rho T + k)^{d-1} + T + k] e^{-\rho T-k} \\
&\leq c(n^d + T^d + k^d)e^{-T-k},
\end{aligned} \tag{4.9}$$

by further increasing the value of T if necessary.

In particular,

$$P_\rho[\eta \in A] \leq \sum_{k=1}^{\infty} \mathbb{P}_\rho[\eta \in A(k)] \leq c(n^d + T^d)e^{-T}. \tag{4.10}$$

Combining the equation above with (4.5) concludes the proof. \square

4.1 Coupling

In this subsection we present the proof of Proposition 4.1. The proof follows the same general steps from [4] and [3]. For this reason, we omit some simple computations.

The idea for constructing the coupling is to start with two independent configurations η_0 and η_0^* and evolve them simultaneously in order to obtain the domination at time T . We first observe that we can restrict ourselves to a larger box H^* around H and assume that all particles that end up inside H at time T never leave H^* . Now, to obtain the domination, we fix a deterministic sequence of times $(s_i)_{i \geq 0}$ and, in each of these times, we construct a pairing between particles of η and of η^* that are inside H^* . The evolution is then set in a way that, if a pair of matched particles meets, they continue evolving together. In particular, the probability that there is no domination at time T is bounded by the probability that there exists a particle of η that never meets a pair. This will be easily bounded with the aid of Proposition B.3.

We now proceed to prove Proposition 4.1.

Proof of Proposition 4.1. Fix T large enough so that Proposition B.3 applies. The coupling will use independent initial configurations η_0 and η_0^* with respective densities ρ and ρ^* . Besides, we consider two independent copies \mathcal{S} and \mathcal{S}' of the graphical construction presented in Section 2.

The process $\eta = (\eta_t)_{t \geq 0}$ will follow the walks from \mathcal{S} , while the process $\eta^* = (\eta_s^*)_{s \geq 0}$ will alternate between the two constructions. This implies that η is independent of η_0^* .

Consider the sequence of times $s_k = kT^{\frac{1}{d+2}}$, for $k = 0, \dots, \lfloor T^{\frac{d+1}{d+2}} \rfloor$, and fix the set

$$H^* = \left[-3\rho T, n + 3\rho T \right]^d. \quad (4.11)$$

Proceeding as in (4.9), we can bound the probability of the event

$$A = \left\{ \begin{array}{l} \text{there exists a particle of } \eta \\ \text{that is outside } H^* \text{ for some time } s_k \\ \text{and is inside } H \text{ at time } T \end{array} \right\} \quad (4.12)$$

by

$$\mathbb{P}[A] \leq cT^{\frac{d+1}{d+2}}(n^d + T^d)e^{-T}. \quad (4.13)$$

Let $L = \left\lfloor \frac{c_{20}}{2\sqrt{d}} T^{\frac{1}{\sqrt{d+2}}} \right\rfloor$, and, for $i \in \mathbb{Z}^d$, write

$$H(i) = iL + [0, L]^d. \quad (4.14)$$

Set $I = \{i \in \mathbb{Z}^d : H(i) \cap H^* \neq \emptyset\}$, and notice that

$$|I| \leq \left(\frac{\lceil (6\rho T + n + 1) \rceil}{L} \right)^d \leq c\rho^d (n + T)^d. \quad (4.15)$$

Observe that $H^* \subset \bigcup_{i \in I} H(i)$. If necessary, we increase H^* to coincide with the union $\bigcup_{i \in I} H(i)$.

We will perform the pairing inside each box $H(i)$ with $i \in I$. Concentration bounds on the number of particles yields, for each $i \in \mathbb{Z}^d$,

$$\mathbb{P} \left[\sum_{x \in H(i)} \eta_t(x) \geq \frac{\rho + \rho^*}{2} L^d \right] \leq \exp \left\{ -\frac{1}{24} (\rho^* - \rho)^2 L^d \right\}, \quad (4.16)$$

and

$$\mathbb{P} \left[\sum_{x \in H(i)} \eta_t^*(x) \leq \frac{\rho + \rho^*}{2} L^d \right] \leq \exp \left\{ -\frac{1}{24} (\rho^* - \rho)^2 L^d \right\}, \quad (4.17)$$

In particular, if

$$B_t = \left\{ \sum_{x \in H(i)} \eta_t^*(x) \leq \sum_{x \in H(i)} \eta_t(x), \text{ for some } i \in I \right\}, \quad (4.18)$$

then we can use (4.16) and (4.17) to obtain

$$\mathbb{P}[B_t] \leq 2|I| \exp \left\{ -\frac{1}{24} (\rho^* - \rho)^2 L^d \right\}. \quad (4.19)$$

Here it is important to notice that the bound above does not require η_t and η_t^* to be independent, only that they have the correct marginal distributions.

We now construct the evolution of the process $\eta^* = (\eta_s^*)_{s \geq 0}$. If we are in the event $A \cup B_0$, then η^* evolves using the graphical construction given by the paths \mathcal{S}' and, consequently, η and η^* have independent evolutions. Assume we are in $A^c \cap B_0^c$. We perform a pairing between particles of η_0 and η_0^* inside each set $H(i)$, for $i \in I$. This pairing is deterministic and follows the following steps.

1. First pair as many particles as possible of η_0 to particles of η_0^* that are in the same site.
2. Pair the remaining particles of η_0 to particles of η_0^* that are in the same sub-box $H(i)$.

Observe that this pairing is always possible in the event B_0^c .

This pairing between particles will be used to construct the evolution of the process η^* . Particles of this process will use the trajectories of the construction \mathcal{S}' until the time they meet their pair. When this happens, the particle will follow the trajectory from \mathcal{S} that its corresponding pair from η uses. At the times $(s_k)_{k=1}^{\lfloor T^{\frac{d+1}{d+2}} \rfloor}$, these pairings are remade, following the rules mentioned above. These rules imply that the number of particles that meet a pair cannot decrease when the pairings are remade.

In view of Proposition B.3, the probability that a particle meets its couple between times s_k and s_{k+1} is at least $c_{22} T^{-\frac{d}{2(d+2)}} > 0$. Furthermore, each particle has $\lfloor T^{\frac{d+1}{d+2}} \rfloor \geq \lfloor T^{\frac{2}{3}} \rfloor$ attempts to find a pair. We obtain the bound

$$\begin{aligned} \mathbb{P} \left[\begin{array}{l} \text{a given particle does not find any of its pairs} \\ \text{in any of its allowed attempts, } A^c \cap \bigcap_{i=0}^{\lfloor T^{\frac{d+1}{d+2}} \rfloor - 1} B_{s_i} \end{array} \right] &\leq (1 - c_{22} T^{-\frac{d}{2(d+2)}})^{\lfloor T^{\frac{2}{3}} \rfloor} \\ &\leq e^{-c_{22} T^{\frac{d+8}{6(d+2)}}} \leq e^{-c_{22} T^{\frac{1}{2}}}. \end{aligned} \quad (4.20)$$

In particular, we can use union bounds to obtain

$$\begin{aligned}
\mathbb{P}[\eta_T(x) > \eta_T^*(x), \text{ for some } x \in H] &\leq \mathbb{P}[A] + \sum_{i=0}^{\lfloor T^{\frac{d-1}{d+2}} \rfloor - 1} \mathbb{P}[B_{s_i}] \\
&+ \rho(6\rho T + n + 1)^d \mathbb{P} \left[\begin{array}{l} \text{a given particle does not find any of its pairs} \\ \text{in any of its allowed attempts, } A^c, \cap_{i=1}^{\lfloor T^{\frac{d+1}{d+2}} \rfloor - 1} B_{s_i} \end{array} \right] \\
&\leq cT^{\frac{3}{4}}(n^d + T^d)e^{-T} + 2T^{\frac{d+1}{d+2}}|I|e^{-\frac{1}{24}\rho^2 T^{-\frac{d}{2\sqrt{d+2}}}L^d} + \rho(6\rho T + n + 1)^d e^{-c_{22}T^{\frac{1}{2}}} \\
&\leq \rho^{d+1}(n + T)^{d+1}e^{-c_{11}T^\delta},
\end{aligned} \tag{4.21}$$

concluding the proof. \square

5 The large density case

We now focus on the proof of Theorem 1.3. Here, we develop a renormalization structure for the infection front. We start by providing elementary bounds for the probability that the environment behaves exceptionally bad, and for events where the infection act abnormally. In Subsection 5.2, we establish the main notations used for the renormalization scheme presented in Subsection 5.3. Finally, Subsection 5.4 contains the proof of Theorem 1.3.

5.1 Elementary bounds

In this subsection, we present some rough initial estimates that will be used to bound the events where the infection process behaves exceptionally bad. These estimates are not sharp and rely mostly on union bounds and large deviations for the number of particles or jumps in a given time interval.

Our first lemma bounds the probability that a given vertex has many particles at some moment before a given time.

Lemma 5.1. *There exists a positive constant c_{12} such that, for all $L \geq 1$ and density $\rho \leq L^2$,*

$$\mathbb{P}_\rho \left[\begin{array}{l} \eta_t(0) \geq L^{d+4}, \\ \text{for some } t \in [0, L] \end{array} \right] \leq c_{12}e^{-c_{12}^{-1}L}. \tag{5.1}$$

Proof. Notice that, in order for the origin to have many particles before time L , it is necessary that a large ball around it starts with many particles, or there exists a particle

that performs many jumps before time L . Proceeding as in (4.9), we can bound

$$\begin{aligned} \mathbb{P}_\rho \left[\begin{array}{l} \eta_t(0) \geq L^{d+4}, \\ \text{for some } t \in [0, L] \end{array} \right] &\leq \mathbb{P}_\rho \left[\sum_{x \in B(0, 3L)} \eta_0(x) \geq L^{d+4} \right] \\ &+ \mathbb{P}_\rho \left[\begin{array}{l} \text{there exists some particle that} \\ \text{starts outside } B(0, 3L) \\ \text{and reaches the origin before time } L \end{array} \right] \\ &\leq c_{12} e^{-c_{12}^{-1} L}, \end{aligned} \quad (5.2)$$

by choosing c_{12} appropriately. \square

The next lemma bounds the probability that the infection process travels abnormally fast during a given time interval.

Lemma 5.2. *There exists a positive constant c_{13} such that, for all $L \geq 2$ and density $\rho \leq L^2$,*

$$\mathbb{P}_\rho \left[\begin{array}{l} \text{there exists an infected particle} \\ \text{outside } [-L^{d+6}, L^{d+6}]^d \text{ before time } L \end{array} \right] \leq c_{13} e^{-c_{13}^{-1} L}. \quad (5.3)$$

Proof. We can bound the probability of the event above by the probability that some vertex inside $[-L^{d+6}, L^{d+6}]^d$ has many particles at some moment before time L or the infection process travels fast through a field of vertices that are typical. Let A denote the event in the statement of the lemma, and define the events

$$B = \left\{ \begin{array}{l} \eta_t(x) \geq L^{d+4}, \text{ for some} \\ t \in [0, L] \text{ and } x \in [-L^{d+6}, L^{d+6}]^d \end{array} \right\}, \quad (5.4)$$

and

$$C = \left\{ \begin{array}{l} \text{there exists a path of size } L^{d+6} \text{ that starts} \\ \text{at the origin and such that one particle jumps} \\ \text{at each site in order before time } L \end{array} \right\}. \quad (5.5)$$

Notice that

$$\mathbb{P}_\rho[A] \leq \mathbb{P}_\rho[B] + \mathbb{P}_\rho[C \cap B^c]. \quad (5.6)$$

Lemma 5.1 gives a bound on the probability of B . As for the probability of $C \cap B^c$, we use union bound on the possible choices of paths to obtain

$$\begin{aligned} \mathbb{P}_\rho[C \cap B^c] &\leq (2d)^{L^{d+6}} \mathbb{P}[\text{Poisson}(L^{d+5}) \geq L^{d+6}] \\ &\leq (2d)^{L^{d+6}} e^{-c_{16} L^{d+6} \log \frac{L^{d+6}}{L^{d+5}}}. \end{aligned} \quad (5.7)$$

The proof is completed by combining the bounds above and choosing the constants appropriately. \square

Recall that we defined the front of the infection towards the direction e_1 as

$$r_t = \sup\{\langle x, e_1 \rangle : \xi_t(x) > 0\}. \quad (5.8)$$

The next lemma states that, provided the density is large enough depending on the time parameter, there is a big probability that r_t is large.

Lemma 5.3. *There exists a positive constant $c_{14} > 0$ such that, for any positive integer $L \geq 1$*

$$\mathbb{P}_{L^{\frac{1}{2}}}[r_L < 8L] \leq c_{14} L e^{-c_{14}^{-1} L^{\frac{1}{2}}}. \quad (5.9)$$

Proof. Consider the event $A_{k,i}$ defined as

$$A_{k,i} = \left\{ \begin{array}{l} \text{between times } k + \frac{i}{8} \text{ and } k + \frac{i+1}{8}, \text{ there exists a particle} \\ \text{at position } (k+i)e_1 \text{ that jumps to position } (k+i+1)e_1, \\ \text{and there is a particle at position } (k+i+1)e_1 \text{ that does not jump} \end{array} \right\}, \quad (5.10)$$

and notice that, in $\cap_{k=0}^{L-1} \cap_{i=0}^7 A_{k,i}$, we have $r_t \geq 8L$.

Furthermore, we can bound the probability

$$\begin{aligned} \mathbb{P}_{L^{\frac{1}{2}}}[A_{k,i}^c] &\leq \mathbb{P} \left[\text{Poisson}((1 - e^{-\frac{1}{8}})p(e_1)L^{\frac{1}{2}}) = 0 \right] + \mathbb{P} \left[\text{Poisson}(e^{-\frac{1}{8}}L^{\frac{1}{2}}) = 0 \right] \\ &= e^{(1-e^{-\frac{1}{8}})p(e_1)L^{\frac{1}{2}}} + e^{-e^{-\frac{1}{8}}L^{\frac{1}{2}}}. \end{aligned} \quad (5.11)$$

Combining this with union bound yields

$$\mathbb{P}_{L^{\frac{1}{2}}}[r_L < L] \leq \sum_{k=0}^{L-1} \sum_{i=0}^7 \mathbb{P}_{L^{\frac{1}{2}}}[A_{k,i}^c] \leq 8L e^{-c_{14} L^{\frac{1}{2}}}, \quad (5.12)$$

for some positive constant c_{14} , and concludes the proof. \square

5.2 The box notation

Let us now introduce the notation in order to develop our renormalization analysis. We first fix a sequence of scales by setting

$$L_0 = L > 0 \quad \text{and} \quad L_{k+1} = L_k^{d+7}. \quad (5.13)$$

The initial value L_0 will be chosen large enough later.

For each $k \in \mathbb{N}_0$, define the space-time box

$$B_k = [-L_k^{d+6}, L_k^{d+6}]^d \times [0, L_k], \quad (5.14)$$

and for $m \in \mathbb{Z} \times L_k \mathbb{N}_0$, let $B_k(m)$ denote the translated box $B_k(m) = m + B_k$.

Define also the sequence of velocities

$$\vartheta_0 = 8 \quad \text{and} \quad \vartheta_{k+1} = \vartheta_k - \frac{6}{\pi^2(k+1)^2}. \quad (5.15)$$

Notice that $\lim \vartheta_k = 7$.

Our goal is to bound the probability that the infection process travels very slowly towards the e_1 direction. The continuous time nature of the process implies that events of this form do not have a bounded support. For this reason, we introduce events that approximate these and have support contained in the box B_k . For each $k \in \mathbb{N}_0$, define

$$R_k = \{(x, t) \in B_k : \|x\|_\infty = L_k^{d+6} \text{ or } x_1 \geq \vartheta_k L_k \text{ and } t = L_k\}, \quad (5.16)$$




Figure 3: A representation for the sets B_k and R_k . We represent the event E_k by considering the shaded area as the infection process.

and, for $m \in \mathbb{Z} \times L_k \mathbb{N}_0$, let $R_k(m)$ denote the translated set $R_k(m) = m + R_k$. Figure 3 contains a representation of the sets B_k and R_k for $d = 1$.

For $m = (x, t) \in \mathbb{Z} \times \mathbb{N}_0$, we denote by $(\xi_s^m)_{s \geq t}$ for the infection process starting with initial configuration η_t and initial collection of infected particles the ones in x .

Consider the events

$$E_k(m) = \{\xi_t^m(x) = 0, \text{ for all } (x, t) \in R_k(m)\}. \quad (5.17)$$

See Figure 3 for a representation of the event above for $d = 1$. Observe that the events $E_k(m)$ are non-increasing and have support inside $B_k(m)$. When $m = (0, 0)$, we omit this in the definition and denote $E_k(0, 0)$ simply by E_k .

We introduce the sequence of densities

$$\rho_0 = \sqrt{L_0} \quad \text{and} \quad \rho_{k+1} = \rho_k(1 + L_k^{-d/4\sqrt{d+2}}), \quad (5.18)$$

and notice that, for each k , we have $\rho_{k+1} \leq L_k^2$. Furthermore the sequence $(\rho_k)_{k \in \mathbb{N}_0}$ is monotone increasing and $\rho_\infty = \lim \rho_k$ exists and is finite. For each k , define also

$$p_k = \mathbb{P}_{\rho_k}[E_k(m)], \quad (5.19)$$

which does not depend on m , by translation invariance of the process.

Finally, let M_k denote the set of indices m in scale k in the box B_{k+1} , namely

$$M_k = (\mathbb{Z}^d \times L_k \mathbb{N}_0) \cap B_{k+1}, \quad (5.20)$$

and observe that

$$|M_k| \leq L_{k+1}^{d+1}. \quad (5.21)$$

5.3 Estimates on p_k

In this subsection, we provide estimates on p_k , by proving that, provided L_0 is large enough, the sequence $(p_k)_{k \in \mathbb{N}_0}$ decays quickly.

The first lemma we prove here states a recursive inequality that relates p_k to p_{k+1} .

Lemma 5.4. *There exist positive constants ℓ_0 and A such that, if $L_0 \geq \ell_0$, then, for all $k \in \mathbb{N}_0$,*

$$p_{k+1} \leq L_{k+1}^A \left(p_k^{d+8} + e^{-c_{15} L_k^\delta} \right), \quad (5.22)$$

where $\delta = \min \left\{ \frac{1}{2}, \frac{d}{2\sqrt{d+2}} \right\}$.

Proof. Choose ℓ_0 is large enough such that, if $L_0 \geq \ell_0$, then

$$L_k > \frac{8\pi^2}{3}(2d+14)(k+1)^2, \text{ for all } k \in \mathbb{N}_0. \quad (5.23)$$

Consider, for each $m = (y, jL_k) \in M_k$,

$$D_k(m) = D_k(y, jL_k) = \left\{ \begin{array}{l} \xi_t^m(x) > 0, \text{ for some } (x, t) \text{ with} \\ \|x - y\|_\infty = L_k^{d+6}, (t - jL_k) \leq L_k \end{array} \right\}, \quad (5.24)$$

the event where the infection ξ^m travels abnormally fast before time L_k .

We first claim that, on E_{k+1} , one of the following two conditions hold:

1. For some $m \in M_k$, the event $D_k(m)$ holds;
2. There are $2d+15$ values $(m_i)_{i=1}^{2d+15}$ in M_k with different time coordinates such that $E_k(m_i)$ holds for all $i \leq 2d+15$.

In order to verify the claim, suppose we are in the event E_{k+1} , that the first condition does not hold, and that the second one holds for at most $2d+14$ values of m with different time coordinates.

For each $m \in M_k$, let $X(m)$ denote a point in \mathbb{Z}^d such that

$$\langle X(m), e_1 \rangle = \max \{ \langle x, e_1 \rangle : \xi_{L_k}^m(x) > 0 \}. \quad (5.25)$$

By assuming that the first condition above does not hold, we have

$$-L_k^{d+6} \leq \langle X(m) - m, e_1 \rangle \leq L_k^{d+6}. \quad (5.26)$$

Furthermore, using that the second condition does not hold, the lower bound can be improved to

$$\langle X(m) - m, e_1 \rangle \geq \vartheta_k L_k \quad (5.27)$$

for all but at most $2d+14$ different time coordinates of m .

Define the sequence of sites in \mathbb{Z}^d as

$$Y_0 = 0 \quad \text{and} \quad Y_{\ell+1} = X((Y_\ell, \ell L_k)). \quad (5.28)$$

Notice that

$$\xi_{L_{k+1}} \left(\frac{Y_{L_{k+1}}}{L_k} \right) > 0. \quad (5.29)$$

We now estimate

$$\begin{aligned} r_{L_{k+1}} &\geq \langle Y_{\frac{L_{k+1}}{L_k}}, e_1 \rangle = \sum_{\ell=1}^{\frac{L_{k+1}}{L_k}} \langle Y_\ell - Y_{\ell-1}, e_1 \rangle \\ &\geq \vartheta_k L_k \left(\frac{L_{k+1}}{L_k} - 2d - 14 \right) - (2d+14)L_k^{d+6} \\ &\geq \vartheta_{k+1} L_{k+1} + \frac{6}{\pi^2(k+1)^2} L_{k+1} - (2d+14)(\vartheta_k L_k + L_k^{d+6}) \\ &\geq \vartheta_{k+1} L_{k+1} + \frac{6}{\pi^2(k+1)^2} L_{k+1} - 8(2d+14)(L_k + L_k^{d+6}) \\ &> \vartheta_{k+1} L_{k+1}, \end{aligned} \quad (5.30)$$




Figure 4: The first application of the decoupling estimate. Small boxes represent the supports of the events $E_k(m_i)$ and the bold boxes are the supports of the events considered when applying the decoupling.

which is a contradiction with the fact that we are in the event E_{k+1} .

We just concluded that, on E_{k+1} , there are two possible outcomes: either $D_k(m)$ holds for some $m \in M_k$ or there exists a choice of indices $(m_i)_{i=1}^{2d+15}$ in M_k with different time coordinates such that $E_k(m_i)$ holds for all $i \leq 2d + 15$.

Suppose we are in the last case described above, and fix one possible choice of indices $(m_i = (x_i, s_i))_{i=1}^{2d+15}$. Observe that, for all i , we have $L_k \leq s_{i+2} - s_i \leq L_{k+1}$.

By possibly further increasing the value of ℓ_0 , we can apply Theorem 1.4 a few times (see Figure 4) to conclude that

$$\begin{aligned} \mathbb{P}_{\rho_{k+1}} \left[\bigcap_{i=1}^{2d+15} E_k(m_i) \right] &\leq \mathbb{P}_{\rho_{k+1}} \left[\bigcap_{i=1}^{2d+13} E_k(m_i) \right] \mathbb{P}_{\rho_k} [E_k(m_{15})] \\ &\quad + L_k^{4(d+1)} ((2L_k^{d+6} + 1)^d + L_{k+1}^d) e^{-c_4^{-1} L_k^\delta} \\ &\leq \mathbb{P}_{\rho_k} [E_k]^{d+8} + (d+8) L_k^{4(d+1)} ((2L_k^{d+6} + 1)^d + L_{k+1}^d) e^{-c_4^{-1} L_k^\delta}. \end{aligned} \quad (5.31)$$

Union bounds, Lemma 5.2 and a change of constants yields

$$\begin{aligned} \mathbb{P}_{\rho_{k+1}} [E_{k+1}] &\leq |M_k|^{2d+17} \left(\mathbb{P}_{\rho_k} [E_k]^{d+8} + (d+8) c_4 ((2L_k^{d+6} + 1)^d + L_{k+1}^d) e^{-c_4^{-1} L_k^\delta} \right) \\ &\quad + |M_k| \mathbb{P}_{\rho_{k+1}} [D_k] \\ &\leq L_{k+1}^A \left(p_k^{d+8} + e^{-c_{15} L_k^\delta} \right), \end{aligned} \quad (5.32)$$

concluding the proof. \square

In the next lemma, we prove that, provided L_0 is large enough, the probabilities p_k decay fast.

Lemma 5.5. *There exist constants $\ell_1 \geq \ell_0$ and $\Delta > 0$ such that, if $L_0 \geq \ell_1$, then*

$$p_k \leq e^{-(\log L_k)^{1+\Delta}}, \quad \text{for all } k \in \mathbb{N}_0. \quad (5.33)$$

Proof. Fix Δ such that $(d+7)^{1+\Delta} < d+8$. Choose $\ell_1 \geq \ell_0$ large enough such that, for all $L \geq \ell_0$,

$$L^{A(d+7)} e^{(d+7)^{1+\Delta} (\log L)^{1+\Delta}} \left(e^{-(d+8) (\log L)^{1+\Delta}} + e^{-c_{15} L^\delta} \right) \leq 1, \quad (5.34)$$

where A , δ and c_{15} are given by Lemma 5.4. By further increasing ℓ_1 , Lemma 5.3 implies

$$p_0 \leq e^{-(\log L_0)^{1+\Delta}}. \quad (5.35)$$

Assume now that (5.33) holds for some value of k . Let us verify that it remains true for $k + 1$.

By Lemma 5.4, we have

$$\begin{aligned} e^{(\log L_{k+1})^{1+\Delta}} p_{k+1} &\leq L_{k+1}^A e^{(\log L_{k+1})^{1+\Delta}} \left(p_k^{d+8} + e^{-c_{15} L_k^\delta} \right) \\ &\leq L_{k+1}^A e^{(d+7)^{1+\Delta} (\log L_k)^{1+\Delta}} \left(e^{-(d+8)(\log L_k)^{1+\Delta}} + e^{-c_{15} L_k^\delta} \right) \\ &\leq 1, \end{aligned} \quad (5.36)$$

concluding the proof. \square

5.4 Proof of Theorem 1.3

In this subsection, we combine the lemmas provided so far to conclude the proof of Theorem 1.3.

Since the event in (1.3) is non-increasing, it suffices to verify the statement for one value of ρ .

Define $\bar{E}_k(m)$ analogously to $E_k(m)$, but with $R_k(m)$ replaced by $\bar{R}_k(m)$, an m -translation of

$$\bar{R}_k = \{(x, t) \in B_k : \|x\|_\infty = L_k^{d+6} \text{ or } x_1 \geq 7L_k \text{ and } t = L_k\}, \quad (5.37)$$

and notice that this is a non-increasing sequence of events. In particular, for $m = (x, t)$, we have

$$\mathbb{P}_{\rho_\infty}[\bar{E}_k(m)] \leq \mathbb{P}_{\rho_\infty}[E_k(m)] \leq p_k \leq e^{-(\log L_k)^{1+\Delta}}, \quad (5.38)$$

for all $k \geq 0$ and all $m \in M_k$, provided L_0 is large enough. We can now proceed with the proof of Theorem 1.3.

Proof of Theorem 1.3. Choose k_0 large enough such that $\rho_\infty \leq L_{k_0}^2$, and set $t_0 = L_{k_0}$. For $t \geq t_0$, let $k \geq k_0$ be the only value such that

$$L_k \leq t < L_{k+1}, \quad (5.39)$$

and choose $\bar{\ell} \in \left[1, \frac{L_{k+1}}{L_k}\right]$ such that

$$\bar{\ell} L_k \leq t < (\bar{\ell} + 1) L_k. \quad (5.40)$$

Define the event

$$A_k = \bigcup_{m=(x,t) \in M_k} \bar{E}_k(m) \cap D_k(m), \quad (5.41)$$

where $D_k(m)$ is given by (5.24). Notice that

$$\mathbb{P}_{\rho_\infty}[A_k] \leq L_{k+1}^{d+1} (\mathbb{P}_{\rho_\infty}[\bar{E}_k] + \mathbb{P}_{\rho_\infty}[D_k]) \leq 2L_{k+1}^{d+1} e^{-(\log L_k)^{1+\Delta}}. \quad (5.42)$$

On A_k^c , by applying the same concatenation argument of (5.30), we can easily obtain that

$$r_{\ell L_k} \geq 7\ell L_k, \quad \text{for all } 1 \leq \ell \leq \frac{L_{k+1}}{L_k}. \quad (5.43)$$

In particular, if the above holds simultaneously with $r_t \leq t$, then all particles that are at a position that realizes $r_{\bar{\ell}L_k}$ at time $\bar{\ell}L_k$ must jump at least $7\bar{\ell}L_k - t \geq 6L_k$ times between times $\bar{\ell}L_k$ and t . This can be easily bound with concentration on the number of jumps of any given particle. In conclusion, we obtain

$$\begin{aligned} \mathbb{P}_{\rho_\infty}[r_t < t, \eta_0(0) > 0] &\leq \mathbb{P}_{\rho_\infty}[A_k] + \mathbb{P}[\text{Poisson}(L_k) \geq 6K_k] \\ &\leq 2L_{k+1}^{d+1} e^{-(\log L_k)^{1+\Delta}} + e^{-L_k} \\ &\leq 2t^{(d+7)(d+1)} \exp\left\{-\left(\log t^{\frac{1}{d+7}}\right)^{1+\Delta}\right\} + e^{-t^{\frac{1}{d+7}}}, \end{aligned} \tag{5.44}$$

which concludes the proof by possibly changing constants. \square

A Concentration

We collect here some basic facts about concentration of Poisson random variables and biased random walks we use in the text.

Proposition A.1. *There exists $c_{16} > 0$ such that, for any $\rho > 0$ and $A \geq 2\rho$ integer, if $X \sim \text{Poisson}(\rho)$ then*

$$\mathbb{P}[X \geq A] \leq \exp\left\{-c_{16}A \log \frac{A}{\rho}\right\}. \tag{A.1}$$

Remark A.2. In particular we obtain the bound

$$\mathbb{P}[X \geq A] \leq \exp\left\{-c_{16}A \mathbf{1}_{\{A \geq 10\rho\}} \log 10\right\}, \tag{A.2}$$

for any positive integer $A \geq 0$.

Proof. Use Markov's inequality with $\lambda = \log \frac{A}{\rho}$ to obtain

$$\begin{aligned} \mathbb{P}[X \geq A] &= \mathbb{P}[e^{\lambda X} \geq e^{\lambda A}] \\ &\leq \exp\left\{-\lambda A + \rho(e^\lambda - 1)\right\} \\ &\leq \exp\left\{-A \log \frac{A}{\rho} + \rho\left(\frac{A}{\rho} - 1\right)\right\} \\ &\leq \exp\left\{-c_{16}A \log \frac{A}{\rho}\right\}. \end{aligned} \tag{A.3}$$

\square

Our next lemma regards linear deviations of the biased random walk from its mean.

Lemma A.3. *For any $\epsilon > 0$, there exists a positive constant $c_{17} = c_{17}(p(\cdot), d, \epsilon)$ such that, for any $u \geq 0$,*

$$\mathbb{P}[||X_t - \bar{v}t|| \geq \epsilon u, \text{ for some } t \in [0, u]] \leq c_{17}^{-1} e^{-c_{17}u}. \tag{A.4}$$

Proof. First notice that union bound allows us to write

$$\mathbb{P} [|X_t - \bar{v}t| \geq \epsilon u, \text{ for some } t \in [0, u]] \leq \sum_{i=1}^d \mathbb{P} \left[|X_t^i - v_i t| \geq \frac{\epsilon}{d} u, \text{ for some } t \in [0, u] \right], \quad (\text{A.5})$$

where $(X_t^i)_{t \geq 0}$ is a continuous time random walk in \mathbb{Z} that jumps to the right with rate $p(e_i)$ and to the left with rate $p(-e_i)$, and $v_i = p(e_i) - p(-e_i)$. Alternatively, we can construct X^i from a standard biased random walk $(Y_t)_{t \geq 0}$ that jumps with rate one and with distribution $q(1) = 1 - q(-1) = \frac{p(e_i)}{p(e_i) + p(-e_i)}$. Even though the values $q(1)$, $q(-1)$ (and, by consequence, $(Y_t)_{t \geq 0}$) depend on $i \in [d]$, we omit this dependence. With this construction, if $\bar{v}_i = q(1) - q(-1)$, we obtain, for all $i \in [d]$,

$$\begin{aligned} & \mathbb{P} \left[|X_t^i - v_i t| \geq \frac{\epsilon}{d} u, \text{ for some } t \in [0, u] \right] \\ &= \mathbb{P} \left[|Y_t - \bar{v}_i t| \geq \frac{\epsilon}{d} u, \text{ for some } t \in [0, (p(e_i) + p(-e_i))u] \right]. \end{aligned} \quad (\text{A.6})$$

We now work with the last quantity. We will prove that

$$\mathbb{P} [|Y_t - \bar{v}_i t| \geq \epsilon u, \text{ for some } t \in [0, u]] \leq c^{-1} e^{-c\epsilon u}, \quad (\text{A.7})$$

for some suitable choice of $c > 0$ that depends only on $p(\cdot)$ and $\epsilon > 0$. Combining this with (A.5) and (A.6) concludes the proof of the lemma.

In order to prove (A.7), notice first that $t \mapsto Y_t^i - \bar{v}_i t$ is a continuous time martingale. Hence, Doob's maximal Inequality allows us to conclude that, for any $\lambda > 0$,

$$\begin{aligned} & \mathbb{P} [|Y_t - \bar{v}_i t| \geq \epsilon u, \text{ for some } t \in [0, u]] \leq \mathbb{E} [\exp \{ \lambda |Y_u - \bar{v}_i u| \}] e^{-\lambda \epsilon u} \\ & \leq e^{-\lambda \epsilon u} \left(\mathbb{E} [\exp \{ \lambda (Y_u - \bar{v}_i u) \}] + \mathbb{E} [\exp \{ -\lambda (Y_u - \bar{v}_i u) \}] \right). \end{aligned} \quad (\text{A.8})$$

By writing Y_u^i as the sum of a Poissonian number of Bernoulli random variables, one obtains, for every $\lambda \in \mathbb{R}$,

$$\mathbb{E} [e^{\lambda Y_u}] = \exp \{ u (q(1)e^\lambda + q(-1)e^{-\lambda}) \}. \quad (\text{A.9})$$

Combining (A.8) and (A.9), we can bound

$$\begin{aligned} & \mathbb{P} [|Y_t - \bar{v}_i t| \geq \epsilon u, \text{ for some } t \in [0, u]] \\ & \leq e^{-\lambda \epsilon u} \left(\mathbb{E} [\exp \{ \lambda (Y_u - \bar{v}_i u) \}] + \mathbb{E} [\exp \{ -\lambda (Y_u - \bar{v}_i u) \}] \right) \\ & \leq \exp \{ u (q(1)e^\lambda + q(-1)e^{-\lambda} - \lambda \bar{v}_i - \lambda \epsilon) \} \\ & \quad + \exp \{ u (q(1)e^{-\lambda} + q(-1)e^\lambda + \lambda \bar{v}_i - \lambda \epsilon) \} \\ & \leq 2e^{-c\epsilon u}, \end{aligned} \quad (\text{A.10})$$

if $\lambda > 0$ is taken small enough, depending of the values of $q(1)$, $q(-1)$ and ϵ . This yields (A.7) and, together with (A.5) and (A.6), concludes the proof. \square

B Sampling of random walks

In this section, we provide bounds for transition probabilities for biased random walks and collect some consequences of these bounds.

We first consider zero-mean random walks. The following lemma is a consequence of an analogous result for discrete time balanced random walks.

Lemma B.1. *Let $(X_s)_{s \geq 0}$ denote a nearest-neighbor continuous-time random walk with transition probability that has mean zero. There exists a positive constant $c_{18} > 0$ such that, if all $t \geq 1$ and $x \in \mathbb{Z}^d$ satisfying $\|x\| \leq \sqrt{t}$, then*

$$\mathbb{P}_0[X_t = x] \geq \frac{c_{18}}{t^{d/2}}. \quad (\text{B.1})$$

Proof. First notice that $(X_s)_{s \geq 0}$ may be realized as a discrete-time zero-mean lazy random walk $(\tilde{X}_n)_{n \geq 0}$ together with a Poisson process $(P_s)_{s \geq 0}$ in \mathbb{R}_+ with intensity 2 that controls the jump times.

According to the remark after Proposition 2.1.2 from [13], there exists a constant $c > 0$ such that, if $\|x\| \leq \sqrt{n}$, then

$$\mathbb{P}_x[\tilde{X}_n = x] \geq \frac{c}{n^{d/2}}. \quad (\text{B.2})$$

We now bound

$$\begin{aligned} \mathbb{P}_x[X_t = 0] &\geq \sum_{k: |k-2t| \leq t} \mathbb{P}[P_t = k] \mathbb{P}_x[\tilde{X}_n = x] \geq \sum_{k: |k-2t| \leq t} \mathbb{P}[P_t = k] \frac{c}{t^{d/2}} \\ &\geq \frac{c}{t^{d/2}} \mathbb{P}[|P_t - 2t| \leq t] \geq \frac{c_{18}}{t^{d/2}}, \end{aligned} \quad (\text{B.3})$$

concluding the proof. \square

Our next goal is to obtain an analogous result to Lemma B.1 for biased random walks.

Lemma B.2. *Let $(X_s)_{s \geq 0}$ denote a biased nearest-neighbor random walk with transition probability $p(\cdot)$. Assume $p(e_i) > 0$ and $p(-e_i) > 0$, for all $i \in [d]$, and set*

$$v = (v_1, \dots, v_d) = \sum_{y \sim 0} p(y)y \in \mathbb{R}^d. \quad (\text{B.4})$$

There exist positive constants c_{19} , c_{20} , and c_{21} such that, if $t \geq c_{19}$ and $x \in \mathbb{Z}^d$ is such that $\|x - vt\| \leq c_{20}\sqrt{t}$, then

$$\mathbb{P}_0[X_t = x] \geq \frac{c_{21}}{t^{d/2}}. \quad (\text{B.5})$$

The proof is based in writing the biased random walk as the sum of drift terms and a zero-mean continuous time random walk, and using the estimate provided by Lemma B.1.

Proof. For each $i \in [d]$, let

$$p_i = \min\{p(e_i), p(-e_i)\}, \quad (\text{B.6})$$

and write $Z = 2 \sum_{i=1}^d p_i$. Let $(\tilde{X}_s)_{s \geq 0}$ be a continuous-time random walk with transition probability $q(\cdot)$ given by $q(e_i) = q(-e_i) = \frac{p_i}{Z}$.

We can write

$$X_t = \sum_{i=1}^d \text{Sign}(v_i) Y_t^i e_i + \tilde{X}_{t(1 - \sum_{i=1}^d |v_i|)}, \quad (\text{B.7})$$

where $Y_t^i \sim \text{Poisson}(t|v_i|)$ are independent.

We now split the probability of $X_t = x$ according to the value of the sum $\sum_{i=1}^d \text{Sign}(v_i) Y_t^i e_i$. This yields

$$\mathbb{P}_0[X_t = x] = \sum_{y \in \mathbb{Z}^d} \mathbb{P} \left[\sum_{i=1}^d \text{Sign}(v_i) Y_t^i e_i = y \right] \mathbb{P}_0 \left[\tilde{X}_{t(1 - \sum_{i=1}^d |v_i|)} = x - y \right]. \quad (\text{B.8})$$

Set $c_{20} = \frac{1}{2} \sqrt{1 - \sum_{i=1}^d |v_i|}$, and observe that, if $\|y - vt\| \leq c_{20} \sqrt{t}$, then

$$\|x - y\| \leq \|x - vt\| + \|y - vt\| \leq \sqrt{t \left(1 - \sum_{i=1}^d |v_i| \right)}, \quad (\text{B.9})$$

so that Lemma B.1 implies

$$\begin{aligned} \mathbb{P}_0[X_t = x] &\geq \sum_{y: \|y - vt\| \leq c_{20} \sqrt{t}} \mathbb{P} \left[\sum_{i=1}^d \text{Sign}(v_i) Y_t^i e_i = y \right] \frac{c_{18}}{t^{d/2}} \\ &\geq \frac{c_{18}}{t^{d/2}} \mathbb{P} \left[\left\| \sum_{i=1}^d \text{Sign}(v_i) Y_t^i e_i - vt \right\| \leq c_{20} \sqrt{t} \right]. \end{aligned} \quad (\text{B.10})$$

The central limit theorem implies that, if t is large enough, the last probability above is uniformly bounded from below by some positive constant. This concludes the proof. \square

The following proposition states that, provided two random walks do not start very far away, the probability that they meet at a given time t does not decay very fast.

Proposition B.3. *Let $p(\cdot)$ be a transition probability satisfying all hypotheses from Lemma B.2. There exists a positive constant $c_{\gamma\gamma} > 0$ such that, for all $t \geq c_{19}$ and $x, y \in \mathbb{Z}^d$ such that $\|x - y\| \leq \frac{c_{20}}{2} \sqrt{t}$, the following holds. If $(X_s)_{s \geq 0}$ and $(Y_s)_{s \geq 0}$ are independent random walks with jump distribution $p(\cdot)$ and initial positions $X_0 = x$ and $Y_0 = y$, then*

$$\mathbb{P}[X_t = Y_t^y] \geq \frac{c_{22}}{t^{d/2}}. \quad (\text{B.11})$$

Proof. Recall Lemma B.2 and fix $\delta > 0$ such that

$$\delta \leq \frac{c_{21}}{t^{d/2}} \left| B \left(vt, \frac{c_{20}}{2} \sqrt{t} \right) \cap \mathbb{Z}^d \right|, \quad (\text{B.12})$$

where $B(a, r)$ denotes the L^∞ -ball of \mathbb{R}^d with center a and radius r . Using that there exists $c > 0$ such that $|B(vt, \frac{c_{20}}{2} \sqrt{t}) \cap \mathbb{Z}^d| \geq ct^{d/2}$, for all $t \geq c_{19}$, we obtain that δ can be chosen uniformly positive, for all t large enough.

By Lemma B.2, we have

$$\mathbb{P}[X_t = z] \geq \frac{c_{21}}{t^{d/2}}, \quad (\text{B.13})$$

for all $z \in B(x + vt, \frac{c_{20}}{2}\sqrt{t}) \cap \mathbb{Z}^d$. The same holds for the random walk Y . From this, we conclude

$$P[X_t = Y_t] \geq \sum_{z \in B(x+vt, \frac{c_{20}}{2}\sqrt{t}) \cap \mathbb{Z}^d} \mathbb{P}[X_t = z] \mathbb{P}[Y_t = z] \geq \frac{c_{21}}{t^{d/2}}, \quad (\text{B.14})$$

which concludes the proof. \square

References

- [1] Oswaldo Alves, Fabio Machado, and Serguei Popov. The shape theorem for the frog model. *The Annals of Applied Probability*, 12(2):533–546, 2002.
- [2] Oswaldo Alves, Fabio Machado, Serguei Popov, and Krishnamurthi Ravishankar. The shape theorem for the frog model with random initial configuration. *Markov Processes Relat. Fields*, 7(4):525–539, 2001.
- [3] Rangel Baldasso and Augusto Teixeira. How can a clairvoyant particle escape the exclusion process? *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 54(4):2177–2202, 2018.
- [4] Rangel Baldasso and Augusto Teixeira. Spread of an infection on the zero range process. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 56(3):1898–1928, 2020.
- [5] Itai Benjamini and Alexandre Stauffer. Perturbing the hexagonal circle packing: a percolation perspective. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 49(4):1141–1157, 2013.
- [6] Peter Gracar and Alexandre Stauffer. Multi-scale lipschitz percolation of increasing events for poisson random walks. *The Annals of Applied Probability*, 29(1):376–433, 2019.
- [7] Peter Gracar and Alexandre Stauffer. Random walks in random conductances: Decoupling and spread of infection. *Stochastic Processes and their Applications*, 129(9):3547–3569, 2019.
- [8] Marcelo Hilário, Frank Den Hollander, Vladas Sidoravicius, Renato Soares dos Santos, and Augusto Teixeira. Random walk on random walks. *Electronic Journal of Probability*, 20, 2015.
- [9] Milton Jara, Gregorio Moreno, and Alejandro F. Ramírez. Front propagation in an exclusion one-dimensional reactive dynamics. *Markov Processes and Related Fields*, 14(2):185–206, 2008.
- [10] Harry Kesten and Vladas Sidoravicius. The spread of a rumor or infection in a moving population. *The Annals of Probability*, pages 2402–2462, 2005.

- [11] Harry Kesten and Vladas Sidoravicius. A phase transition in a model for the spread of an infection. *Illinois J. Math.*, 50(1-4):547–634, 2006.
- [12] Harry Kesten and Vladas Sidoravicius. A shape theorem for the spread of an infection. *Annals of mathematics*, pages 701–766, 2008.
- [13] Gregory F Lawler and Vlada Limic. *Random walk: a modern introduction*, volume 123. Cambridge University Press, 2010.
- [14] Yuval Peres, Alistair Sinclair, Perla Sousi, and Alexandre Stauffer. Mobile geometric graphs: detection, coverage and percolation. *Proceedings of the twenty-second annual ACM-SIAM symposium on Discrete Algorithms*, pages 412–428, 2011.
- [15] Serguei Popov and Augusto Teixeira. Soft local times and decoupling of random interlacements. *Journal of the European Mathematical Society*, 17(10):2545–2593, 2015.
- [16] Serguei Yu. Popov. Frogs and some other interacting random walks models. *Discrete Mathematics and Theoretical Computer Science*, AC:277–288, 2003.
- [17] Alejandro F. Ramírez and Vladas Sidoravicius. Asymptotic behavior of a stochastic combustion growth process. *Journal of the European Mathematical Society*, 6(3):293–334, 2004.
- [18] Alistair Sinclair and Alexandre Stauffer. Mobile geometric graphs, and detection and communication problems in mobile wireless networks, 2010.
- [19] Alexandre Stauffer. Space-time percolation and detection by mobile nodes. *The Annals of Applied Probability*, 25(5):2416–2461, 2015.
- [20] Alain-Sol Sznitman. Vacant set of random interlacements and percolation. *Annals of mathematics*, pages 2039–2087, 2010.