



*Citation for published version:*

Tian, J, McGrogan, A & Jones, M 2022, 'Low carbon footprint inhalers in England: a review of dispensing data'.  
<https://doi.org/10.1093/ijpp/riac019.053>

*DOI:*

[10.1093/ijpp/riac019.053](https://doi.org/10.1093/ijpp/riac019.053)

*Publication date:*

2022

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication](#)

## University of Bath

### Alternative formats

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Low carbon footprint inhalers in England: a review of dispensing data

## Background

Metered dose inhalers (pMDIs) have a higher carbon footprint than low carbon inhalers (LCIs), such as dry powder inhalers. pMDIs contribute 3.5% of the NHS's CO<sub>2</sub> equivalent emissions. In 2019, NICE and BTS/SIGN guidelines attempted to increase use of LCIs, but their effects & factors influencing success are unknown.

**Aim:** To investigate temporal & geographical variation in LCI prescribing in England over 5 years.


## Methods

**Data**  
CCG dispensed items (March 2016-Feb 2021) from openprescribing.net  
CCG population characteristics from ONS, PHE & CCG websites

**Key measure**  
Low carbon inhaler % = LCI items relative to total inhaler items (pMDI + LCI)


**Statistical analysis**  
Interrupted time series analysis to investigate temporal variation. Multivariate regression models to investigate geographical variation

## Low carbon inhaler % by CCG



SABA & ICS: both advice on climate change in CCG guidelines & asthma prevalence associated with higher %LCI  
ICS & ICS+LABA: CCG population <15 years associated with lower %LCI

# Despite current initiatives, use of low carbon inhalers in England has fallen since 2019



Only a small increase in low carbon inhaler % seen after publication of NICE & British Thoracic Society guidelines, which was soon erased by the long-term trend

JT acknowledges the EPSRC Centre for Doctoral Training in Aerosol Science (EP/S023593/1) for financial support and the award of a China Scholarship Council (CSC)-University of Bristol joint-funded scholarship for PhD research

Presenter: **Jiangnan Tian**, **Anita McGrogan**, **Matthew Jones**  
@MatthewJonesUoB M.D.Jones@bath.ac.uk



Asthma icons created by Smashicons - Flaticon: <https://www.flaticon.com/free-icons/asthma>