



*Citation for published version:*

Bahia, PK, Pugh, V, Hoyland, K, Hensley, V, Rattray, M & Williams, RJ 2012, 'Neuroprotective effects of phenolic antioxidant tBHQ associate with inhibition of FoxO3a nuclear translocation and activity', *Journal of Neurochemistry*, vol. 123, no. 1, pp. 182-191. <https://doi.org/10.1111/j.1471-4159.2012.07877.x>

*DOI:*

[10.1111/j.1471-4159.2012.07877.x](https://doi.org/10.1111/j.1471-4159.2012.07877.x)

*Publication date:*

2012

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication](#)

© 2012 The Authors Journal of Neurochemistry © 2012 International Society for Neurochemistry

**University of Bath**

**Alternative formats**

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ORIGINAL  
ARTICLENeuroprotective effects of phenolic antioxidant  
tBHQ associate with inhibition of FoxO3a nuclear  
translocation and activityParmvir K. Bahia,<sup>\*,1</sup> Victoria Pugh,<sup>†</sup> Kimberley Hoyland,<sup>\*</sup>  
Victoria Hensley,<sup>\*</sup> Marcus Rattray<sup>†</sup> and Robert J. Williams<sup>\*,‡</sup><sup>\*</sup>Wolfson Centre for Age-Related Diseases, King's College London, London, UK<sup>†</sup>Reading School of Pharmacy, University of Reading, UK<sup>‡</sup>Department of Biology and Biochemistry, University of Bath, UK**Abstract**

The Forkhead transcription factor, FoxO3a induces genomic death responses in neurones following translocation from the cytosol to the nucleus. Nuclear translocation of FoxO3a is triggered by trophic factor withdrawal, oxidative stress and the stimulation of extrasynaptic NMDA receptors. Receptor activation of phosphatidylinositol 3-kinase (PI3K)–Akt signalling pathways retains FoxO3a in the cytoplasm, thereby inhibiting the transcriptional activation of death-promoting genes. We hypothesized that phenolic antioxidants such as tert-Butylhydroquinone (tBHQ), which is known to stimulate PI3K–Akt signalling, would inhibit FoxO3a translocation and activity. Treatment of cultured cortical neurones with NMDA increased the nuclear localization of FoxO3a, reduced the phosphorylation of FoxO3a, increased caspase activity and up-regulated Fas ligand expression. In contrast the phenolic antioxidant,

tBHQ, caused retention of FoxO3a in the cytosol coincident with enhanced PI3K- dependent phosphorylation of FoxO3a. tBHQ-induced nuclear exclusion of FoxO3a was associated with reduced FoxO-mediated transcriptional activity. Exposure of neurones to tBHQ inhibited NMDA-induced nuclear translocation of FoxO3a, prevented NMDA-induced up-regulation of FoxO-mediated transcriptional activity, blocked caspase activation and protected neurones from NMDA-induced excitotoxic death. Collectively, these data suggest that phenolic antioxidants such as tBHQ oppose stress-induced activation of FoxO3a and therefore have potential neuroprotective utility in neurodegeneration.

**Keywords:** caspase, dietary antioxidant, Fas ligand, motor neuron, NMDA receptor, PI3-kinase.

*J. Neurochem.* (2012) **123**, 182–191.

The mammalian homologue, class O of forkhead box transcription factors (FoxO) comprises four members (Fukunaga *et al.* 2005): *FoxO1*, *FoxO3*, *FoxO4* and *FoxO6*, and of these *FoxO1*, *FoxO3* and *FoxO6* expression has been detected in the brain (Furuyama *et al.* 2000; Biggs *et al.* 2001; Hoekman *et al.* 2006). FoxO binds to a core DNA consensus sequence, referred to as the Forkhead-response element (FHRE) to increase expression of pro-apoptotic genes (Furuyama *et al.* 2000). While FoxO activity is essential to normal programmed cell death during development, it has also been implicated in initiating apoptosis in cases of brain injury and in the progression of neurodegenerative disorders in the mature nervous system (Chong *et al.* 2005; Fukunaga *et al.* 2005; Su *et al.* 2009). There are several stressful triggers that stimulate nuclear translocation of FoxO in neurones including trophic factor withdrawal (Gilley *et al.* 2003; Barthelemy *et al.* 2004; Biswas *et al.*

2007), oxidative stress (Lehtinen *et al.* 2006), ischaemia (Fukunaga *et al.* 2005) and the activation of extrasynaptic NMDA receptors (Dick and Bading 2010). FoxO subsequently increases expression of genes encoding death

Received October 20, 2011; revised manuscript received July 11, 2012; accepted July 12, 2012.

Address correspondence and reprint requests to Dr Robert Williams, Department of Biology and Biochemistry, University of Bath, Claverton Down, Bath, BA2 7AY, UK. E-mail: r.j.williams@bath.ac.uk

<sup>1</sup>Present address: Department of Molecular Pharmacology and Physiology, University of South Florida, Tampa, USA.

Re-use of this article is permitted in accordance with the Terms and Conditions set out at [http://wileyonlinelibrary.com/onlineopen/OnlineOpen\\_Terms](http://wileyonlinelibrary.com/onlineopen/OnlineOpen_Terms)

**Abbreviations used:** BSA, bovine serum albumin; FHRE, Forkhead-response element; FoxO, Forkhead transcription factor; PBS, phosphate-buffered saline; tBHQ, tert-Butylhydroquinone.

proteins such as Bim (Bcl-2-interacting mediator of cell death) and Fas ligand (Brunet *et al.* 1999; Gilley *et al.* 2003; Barthelemy *et al.* 2004; Biswas *et al.* 2007).

Transcriptional activity of FoxO is partly determined by its subcellular location, which in turn is regulated by phosphorylation of three key residues at Thr-24, Ser-256 and Ser-319 (Nakae *et al.* 1999) lying downstream of PI3K–Akt signalling. FoxO3a phosphorylation at these sites causes it to be exported from the nucleus and retained in the cytoplasm in a Crm-1- and 14-3-3-dependent manner, thus rendering FoxO3a incapable of transcribing target apoptotic genes (Brunet *et al.* 1999). Neuroprotective phenolic antioxidants such as tBHQ and dietary flavonoids stimulate PI3K–Akt signalling and up-regulate Nrf2-dependent antioxidant response element activity (Johnson *et al.* 2002; Kraft *et al.* 2004; Schroeter *et al.* 2007; Bahia *et al.* 2008; Arredondo *et al.* 2010; Williams and Spencer 2012), but it is not known if they also inhibit FoxO3a activity to exert neuroprotective effects in neurones. The aim of this study, therefore, was to establish a link between FoxO3a activity and NMDA receptor-evoked neuronal cell death and to determine if phenolic antioxidants such as tBHQ could exert neuroprotection by opposing activation of this pro-apoptotic pathway. The work described here demonstrates that exposure of neurones to tBHQ inhibited NMDA-induced nuclear translocation of FoxO3a, prevented NMDA-induced up-regulation of FoxO-mediated transcriptional activity, blocked caspase activation and protected neurones from NMDA-induced excitotoxic death.

## Materials and methods

### Materials

NMDA and kainate were purchased from Tocris Bioscience (Bristol, UK). Tert-butyl hydroquinone (tBHQ) was purchased from Sigma-Aldrich (Poole, UK) and wortmannin from Calbiochem Corporation (La Jolla, CA, USA). Primary antibodies used were anti-FoxO3a and anti-phospho FoxO3a (Cell Signaling Technology, Danvers, MA, USA), anti-Fas ligand from Millipore (Watford, UK), anti-activated caspase 3 (Asp175) from New England Biolabs (Hitchin, UK), anti-beta Tubulin from Abcam (Cambridge, UK), anti-SMI-32R from Covance (Princeton, NJ, USA). Fluorescently labelled secondary antibodies, chicken anti-rabbit Alexa Fluor<sup>®</sup> 488 and goat anti-mouse Alexa Fluor<sup>®</sup> 568 were obtained from Invitrogen (Paisley, UK) and horseradish peroxidase-conjugated goat anti-rabbit antibody from Millipore. Cells were mounted in Mowiol<sup>®</sup> mounting medium with Hoechst 33342 (Calbiochem Corporation). The Dual-Glo luciferase assay system, pGL3 basic, pRL-TK renilla and restriction enzymes were from Promega (Madison, WI, USA). Cell culture reagents were obtained from Invitrogen and all other reagents from Sigma-Aldrich or Merck.

### Cell culture

Animals were maintained and used according to the U.K. Animals (Scientific Procedures) Act, 1986 and institutional ethical approval.

Animals were killed using cervical dislocation (Schedule 1 procedure) according to Home Office guidelines. Primary cultured neurones were prepared from 15 to 16-day-old Swiss mouse embryos (NIH, Harlan, UK).

### Cortical neurone culture

Cortical neurones were prepared essentially as described previously (Hoey *et al.* 2009). Briefly, dissociated cortical neurones were plated at 10<sup>6</sup> cells/mL into 24-well or six-well Nunc multiwell plates (Nunc, Naperville, IL, USA) that had been coated previously overnight with 15 µg/mL poly l-ornithine (Sigma-Aldrich). Neurones were seeded in phenol red-free Neurobasal B27-supplemented medium (Invitrogen). Cells were cultured at 37°C in a humidified atmosphere of 95% air and 5% CO<sub>2</sub> and experiments were performed at 7–9 DIV. Neurones for immunocytochemistry were cultured on 13-mm coverslips pre-coated with poly l-ornithine and 2 µg/mL laminin (Sigma-Aldrich).


### Motor neurone culture

Motor neurones were prepared essentially as described previously (Rainey-Smith *et al.* 2010). Spinal cords from embryos were dissected out into phosphate-buffered saline containing 0.6% glucose (PBS-gluc). The meninges were removed and the cords cut into ~5 mm<sup>3</sup> pieces. Up to five cords in 1.5 mL PBS-gluc were transferred to 15 mL centrifuge tubes. Trypsin (Invitrogen) was added to a final concentration of 0.05%. The tubes were then incubated at 37°C for 10 min with gentle agitation every 2 min. The supernatant was removed and the tissue resuspended in 1 mL L-15 medium containing 0.4% w/v bovine serum albumin (BSA) and 0.1 mg/mL DNase (D2; Worthington Biochemical Corporation, Lakewood, NJ, USA). Tissue was then agitated for 3 min, triturated and the supernatant transferred to a clean centrifuge tube. The pellet was then resuspended in 1 mL L-15 containing 0.4% w/v BSA and 20 µg/mL Dnase and triturated again and the supernatant collected. This process was repeated and the pooled supernatants were then transferred onto a BSA cushion (1.5 mL of L-15 containing 4% w/v BSA for the equivalent of five spinal cords) and spun for 5 min at 500 g in a Heraeus Biofuge Primo centrifuge (DJB Labcare, Bucks, UK). The supernatant was then removed and the pellet resuspended in 1 mL of culture medium consisting of phenol red-free Neurobasal<sup>™</sup> Medium supplemented with glutamine (2 mM), streptomycin (100 µg/mL and penicillin 100 u/mL) and B27 diluted 1 : 50. This was carefully layered onto 1.5 mL OptiPrep<sup>®</sup> density gradient medium (Axis-Shield UK, Cambridge, UK), diluted in L-15 to contain 10.4% w/v iodixanol, in a 15 mL centrifuge tube. The cells were then spun at 1000 g for 15 min with the centrifuge brake switched off. Cells visible at the medium/OptiPrep<sup>®</sup> interface were collected and spun through a second BSA cushion for 5 min at 1500 rpm. Between 20 000 and 30 000 cells were resuspended per millilitre of culture medium and strained through a 40 µm nylon strainer (VWR International Ltd, Leicestershire, UK). Cells were then plated out onto 13 mm coverslips coated with poly l-ornithine (1 h at 20°C) followed by a second incubation with culture medium containing 2 µg/mL laminin (Sigma-Aldrich).

### Plasmid constructs

*pFHRE-luc*: Synthesized oligos containing three repeats of the insulin response element (as described by Guo *et al.* 1999) were






**Fig. 1** NMDA inhibits FoxO3a phosphorylation, stimulates FoxO3a nuclear translocation and up-regulates FasL. Cortical neurones were treated with NMDA (50 μM) for 2 h, and FoxO3a phosphorylation and FoxO3a translocation measured by immunoblotting (b, d) and immunocytochemistry, respectively, (a) and FasL expression by immunoblotting only (c, e). Reduced dephosphorylation of both pFoxO3a ser<sup>253</sup> and pFoxO3a ser<sup>319/321</sup> was evident following exposure to NMDA (b, d) with no change in total levels of FoxO3a or β-tubulin, but was associated with an increase in the levels of FasL (c, e). Blots shown are representative of three independent experiments. (d) Data presented as mean band intensity (mean ± SEM, *n* = 3) expressed as pFoxO3a relative to total FoxO3a following exposure to Vehicle control

or NMDA (\**p* < 0.05, \*\**p* < 0.01). (e) Data presented as mean band intensity (mean ± SEM, *n* = 3) expressed as FasL relative to β-tubulin following exposure to Vehicle control or NMDA. To determine whether NMDA-mediated dephosphorylation of FoxO3a resulted in the nuclear accumulation of FoxO3a, cells were probed with an anti-FoxO3a antibody [(a), left and right panels] and counter-stained with Hoe 33342 (right panel only) to label nuclei. Control represents staining because of secondary antibody only. Under basal conditions, FoxO3a was distributed throughout the cytoplasm of cortical neurones with some diffuse staining evident in the nucleus (a, Vehicle). Following treatment with NMDA, FoxO3a staining was almost exclusively nuclear (a, NMDA). Scale bar = 20 μm.

phosphorylation at the other sites (Nakae *et al.* 2000) and the absence of S1 phosphorylation results in unmasking of the FoxO3a nuclear localization signal (Fukunaga *et al.* 2005). We first determined whether excitotoxic stimuli reduce FoxO3a phosphorylation and drive nuclear accumulation of FoxO3a in neurones. Cortical neurones were treated with NMDA (50 μM) for 2 h, and FoxO3a phosphorylation and FoxO3a translocation measured by immunoblotting and immunocytochemistry. There was significant dephosphorylation of both the S1 (pFoxO3a ser<sup>253</sup>) and S2 (pFoxO3a ser<sup>319/321</sup>) sites by 80% and 73%, respectively, relative to total FoxO3a levels following exposure to NMDA (Fig 1b and d) with no change in total levels of FoxO3a or β-tubulin (Fig 1b). To determine whether NMDA-mediated dephosphorylation of FoxO3a resulted in the nuclear accumulation of FoxO3a cells were probed with an anti-FoxO3a antibody (Fig 1a). Under control conditions FoxO3a was distributed

throughout the cytoplasm of cortical neurones with some diffuse staining evident in the nucleus (Fig 1a, Vehicle). Following treatment with NMDA, FoxO3a staining was almost exclusively nuclear (Fig 1a, NMDA). Kainate induced a similar redistribution of FoxO3 from cytoplasm to nucleus in cortical neurones (not shown) and also in motor neurones (Figure S1).

We next examined Fas ligand (FasL) expression. FasL is a member of the tumour necrosis factor (TNF) family, which binds to the Fas receptor to initiate cell death via caspase signalling cascades (e.g. Raoul *et al.* 1999). The FasL gene has three FHRE (Brunet *et al.* 1999), and induction of FasL by FoxO3A has been linked directly with neuronal death (Barthelemy *et al.* 2004). To determine whether excitotoxic signalling to FoxO3a resulted in increased expression of FasL, immunoblotting was performed on lysates from cortical neurones treated with NMDA. Low levels of FasL



**Fig. 2** tert-Butylhydroquinone (tBHQ) stimulates FoxO3a phosphorylation and promotes cytosolic accumulation of FoxO3a. Cortical neurones were treated with tBHQ (30  $\mu$ M; 2 h), or wortmannin (150 nM; 2 h) and FoxO3a phosphorylation and FoxO3a translocation were measured by immunoblotting and immunocytochemistry. tBHQ increased the phosphorylation of both the S1 (pFoxO3a ser<sup>253</sup>) and S2 (pFoxO3a ser<sup>319/321</sup>) sites without changing the total levels of FoxO3a or  $\beta$ -tubulin (b, d). Blots shown are representative of three independent experiments. (d) Data presented as mean band intensity (mean  $\pm$  SEM,  $n = 3$ ) expressed as pFoxO3a relative to total FoxO3a following exposure to Vehicle control or tBHQ ( $*p < 0.05$ ). Basal (vehicle) and tBHQ-induced phosphorylation of pFoxO3a ser<sup>253</sup> was abolished by wortmannin (c). To determine whether tBHQ-induced


phosphorylation of FoxO3a affected the cellular distribution of FoxO3a cells were probed with an anti-FoxO3a antibody [(a) left and right panels] and counter-stained with Hoechst 33342 (right panel only) to label nuclei. Under control conditions, FoxO3a was distributed throughout the cytoplasm of cortical neurones with some low-level diffuse staining evident in the nucleus (Fig 2, Vehicle). Following treatment with tBHQ, FoxO3a staining was strongly cytoplasmic (Fig 2, tBHQ). This was particularly evident when compared with cells treated with the PI3K inhibitor wortmannin, which resulted in pronounced nuclear accumulation of FoxO3a (Fig 2, Wortmannin). Images shown are representative of 100 individual cells. tBHQ caused strong cytoplasmic accumulation of FoxO3a in 45% of stained cells. Scale bar = 20  $\mu$ m.

protein could be detected under control conditions and exposure to NMDA (50  $\mu$ M, 2 h) caused a two-fold increase in FasL (Fig 1c and e), suggesting that following translocation to the nucleus, FoxO3a up-regulates FasL in cortical neurones.

#### tBHQ enhances phosphorylation and cytoplasmic retention of FoxO3a

Having demonstrated that NMDA caused FoxO3a dephosphorylation and translocation to the nucleus, and induced FasL, we wished to determine whether neuroprotective phenolic antioxidants such as tBHQ could suppress FoxO3a

activation. It was not possible to demonstrate tBHQ inhibition of FasL expression as the basal levels in neurones were too low (see Fig 1c). Cortical neurones were treated with tBHQ (30  $\mu$ M, 2 h), and FoxO3a phosphorylation and FoxO3a translocation were measured by immunoblotting and immunocytochemistry. tBHQ induced a clear trend towards increased phosphorylation of both the S1 (pFoxO3a ser<sup>253</sup>) and S2 (pFoxO3a ser<sup>319/321</sup>) sites by 149% and 151%, respectively, relative to total foxO3a levels without changing the total levels of FoxO3a or  $\beta$ -tubulin (Fig 2b). To determine whether tBHQ-mediated phosphorylation of FoxO3a affected the cellular distribution of FoxO3a, cells



**Fig. 3** Inhibition of FKHD-mediated gene expression by tert-Butylhydroquinone (tBHQ). Cortical neurones (8 DIV) were transfected with pARE-luc (a, b) or FHRE-luc (c, d) and then treated with, 10 μM or 30 μM tBHQ for 24 h (a and c) or 30 μM tBHQ for different time periods as shown (b and d). Data represent mean ± SEM, analysed using one-way ANOVA and Bonferroni *post hoc* test ( $n = 3$ , each concentration performed in triplicate) where \* $p < 0.05$ , \*\* $p < 0.01$  \*\*\* $p < 0.001$  compared to vehicle control.

were probed with an anti-FoxO3a antibody (Fig 2a). Under control conditions, FoxO3a was distributed throughout the cytoplasm of cortical neurones with some low-level diffuse staining evident in the nucleus (Fig 2, Vehicle). Following treatment with tBHQ, FoxO3a staining was strongly cytoplasmic (Fig 2, tBHQ). This was particularly evident when compared with cells treated with the PI3K inhibitor, wortmannin, which resulted in pronounced nuclear accumulation of FoxO3a (Fig 2, Wortmannin). Wortmannin abolished basal (vehicle) and tBHQ-mediated FoxO3a phosphorylation (Fig 2c) and prevented tBHQ-induced cytoplasmic retention of FoxO3a (not shown).


#### tBHQ enhances antioxidant response element- and inhibits forkhead-response element- activity in neurones

To ascertain whether the cytoplasmic retention of FoxO3a by tBHQ is accompanied by a decrease in FoxO activity, a luciferase assay was employed to examine changes in gene expression downstream of a forkhead-response element (FHRE-luc). To first define the experimental parameters, the effect of tBHQ on ARE activity were examined, as ARE is a well-characterized target of tBHQ (Kraft *et al.* 2004; Shih *et al.* 2005; Bahia *et al.* 2008). tBHQ caused a concentration- (Fig 3a) and time-dependent (Fig 3b) increase in ARE-mediated luciferase expression in neurones with maximal effects observed at 30 μM tBHQ for 24 h. In contrast, tBHQ

caused a significant decrease in basal levels of FHRE activity with maximum effects again observed at 30 μM tBHQ for 24 h (Fig 3c and d).

#### tBHQ inhibits NMDA-induced nuclear translocation of FoxO3a and reduces caspase 3A activation

We next tested if the induction of apoptotic markers following treatment with NMDA correlated with increase in nuclear FoxO3a and also whether NMDA-dependent caspase 3 activation and FoxO3a nuclear localization could be abrogated by tBHQ. As expected, NMDA (50 μM, 2–18 h) caused a time-dependent elevation in neurones containing activated caspase 3 (Fig 4a, top panel). Relative to vehicle controls, the number of cells containing activated caspase after NMDA treatment were  $291 \pm 93\%$ ,  $290 \pm 82\%$ ,  $211 \pm 35\%$  and  $151 \pm 42\%$  at 2, 4, 6 and 18 h, respectively ( $n = 6–9$ ), which was statistically significant at the 6-h time point (Fig 4b). At all time points, the number of activated caspase-positive cells in the presence of tBHQ (30 μM) and NMDA (50 μM) was lower than the number of activated caspase-positive cells in the presence of NMDA alone, and at 6 h, the tBHQ reduction of NMDA-induced caspase activation was highly significant ( $40 \pm 10\%$  of NMDA-treated numbers,  $n = 9$ ). The phenolic antioxidant tBHQ in its own right did not significantly decrease the number of caspase-positive cells at any time point. We note



**Fig. 4** NMDA induction of activated caspase is accompanied by accumulation of nuclear FoxO3A and blocked by tert-Butylhydroquinone (tBHQ). (a) Representative photomicrograph showing cortical neurons (8DIV) after 6-h treatment with 0.1% ethanol vehicle (C), NMDA (50 μM) (N), tBHQ (30 μM) (T), or NMDA and tBHQ combined (N + T). Top panel shows staining for activated caspase 3 (green), and nuclear Hoe 33342 (blue). Bottom panel shows staining for foxO3A (green) and Hoe 33342 (blue), arrowheads indicate cells with strongly foxO3A-positive nuclei. Scale bar = 10 μm. (b) shows quan-


tification of nuclei associated with activated caspase 3 at different time points after application of 0.1% ethanol vehicle (C), NMDA (50 μM) (N), tBHQ (30 μM) (T) or tBHQ and NMDA combined (T + N). (c) shows quantification of foxO3A-positive nuclei after the same treatments. In B + C, Bars show means and SEM derived from counts of six to nine fields from three experiments. Analysis was by ANOVA and Bonferroni's multiple comparison test: \*, \*\*, \*\*\* =  $p < 0.05$ ,  $p < 0.01$ ,  $p < 0.001$  compared to vehicle control. #, ##, ### =  $p < 0.05$ ,  $p < 0.01$ ,  $p < 0.001$  compared to NMDA treatment only.

that, in these experiments, longer incubation times resulted in a time-dependent increase in the number of activated caspase 3-positive cells, even in the absence of any stimulation, suggesting some cell death in these cultures. For example, at 18 h, the vehicle control cultures had  $361 \pm 85\%$  ( $n = 9$ ) more caspase-positive cells compared with the vehicle control cultures at 2 h. In parallel with induction of the apoptotic marker, activated caspase 3, NMDA-induced nuclear foxO3A (Fig. 4a). The NMDA-induced increase in cells with nuclear foxO3A were  $239 \pm 27\%$ ,  $188 \pm 26\%$ ,  $295 \pm 32\%$  and  $312 \pm 70\%$  to the level in vehicle-treated controls, at 2, 4, 6 and 18 h respectively (Fig 4c,  $n = 9$ ). This increase was statistically significant at all time points tested. At all time points, tBHQ prevented the NMDA-induced increase in foxo3A. Co-administration of tBHQ with NMDA significantly reduced in the number of cells with nuclear foxO3A to  $51 \pm 12\%$ ,  $49 \pm 16\%$ ,  $39 \pm 6\%$  and  $30 \pm 6\%$  to the levels with NMDA alone at 2, 4, 6 and 18 h respectively (Fig 4c,  $n = 6-9$ ) tBHQ in its own right did not cause a significant change in the number of cells with nuclear foxO3A.

#### tBHQ opposes NMDA-induced increase in forkhead-response element-activity and protects against NMDA-evoked toxicity

We further tested if tBHQ opposed NMDA effects via regulation of FHRE activity. Neurons were treated with NMDA (50 μM; 6 h) in the presence and absence of tBHQ (30 μM). NMDA increased FHRE activity consistent with the observed increase in nuclear FoxO3a and this increase was attenuated by co-treatment with tBHQ (30 μM) (Fig 5a). We note that longer term exposure (24 h) to NMDA (50 μM) induced neuronal loss, which confounded the luciferase expression data. Short-term exposure (20 min) to NMDA inhibited FHRE activity (data not shown), consistent with previous reports of pre-conditioning-dependent neuroprotection (Soriano *et al.* 2006).

Finally, to determine whether tBHQ actions can confer protection against excitotoxicity, neuronal viability was examined following exposure to NMDA (3–300 μM, 24 h) in the presence or absence of tBHQ (30 μM). Exposure to NMDA caused concentration-dependent neuronal cell damage as assessed by a reduction in MTT turnover.



**Fig. 5** tert-Butylhydroquinone (tBHQ) opposes NMDA-induced up-regulation of FKHD-mediated gene expression and protects against NMDA-evoked neuronal damage. (a) Cortical neurones (8 DIV) were transfected with FHRE-luc and were then treated with NMDA (50  $\mu$ M; 6 h; ■) or vehicle control (□) in the presence and absence of tBHQ (30  $\mu$ M). Data represent mean  $\pm$  SEM, analysed using one-way ANOVA and Bonferroni *post hoc* test ( $n = 3$ , each concentration performed in triplicate). NMDA increased FHRE activity compared with vehicle control \* $p < 0.05$  and tBHQ inhibited FKHR compared to vehicle control # $p < 0.05$ . (b) Cultured cortical neurones (8 DIV) were incubated with increasing concentrations of NMDA in the presence of 30  $\mu$ M tBHQ (▲) or vehicle control (■) for 24 h. Cell viability was assessed using the MTT assay where the absorbance of reduced MTT was read at 505 nm. Neurones co-treated with tBHQ were significantly protected against 100 and 300  $\mu$ M NMDA insults. Data represent mean  $\pm$  SEM, analysed using one-way ANOVA and the Bonferroni *post hoc* test ( $n = 4$ ) where \*\* $p < 0.01$  for individual NMDA concentrations versus control.

Co-treatment with tBHQ significantly inhibited NMDA-induced reduction in MTT turnover, consistent with tBHQ eliciting a neuroprotective effect. (Fig 5b).

## Discussion

FoxO3a activity is enhanced in neurones principally by a loss of trophic support (Gilley *et al.* 2003), but here we have shown that prolonged exposure of cortical neurones to NMDA also appears to enhance nuclear accumulation of

FoxO3a. This is in general agreement with a recent report that extrasynaptic NMDA receptor stimulation enhances FoxO3a nuclear translocation in hippocampal neurones (Dick and Bading 2010). Enhanced nuclear translocation of FoxO3a is consistent with an NMDA-stimulated loss of PI3-kinase-dependent phosphorylation of FoxO3a, which has also been observed following exposure to high concentrations of glutamate (Zheng and Quirion 2009). Here, we show a correlation between NMDA-induced loss of FoxO3a phosphorylation and FoxO3 nuclear localization, which is reflected in an increase in FHRE-mediated gene expression, induction of a pro-apoptotic protein known to be regulated directly by FoxO proteins, FasL, and NMDA-induced cell death. In contrast, calcium-signalling evoked by stimulation of synaptic NMDA receptors has been shown to protect against FoxO3a translocation (Dick and Bading 2010), indeed, we observed inhibition of FHRE-mediated gene expression upon short-term exposure to NMDA (not shown), which is consistent with neuroprotective PI3-kinase-dependent Akt signalling activity associated with acute activation of this receptor population (Perkinton *et al.* 2002; Soriano *et al.* 2006). It is likely, therefore, that the switch to increased FHRE-mediated gene expression and pro-death signalling observed following prolonged exposure to NMDA results from down-regulation of the protective PI3-kinase-dependent element of the pathway and recruitment of the extrasynaptic NMDA receptor population in a similar manner to the well-characterized CREB shut-off pathway (Hardingham and Bading 2010).

As tBHQ as well as various dietary polyphenolics stimulate PI3-kinase-dependent Akt phosphorylation and up-regulate Nrf2-mediated antioxidant response element activity (Kang *et al.* 2002; Kraft *et al.* 2004; Schroeter *et al.* 2007; Bahia *et al.* 2008; Arredondo *et al.* 2010), and tBHQ has been reported to protect motor neurones against FasL-induced neurodegeneration (Pehar *et al.* 2007), we considered that tBHQ might also suppress FoxO3a nuclear translocation, which might underlie some of its neuroprotective efficacy. Accordingly, we found that tBHQ stimulated PI3-kinase-dependent phosphorylation of FoxO3a in neurones and this correlated with its retention in the cytosol against a basal low-level accumulation of FoxO3a in the nucleus. Furthermore, tBHQ effectively suppressed NMDA-dependent FoxO3a nuclear accumulation and FHRE activity, attenuated NMDA-dependent activation of caspase 3 in neurones and reduced NMDA-induced neuronal death. Although these observations are strictly correlative, they are strongly suggestive of a very close relationship between regulation of FoxO3a activity and neuronal cell death and identify tBHQ and compounds of this type, for example dietary polyphenolic compounds, as potential suppressors of FoxO-dependent cell death. Our findings have been carried out in neuronal cultures with very few astrocytes present (< 5%), suggesting that antioxidant signalling to prevent

FoxO activation and neuronal death are mediated by pathways intrinsic to neurones, nevertheless we cannot completely exclude at this time that tBHQ is acting on astrocytes to confer neuronal protection. The precise mechanisms underlying these protective effects are not yet clear, but there is emerging evidence that deacetylation and methylation regulate the stability and activity of FoxO3 (Wang *et al.* 2012). Most notably, oxidative stress-evoked neuronal apoptosis is inhibited by Set9-mediated lysine methylation of FoxO3 (Xie *et al.* 2012) and this therefore represents a potential target for the protective actions of tBHQ and dietary antioxidants.

Our study supports emerging data, which suggest that FoxO3A is a key mediator of the progressive neuronal cell loss associated with neurodegeneration and with other disorders of the nervous system. For example, increased FoxO3a has been found localized to Lewy bodies and Lewy neurites in Parkinson's disease brain tissue (Su *et al.* 2009) and activation of FoxO by LRRK2, associated with autosomal-dominant late-onset Parkinson's disease, enhances neuronal cell death in *Drosophila* (Kanao *et al.* 2010). With respect to Alzheimer's disease, inactivation of FoxO3a activity is correlated with reduced Alzheimer's disease-like pathology and with preservation of spatial reference memory in Tg2576 mice (Qin *et al.* 2008). This implies that APP over-expression activates FoxO3a in brain regions associated with spatial memory, which is consistent with the reduced levels of Akt phosphorylation observed in the hippocampus of APP<sup>swe</sup> mutant mice (Abbott *et al.* 2008). In motor neurons, FoxO3a activity is thought to contribute to neuronal cell death (Brunet *et al.* 1999; Barthelemy *et al.* 2004) and similarly we interpret our observed increase in nuclear FoxO3a in motor neurons as a contributing factor to apoptosis; however, this is not consistent with findings that stimulating FoxO3a activity is broadly neuroprotective in *C-elegans*, *Drosophila* and in mouse motor neurons (Mojsilovic-Petrovic *et al.* 2009). It is therefore possible that the enhanced nuclear levels of FoxO3a observed in our motor neurons represent a failed neuroprotective response. Generally, differences in beneficial versus harmful effects of FoxO3a are more likely to relate to cell-circumstance-specific level of expression, activation and post-translational modification (Mojsilovic-Petrovic *et al.* 2009).

On the basis of our data, we propose, therefore, that tBHQ and perhaps other dietary polyphenolics, in addition to their well-documented effects on antioxidant response element-mediated gene expression, exert neuroprotection by regulating the localization and activity of FoxO3a, thereby opposing apoptosis. In this context, we note that a number of flavonoids are also capable of modestly reducing FHRE activity (data not shown), suggesting that regulation of nuclear translocation of FoxO3A might be a common feature of molecules possessing antioxidant activity. Indeed, our data predict that activation of the antioxidant response element is

accompanied by reciprocal regulation of FHRE. We do not yet know if these observations are mechanistically related, but hypoxic pre-conditioning which up-regulates an antioxidant response protects against transient global ischaemia by increasing FoxO phosphorylation (Zhan *et al.* 2010) and signalling pathways linking oxidative stress to neuronal cell death have previously been attributed to FoxO activation (Lehtinen *et al.* 2006; Yuan *et al.* 2009), so the link between these responses is compelling. In summary, our data support a role for FoxO3a in NMDA-evoked neuronal apoptosis and suggest that phenolic antioxidants such as tBHQ oppose stress-induced activation of FoxO3a and therefore may have potential neuroprotective utility in neurodegeneration.

## Acknowledgements

This study was supported by the Motor Neurone Disease Association (Williams 6030). MR acknowledges the support of the ALS Association (Grant Z6BBIO). KH and VH were recipients of Wellcome Trust Summer Studentships. The authors declare no conflicts of interest.

## Supporting information

Additional supporting information may be found in the online version of this article:

**Figure S1.** Kainate drives nuclear accumulation of Foxo3a in motor neurones.

As a service to our authors and readers, this journal provides supporting information supplied by the authors. Such materials are peer-reviewed and may be re-organized for online delivery, but are not copy-edited or typeset. Technical support issues arising from supporting information (other than missing files) should be addressed to the authors.

## References

- Abbott J. J., Howlett D. R., Francis P. T. and Williams R. J. (2008) Abeta(1-42) modulation of Akt phosphorylation via alpha7 nAChR and NMDA receptors. *Neurobiol. Aging* **29**, 992–1001.
- Arredondo F., Echeverry C., Abin-Carriquiry J. A., Blasina F., Antúnez K., Jones D. P., Go Y. M., Liang Y. L. and Dajas F. (2010) After cellular internalization, quercetin causes Nrf2 nuclear translocation, increases glutathione levels, and prevents neuronal death against an oxidative insult. *Free Radic. Biol. Med.* **49**, 738–747.
- Bahia P. K., Rattray M. and Williams R. J. (2008) Dietary flavonoid (-)epicatechin stimulates phosphatidylinositol 3-kinase-dependent anti-oxidant response element activity and up-regulates glutathione in cortical astrocytes. *J. Neurochem.* **106**, 2194–2204.
- Barthelemy C., Henderson C. and Pettmann B. (2004) Foxo3a induces motoneuron death through the Fas pathway in cooperation with JNK. *BMC Neurosci.* **5**, 48.
- Biggs III W. H., Cavenee W. K. and Arden K. C. (2001) Identification and characterization of members of the FKHR (FOX O) subclass of winged-helix transcription factors in the mouse. *Mamm. Genome* **12**, 416–425.
- Biswas S. C., Shi Y., Sproul A. and Greene L. A. (2007) Pro-apoptotic Bim induction in response to nerve growth factor deprivation requires simultaneous activation of three different death signaling pathways. *J. Biol. Chem.* **282**, 29368–29374.

- Brunet A., Bonni A., Zigmond M. J., Lin M. Z., Juo P., Hu L. S., Anderson M. J., Arden K. C., Blenis J. and Greenberg M. E. (1999) Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* **96**, 857–868.
- Chong Z. Z., Li F. and Maiese K. (2005) Oxidative stress in the brain: novel cellular targets that govern survival during neurodegenerative disease. *Prog. Neurobiol.* **75**, 207–246.
- Dick O. and Bading H. (2010) Synaptic activity and nuclear calcium signaling protect hippocampal neurons from death signal-associated nuclear translocation of FoxO3a induced by extrasynaptic N-methyl-D-aspartate receptors. *J. Biol. Chem.* **285**, 19354–19361.
- Fukunaga K., Ishigami T. and Kawano T. (2005) Transcriptional regulation of neuronal genes and its effect on neural functions: expression and function of forkhead transcription factors in neurons. *J. Pharmacol. Sci.* **98**, 205–211.
- Furuyama T., Nakazawa T., Nakano I. and Mori N. (2000) Identification of the differential distribution patterns of mRNAs and consensus binding sequences for mouse DAF-16 homologues. *Biochem. J.* **349**, 629–634.
- Gilley J., Coffer P. J. and Ham J. (2003) FOXO transcription factors directly activate bim gene expression and promote apoptosis in sympathetic neurons. *J. Cell Biol.* **162**, 613–622.
- Guo S., Rena G., Cichy S., He X., Cohen P. and Unterman T. (1999) Phosphorylation of serine 256 by protein kinase B disrupts transactivation by FKHR and mediates effects of insulin on insulin-like growth factor-binding protein-1 promoter activity through a conserved insulin response sequence. *J. Biol. Chem.* **274**, 17184–17192.
- Hardingham G. E. and Bading H. (2010) Synaptic versus extrasynaptic NMDA receptor signalling: implications for neurodegenerative disorders. *Nat. Rev. Neurosci.* **10**, 682–696.
- Hoekman M. F., Jacobs F. M., Smidt M. P. and Burbach J. P. (2006) Spatial and temporal expression of FoxO transcription factors in the developing and adult murine brain. *Gene Expr. Patterns* **6**, 134–140.
- Hoey S. E., Williams R. J. and Perkinson M. S. (2009) Synaptic NMDA receptor activation stimulates alpha-secretase amyloid precursor protein processing and inhibits amyloid-beta production. *J. Neurosci.* **29**, 4442–4460.
- Johnson D. A., Andrews G. K., Xu W. and Johnson J. A. (2002) Activation of the antioxidant response element in primary cortical neuronal cultures derived from transgenic reporter mice. *J. Neurochem.* **81**, 1233–1241.
- Kanao T., Venderova K., Park D. S., Unterman T., Lu B. and Imai Y. (2010) Activation of FoxO by LRRK2 induces expression of proapoptotic proteins and alters survival of postmitotic dopaminergic neuron in *Drosophila*. *Hum. Mol. Genet.* **19**, 3747–3758.
- Kang K. W., Lee S. J., Park J. W. and Kim S. G. (2002) Phosphatidylinositol 3-kinase regulates nuclear translocation of NF-E2-related factor 2 through actin rearrangement in response to oxidative stress. *Mol. Pharmacol.* **62**, 1001–1010.
- Kraft A. D., Johnson D. A. and Johnson J. A. (2004) Nuclear factor E2-related factor 2-dependent antioxidant response element activation by tert-butylhydroquinone and sulforaphane occurring preferentially in astrocytes conditions neurons against oxidative insult. *J. Neurosci.* **24**, 1101–1112.
- Lehtinen M. K., Yuan Z., Boag P. R. *et al.* (2006) A conserved MST-FOXO signaling pathway mediates oxidative-stress responses and extends life span. *Cell* **125**, 987–1001.
- Mojisilovic-Petrovic J., Nedelsky N., Boccitto M. *et al.* (2009) FOXO3a is broadly neuroprotective in vitro and in vivo against insults implicated in motor neuron diseases. *J. Neurosci.* **29**, 8236–8247.
- Nakae J., Park B. C. and Accili D. (1999) Insulin stimulates phosphorylation of the forkhead transcription factor FKHR on serine 253 through a Wortmannin-sensitive pathway. *J. Biol. Chem.* **274**, 15982–15985.
- Nakae J., Barr V. and Accili D. (2000) Differential regulation of gene expression by insulin and IGF-1 receptors correlates with phosphorylation of a single amino acid residue in the forkhead transcription factor FKHR. *EMBO J.* **19**, 989–996.
- Peihar M., Vargas M. R., Robinson K. M., Cassina P., Díaz-Amarilla P. J., Hagen T. M., Radi R., Barbeito L. and Beckman J. S. (2007) Mitochondrial superoxide production and nuclear factor erythroid 2-related factor 2 activation in p75 neurotrophin receptor-induced motor neuron apoptosis. *J. Neurosci.* **27**, 7777–7785.
- Perkinson M. S., Ip J. K., Wood G. L., Crosssthaite A. J. and Williams R. J. (2002) Phosphatidylinositol 3-kinase is a central mediator of NMDA receptor signalling to MAP kinase (Erk1/2), Akt/PKB and CREB in striatal neurones. *J. Neurochem.* **80**, 239–254.
- Qin W., Zhao W., Ho L., Wang J., Walsh K., Gandy S. and Pasinetti G. M. (2008) Regulation of forkhead transcription factor FoxO3a contributes to calorie restriction-induced prevention of Alzheimer's disease-type amyloid neuropathology and spatial memory deterioration. *Ann. N.Y. Acad. Sci.* **1147**, 335–347.
- Rainey-Smith S. R., Andersson D. A., Williams R. J. and Rattray M. (2010) Tumour necrosis factor alpha induces rapid reduction in AMPA receptor-mediated calcium entry in motor neurones by increasing cell surface expression of the GluR2 subunit: relevance to neurodegeneration. *J. Neurochem.* **113**, 692–703.
- Raoul C., Henderson C. E. and Pettmann B. (1999) Programmed cell death of embryonic motoneurons triggered through the Fas death receptor. *J. Cell Biol.* **147**, 1049–1062.
- Schroeter H., Bahia P., Spencer J. P., Sheppard O., Rattray M., Cadenas E., Rice-Evans C. and Williams R. J. (2007) (-)Epicatechin stimulates ERK-dependent cyclic AMP response element activity and up-regulates GluR2 in cortical neurons. *J. Neurochem.* **101**, 1596–1606.
- Shih A. Y., Li P. and Murphy T. H. (2005) A small-molecule-inducible Nrf2-mediated antioxidant response provides effective prophylaxis against cerebral ischemia in vivo. *J. Neurosci.* **25**, 10321–10335.
- Soriano F. X., Papadia S., Hofmann F., Hardingham N. R., Bading H. and Hardingham G. E. (2006) Preconditioning doses of NMDA promote neuroprotection by enhancing neuronal excitability. *J. Neurosci.* **26**, 4509–4518.
- Su B., Liu H., Wang X. *et al.* (2009) Ectopic localization of FOXO3a protein in Lewy bodies in Lewy body dementia and Parkinson's disease. *Mol. Neurodegener.* **4**, 32.
- Wang F., Chan C. H., Chen K., Guan X., Lin H. K. and Tong Q. (2012) Deacetylation of FOXO3 by SIRT1 or SIRT2 leads to Skp2-mediated FOXO3 ubiquitination and degradation. *Oncogene* **31**, 1546–1557.
- Williams R. J. and Spencer J. P. (2012) Flavonoids, cognition, and dementia: actions, mechanisms, and potential therapeutic utility for Alzheimer disease. *Free Radic. Biol. Med.* **52**, 35–45.
- Xie Q., Hao Y., Tao L., Peng S., Rao C., Chen H., You H., Dong M. Q. and Yuan Z. (2012) Lysine methylation of FOXO3 regulates oxidative stress-induced neuronal cell death. *EMBO Rep.* **13**, 371–377.
- Yuan Z., Lehtinen M. K., Merlo P., Villén J., Gygi S. and Bonni A. (2009) Regulation of neuronal cell death by MST1-FOXO1 signaling. *J. Biol. Chem.* **284**, 11285–11292.
- Zhan L., Wang T., Li W., Xu Z. C., Sun W. and Xu E. (2010) Activation of Akt/FoxO signaling pathway contributes to induction of neuroprotection against transient global cerebral ischemia by hypoxic pre-conditioning in adult rats. *J. Neurochem.* **114**, 897–908.
- Zheng W. H. and Quirion R. (2009) Glutamate acting on N-methyl-D-aspartate receptors attenuates insulin-like growth factor-1 receptor tyrosine phosphorylation and its survival signaling properties in rat hippocampal neurons. *J. Biol. Chem.* **284**, 855–861.