



Citation for published version:

Ignat, R & Moser, R 2022, 'Separation of domain walls with nonlocal interaction and their renormalised energy by Γ -convergence in thin ferromagnetic films', *Journal of Differential Equations*, vol. 339, pp. 395-475.
<https://doi.org/10.1016/j.jde.2022.08.024>

DOI:

[10.1016/j.jde.2022.08.024](https://doi.org/10.1016/j.jde.2022.08.024)

Publication date:

2022

Document Version

Early version, also known as pre-print

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Separation of domain walls with nonlocal interaction and their renormalised energy by Γ -convergence in thin ferromagnetic films

Radu Ignat * Roger Moser†

July 13, 2020

Abstract

We analyse two variants of a nonconvex variational model from micromagnetics with a nonlocal energy functional, depending on a small parameter $\epsilon > 0$. The model gives rise to transition layers, called Néel walls, and we study their behaviour in the limit $\epsilon \rightarrow 0$. The analysis has some similarity to the theory of Ginzburg-Landau vortices. In particular, it gives rise to a renormalised energy that determines the interaction (attraction or repulsion) between Néel walls to leading order. But while Ginzburg-Landau vortices show attraction for degrees of the same sign and repulsion for degrees of opposite signs, the pattern is reversed in this model.

In a previous paper, we determined the renormalised energy for one of the models studied here under the assumption that the Néel walls stay separated from each other. In this paper, we present a deeper analysis that in particular removes this assumption. The theory gives rise to an effective variational problem for the positions of the walls, encapsulated in a Γ -convergence result. In the second part of the paper, we turn our attention to another, more physical model, including an anisotropy term. We show that it permits a similar theory, but the anisotropy changes the renormalised energy in unexpected ways and requires different methods to find it.

Keywords: domain walls, repulsive/attractive interaction, renormalised energy, compactness, Γ -convergence.

Contents

1	Introduction	2
1.1	Background	2
1.2	The confined problem	2
1.3	The unconfined problem	9
1.4	Comparison between the confined and the unconfined model	11
1.5	Comparison with other Ginzburg-Landau models	12
1.6	Representations of the stray field energy	12
2	Analysis for the confined problem	13
2.1	Compactness	13
2.2	Blow-up of the renormalised energy	16
2.3	Energy estimates	21
2.4	Proofs of the main results	29

*Institut de Mathématiques de Toulouse & Institut Universitaire de France, UMR 5219, Université de Toulouse, CNRS, UPS IMT, F-31062 Toulouse Cedex 9, France. Email: Radu.Ignat@math.univ-toulouse.fr

†Department of Mathematical Sciences, University of Bath, Bath BA2 7AY, UK. Email: r.moser@bath.ac.uk

3	Analysis for the unconfined problem	33
3.1	The function I	33
3.2	The limiting stray field potential: construction	34
3.3	The limiting stray field potential: further properties	40
3.4	Preliminary estimates	46
3.5	Proof of Theorem 8	50
	3.5.1 Preparation	50
	3.5.2 A lower bound for the interaction energy	51
	3.5.3 An upper bound for the interaction energy	54
3.6	Renormalised energy, separation and Γ -convergence	57

1 Introduction

1.1 Background

According to the theory of micromagnetics, the magnetisation in ferromagnetic materials is given by a vector field of constant length that obeys a variational principle. The underlying energy functional contains several competing terms of rather different natures. In many contexts, it gives rise to regions with approximately constant or slowly varying magnetisation, separated by transition layers where the magnetisation varies rapidly. Néel walls are transition layers of this type found in soft thin films and are one of the most important features there.

We study a model for Néel walls derived from the micromagnetic energy functional (discussed in detail, e.g., by Hubert-Schäfer [12] or by De Simone-Kohn-Müller-Otto [10]), but including some simplifications. Most significantly, our model depends on the assumption that all the Néel walls are parallel to each other, and they are one-dimensional transition layers where the magnetisation varies in-plane. As a result, the magnetisation is described by a map

$$m: J \rightarrow \mathbb{S}^1,$$

where J is an interval. We consider two different, but closely related, problems in this paper. The crucial difference between them is that $J = (-1, 1)$ in one case and $J = \mathbb{R}$ in the other case. For reasons that are explained below, the different domains will necessitate a somewhat different set-up of the problem, and therefore we explain the two problems separately.

1.2 The confined problem

We begin with the case where $J = (-1, 1)$, i.e., the whole magnetisation profile is confined to the fixed interval $(-1, 1)$. Then $m: (-1, 1) \rightarrow \mathbb{S}^1$ represents the in-plane magnetisation vector field, renormalised to unit length. As we have a boundary, our problem requires boundary conditions. For this purpose, we fix the angle

$$\alpha \in (0, \pi)$$

and require that

$$m_1(-1) = m_1(1) = \cos \alpha.$$

This will fix $m_2(\pm 1)$ up to the sign as well, but we allow either sign. It is convenient to extend m_1 by $\cos \alpha$ outside of $(-1, 1)$. The energy functional depends on a parameter $\epsilon > 0$ and is of the form

$$E_\epsilon(m) = \frac{\epsilon}{2} \int_{-\infty}^{\infty} |m'|^2 dx_1 + \frac{1}{2} \|m_1 - \cos \alpha\|_{\dot{H}^{1/2}(\mathbb{R})}^2, \quad (1)$$

where $\dot{H}^{1/2}(\mathbb{R})$ is the homogeneous Sobolev space of fractional order 1/2 and we write x_1 for the independent variable for reasons that will become clear later. For details about the background of

this model, and how it is derived from the full micromagnetic energy, we refer to previous work [8, 9, 30, 31, 7, 21, 16, 5, 29, 32, 18, 19, 20]. We note, however, that the first term on the right hand side in (1) is called exchange energy and the second term is called stray field energy or magnetostatic energy in the theory of micromagnetics.

We emphasise that the second term in E_ϵ is of nonlocal nature. This makes the analysis of the problem challenging, but the nonlocal nature is also responsible for the structure of the transition layers. When ϵ is small, the energy favours magnetisations such that $m_1 - \cos \alpha$ is small in the $\dot{H}^{1/2}$ -sense, but for topological reasons, it may not be able to vanish identically. Then we expect transitions between the points $(\cos \alpha, \pm \sin \alpha)$ on \mathbb{S}^1 , and these are the Néel walls that we study.

Since any continuous transition between $(\cos \alpha, \pm \sin \alpha)$ requires that one of the points $(\pm 1, 0)$ on \mathbb{S}^1 be attained, we may use the preimages of these points under m (i.e., the zeros of m_2) as a proxy for the locations of the Néel walls. There are two different types of Néel walls, depending on the sign of m_1 at such a point. We are interested mainly in configurations with several Néel walls, located at certain points a_1, \dots, a_N with $a_1 < \dots < a_N$. Thus given $N \in \mathbb{N} \cup \{0\}$, we define¹

$$A_N = \{a \in (-1, 1)^N : -1 < a_1 < \dots < a_N < 1\}.$$

An element $a = (a_1, \dots, a_N) \in A_N$ will typically be accompanied by $d \in \{\pm 1\}^N$, which indicates the type of the Néel walls. That is, the pair (a, d) gives rise to the condition $m_1(a_n) = d_n$ for $n = 1, \dots, N$. We define

$$M(a, d) = \{m \in H^1((-1, 1); \mathbb{S}^1) : m_1(a_n) = d_n \text{ for } n = 1, \dots, N \text{ and } m_1(-1) = m_1(1) = \cos \alpha\}.$$

A natural question is how much energy it takes to form Néel walls of given types at given positions. In terms of the above notation, we may pose the question as follows: what is

$$\inf_{M(a, d)} E_\epsilon \tag{2}$$

for given $a \in A_N$ and $d \in \{\pm 1\}^N$? The question has previously been studied by DeSimone-Kohn-Müller-Otto [9], whose results show that if we set

$$\gamma_n = d_n - \cos \alpha \tag{3}$$

for $n = 1, \dots, N$, then

$$\inf_{M(a, d)} E_\epsilon = \frac{\pi \sum_{n=1}^N \gamma_n^2}{2 \log \frac{1}{\epsilon}} + O\left(\frac{\log \log \frac{1}{\epsilon}}{(\log \epsilon)^2}\right)$$

as $\epsilon \searrow 0$. The result was improved in our previous paper [18]. The first observation is that the asymptotic behaviour of this quantity is more easily expressed in

$$\delta = \epsilon \log \frac{1}{\epsilon}$$

than in ϵ itself because δ is the appropriate size of the core of a Néel wall. (A Néel wall is a two-scale object, containing a core and two logarithmically decaying tails [30, 31, 18].) For this reason, we will assume this relationship between ϵ and δ throughout the paper and work mostly with δ henceforth. Another observation is that the problem behaves similarly to Ginzburg-Landau vortices in some respects. In particular, similarly to the theory of Ginzburg-Landau vortices (see the seminal book of Bethuel-Brezis-Hélein [2]), we can compute a renormalised energy that gives a lot of information about the energy required for a certain collection of Néel walls [18]. It is given by the function²

¹We include $N = 0$ here to avoid special treatment of this case later on. The definition should then be interpreted as follows: as $(-1, 1)^N$ is the set of maps $\{1, \dots, N\} \rightarrow (-1, 1)$, for $N = 0$, we have the unique map $\emptyset \rightarrow (-1, 1)$. The additional constraints become vacuous. So A_0 is *not* empty.

²For $N = 0$, equation (4) reads $W = 0$, while for $N = 1$, the definition of $W(a_1, d_1)$ contains only the first term.

$$W(a, d) = -\frac{\pi}{2} \sum_{n=1}^N \gamma_n^2 \log(2 - 2a_n^2) - \frac{\pi}{2} \sum_{n=1}^N \sum_{k \neq n} \gamma_k \gamma_n \log \left(\frac{1 + \sqrt{1 - \varrho(a_k, a_n)^2}}{\varrho(a_k, a_n)} \right), \quad (4)$$

where

$$\varrho(b, c) = \frac{|b - c|}{1 - bc} \in [0, 1] \quad (5)$$

for $b, c \in (-1, 1)$. Besides the above interaction energy $W(a, d)$, the asymptotic behaviour of E_ϵ as $\epsilon \searrow 0$ also generates an intrinsic renormalised energy $e(d_n)$, characterising the energy of the core of a Néel wall of sign $d_n \in \{\pm 1\}$ (see [18, Definition 26]).

Theorem 1 (Renormalised energy [18]). *Let $N \geq 0$. Then there exists a function $e : \{\pm 1\} \rightarrow \mathbb{R}$ such that for any $a \in A_N$ and $d \in \{\pm 1\}^N$,*

$$\inf_{M(a, d)} E_\epsilon = \frac{\pi \sum_{n=1}^N \gamma_n^2}{2 \log \frac{1}{\delta}} + \frac{W(a, d) + \sum_{n=1}^N e(d_n)}{(\log \frac{1}{\delta})^2} + o\left(\frac{1}{(\log \frac{1}{\delta})^2}\right) \quad \text{as } \epsilon \searrow 0.$$

While this theorem is formulated for fixed Néel walls, the quantity $\inf_{M(a, d)} E_\epsilon$ is continuous with respect to the positions of the walls encoded in a . It is not difficult to see (and is proved nevertheless in Section 2.4 below) that the following is true.

Proposition 2. *Let $a \in A_N$ and $d \in \{\pm 1\}^N$. Then for any $c_0 > 0$ there exists $r > 0$ such that for any $b \in A_N$, the following holds true. If $|b_n - a_n| \leq r$ for $n = 1, \dots, N$, then for any $\epsilon \in (0, \frac{1}{4}]$,*

$$\left| \inf_{M(a, d)} E_\epsilon - \inf_{M(b, d)} E_\epsilon \right| \leq \frac{c_0}{(\log \delta)^2}.$$

Examining the function W can now give some information on how the energy depends on the positions of these walls, and therefore on the interaction between them. In particular, for every pair a_k, a_ℓ of Néel walls, we have an interaction term involving $\varrho(a_k, a_\ell)$, and here ϱ may be regarded as a distance function. The interaction term in (4) may be increasing or decreasing in $\varrho(a_k, a_\ell)$, depending on the sign of $\gamma_k \gamma_\ell$, and accordingly, we can interpret the interaction as repulsive or attractive. Examining the expression more closely,³ we see that we have attraction between Néel walls of the same sign and repulsion for different signs.

But such considerations are supported by the existing theory only as long as a lower bound for their distances is assumed, because the estimates from our paper [18] give no control of the remainder term when this assumption is removed. This is one of the gaps that we fill in this paper, provided that the transition angle α is such that $|\frac{\pi}{2} - \alpha|$ is sufficiently small. In order to quantify this statement, we will often assume in the following that $\alpha \in (\theta_N, \pi - \theta_N)$, where $\theta_N \in [0, \frac{\pi}{2}]$ is defined as follows: $\theta_0 = \theta_1 = \theta_2 = 0$, and for $N \geq 3$,

$$\theta_N = \begin{cases} \arccos\left(\frac{\sqrt{N+1}-1}{N}\right) & \text{if } N \text{ is odd,} \\ \arccos\left(\frac{\sqrt{N}-1}{N-1}\right) & \text{if } N \text{ is even.} \end{cases}$$

Obviously, θ_N is nondecreasing in N and $\theta_N \rightarrow \frac{\pi}{2}$ as $N \rightarrow \infty$.

Roughly speaking, our first new result states that families of profiles with energy of order $O(1/|\log \delta|)$ subconverge to a limiting profile. Moreover, if the energy is even bounded by

$$\frac{\pi}{2|\log \delta|} \sum_{n=1}^N \gamma_n^2 + O\left(\frac{1}{(\log \delta)^2}\right)$$

³The function $t \mapsto \log((1 + \sqrt{1 - t^2})/t)$ is decreasing in $[0, 1)$ and behaves like $\log(1/t)$ as $t \searrow 0$. Therefore, the leading order term in (4) is $\frac{\pi}{2} \sum_{n=1}^N \sum_{k \neq n} \gamma_k \gamma_n \log |a_k - a_n|$.

(for the numbers γ_n as in (3), determined by the limiting profile), then all the Néel walls stay separated from one another, provided that $\theta_N < \alpha < \pi - \theta_N$.⁴ But in order to formulate this rigorously, we need to introduce some additional notation first.

In the following result, we represent a continuous map $m: (-1, 1) \rightarrow \mathbb{S}^1$ in terms of its lifting $\phi: (-1, 1) \rightarrow \mathbb{R}$, determined up to multiples of 2π by the condition $m = (\cos \phi, \sin \phi)$. We abuse notation and write $E_\epsilon(\phi)$ instead of $E_\epsilon(m)$ sometimes. In the limit $\epsilon \searrow 0$, the magnetisation is expected to satisfy $m_1 = \cos \alpha$ almost everywhere. In terms of the lifting, this means that ϕ takes values in $2\pi\mathbb{Z} \pm \alpha$ almost everywhere. We also expect that the limiting lifting will be in $\text{BV}(-1, 1)$. We therefore define the set of piecewise constant functions

$$\Phi = \{\phi \in \text{BV}(-1, 1) : \phi(x_1) \in 2\pi\mathbb{Z} \pm \alpha \text{ for almost all } x_1 \in (-1, 1)\}.$$

We define the two functions

$$\iota: \Phi \rightarrow \mathbb{N} \cup \{0\} \quad \text{and} \quad \eta: \Phi \rightarrow [0, \infty)$$

as follows. For $b \in \mathbb{R}$, let δ_b denote the Dirac measure in \mathbb{R} concentrated at b . Observe that for any $\phi \in \Phi$, there exists a unique representation of the distributional derivative ϕ' of the form

$$\phi' = \sum_{\ell=1}^L \sigma_\ell \delta_{b_\ell},$$

where $b_1, \dots, b_L \in (-1, 1)$ with $b_1 \leq \dots \leq b_L$ and $\sigma_1, \dots, \sigma_L \in \{\pm 2\alpha, \pm 2(\pi - \alpha)\}$, and where $|\sigma_{\ell-1} + \sigma_\ell| = 2\pi$ whenever $b_{\ell-1} = b_\ell$ for $\ell = 1, \dots, L$.⁵ Given this representation, we define

$$\iota(\phi) = L \quad \text{and} \quad \eta(\phi) = \frac{\pi}{2} \sum_{\ell=1}^L \left(1 - \cos \frac{\sigma_\ell}{2}\right)^2.$$

Thus, $\iota(\phi)$ represents the number of what we think of as *individual* Néel walls encoded by ϕ (each corresponding to a simple transition of angle $\pm\alpha$ or $\pm 2(\pi - \alpha)$ between the points $(\cos \alpha, \pm \sin \alpha)$ on \mathbb{S}^1). The number $\eta(\phi)$ corresponds to the energy required for these Néel walls to leading order, as identified in Theorem 1.

Our theory gives special relevance to functions $\phi \in \Phi$ such that every jump has size $\pm 2\alpha$ or $\pm 2(\pi - \alpha)$, and thus every jump corresponds to exactly *one* transition between the points $(\cos \alpha, \pm \sin \alpha)$ along \mathbb{S}^1 . This is the case exactly when $\iota(\phi)$ coincides with the number of jumps and is equivalent to the following condition.

Definition 3. *We say that $\phi \in \Phi$ is simple if there exist $N \in \mathbb{N} \cup \{0\}$, $a \in A_N$, and $\omega \in \{\pm 2\alpha, \pm 2(\pi - \alpha)\}^N$ such that⁶*

$$\phi' = \sum_{n=1}^N \omega_n \delta_{a_n}. \tag{6}$$

If so, define $d \in \{\pm 1\}^N$ by

$$d_n = \begin{cases} 1 & \text{if } |\omega_n| = 2\alpha \\ -1 & \text{if } |\omega_n| = 2\pi - 2\alpha, \end{cases} \tag{7}$$

provided that $\alpha \neq \pi/2$, for $n = 1, \dots, N$. For $\alpha = \pi/2$, the jump sizes ω_n are not sufficient to determine d . Instead, we then define $d_n = 1$ if either $\lim_{x_1 \nearrow a_n} \phi(x_1) \in 2\pi\mathbb{Z} - \pi/2$ and $\omega_n = \pi$ or $\lim_{x_1 \nearrow a_n} \phi(x_1) \in 2\pi\mathbb{Z} + \pi/2$ and $\omega_n = -\pi$, and we define $d_n = -1$ otherwise. Then (a, d) is called the transition profile of ϕ . For simple functions ϕ , we have $N = \iota(\phi)$.

⁴This condition means that the neighbouring transitions have comparable sizes.

⁵This condition implies that σ_ℓ and $\sigma_{\ell+1}$ have the same sign; thus, the jumps of ϕ are decomposed in a (strictly) monotone way and the representation of ϕ' is unique.

⁶If $N = 0$, then (6) reads $\phi' = 0$, i.e., constant functions are simple.

We note that for any $a \in A_N$ and $d \in \{\pm 1\}^N$, the pair (a, d) is the transition profile of a simple function $\phi \in \Phi$. Moreover, this function is unique up to a constant in $2\pi\mathbb{Z}$.⁷ If $\phi \in \Phi$ is simple with transition profile (a, d) , then we may define $\gamma_n = d_n - \cos \alpha$ for $n = 1, \dots, N$, and we observe that

$$\eta(\phi) = \frac{\pi}{2} \sum_{n=1}^N \gamma_n^2.$$

This corresponds to the leading order term in the expansion of Theorem 1. In other words, we see here, as already mentioned, that η represents the energy of the Néel walls encoded in ϕ . This remains true if ϕ is not simple.

We now have a compactness result for profiles with suitably bounded energy. Under additional assumptions, we can rule out that several Néel walls collide as $\epsilon \searrow 0$, and thus the limiting profile stays simple.

Theorem 4 (Compactness and Separation). *Suppose that $(\phi_\epsilon)_{\epsilon>0}$ is a family of functions in $H^1(-1, 1)$ such that $\phi_\epsilon(-1) = \pm\alpha$ and $\phi_\epsilon(1) \in 2\pi\mathbb{Z} \pm \alpha$ for all $\epsilon > 0$. Suppose that*

$$\limsup_{\epsilon \searrow 0} |\log \epsilon| E_\epsilon(\phi_\epsilon) < \infty.$$

1. *Then there exist a sequence of numbers $\epsilon_k \searrow 0$ and a function $\phi_0 \in \Phi$ such that $\phi_{\epsilon_k} \rightarrow \phi_0$ in $L^2(-1, 1)$.*
2. *For the limit ϕ_0 from statement 1, let $N = \iota(\phi_0)$. If $\delta = \epsilon \log \frac{1}{\delta}$, $\theta_N < \alpha < \pi - \theta_N$, and*

$$E_\epsilon(\phi_\epsilon) \leq \frac{\eta(\phi_0)}{\log \frac{1}{\delta}} + O\left(\frac{1}{(\log \frac{1}{\delta})^2}\right), \quad \text{as } \epsilon \searrow 0, \quad (8)$$

then ϕ_0 is simple.

In other words, under condition (8), the jump points $a_1 < a_2 < \dots < a_N$ of ϕ_0 are distinct and the size of each jump lies in $(-2\pi, 2\pi)$. Without condition (8), the limit ϕ_0 may have jumps of any size in $2\pi\mathbb{Z} + \{0, \pm 2\alpha\}$. In general, this will of course amount to several transitions between the points $(\cos \alpha, \pm \sin \alpha)$, so we think of this as an accumulation of several Néel walls, or a composite Néel wall. According to the last statement of the theorem, however, a composite Néel wall requires more energy than the sum of its individual parts if $\theta_N < \alpha < \pi - \theta_N$. In contrast, if (8) is satisfied, then the walls stay separated and during the transition, ϕ_ϵ passes the set $\pi\mathbb{Z}$ only *once*.

For simple limit configurations, the results from our previous paper [18] then apply, and we can prove the following Γ -convergence result, which improves Theorem 1 insofar as no assumption of the position of the Néel walls is required. For the Γ -convergence result, there is no need of the angle constraint $\theta_N < \alpha < \pi - \theta_N$ because we restrict ourselves to simple functions ϕ_0 . This restriction corresponds to the assumption in Ginzburg-Landau models that only vortices of degree ± 1 are present, which is a common feature in Γ -convergence results in such theories (see e.g. [1, 15]).

Corollary 5 (Γ -convergence). *Let $\alpha \in (0, \pi)$.*

1. *(Lower bound) Let $(\phi_\epsilon)_{\epsilon>0}$ be a family of functions in $H^1(-1, 1)$ such that $\phi_\epsilon(-1) = \pm\alpha$, $\phi_\epsilon(1) \in 2\pi\mathbb{Z} \pm \alpha$ for all $\epsilon > 0$, and such that $\phi_\epsilon \rightarrow \phi_0$ in $L^2(-1, 1)$ for a simple $\phi_0 \in \Phi$. Suppose that (a, d) is the transition profile of ϕ_0 and $N = \iota(\phi_0)$. Then*

$$E_\epsilon(\phi_\epsilon) \geq \frac{\eta(\phi_0)}{\log \frac{1}{\delta}} + \frac{\sum_{n=1}^N e(d_n) + W(a, d)}{(\log \frac{1}{\delta})^2} - o\left(\frac{1}{(\log \frac{1}{\delta})^2}\right) \quad \text{as } \epsilon \searrow 0.$$

⁷If $m = (\cos \phi, \sin \phi)$, then one can easily find an approximation sequence $m_\epsilon \in M(a, d)$ of m (i.e., $m_\epsilon \rightarrow m$ in $L^2(-1, 1)$).

2. (Upper bound) If $\phi_0 \in \Phi$ is simple with transition profile (a, d) and if $N = \iota(\phi_0)$, then there exists a family $(\phi_\epsilon)_{\epsilon > 0}$ in $H^1(-1, 1)$ such that $\phi_\epsilon(-1) = \pm\alpha$, $\phi_\epsilon(1) \in 2\pi\mathbb{Z} \pm \alpha$ for all $\epsilon > 0$, $\phi_\epsilon \rightarrow \phi_0$ in $L^2(-1, 1)$, and

$$E_\epsilon(\phi_\epsilon) \leq \frac{\eta(\phi_0)}{\log \frac{1}{\delta}} + \frac{\sum_{n=1}^N e(d_n) + W(a, d)}{(\log \frac{1}{\delta})^2} + o\left(\frac{1}{(\log \frac{1}{\delta})^2}\right) \quad \text{as } \epsilon \searrow 0.$$

This result amounts to an asymptotic expansion by Γ -convergence of the energy E_ϵ at second order (see [3] for more details about Γ -expansions). The first order terms tell us that η corresponds to the quantised energy of the individual Néel walls and gives rise to the Γ -convergence $|\log \delta| E_\epsilon \xrightarrow{\Gamma} \eta$ over the space of admissible liftings ϕ with the topology of $L^2(-1, 1)$. The second order terms include (apart from the contribution $e(d)$ depending only on the number and types of the Néel walls) the function $W(a, d)$, which encapsulates the interaction between the Néel walls and governs their optimal positions. When the energy is rescaled by $(\log \delta)^2$, the second order terms become the dominant, asymptotically finite contributions in the expansion. We call them ‘renormalised energy’, which is standard terminology in Ginzburg-Landau type theories. Sometimes, however, we apply this name to $W(a, d)$ alone.

A related Γ -convergence result has been proved in [13], but it is at the first order and weaker than Corollary 5 because it makes a statement about \mathbb{S}^1 -valued maps rather than their liftings.

Néel walls may arise as the result of topological constraints. If we have a continuous function $\phi: [-1, 1] \rightarrow \mathbb{R}$ with fixed values on the boundary, say $\phi(-1) = \alpha$ (or $\phi(-1) = -\alpha$) and $\phi(1) = 2\pi\ell + \alpha$ (or $\phi(1) = 2\pi\ell - \alpha$) for a fixed number $\ell \in \mathbb{Z}$, then the magnetisation $m = (\cos \phi, \sin \phi): [-1, 1] \rightarrow \mathbb{S}^1$ will have at least N Néel walls (transitions between $(\cos \alpha, \pm \sin \alpha)$), where $N = 2|\ell|$ or $N = 2|\ell| \pm 1$, depending on the signs in the boundary data. If the number of Néel walls is exactly N , then their signs will alternate. In this situation, we speak of *topological* Néel walls. Thus if we define

$$d_N^+ = (1, -1, 1, \dots, \pm 1) \in \{\pm 1\}^N \quad \text{and} \quad d_N^- = (-1, 1, -1, \dots, \pm 1) \in \{\pm 1\}^N,$$

then we expect a renormalised energy given by $W(a, d_N^\pm)$ for some $a \in A_N$. Topological Néel walls are expected when we minimise E_ϵ subject to the above boundary data (which amounts to prescribing the topological degree). Indeed, it can be shown, with arguments similar to a related model [19, Lemma 3.1], that there are exactly as many Néel walls for minimisers as the topology requires and thus their signs necessarily alternate. When passing to the limit, we expect that the positions of the Néel walls converge to a minimiser of $W(\cdot, d_N^\pm)$ if such a minimiser exists.

The following result tells us when this is the case. The findings are consistent with the expectation that topological Néel walls stay separated for α close to $\pi/2$, while they tend to collide in the limit $\epsilon \searrow 0$ if $|\cos \alpha|$ is too large. We will shortly discuss another result which makes this more precise.

Proposition 6. *Given $N \geq 2$ and $d = d_N^+$ or d_N^- , let*

$$\Theta = \begin{cases} (\theta_N, \pi - \theta_N) & \text{if } N \text{ is even,} \\ (\theta_{N-2}, \pi - \theta_N) & \text{if } N \text{ is odd and } d_1 = 1, \\ (\theta_N, \pi - \theta_{N-2}) & \text{if } N \text{ is odd and } d_1 = -1. \end{cases} \quad (9)$$

Then

1. $W(\cdot, d)$ admits a minimizer over A_N if $\alpha \in \Theta$;
2. $\inf_{a \in A_N} W(a, d) = -\infty$ if $N \geq 3$ and $\alpha \notin \overline{\Theta}$.

In the specific situation of topological Néel walls, the information provided by Theorem 4 can be improved. We formulate the following statement for $\phi(-1) = -\alpha$ and $\phi(1) = 2\pi\ell + \alpha$ with $\ell \in \mathbb{N} \cup \{0\}$ only (for simplicity), but the other cases allow similar conclusions, too. Here we denote

$$e_+ := e(1) \quad \text{and} \quad e_- := e(-1)$$

for the function $e: \{\pm 1\} \rightarrow \mathbb{R}$ from Theorem 1.

Corollary 7 (Topological Néel walls). *Let $\ell \in \mathbb{N} \cup \{0\}$ and $N = 2\ell + 1$. Suppose that $\theta_N < \alpha < \pi - \theta_N$. Suppose that $(\phi_\epsilon)_{\epsilon > 0}$ is a family of functions in $H^1(-1, 1)$ such that $\phi_\epsilon(-1) = -\alpha$ and $\phi_\epsilon(1) = 2\pi\ell + \alpha$ for all $\epsilon > 0$. Let⁸*

$$\mathcal{E}_0 := \pi\ell(1 + \cos^2 \alpha) + \frac{\pi}{2}(1 - \cos \alpha)^2 \quad \text{and} \quad \mathcal{E}_1 := (\ell + 1)e_+ + \ell e_-.$$

1. Then

$$E_\epsilon(\phi_\epsilon) \geq \frac{\mathcal{E}_0}{\log \frac{1}{\delta}} + \frac{\mathcal{E}_1 + \inf_{\tilde{a} \in A_N} W(\tilde{a}, d_N^+)}{(\log \frac{1}{\delta})^2} - o\left(\frac{1}{(\log \frac{1}{\delta})^2}\right) \quad \text{as } \epsilon \searrow 0.$$

2. If

$$E_\epsilon(\phi_\epsilon) \leq \frac{\mathcal{E}_0}{\log \frac{1}{\delta}} + O\left(\frac{1}{(\log \frac{1}{\delta})^2}\right) \quad \text{as } \epsilon \searrow 0,$$

then there exist $a \in A_N$, a sequence $\epsilon_k \searrow 0$, and a simple $\phi_0 \in \Phi$ with $\phi_0(-1) = -\alpha$ and $\phi_0(1) = 2\pi\ell + \alpha$ such that $\phi_{\epsilon_k} \rightarrow \phi_0$ in $L^2(-1, 1)$ and

$$\phi_0' = \sum_{n=1}^N \omega_n \delta_{a_n},$$

where $\omega_n = 2\alpha$ for n odd and $\omega_n = 2\pi - 2\alpha$ for n even.

3. If every ϕ_ϵ is a minimiser of E_ϵ for its boundary data, then all of the above holds and

$$W(a, d_N^+) = \inf_{\tilde{a} \in A_N} W(\tilde{a}, d_N^+).$$

Combining Corollary 5 and statements 1 and 2 of Corollary 7, we deduce that

$$\min_{\mathcal{T}_N} |\log \delta| (|\log \delta| E_\epsilon - \mathcal{E}_0) \rightarrow \mathcal{E}_1 + \min_{A_N} W(\cdot, d_N^+) \quad \text{as } \epsilon \rightarrow 0,$$

where \mathcal{T}_N is the set of topological Néel walls with signs d_N^+ . Statement 3 in Corollary 7 determines (asymptotically as $\epsilon \searrow 0$) the optimal positions of Néel walls by minimising the renormalised energy $W(\cdot, d_N^+)$ when a winding number is prescribed by the boundary data. For any result of this sort, we need to exclude neighbouring walls of the same sign (which is guaranteed by the topology in Corollary 7), because they will attract each other. This is a feature of the problem and not a shortcoming of the theory. The question of attraction and repulsion of Néel walls is discussed from a physical point of view by Hubert and Schäfer [12, Section 3.6. (C)].

⁸The quantity \mathcal{E}_0 corresponds to the leading order term in the energy expansion of ϕ_ϵ . For example, a pair of Néel walls of signs $(+1, -1)$ generates to leading order the energy

$$\frac{\pi}{2|\log \delta|} [(1 - \cos \alpha)^2 + (1 + \cos \alpha)^2] = \frac{\pi}{|\log \delta|} (1 + \cos^2 \alpha).$$

We also need to exclude the case of small transition angles α . This is because otherwise, the attraction of next-to-neighbouring walls may dominate the (comparatively small) repulsion of neighbouring walls. This is a phenomenon that was exploited in other papers of the authors [19, 20] (in a somewhat different model with fixed ϵ) to construct energy minimisers comprising several Néel walls. For the problem studied here, a plausible consequence (not proved here) is that for small transition angles, several Néel walls may in the limit $\epsilon \searrow 0$ collapse to a single *composite* Néel wall, corresponding to a jump of more than 2π in the lifting (i.e., if $\alpha \in (0, \pi)$ is small, under the condition (8), we may have a jump of size $|\omega_n| \geq 2\pi$).

1.3 The unconfined problem

If we consider $m: \mathbb{R} \rightarrow \mathbb{S}^1$, rather than restricting the domain to a bounded interval, then a functional as in the preceding section will not have any nontrivial critical points; indeed, as simple dilation of the form $\tilde{m}(x_1) = m(\lambda x_1)$ for $\lambda \in (0, 1)$ will decrease the energy unless m is constant. Any nontrivial structure, including a Néel wall, is therefore inherently unstable. The situation changes, however, if the energy functional contains an additional term modelling anisotropy.

For $\epsilon > 0$ and $\alpha \in (0, \pi)$, we now study the functionals

$$E_\epsilon(m) = \frac{1}{2} \int_{-\infty}^{\infty} (\epsilon |m'|^2 + (m_1 - \cos \alpha)^2) dx + \frac{1}{2} \|m_1 - \cos \alpha\|_{\dot{H}^{1/2}(\mathbb{R})}^2$$

for $m \in H_{\text{loc}}^1(\mathbb{R}; \mathbb{S}^1)$. The additional term arises from the combination of a crystalline anisotropy with easy axis parallel to the m_2 -axis and an external magnetic field parallel to the m_1 -axis. For simplicity, we call it the anisotropy energy.

We redefine several quantities in this section, including E_ϵ , but as we will treat the two problems separately, this will not cause any problems. Does this situation still allow results similar to our previous paper [18] and the extension from the preceding section? The answer is yes, and we will give some of the corresponding results here. Many of the arguments are in fact similar to the confined case. What may seem somewhat surprising, however, is the form of the renormalised energy for the new problem. There is now an interaction between all three energy terms (exchange, stray field, and anisotropy energy), and while the anisotropy energy contributes directly to the renormalised energy, it also does so indirectly by modifying the contribution from the stray field energy.

For $N \in \mathbb{N}$, we now define

$$A_N = \{a = (a_1, \dots, a_N) \in \mathbb{R}^N : a_1 < \dots < a_N\},$$

whereas for $a \in A_N$ and $d \in \{\pm 1\}^N$, we define

$$M(a, d) = \{m \in H_{\text{loc}}^1(\mathbb{R}; \mathbb{S}^1) : E_1(m) < \infty \text{ and } m_1(a_n) = d_n \text{ for } n = 1, \dots, N\}.$$

Let

$$I(t) = \int_0^\infty \frac{se^{-s}}{s^2 + t^2} ds, \quad t > 0.$$

Also define

$$I_0 = \int_0^\infty e^{-s} \log s ds. \tag{10}$$

This is minus the Euler-Mascheroni constant⁹ [26]. (The latter is traditionally denoted by γ , but we use this symbol extensively for something else.) Then the renormalised energy turns out to be

$$W(a, d) = -\frac{\pi}{2} I_0 \sum_{n=1}^N \gamma_n^2 - \frac{\pi}{2} \sum_{n=1}^N \sum_{k \neq n}^N \gamma_k \gamma_n I(|a_k - a_n|). \tag{11}$$

⁹Recalling the gamma function $\Gamma(t) = \int_0^\infty s^{t-1} e^{-s} ds$ for $t > 0$, we compute $I_0 = \Gamma'(1)$, which is known to be $\lim_{n \rightarrow \infty} (\log n - \sum_{k=1}^n \frac{1}{k}) = -\gamma = -0.577\dots$

Here again, given $a \in A_N$ and $d \in \{\pm 1\}^N$, we set $\gamma_n = d_n - \cos \alpha$.

It is worth pointing out that $I(t)$ is logarithmic to leading order again. This is not immediately obvious here, but we will prove in Lemma 20 below that $|I(t) + \log t - I_0| \leq \pi t/2$ for all $t > 0$. On the other hand, the function I decays at the rate $1/t^2$ as $t \rightarrow \infty$. The extra term $-\frac{\pi}{2} I_0 \gamma_n^2$ for $n = 1, \dots, N$ in (11) may be thought of as a ‘self-interaction’ contribution from each Néel wall. As $\lim_{t \searrow 0} (I(t) + \log t) = I_0$ (by Lemma 20), there is some analogy to the definition of W in the confined case. Indeed, the terms involving $\log(2 - 2a_n^2)$ in the renormalised energy in (4) may be interpreted the same way, although it is also identified as a ‘tail-boundary interaction’ term in our previous paper [18]. As in the confined model, next to the interaction energy $W(a, d)$ we also have the terms $e(d_n)$ for every Néel wall of sign $d_n \in \{\pm 1\}$. In the following, we prove a result on the expansion of the minimal energy in the unconfined case when the transition profile (a, d) is fixed, similar to Theorem 1 for the confined case.

Theorem 8 (Renormalised energy). *Let $e: \{\pm 1\} \rightarrow \mathbb{R}$ be the function from Theorem 1. For any $a \in A_N$ and $d \in \{\pm 1\}^N$, the following expansion holds as $\epsilon \searrow 0$:*

$$\inf_{M(a,d)} E_\epsilon = \frac{\pi \sum_{n=1}^N \gamma_n^2}{2 \log \frac{1}{\delta}} + \frac{1}{(\log \delta)^2} \left(W(a, d) + \sum_{n=1}^N e(d_n) \right) + o\left(\frac{1}{(\log \delta)^2} \right).$$

Most of the new arguments in the proof of this result concern the effects of the anisotropy term on the limiting stray field potential. When proving the corresponding result for the confined problem [18], we represented the stray field energy $\frac{1}{2} \|m_1 - \cos \alpha\|_{H^{1/2}(\mathbb{R})}^2$ in terms of the harmonic extension of $m_1 - \cos \alpha$ to the half-plane $\mathbb{R}_+^2 = \mathbb{R} \times (0, \infty)$ and made extensive use of the fact that a harmonic function $u: \mathbb{R}_+^2 \rightarrow \mathbb{R}$ remains harmonic under composition with a conformal map (especially a Möbius transform). In other words, we used the underlying symmetry of the stray field energy. The anisotropy term does not share this symmetry, and therefore, this approach will not work any more. Instead, we now use the Fourier transform in x_1 in order to turn a harmonic function on \mathbb{R}_+^2 into a solution of an ordinary differential equation. This will allow us to represent the solutions as oscillatory integrals. Of course, some of the previous arguments can still be used here, and therefore, parts of the proof of this result are given as a sketch only, emphasising the changes relative to our previous work [18].

The results from section 2 (in particular, Theorem 4 and Corollaries 5 and 7) will have almost identical counterparts for the new situation; see Theorem 31 below. But this now requires nothing more than a combination of the arguments from the confined case and from the proof of Theorem 8. For the convenience of the reader, we give a very brief sketch of the proof in Section 3.6.

Having found a new renormalised energy W , we may now wish to use it to determine the (asymptotically) optimal positions of a set of Néel walls. As before, neighbouring walls of the same sign will attract each other, and therefore, no minimisers of $W(\cdot, d)$ in A_N are to be expected unless $d_{n+1} = -d_n$ for $n = 1, \dots, N-1$. But in contrast to the confined model, we now have to consider the possibility of arbitrarily large distances between the Néel walls as well. We find that in some cases, no minimum exists even though the infimum is finite. More precisely, we know the following.

Proposition 9. *Let $N \geq 2$ and $d = d_N^+$ or d_N^- . Suppose that Θ is defined as in (9).*

1. *If $\alpha \in \Theta$, then $\inf_{A_N} W(\cdot, d) > -\infty$.*
2. *If $N \geq 3$ and $\alpha \notin \bar{\Theta}$, then $\inf_{A_N} W(\cdot, d) = -\infty$.*
3. *Suppose that $N = 3$. If $d_1 \cos \alpha \in (-\frac{7}{9}, -\frac{1}{3})$, then $W(\cdot, d)$ has a unique critical point, which is not a minimiser. If $d_1 \cos \alpha \notin (-\frac{7}{9}, -\frac{1}{3})$, then $W(\cdot, d)$ has no critical point.*
4. *Suppose that $N \geq 2$ and $d_1 \cos \alpha \geq 0$ or $d_N \cos \alpha \geq 0$. Then $W(\cdot, d)$ has no critical point (and thus no minimisers). In particular, this is always the case if N is even.*

The proposition gives only partial information for $N \geq 5$ odd. The question of existence/non-existence of minimisers is open here, except for the cases covered in statements 2 and 4.

Some of these observations are obviously consistent with our results [19, 20] about the existence/nonexistence of minimisers of the functional E_1 for a prescribed winding number. In particular, for winding numbers giving rise to the situation of statement 4, we find that E_1 has no minimiser. Statement 3, however, shows an apparent contrast. In this situation, the functional E_1 *has* a minimiser, provided that α is sufficiently small (and by symmetry of the model, this also applies when $\pi - \alpha$ is small). Although our previous papers only consider the case $\epsilon = 1$, we expect that similar results apply to every fixed $\epsilon > 0$, but no analysis has been carried out studying how the threshold for α or the shape of the minimisers depend on ϵ .

1.4 Comparison between the confined and the unconfined model

As we have seen, by and large, our two models permit the same sort of results. There are, however, some subtle differences both on a technical level and in the consequences.

As mentioned previously, the confined and the unconfined model have different mechanisms to stabilise the Néel walls. The same mechanisms also determine the transition angle α . In the confined model, this is the result of the steric interaction with the sample edges (represented by the boundary of the interval J). In the unconfined model, we have a combination of the anisotropy effect and an external magnetic field instead.

The two models lead to renormalised energies with a lot of similarities but also some qualitative differences. At close range, the interaction between any two Néel walls is essentially the same and logarithmic at leading order for both models. But in the unconfined model, the walls can be arbitrarily far apart, and then the long-range interaction decays quadratically to 0. The confined model prevents this by the set-up, but the confinement is also visible in the renormalised energy in the form of a boundary interaction term (also logarithmic).

As a consequence, the global energy landscapes of the two renormalised energies look quite different. This is most easily demonstrated in the context of Corollary 7, where we consider $m = (\cos \phi, \sin \phi)$ with a prescribed winding number for ϕ . In this situation, the number and signs of the Néel walls are essentially given. In the confined model, according to Proposition 6, the renormalised energy W attains its minimum among all admissible configurations for some values of α and is unbounded below otherwise. (Roughly speaking, we have minimisers when $|\pi/2 - \alpha|$ is small.) For the unconfined model, the question is less clear, but is discussed in Proposition 9. While the result does not cover all possible cases, it does give a complete overview for arrays of up to four Néel walls, and it turns out that W *never* has a minimiser in all cases where we know the answer.

For both models, given $a \in A_N$ and $d \in \{\pm 1\}^N$, we identify a limiting stray field potential $u_{a,d}^*: \mathbb{R}_+^2 \rightarrow \mathbb{R}$ in the proofs below, the understanding of which is crucial for all the results discussed in this paper. Near every Néel wall, this function behaves like the phase of a vortex (which is why the theory has some connections to Ginzburg-Landau vortices), and at leading order, this is the same for both models. The behaviour at ∞ is similar, too, but only in the unconfined model does it have immediate consequences. This is what determines the decay of W in the unconfined model as the distance between the Néel walls increases. Furthermore, the way that the limiting stray field potential arises from E_ϵ is somewhat different for the two models. The exchange energy has only an indirect effect on it, and thus in the confined model, where E_ϵ consists only of the exchange energy and the stray field energy, we mostly need to study the latter. In the unconfined model, we also have the anisotropy energy, which has a fundamental effect on $u_{a,d}^*$. Indeed, in many of the proofs in Section 3, we have to consider $u_{a,d}^*$ and a limiting anisotropy energy jointly. This is somewhat surprising, as a variant of the Pohozaev identity [19, Proposition 1.1] implies that for critical points of E_ϵ , the exchange energy and the anisotropy energy are of the same magnitude, while the stray field energy is larger by a factor of order $|\log \epsilon|$.

1.5 Comparison with other Ginzburg-Landau models

The connection between our theory and Ginzburg-Landau vortices is not obvious, but can be seen when studying the stray field energy in terms of a stray field potential. This is a function on $\mathbb{R}_+^2 = \mathbb{R} \times (0, \infty)$ that is obtained from the magnetisation m by means of a boundary value problem (see (14) and (15) below) and in the limit $\epsilon \searrow 0$ gives rise to the aforementioned function $u_{a,d}^*$. Its physical relevance is that its gradient corresponds to the magnetic stray field induced by m . Near a Néel wall, the stray field potential behaves like the phase of a vortex in \mathbb{R}_+^2 . As the energy carried by the stray field potential turns out to be the dominant term in the limit, the interaction between these ‘vortices’ matters a lot.

In the context of Ginzburg-Landau type models, computing the renormalised energy between topological singularities has become a topic extensively studied in last three decades, see, e.g., the seminal book of Bethuel-Brezis-Hélein [2] and the further developments discussed by Sandier-Serfaty [36]. In two-dimensional Ginzburg-Landau models, one of the key tools is the Jacobian of the order parameter, which detects the point singularities in the shape of vortices. Many techniques have been developed based on the Jacobian, yielding compactness results, lower bounds, and Γ -convergence (at the first and second order), see e.g. [1, 6, 14, 15, 23, 24, 22, 28, 34, 35].

However, in our models, there is no topological invariant playing the role of the Jacobian, even though we have a stray field potential that behaves like a vortex angle around a domain wall. In fact, our problem is phenomenologically different from the standard Ginzburg-Landau model. First, a Néel wall is a two length-scale topological object, while the vortex has only one scale. Second, while the renormalised energy between Ginzburg-Landau vortices is generated solely by the tail-tail interaction of neighbouring vortices, in our problem, there is a core-tail interaction as well. In fact, the latter even dominates the tail-tail interaction between Néel walls, which is a new feature in the context of Ginzburg-Landau problems. It also gives rise to a new phenomenon: we have attraction for walls of the same sign and repulsion for walls of different signs, which is exactly the opposite in the context of Ginzburg-Landau vortices.

As a result of all of this, we can use the well-known ideas for Ginzburg-Landau vortices as a motivation for much of our theory, but they are generally not sufficient and can occasionally be misleading. The different attraction/repulsion pattern also gives rise to an energy landscape in the renormalised energy that is quite different from Ginzburg-Landau vortices. Some of this energy landscape is explored in Propositions 6 and 9 above, but more work is required here.

1.6 Representations of the stray field energy

A good understanding of the nonlocal term in the energy functional, the stray field energy, is essential for the analysis of both problems. In both cases, it is given in terms of the seminorm $\|\cdot\|_{\dot{H}^{1/2}(\mathbb{R})}$, and there are several ways to characterise this quantity. Possibly the best-known representation involves the Fourier transform $\hat{f}(\xi) = \int_{\mathbb{R}} e^{-i\xi x_1} f(x_1) dx_1$ for $\xi \in \mathbb{R}$. Then

$$\|f\|_{\dot{H}^{1/2}(\mathbb{R})}^2 = \frac{1}{2\pi} \int_{-\infty}^{\infty} |\xi| |\hat{f}|^2 d\xi.$$

Our arguments mostly rely on different representations, however, one of which has a physical interpretation as well. (This is the reason why $\dot{H}^{1/2}(\mathbb{R})$ appears in the problem in the first place.)

Given $f \in \dot{H}^{1/2}(\mathbb{R})$, consider the boundary value problem

$$\Delta u = 0 \quad \text{in } \mathbb{R}_+^2, \tag{12}$$

$$\frac{\partial u}{\partial x_2} = -f' \quad \text{on } \mathbb{R} \times \{0\}. \tag{13}$$

Let $C_0^\infty(\overline{\mathbb{R}_+^2})$ denote the set of smooth functions in \mathbb{R}_+^2 with compact support in $\mathbb{R} \times [0, \infty)$. If $\dot{H}^1(\mathbb{R}_+^2)$ denotes the completion of $C_0^\infty(\overline{\mathbb{R}_+^2})$ with respect to the inner product

$$\langle u, v \rangle_{\dot{H}^1(\mathbb{R}_+^2)} = \int_{\mathbb{R}_+^2} \nabla u \cdot \nabla v \, dx,$$

then there exists a unique solution u of (12), (13) in $\dot{H}^1(\mathbb{R}_+^2)$. (If interpreted as a function, however, then u is unique only up to a constant.) If $f = m_1 - \cos \alpha$ for a given magnetisation m , then u should be regarded as a potential for the magnetic stray field induced by m . Then

$$\|f\|_{\dot{H}^{1/2}(\mathbb{R})}^2 = \int_{\mathbb{R}_+^2} |\nabla u|^2 \, dx.$$

There exists a dual representation, which may be more familiar but has no physical interpretation. As $\Delta u = 0$, the vector field $\nabla^\perp u = (-\frac{\partial u}{\partial x_2}, \frac{\partial u}{\partial x_1})$ is curl free in \mathbb{R}_+^2 . Hence there exists $v: \mathbb{R}_+^2 \rightarrow \mathbb{R}$ such that $\nabla v = \nabla^\perp u$. Because $\frac{\partial v}{\partial x_1} = -\frac{\partial u}{\partial x_2} = f'$ on $\mathbb{R} \times \{0\}$, we may further choose v such that $v = f$ on $\mathbb{R} \times \{0\}$. We then compute $\Delta v = 0$ in \mathbb{R}_+^2 , so v is a harmonic extension of f . Indeed it is the unique harmonic extension with finite Dirichlet energy. Now we may also write

$$\|f\|_{\dot{H}^{1/2}(\mathbb{R})}^2 = \int_{\mathbb{R}_+^2} |\nabla v|^2 \, dx.$$

Finally, we mention one more representation of the stray field energy that does not play a significant role here, but has been used extensively for other questions about the model [19, 20]. It can be shown that [27, Chapter 7]

$$\|f\|_{\dot{H}^{1/2}(\mathbb{R})}^2 = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{(f(s) - f(t))^2}{(s - t)^2} \, ds \, dt.$$

When studying the functional E_ϵ , we apply these formulas to the function $m_1 - \cos \alpha$. For the problem described in Section 1.3, this is a function defined on \mathbb{R} , but in Section 1.2, it is initially defined in the interval $(-1, 1)$. We therefore extend m_1 by $\cos \alpha$ everywhere else. Then $\|m_1 - \cos \alpha\|_{\dot{H}^{1/2}(\mathbb{R})}$ can be represented in terms of the boundary value problem

$$\Delta u = 0 \quad \text{in } \mathbb{R}_+^2, \tag{14}$$

$$\frac{\partial u}{\partial x_2} = -m'_1 \quad \text{on } \mathbb{R} \times \{0\}. \tag{15}$$

for both problems. This problem has a unique solution $u \in \dot{H}^1(\mathbb{R}_+^2)$, which is called the *stray field potential* induced by the magnetisation m .

2 Analysis for the confined problem

The purpose of this section is to prove Theorem 4, Corollaries 5 and 7, and Propositions 2 and 6. Thus we consider the energy functional E_ϵ defined in (1) and the renormalised energy defined in (4). Furthermore, both A_N and $M(a, d)$ (for $a \in A_N$ and $d \in \{\pm 1\}^N$) are defined as in Section 1.2.

2.1 Compactness

We first address the compactness result, i.e., statement 1 in Theorem 4. For comparison, we mention that there are other compactness results for \mathbb{S}^1 -valued magnetisations in terms of their liftings in various ferromagnetic thin-film regimes in [33, 25, 17].

Even though we obtain accumulation points in $\Phi \subset \text{BV}(-1, 1)$ here, there is no compactness with respect to the BV-topology [21, Theorem 3]. We use the L^2 -topology for convenience, but obviously it can be replaced by any other L^p -topology with $p < \infty$.

Proposition 10. *Consider a sequence $(\epsilon_k)_{k \in \mathbb{N}}$ of positive numbers with $\epsilon_k \searrow 0$ as $k \rightarrow \infty$. Let $(\phi_k)_{k \in \mathbb{N}}$ be a sequence in $H^1(-1, 1)$ such that every ϕ_k satisfies the boundary conditions $\phi_k(-1) = \pm \alpha$ and $\phi_k(1) \in 2\pi\mathbb{Z} \pm \alpha$. Suppose that*

$$\limsup_{k \rightarrow \infty} |\log \epsilon_k| E_{\epsilon_k}(\phi_k) < \infty. \quad (16)$$

Then $\{\phi_k : k \in \mathbb{N}\}$ is uniformly bounded in $L^\infty(-1, 1)$ and relatively compact in $L^2(-1, 1)$. Moreover, if ϕ_0 is any accumulation point, then $\phi_0 \in \Phi$.

Proof. This statement is an improvement of a result of the first author [13, Theorem 1]. In fact, only the compactness of the \mathbb{S}^1 -valued maps $m_\epsilon = (\cos \phi_\epsilon, \sin \phi_\epsilon)$ is proved in [13], while here we give a more precise statement in terms of the lifting ϕ_ϵ . Nevertheless, we can follow some of the same steps here, referring to the earlier paper for some of the details.

First recall that for $m \in H^1((-1, 1); \mathbb{S}^1)$, the $\dot{H}^{1/2}$ -seminorm of $m_1 - \cos \alpha$ is controlled by the stray field energy. More precisely,

$$\|m_1 - \cos \alpha\|_{\dot{H}^{1/2}(\mathbb{R})}^2 \leq 2E_\epsilon(m).$$

Consider $m_k = (\cos \phi_k, \sin \phi_k)$. Since $m_{k1}(-1) = \cos \alpha$, inequality (16) is therefore enough to conclude that $m_{k1} \rightarrow \cos \alpha$ in $L^2(-1, 1)$.

Next we localise the large variations of m_{k2} . Since each m_k belongs to $H^1((-1, 1); \mathbb{S}^1)$, m_k is a uniformly continuous function (for fixed k). Therefore, the set

$$m_{k2}^{-1}\left(\left(-\frac{1}{2} \sin \alpha, \frac{1}{2} \sin \alpha\right)\right)$$

is open and every connected component is an open interval. If (a, b) is one of these connected components, then $|m_{k2}(a)| = |m_{k2}(b)| = \frac{1}{2} \sin \alpha$. If they have opposite signs, then

$$b - a \geq \frac{\sin^2 \alpha}{\|m'_k\|_{L^2(-1, 1)}^2}$$

by the fundamental theorem of calculus and the Cauchy-Schwarz inequality. Thus, discarding all the other connected components, we get a finite number N_k of intervals (a_n^k, b_n^k) , $1 \leq n \leq N_k$, such that

1. $-1 < a_1^k < b_1^k \leq a_2^k < b_2^k \leq \dots \leq a_{N_k}^k < b_{N_k}^k < +1$,
2. $m_{k2}(a_n^k) = \pm \frac{1}{2} \sin \alpha$ and $m_{k2}(b_n^k) = \mp \frac{1}{2} \sin \alpha$ (with opposite sign),
3. $-\frac{1}{2} \sin \alpha < m_{k2} < \frac{1}{2} \sin \alpha$ in (a_n^k, b_n^k) for $n = 1, \dots, N_k$, and
4. $m_{k2} < \frac{1}{2} \sin \alpha$ or $m_{k2} > -\frac{1}{2} \sin \alpha$ in (b_n^k, a_{n+1}^k) for $n = 0, \dots, N_k$, where $b_0^k = -1$ and $a_{N_k+1}^k = 1$

(see Figure 1). Then, using a duality argument and an interpolation inequality for a well-chosen cut-off function, we can prove that

$$\sup_{k \in \mathbb{N}} N_k < \infty.$$

The details for these arguments can be found in the aforementioned paper [13].

In order to prove the relative compactness of $(\phi_k)_{k \in \mathbb{N}}$, we construct step functions $\psi_k : \mathbb{R} \rightarrow 2\pi\mathbb{Z} \pm \alpha$ approximating ϕ_k . To this end, if $N_k > 0$, choose $t_n^k \in (a_n^k, b_n^k)$ such that $\phi_k(t_n^k) \in \pi\mathbb{Z}$ (so




Figure 1: The variations of m_2 .

$m_{k2}(t_n^k) = 0$) for $n = 1, \dots, N_k$. Furthermore, set $t_0^k = -1$ and $t_{N_k+1}^k = 1$. By properties 3 and 4 of the intervals $(a_1^k, b_1^k), \dots, (a_{N_k}^k, b_{N_k}^k)$, the values of m_k are restricted to one of the arcs $\mathcal{C}^\pm = \{m \in \mathbb{S}^1 : \pm m_2 \geq -\frac{1}{2} \sin \alpha\}$ in $[t_{n-1}^k, t_n^k]$ for $n = 1, \dots, N_k + 1$. Hence if we choose $\phi_n^k \in 2\pi\mathbb{Z} \pm \alpha$ such that $(\cos \phi_n^k, \sin \phi_n^k) \in \mathcal{C}^\pm$ as well and $|\phi_n^k - \phi_k(t_n^k)| < \pi$, then

$$|\phi_k - \phi_n^k| < \pi \quad (17)$$

throughout $[t_{n-1}^k, t_n^k]$. Let χ_n^k be the characteristic function of $(t_{n-1}^k, t_n^k]$ and set

$$\psi_k = \sum_{n=1}^{N_k+1} \phi_n^k \chi_n^k.$$

If $N_k = 0$, then $m_k(x_1) \in \mathcal{C}^\pm$ for all $x_1 \in (-1, 1)$ for one of these arcs. It follows that $\phi_k(-1) = \phi_k(1)$ and we set $\psi_k \equiv \phi_k(1)$. Since $\phi_k(-1) = \pm\alpha$, it follows that $|\phi_k| \leq \pi(N_k + 2)$ in $(-1, 1)$. Since $(N_k)_{k \in \mathbb{N}}$ is bounded, we deduce that (ϕ_k) is uniformly bounded in $L^\infty(-1, 1)$. Moreover, inequality (17) implies that there exists a universal constant $C > 0$ with

$$(\phi_k - \psi_k)^2 \leq C |e^{i\phi_k} - e^{i\psi_k}|^2 \quad (18)$$

in $(-1, 1)$. As a consequence of properties 3 and 4, we further have the inequality

$$|\sin \phi_k + \sin \psi_k| = |m_{k2} + \sin \psi_k| \geq \frac{1}{2} \sin \alpha$$

everywhere in $(-1, 1)$. Thus (18) implies that

$$\begin{aligned} \int_{-1}^1 (\phi_k - \psi_k)^2 dx_1 &\leq C \int_{-1}^1 (\cos \phi_k - \cos \psi_k)^2 dx_1 + C \int_{-1}^1 (\sin \phi_k - \sin \psi_k)^2 dx_1 \\ &\leq C \int_{-1}^1 (\cos \phi_k - \cos \alpha)^2 dx_1 + \frac{4C}{\sin^2 \alpha} \int_{-1}^1 (\sin^2 \phi_k - \sin^2 \psi_k)^2 dx_1 \\ &\leq C \left(1 + \frac{16}{\sin^2 \alpha}\right) \int_{-1}^1 (\cos \phi_k - \cos \alpha)^2 dx_1 \rightarrow 0 \end{aligned}$$

as $k \rightarrow \infty$. The last step is due to the convergence $m_{k1} \rightarrow \cos \alpha$ in $L^2(-1, 1)$ established earlier in the proof. It now follows that any accumulation point of $\{\phi_k : k \in \mathbb{N}\}$ will also be an accumulation point of $\{\psi_k : k \in \mathbb{N}\}$ and vice versa.

The uniform bound for N_k implies that the sequence $(\psi_k)_{k \in \mathbb{N}}$ is uniformly bounded in $BV(-1, 1)$ and in $L^\infty(-1, 1)$. Therefore, it is relatively compact in $L^2(-1, 1)$ and any accumulation point belongs to $BV(-1, 1)$ and takes values in $2\pi\mathbb{Z} \pm \alpha$ almost everywhere. \square

2.2 Blow-up of the renormalised energy

In this section we examine the behaviour of the renormalised energy W defined in (4) when two or more Néel walls approach each other or the boundary. This question is more subtle than it may appear, because there are attractive and repulsive terms in the definition of W .

We will work with the quantity

$$\rho(a) = \frac{1}{2} \min\{2a_1 + 2, a_2 - a_1, \dots, a_N - a_{N-1}, 2 - 2a_N\}, \quad (19)$$

defined for $N \geq 1$ (although for the moment we assume that $N \geq 2$).

Lemma 11. *Let $N \geq 2$, $B_1, \dots, B_N < 0$, and $A_{k\ell} \in \mathbb{R}$ for $1 \leq k < \ell \leq N$. Let $f: A_N \rightarrow \mathbb{R}$ be defined by the formula*

$$f(a) = \sum_{k=1}^N B_k \log(1 - a_k^2) + \sum_{1 \leq k < \ell \leq N} A_{k\ell} \log \varrho(a_\ell, a_k), \quad \text{for } a \in A_N,$$

where ϱ is given in (5). Then the following holds true.

1. If $\sum_{K \leq k < \ell \leq L} A_{k\ell} < 0$ for all $K, L \in \{1, \dots, N\}$ with $K < L$, then $f(a) \rightarrow \infty$ as $\rho(a) \rightarrow 0$. As a consequence, f is bounded below.
2. If there exist $K < L$ such that $\sum_{K \leq k < \ell \leq L} A_{k\ell} > 0$, then f is unbounded below, i.e., there exists a sequence $(a^{(i)})_{i \in \mathbb{N}}$ in A_N such that $\rho(a^{(i)}) \rightarrow 0$ and $f(a^{(i)}) \rightarrow -\infty$ as $i \rightarrow \infty$.

We will apply this lemma for numbers of the form $A_{k\ell} = \gamma_k \gamma_\ell$, where γ_n is as in (3). In order to verify the hypothesis in some cases, we use the following result.

Lemma 12. *Let $N \geq 2$ and $d = d_N^+$ or d_N^- , and let $\gamma_n = d_n - \cos \alpha$ for $n = 1, \dots, N$. Suppose that $\Theta \subset (0, \pi)$ is defined as in (9).*

1. If $\alpha \in \Theta$, then for any $K, L \in \{1, \dots, N\}$ with $K < L$:

$$\sum_{K \leq k < \ell \leq L} \gamma_k \gamma_\ell < 0.$$

2. If $N \geq 3$ and $\alpha \notin \bar{\Theta}$, then there exist $K < L$ such that $\sum_{K \leq k < \ell \leq L} \gamma_k \gamma_\ell > 0$.

We prove the two statements in Lemma 11 independently.

Proof of Lemma 11, statement 1. We argue by induction over N . If $N = 2$ and $A_{12} < 0$, then the three terms in $f(a)$ are positive because $a_1, a_2 \in (-1, 1)$ and $\varrho(a_1, a_2) \in (0, 1)$. Assuming that $\rho(a) \rightarrow 0$, we conclude that either one of the points a_1 or a_2 approaches the boundary, or $\varrho(a_1, a_2) \rightarrow 0$. In both cases, the statement is obvious.

Now suppose that the statement is true for all integers between 2 and $N - 1$. Consider a sequence $(a^{(i)})_{i \in \mathbb{N}}$ in A_N with $\rho(a^{(i)}) \rightarrow 0$. We may assume without loss of generality that $a_n^{(i)} \rightarrow a_n$ for some $a_n \in [-1, 1]$ for every $n = 1, \dots, N$. (If not, we choose a subsequence with this property.) By the assumption that $\rho(a^{(i)}) \rightarrow 0$, either one of the limit points a_1 or a_N is on the boundary (i.e., $a_1 = -1$ or $a_N = 1$), or at least two of the limit points a_1, \dots, a_N are equal.

We consider several cases.

Case A. If $a_1 < a_N$, choose a partition $\Lambda_1, \dots, \Lambda_J$ of $\{1, \dots, N\}$ such that $a_k = a_\ell$ whenever $k, \ell \in \Lambda_j$, but $a_k < a_\ell$ if $k \in \Lambda_{j_1}$ and $\ell \in \Lambda_{j_2}$ for $j_1 < j_2$. Note that $|\Lambda_j| < N$ for every $1 \leq j \leq J$.

Case A.1. Suppose first that $J = N$ (i.e., $a_1 < \dots < a_N$) and either $a_1 = -1$ or $a_N = 1$. In this case, the terms in f involving $\log \varrho(a_\ell^{(i)}, a_k^{(i)})$ are uniformly bounded, while at least one boundary interaction term in f (involving $\log(1 - (a_1^{(i)})^2)$ or $\log(1 - (a_N^{(i)})^2)$) blows up to $+\infty$ as $i \rightarrow \infty$.

Case A.2. Now suppose that $J < N$. Then $|\Lambda_j| \geq 2$ for at least one value of j . But for every j such that $|\Lambda_j| \geq 2$, the induction assumption implies that

$$\sum_{k \in \Lambda_j} B_k \log(1 - (a_k^{(i)})^2) + \sum_{\substack{k < \ell \\ k, \ell \in \Lambda_j}} A_{k\ell} \log \varrho(a_\ell^{(i)}, a_k^{(i)}) \rightarrow \infty.$$

All the other terms in the definition of f remain uniformly bounded from below (some boundary interaction term in f might blow up to $+\infty$), and thus, it follows that $f(a^{(i)}) \rightarrow \infty$ as $i \rightarrow \infty$.

Case B. If $a_1 = a_N$, then we distinguish two cases again.

Case B.1. Suppose that $-1 < a_1 = a_N < 1$. We define $\sigma_i = a_N^{(i)} - a_1^{(i)}$ and note that $\sigma_i \rightarrow 0$ as $i \rightarrow \infty$. We set $\tilde{a}^{(i)} = (\tilde{a}_1^{(i)}, \dots, \tilde{a}_N^{(i)})$, where

$$\tilde{a}_n^{(i)} = \frac{a_n^{(i)} - a_1^{(i)}}{2\sigma_i}, \quad 1 \leq n \leq N.$$

Then $\tilde{a}^{(i)} \in A_N$ and $\tilde{a}_1^{(i)} = 0 < \frac{1}{2} = \tilde{a}_N^{(i)}$, so $\tilde{a}^{(i)}$ is consistent with Case A. Since all of the points in $a^{(i)}$ and $\tilde{a}^{(i)}$ stay in an interval of the form $[c - 1, 1 - c]$ for some $c \in (0, 1)$, then all the boundary interaction terms in the formulas for $f(a^{(i)})$ and $f(\tilde{a}^{(i)})$ are uniformly bounded. Furthermore,

$$\log(\tilde{a}_\ell^{(i)} - \tilde{a}_k^{(i)}) - \log 2 \leq \log \varrho(\tilde{a}_\ell^{(i)}, \tilde{a}_k^{(i)}) \leq \log(\tilde{a}_\ell^{(i)} - \tilde{a}_k^{(i)}) - \log(2c - c^2)$$

for every $k < \ell$ and every $i \in \mathbb{N}$, and the same inequalities hold for a . Therefore, there exists $C > 0$ such that for every i ,

$$f(a^{(i)}) \geq f(\tilde{a}^{(i)}) + \sum_{k < \ell} A_{k\ell} \log(2\sigma_i) - C.$$

If $\rho(\tilde{a}^{(i)}) \rightarrow 0$, then the arguments of Case A show that $f(\tilde{a}^{(i)}) \rightarrow \infty$ as $i \rightarrow \infty$. Otherwise, the values $f(\tilde{a}^{(i)})$ will remain bounded. In both cases, we know that $\log(2\sigma_i) \rightarrow -\infty$, while $\sum_{k < \ell} A_{k\ell} < 0$ by the above assumption. It follows that $f(a^{(i)}) \rightarrow \infty$ as $i \rightarrow \infty$.

Case B.2. Finally, assume that $a_1 = a_N \in \{\pm 1\}$. It suffices to consider the case where $a_1 = a_N = 1$ (as the other case is similar). For $b \in (-1, 1)$, the Möbius transform $\Phi_b: \mathbb{R}_+^2 \rightarrow \mathbb{R}_+^2$ is defined by

$$\Phi_b(z) = \frac{z + b}{1 + bz}, \quad z \in \mathbb{C}. \quad (20)$$

It is readily checked that ϱ is invariant under Φ_b .

We set $\tilde{a}_k^{(i)} = \Phi_{-a_1^{(i)}}(a_k^{(i)})$ for $1 \leq k \leq N$. So $\tilde{a}^{(i)} \in A_N$ and $\tilde{a}_1^{(i)} = 0$. If i is sufficiently large, then also $1 - (\tilde{a}_k^{(i)})^2 \geq 1 - (a_k^{(i)})^2$ for $1 \leq k \leq N$, because $a_k^{(i)} \rightarrow 1$ as $i \rightarrow \infty$. As ϱ is invariant under the Möbius transform, we know that $\varrho(\tilde{a}_\ell^{(i)}, \tilde{a}_k^{(i)}) = \varrho(a_\ell^{(i)}, a_k^{(i)})$. Therefore, we deduce that

$$f(a^{(i)}) \geq f(\tilde{a}^{(i)}) + B_1 \log(1 - (a_1^{(i)})^2).$$

If $\rho(\tilde{a}^{(i)}) \rightarrow 0$, then, as $\tilde{a}_1 = 0$, the arguments of Cases A and B.1 apply. Hence $f(\tilde{a}^{(i)}) \rightarrow \infty$ as $i \rightarrow \infty$. Otherwise, the values $f(\tilde{a}^{(i)})$ will remain bounded. In both cases, we know that $B_1 < 0$ and $a_1^{(i)} \rightarrow 1$. Thus $f(a^{(i)}) \rightarrow \infty$ as $i \rightarrow \infty$.

To prove that f is bounded below, we consider a minimising sequence $(a^{(i)})_{i \in \mathbb{N}}$. Then by what we have just proved, $\rho(a^{(i)})$ stays uniformly away from 0. Since f is bounded for $\rho(a) \geq C > 0$, the conclusion follows. \square

Proof of Lemma 11, statement 2. Fix $a \in A_N$ such that $a_K < 0$ and $a_L > 0$. For $\eta \in (0, 1]$, define $a^{(\eta)} = (a_1^{(\eta)}, \dots, a_N^{(\eta)}) \in A_N$ with

$$a_k^{(\eta)} = \begin{cases} a_k & \text{if } k < K \text{ or } k > L \\ \eta a_k & \text{if } K \leq k \leq L. \end{cases}$$

Note that $1 - (a_k^{(\eta)})^2$ stays away from 0 uniformly and $|\log \varrho(a_\ell^{(\eta)}, a_k^{(\eta)}) - \log(a_\ell^{(\eta)} - a_k^{(\eta)})|$ is uniformly bounded for every $k < \ell$ and every $\eta \in (0, 1]$. Thus, there exists a constant $C > 0$ such that

$$\begin{aligned} f(a^{(\eta)}) \leq & C + \sum_{K \leq k < \ell \leq L} A_{k\ell} (\log(a_\ell - a_k) + \log \eta) + \sum_{\substack{k < \ell < K \text{ or} \\ L < k < \ell \text{ or} \\ k < K < L < \ell}} A_{k\ell} \log(a_\ell - a_k) \\ & + \sum_{k < K \leq \ell \leq L} A_{k\ell} \log(\eta a_\ell - a_k) + \sum_{K \leq k \leq L < \ell} A_{k\ell} \log(a_\ell - \eta a_k). \end{aligned}$$

Now we let $\eta \searrow 0$. We note that $\log(\eta a_\ell - a_k) \rightarrow \log |a_k|$ when $k < K$ and $\log(a_\ell - \eta a_k) \rightarrow \log a_\ell$ when $\ell > L$. Thus

$$\limsup_{\eta \searrow 0} \left(f(a^{(\eta)}) - \log \eta \sum_{K \leq k < \ell \leq L} A_{k\ell} \right) < \infty.$$

By the assumption $\sum_{K \leq k < \ell \leq L} A_{k\ell} > 0$, we find that $f(a^{(\eta)}) \rightarrow -\infty$ as $\eta \searrow 0$. \square

Proof of Lemma 12. In order to prove statement 1, we need to show that

$$\sum_{K \leq k < \ell \leq L} (d_k - \cos \alpha)(d_\ell - \cos \alpha) < 0 \tag{21}$$

for all $K, L \in \{1, \dots, N\}$ with $K < L$, provided that $\alpha \in \Theta$.

If $N = 2$, then (21) holds true for every $\alpha \in (0, \pi)$. For $N \geq 3$, we consider the sums

$$S_0^K = \sum_{k=0}^{K-1} \sum_{\ell=k+1}^{K-1} ((-1)^k - \cos \alpha)((-1)^\ell - \cos \alpha)$$

and

$$S_1^K = \sum_{k=1}^K \sum_{\ell=k+1}^K ((-1)^k - \cos \alpha)((-1)^\ell - \cos \alpha)$$

for any $K = 2, \dots, N$. We compute

$$\begin{aligned}
S_0^K &= \frac{1}{2} \left(\sum_{k=0}^{K-1} \sum_{\ell=0}^{K-1} ((-1)^k - \cos \alpha)((-1)^\ell - \cos \alpha) - \sum_{k=0}^{K-1} ((-1)^k - \cos \alpha)^2 \right) \\
&= \frac{1}{2} \sum_{k=0}^{K-1} \sum_{\ell=0}^{K-1} ((-1)^{k+\ell} - 2(-1)^\ell \cos \alpha + \cos^2 \alpha) - \frac{1}{2} \sum_{k=0}^{K-1} (1 - 2(-1)^k \cos \alpha + \cos^2 \alpha) \\
&= \frac{1}{2} \sum_{k=0}^{K-1} \sum_{\ell=0}^{K-1} (-1)^{k+\ell} - K \cos \alpha \sum_{\ell=0}^{K-1} (-1)^\ell + \frac{K^2}{2} \cos^2 \alpha - \frac{K}{2} + \cos \alpha \sum_{k=0}^{K-1} (-1)^k - \frac{K}{2} \cos^2 \alpha \\
&= \frac{1}{2} \left(\sum_{k=0}^{K-1} (-1)^k \right)^2 + (1 - K) \cos \alpha \sum_{k=0}^{K-1} (-1)^k + \frac{K}{2} ((K - 1) \cos^2 \alpha - 1).
\end{aligned}$$

Similarly,

$$S_1^K = \frac{1}{2} \left(\sum_{k=1}^K (-1)^k \right)^2 + (1 - K) \cos \alpha \sum_{k=1}^K (-1)^k + \frac{K}{2} ((K - 1) \cos^2 \alpha - 1).$$

If K is even, then

$$\sum_{k=0}^{K-1} (-1)^k = \sum_{k=1}^K (-1)^k = 0,$$

and thus

$$S_0^K = S_1^K = \frac{K}{2} ((K - 1) \cos^2 \alpha - 1).$$

If K is odd, then

$$\sum_{k=0}^{K-1} (-1)^k = 1 \quad \text{and} \quad \sum_{k=1}^K (-1)^k = -1,$$

and thus

$$S_0^K = \frac{K-1}{2} (K \cos^2 \alpha - 2 \cos \alpha - 1)$$

and

$$S_1^K = \frac{K-1}{2} (K \cos^2 \alpha + 2 \cos \alpha - 1).$$

Hence for any given $N \geq 3$, we conclude that (21) is satisfied under the following conditions.

- If N is even, then (21) reduces to the condition that $S_0^{N-1} < 0$ and $S_1^{N-1} < 0$. This amounts to $(N - 1) \cos^2 \alpha + 2|\cos \alpha| - 1 < 0$, i.e., to the inequalities $\theta_N < \alpha < \pi - \theta_N$.
- If N is odd and $d_1 = 1$, we have the condition that $S_0^N < 0$ and $S_1^{N-2} < 0$, i.e.,

$$\begin{aligned}
N \cos^2 \alpha - 2 \cos \alpha - 1 &< 0, \\
(N - 2) \cos^2 \alpha + 2 \cos \alpha - 1 &< 0.
\end{aligned}$$

If $\cos \alpha \geq 0$, then the second inequality is the strongest, otherwise it is the first. Thus this case amounts to $\theta_{N-2} < \alpha < \pi - \theta_N$.

- if N is odd and $d_1 = -1$, we have the condition that $S_1^N < 0$ and $S_0^{N-2} < 0$, i.e.,

$$\begin{aligned} N \cos^2 \alpha + 2 \cos \alpha - 1 &< 0, \\ (N - 2) \cos^2 \alpha - 2 \cos \alpha - 1 &< 0. \end{aligned}$$

This leads to $\theta_N < \alpha < \pi - \theta_{N-2}$.

The proof of statement 2 is similar. Indeed, if N is even, then the sum can be made positive if, and only if, there exists $K \leq N$ such that $S_0^K > 0$ or $S_1^K > 0$. By the above computations, this is the case if, and only if, $\alpha \notin [\theta_N, \pi - \theta_N]$.

If N is odd, then it is convenient to consider the cases $\cos \alpha \geq 0$ and $\cos \alpha < 0$ separately. Suppose that $\cos \alpha \geq 0$. If $d_1 = 1$ (and thus $d_N = 1$ as well), then we require the existence of $K \leq N$ such that $S_0^K > 0$ or $S_1^{K-1} > 0$. It is readily checked that any of these inequalities will imply that in particular $S_1^{N-2} > 0$. This leads to the condition $\alpha < \theta_{N-2}$. If $d_1 = -1$, then we require $K \leq N$ such that $S_0^{K-1} > 0$ or $S_1^K > 0$. In this case, the term S_1^N is the greatest. It is positive when $\alpha < \theta_N$. The situation for $\cos \alpha < 0$ is similar. \square

We can now answer the question at the beginning of this section. If $\theta_N < \alpha < \pi - \theta_N$, then the repulsion between neighbouring walls of *different signs* will dominate and the renormalised energy will blow up when two such walls approach each other or a wall (of any sign) approaches the boundary. As discussed previously, no such conclusion can be expected when two walls of the same sign approach each other. In fact, by (4), the renormalised energy tends to $-\infty$ when two neighbouring walls of the same sign approach one another. This is consistent with the energy landscape: after the ‘collision’ of two such walls, the number of walls decreases. Thus the total energy, normalised by $(\log \delta)^2$, decreases by $O(|\log \delta|)$. As $\delta \searrow 0$, this should be interpreted as ‘ $-\infty$ ’ in terms of $W(a, d)$.

The above observation can be formulated as follows. Recall that for $a \in A_N$, the quantity $\rho(a)$ is defined in (19).

Proposition 13 (Repulsion of Néel walls). *Let $N \in \mathbb{N}$ such that $\theta_N < \alpha < \pi - \theta_N$. Suppose that $(a^{(i)})_{i \in \mathbb{N}}$ is a sequence in A_N such that $\rho(a^{(i)}) \rightarrow 0$ as $i \rightarrow \infty$ and $(d^{(i)})_{i \in \mathbb{N}}$ is a sequence in $\{\pm 1\}^N$. Suppose further that*

$$\liminf_{i \rightarrow \infty} \min \left\{ a_{k+1}^{(i)} - a_k^{(i)} : 1 \leq k \leq N-1 \text{ and } d_k^{(i)} = d_{k+1}^{(i)} \right\} > 0.$$

Then $W(a^{(i)}, d^{(i)}) \rightarrow \infty$ as $i \rightarrow \infty$.

Proof. By the definition of $W(a, d)$ and $\varrho(a_k, a_\ell)$, there exists a constant $C > 0$ (depending on N) such that

$$W(a, d) \geq -\frac{\pi}{2} \sum_{k=1}^N \gamma_k^2 \log(1 - a_k^2) + \pi \sum_{k < \ell} \gamma_k \gamma_\ell \log \varrho(a_\ell, a_k) - C \quad (22)$$

for any $a \in A_N$ and $d \in \{\pm 1\}^N$, where $\gamma_n = d_n - \cos \alpha$. Under the extra assumption that $d_n^{(i)} = -d_{n+1}^{(i)}$ for $n = 1, \dots, N-1$ and for every $i \in \mathbb{N}$ (i.e., the signs $d_n^{(i)}$ alternate in n), then in view of (22) and Lemma 11 (applied to $A_{k\ell} = \pi \gamma_k \gamma_\ell$ and $B_k = -\frac{\pi}{2} \gamma_k^2$) and Lemma 12, the conclusion follows.

Next we consider the general case. We may assume without loss of generality that there exists a partition $\Lambda_1, \dots, \Lambda_J$ of $\{1, \dots, N\}$ with the property that

$$\liminf_{i \rightarrow \infty} \min \left\{ a_\ell^{(i)} - a_k^{(i)} : 1 \leq k < \ell \leq N \text{ and } \{k, \ell\} \not\subseteq \Lambda_j \text{ for } j = 1, \dots, J \right\} > 0,$$

while

$$\lim_{i \rightarrow \infty} \max \left\{ a_\ell^{(i)} - a_k^{(i)} : 1 \leq k < \ell \leq N \text{ and } k, \ell \in \Lambda_j \right\} = 0$$

for any $j = 1, \dots, J$ such that $|\Lambda_j| \geq 2$. Otherwise, we pass to a subsequence with this property. (We think of each Λ_j as the set of indices of a cluster of points approaching one another as $i \rightarrow \infty$.)

We may further assume that $d_k^{(i)}$ is independent of i .

Fix $j \in \{1, \dots, J\}$. Then for any sufficiently large value of i , it is clear that Λ_j comprises a consecutive set of numbers, i.e., there exist $K_j, L_j \in \{1, \dots, N\}$ such that $\Lambda_j = \{K_j, \dots, L_j\}$, since the points of each $a^{(i)}$ are ordered. The hypothesis of the proposition then implies that for any fixed j , either $d_k^{(i)} = (-1)^k$ for all $k \in \Lambda_j$ or $d_k^{(i)} = -(-1)^k$ for all $k \in \Lambda_j$. Since θ_N is increasing in N , Lemma 11 and Lemma 12 show that

$$-\frac{\pi}{2} \sum_{k=K_j}^{L_j} (\gamma_k^{(i)})^2 \log(1 - (a_k^{(i)})^2) + \pi \sum_{K_j \leq k < \ell \leq L_j} \gamma_k^{(i)} \gamma_\ell^{(i)} \log \varrho(a_\ell^{(i)}, a_k^{(i)}) \rightarrow \infty$$

as $i \rightarrow \infty$ for every $j = 1, \dots, J$ such that $|\Lambda_j| \geq 2$. (Here $\gamma_k^{(i)} = d_k^{(i)} - \cos \alpha$.) Finally, we observe that the condition $\rho(a^{(i)}) \rightarrow 0$ implies that either $a_1^{(i)} \rightarrow -1$ or $a_N^{(i)} \rightarrow 1$ or there exists $j \in \{1, \dots, N\}$ with $|\Lambda_j| \geq 2$. As we have inequality (22), the claim then follows. \square

Now we have all the tools for the proof of Proposition 6.

Proof of Proposition 6. For the proof of statement 1, let $(a^{(i)})_{i \in \mathbb{N}}$ be a minimising sequence of $W(\cdot, d)$ over A_N . (By what we know so far, this might mean that $W(a^{(i)}, d) \rightarrow -\infty$ as $i \rightarrow \infty$.) Assume by contradiction that for a subsequence (still denoted by $(a^{(i)})_{i \in \mathbb{N}}$) we have $\rho(a^{(i)}) \rightarrow 0$ as $i \rightarrow \infty$. Then we use (22) to estimate W from below in terms of quantities that can be controlled with the help of Lemmas 11 and 12. We conclude that $W(a^{(i)}, d) \rightarrow +\infty$, which contradicts the minimising character of $(a^{(i)})_{i \in \mathbb{N}}$. Thus, $\liminf_{i \rightarrow \infty} \rho(a^{(i)}) > 0$. Passing to a subsequence if necessary, we may assume that $a^{(i)} \rightarrow a$ as $i \rightarrow \infty$ with $a \in A_N$. By definition (4) of $W(\cdot, d)$, we deduce that $W(a^{(i)}, d) \rightarrow W(a, d)$ as $i \rightarrow \infty$, i.e., the infimum of $W(\cdot, d)$ over A_N is finite and achieved by a .

For statement 2, we first use Lemma 12, which tells us that for $N \geq 3$ and $\alpha \notin \bar{\Theta}$, there exist $K < L$ such that $\sum_{K \leq k < \ell \leq L} \gamma_k \gamma_\ell > 0$. Therefore, by Lemma 11, there exists a sequence $(a^{(i)})_{i \in \mathbb{N}}$ in A_N such that $\rho(a^{(i)}) \rightarrow 0$ and

$$-\frac{\pi}{2} \sum_{k=1}^N \gamma_k^2 \log(1 - (a_k^{(i)})^2) + \pi \sum_{1 \leq k < \ell \leq N} \gamma_k \gamma_\ell \log \varrho(a_\ell^{(i)}, a_k^{(i)}) \rightarrow -\infty$$

as $i \rightarrow \infty$. Comparing with definition (4), we see that there exists a constant C such that

$$W(a^{(i)}, d) \leq C - \frac{\pi}{2} \sum_{k=1}^N \gamma_k^2 \log(1 - (a_k^{(i)})^2) + \pi \sum_{1 \leq k < \ell \leq N} \gamma_k \gamma_\ell \log \varrho(a_\ell^{(i)}, a_k^{(i)}).$$

Hence $W(a^{(i)}, d) \rightarrow -\infty$ as well. \square

2.3 Energy estimates

In this section we improve some of the energy estimates from our previous paper [18]. In particular, this will remove the need to bound the distance between two Néel walls from below.

We use several of the tools from the previous paper here. Therefore, we discuss them briefly before giving some improved estimates. This includes in particular the construction of a limiting

stray field potential, given in terms of a function on the upper half-plane $\mathbb{R}_+^2 = \mathbb{R} \times (0, \infty)$ satisfying a certain boundary value problem.

We use the following notation. For $r > 0$, we set

$$\Omega_r(a) = \mathbb{R}_+^2 \setminus \bigcup_{n=1}^N B_r(a_n, 0),$$

where $B_r(x)$ denotes the open disk centred at x of radius $r > 0$. Apart from the half-plane \mathbb{R}_+^2 , which is sometimes regarded as a subset of \mathbb{C} by the usual identification, we consider

$$S = \{x_1 + ix_2 \in \mathbb{C} : x_1 > 0, 0 < x_2 < \pi\}$$

and the map $F: S \rightarrow \mathbb{R}_+^2$ given by

$$F(w) = -\frac{1}{\cosh w} = -\frac{2}{e^w + e^{-w}}, \quad w \in S.$$

Furthermore, we consider the function $\hat{u}: S \rightarrow \mathbb{R}$ with

$$\hat{u}(w) = \frac{\pi}{2} - \operatorname{Im} w$$

and $u: \mathbb{R}_+^2 \rightarrow \mathbb{R}$ given by

$$u = \hat{u} \circ F^{-1}.$$

This function solves the boundary value problem

$$\begin{aligned} \Delta u &= 0 && \text{in } \mathbb{R}_+^2, \\ u &= \frac{\pi}{2} && \text{on } \{0\} \times (-1, 0), \\ u &= -\frac{\pi}{2} && \text{on } \{0\} \times (0, 1), \\ \frac{\partial u}{\partial x_2} &= 0 && \text{on } \{0\} \times (-\infty, -1) \text{ and on } \{0\} \times (1, \infty). \end{aligned}$$

Note that $|u| \leq \frac{\pi}{2}$ in \mathbb{R}_+^2 and $u(z) \rightarrow 0$ as $|z| \rightarrow \infty$. For $b \in (-1, 1)$, we recall the Möbius transform Φ_b defined in (20). Observe that $\Phi_b^{-1} = \Phi_{-b}$. We define

$$u_b = u \circ \Phi_{-b}.$$

For $a \in A_N$ and $d \in \{\pm 1\}^N$, set $\gamma_n = d_n - \cos \alpha$ and

$$u_{a,d}^* = \sum_{n=1}^N \gamma_n u_{a_n}.$$

This function plays the role of a limiting stray field potential for an array of Néel walls at the points a_1, \dots, a_N of signs d_1, \dots, d_N . The renormalised energy is related to

$$\frac{1}{2} \lim_{r \searrow 0} \left(\int_{\Omega_r(a)} |\nabla u_{a,d}^*|^2 dx - \pi \sum_{n=1}^N \gamma_n^2 \log \frac{1}{r} \right), \quad (23)$$

which happens to equal $-W(a, d)$. However, the full renormalised energy contains another term, which turns out to be $2W(a, d)$, giving $W(a, d)$ as the sum [18, page 442]. The quantity in (23) can be identified as the contribution of the interaction between all the logarithmically decaying tails of

the Néel walls, whereas the other term corresponds to the interaction between pairs of a tail from one wall and the core of another. This relation between the two terms can surely be no coincidence, and indeed we discover something similar for the unconfined problem below, but the reason is unclear.

In order to improve the results from the previous paper [18], we need above all to refine some estimates for the Dirichlet energy in (23). We begin with a result similar to [18, Lemma 9], but we prove an estimate in the half-disk $B_R^+(b, 0) = B_R(b, 0) \cap \mathbb{R}_+^2$ instead of the half-space \mathbb{R}_+^2 . This is the natural estimate in the context of Ginzburg-Landau theory, because u_b behaves like the phase of a vortex of degree 1.

Lemma 14. *Let $b \in (-1, 1)$ and $0 < r < R \leq 1 - |b|$. Then*

$$\int_{B_R^+(b, 0) \setminus B_r(b, 0)} |\nabla u_b|^2 dx \leq \pi \log \frac{R}{r} + \frac{\pi}{2} \log 76.$$

Proof. By the symmetry, it suffices to consider the case $b \in [0, 1)$. The first step is to examine the set $\Phi_{-b}(B_R^+(b, 0) \setminus B_r(b, 0))$. To this end, we fix $\rho \in (0, 1 - b]$ and we observe that $\Phi_{-b}(\partial^+ B_\rho(b, 0))$ is a semicircle centred on $\mathbb{R} \times \{0\}$ by the standard properties of the Möbius transform (where $\partial^+ B_\rho(b, 0) = \partial B_\rho(b, 0) \cap \mathbb{R}_+^2$). In order to determine this semicircle, it suffices to compute

$$\Phi_{-b}(b + \rho) = \frac{\rho}{1 - b^2 - b\rho} \quad \text{and} \quad \Phi_{-b}(b - \rho) = -\frac{\rho}{1 - b^2 + b\rho}.$$

Observing that $\frac{\rho}{1 - b^2 - b\rho} > \frac{\rho}{1 - b^2 + b\rho}$, we then see that

$$\Phi_{-b}(B_\rho^+(b, 0)) \subseteq B_{\rho/(1 - b^2 - b\rho)}^+(0), \quad \Phi_{-b}(\mathbb{R}_+^2 \setminus B_\rho(b, 0)) \subseteq \mathbb{R}_+^2 \setminus B_{\rho/(1 - b^2 + b\rho)}^+(0).$$

Thus

$$\Phi_{-b}(B_R^+(b, 0) \setminus B_r(b, 0)) \subseteq B_{R/(1 - b^2 - bR)}^+(0) \setminus B_{r/(1 - b^2 + br)}(0).$$

Set $\tilde{R} = R/(1 - b^2 - bR)$ and $\tilde{r} = r/(1 - b^2 + br)$. Using the identity

$$|\cosh w|^2 = \frac{1}{2} \cosh(2 \operatorname{Re} w) + \frac{1}{2} \cos(2 \operatorname{Im} w), \quad w \in \mathbb{C},$$

we see that for any $\rho > 0$,

$$\begin{aligned} F^{-1}(\mathbb{R}_+^2 \setminus B_\rho(0)) &\subseteq \left\{ w \in S : \operatorname{Re} w \leq \frac{1}{2} \operatorname{arcosh} \left(\frac{2}{\rho^2} + 1 \right) \right\}, \\ F^{-1}(B_\rho^+(0)) &\subseteq \left\{ w \in S : \operatorname{Re} w > \frac{1}{2} \operatorname{arcosh} \left(\frac{2}{\rho^2} - 1 \right) \right\}. \end{aligned}$$

As conformal maps leave the Dirichlet energy invariant, it follows that

$$\begin{aligned} \int_{B_R^+(b, 0) \setminus B_r(b, 0)} |\nabla u_b|^2 dx &\leq \int_{B_{\tilde{R}}^+(0) \setminus B_{\tilde{r}}(0)} |\nabla u|^2 dx \\ &\leq \int_{\{w \in S : \operatorname{arcosh}(2/\tilde{R}^2 - 1) < 2 \operatorname{Re} w < \operatorname{arcosh}(2/\tilde{r}^2 + 1)\}} |\nabla \hat{u}|^2 dx \\ &= \frac{\pi}{2} \left(\operatorname{arcosh} \left(\frac{2(1 - b^2 + br)^2}{r^2} + 1 \right) - \operatorname{arcosh} \left(\frac{2(1 - b^2 - bR)^2}{R^2} - 1 \right) \right). \end{aligned}$$

Note that for any $y \geq 1$,

$$|\operatorname{arcosh} y - \log y| = \left| \log \left(y + \sqrt{y^2 - 1} \right) - \log y \right| = \log \left(1 + \sqrt{1 - \frac{1}{y^2}} \right) \leq \log 2.$$

Hence

$$\int_{B_R^+(b,0) \setminus B_r(b,0)} |\nabla u_b|^2 dx \leq \frac{\pi}{2} \log \left(\frac{R^2(2(1-b^2+br)^2+r^2)}{r^2(2(1-b^2-bR)^2-R^2)} \right) + \pi \log 2.$$

Using the inequality $R \leq 1-b$, we obtain $1-b^2-bR \geq 1-b^2-b(1-b) = 1-b$. Similarly, $1-b^2+br \leq 1-b^2+b(1-b) = (1+2b)(1-b)$. Therefore, using $r \leq 1-b$ and $R \leq 1-b$ also to estimate r^2 and R^2 , we conclude that

$$\begin{aligned} \int_{B_R^+(b,0) \setminus B_r(b,0)} |\nabla u_b|^2 dx &\leq \pi \log \frac{R}{r} + \frac{\pi}{2} \log \left(\frac{2(1+2b)^2(1-b)^2 + (1-b)^2}{2(1-b)^2 - (1-b)^2} \right) + \pi \log 2 \\ &= \pi \log \frac{R}{r} + \frac{\pi}{2} \log (2(1+2b)^2 + 1) + \pi \log 2 \\ &\leq \pi \log \frac{R}{r} + \frac{\pi}{2} \log 19 + \pi \log 2 \\ &\leq \pi \log \frac{R}{r} + \frac{\pi}{2} \log 76, \end{aligned}$$

as required. \square

Note that $u_{a,d}^*$ does not belong to $\dot{H}^1(\mathbb{R}_+^2)$ because of the singularities at $(a_n, 0)$. As a consequence of the preceding inequality, however, we can regularise it near the singular points and at the same time obtain good estimates.

Lemma 15. *Let $c > 0$ and $N \in \mathbb{N}$. Then there exists $C > 0$ such that for any $a \in A_N$ and $d \in \{\pm 1\}^N$ and for any $\epsilon \in (0, \frac{1}{2}]$ with $\rho(a) \geq c\delta = c\epsilon|\log \epsilon|$, there exists $\xi \in \dot{H}^1(\mathbb{R}_+^2)$ such that $\xi = u_{a,d}^*$ in $\Omega_{c\delta}(a)$ and such that the inequalities*

$$\sup_{x_1 \in \mathbb{R}} |u_{a,d}^*(x_1, 0) - \xi(x_1, 0)| \leq \pi$$

and¹⁰

$$\int_{\mathbb{R}_+^2} |\nabla \xi|^2 dx \leq \pi \log \frac{1}{\delta} \sum_{n=1}^N \gamma_n^2 - 2W(a, d) + C$$

are satisfied, where $\gamma_n = d_n - \cos \alpha$ for $n = 1, \dots, N$.

Proof. Fix $n \in \{1, \dots, N\}$. Then Lemma 14 implies that

$$\int_{B_{c\delta}^+(a_n, 0) \setminus B_{c\delta/2}(a_n, 0)} |\nabla u_{a_n}|^2 dx \leq \frac{\pi}{2} \log 304.$$

Note that the function

$$f(x) = u_{a_n}(x_1 + a_n, x_2) + \arctan \frac{x_1}{x_2}$$

is harmonic in \mathbb{R}_+^2 and constant on $(-c\delta, c\delta) \times \{0\}$. Moreover, as $|\nabla f|^2(x) \leq 2(|\nabla u_{a_n}|^2(x_1 + a_n, x_2) + 1/|x|^2)$, we estimate

$$\int_{B_{c\delta}^+(0) \setminus B_{c\delta/2}(0)} |\nabla f|^2 dx \leq \pi \log 1216. \quad (24)$$

We may extend the function $\tilde{f}(x) = f(x) - f(0)$ to $B_{c\delta}(0)$ by the odd reflection $\tilde{f}(x_1, -x_2) = -\tilde{f}(x_1, x_2)$ for $(x_1, x_2) \in B_{c\delta}^+(0)$. Then \tilde{f} is harmonic in $B_{c\delta}(0)$ and so are its derivatives. The mean value formula then gives

$$\left| \frac{\partial \tilde{f}}{\partial x_i}(x) \right| \leq \frac{16}{\pi c^2 \delta^2} \int_{B_{c\delta/4}(x)} \left| \frac{\partial \tilde{f}}{\partial x_i}(y) \right| dy$$

¹⁰This inequality provides an upper bound for the quantity (23), because $\xi = u_{a,d}^*$ in $\Omega_{c\delta}(a)$.

for any $x \in \partial B_{3c\delta/4}(0)$. The maximum principle allows us to extend this inequality to all $x \in B_{3c\delta/4}(0)$. Combining the resulting estimate with (24) and Hölder's inequality, we obtain a universal constant C_1 such that

$$\|\nabla f\|_{L^\infty(B_{c\delta/2}^+(0))} \leq \frac{C_1}{c\delta}.$$

That is,

$$\left| \nabla u_{a_n}(x) + \frac{(x_2, a_n - x_1)}{(x_1 - a_n)^2 + x_2^2} \right| \leq \frac{C_1}{c\delta} \quad (25)$$

for every $x \in B_{c\delta/2}^+(a_n, 0)$.

In addition, for $k \neq n$, we obtain the inequality

$$\begin{aligned} \int_{B_{c\delta}^+(a_k, 0)} |\nabla u_{a_n}|^2 dx &\leq \int_{B_{|a_n - a_k| + c\delta}^+(a_n, 0) \setminus B_{|a_n - a_k| - c\delta}(a_n, 0)} |\nabla u_{a_n}|^2 dx \\ &\leq \pi \log 3 + \frac{\pi}{2} \log 76 = \frac{\pi}{2} \log 684. \end{aligned}$$

(Here we have used Lemma 14 and the fact that $|a_n - a_k| \geq 2c\delta$ for $k \neq n$.) From this we conclude, as above, that

$$|\nabla u_{a_n}(x)| \leq \frac{C_2}{c\delta} \quad (26)$$

in $B_{c\delta/2}^+(a_k, 0)$ for a universal constant C_2 .

Now choose $\eta \in C_0^\infty(B_{c\delta/2}(0))$ with $\eta \equiv 1$ in $B_{c\delta/4}(0)$ and with $0 \leq \eta \leq 1$ and $|\nabla \eta| \leq \frac{8}{c\delta}$ everywhere. Set

$$\tilde{u}_{a_n}(x) = (1 - \eta(x_1 - a_n, x_2))u_{a_n}(x) + \eta(x_1 - a_n, x_2) \int_{B_{c\delta/2}^+(a_n, 0) \setminus B_{c\delta/4}(a_n, 0)} u_{a_n} dy.$$

As $|u| \leq \frac{\pi}{2}$, we have $-\frac{\pi}{2} \leq \tilde{u}_{a_n} \leq \frac{\pi}{2}$. Moreover, it is clear that \tilde{u}_{a_n} coincides with u_{a_n} in $\mathbb{R}_+^2 \setminus B_{c\delta/2}(a_n, 0)$. Thus, by Lemma 9 in [18], we find universal constants C_3, C_4 such that

$$\begin{aligned} \int_{\mathbb{R}_+^2} |\nabla \tilde{u}_{a_n}|^2 dx &\leq \int_{\mathbb{R}_+^2 \setminus B_{c\delta}(a_n, 0)} |\nabla u_{a_n}|^2 dx + C_3 \\ &\leq \pi \log \frac{1}{\delta} + \pi \log \frac{1}{c} + \pi \log(2 - 2a_n^2) + \frac{C_4 c \delta |a_n|}{1 - a_n^2} + \frac{C_4 c^2 \delta^2}{(1 - a_n^2 - c\delta |a_n|)^2} + C_3. \end{aligned}$$

Since $1 - a_n^2 = (1 - a_n)(1 + a_n) \geq c\delta$ and $1 - a_n^2 - c\delta |a_n| = (1 - |a_n|)(1 + |a_n|) - c\delta |a_n| \geq c\delta(1 + |a_n|) - c\delta |a_n| = c\delta$, we conclude that

$$\int_{\mathbb{R}_+^2} |\nabla \tilde{u}_{a_n}|^2 dx \leq \pi \log \frac{1}{\delta} + \pi \log(2 - 2a_n^2) + C_5 \quad (27)$$

for a constant C_5 that depends only on c .

Furthermore, for $k \neq n$,

$$\begin{aligned} \int_{\mathbb{R}_+^2} \nabla \tilde{u}_{a_k} \cdot \nabla \tilde{u}_{a_n} dx &= \int_{\mathbb{R}_+^2} \nabla u_{a_k} \cdot \nabla u_{a_n} dx + \int_{\mathbb{R}_+^2} \nabla \tilde{u}_{a_k} \cdot (\nabla \tilde{u}_{a_n} - \nabla u_{a_n}) dx \\ &\quad + \int_{\mathbb{R}_+^2} (\nabla \tilde{u}_{a_k} - \nabla u_{a_k}) \cdot \nabla u_{a_n} dx. \end{aligned}$$

Define $\zeta_n(x) = \arctan \frac{x_1 - a_n}{x_2}$. Then we estimate

$$\begin{aligned} \int_{\mathbb{R}_+^2} \nabla \tilde{u}_{a_k} \cdot (\nabla \tilde{u}_{a_n} - \nabla u_{a_n}) dx &= \int_{B_{c\delta/2}^+(a_n, 0)} \nabla \tilde{u}_{a_k} \cdot (\nabla \tilde{u}_{a_n} - \nabla u_{a_n}) dx \\ &\leq \frac{\pi c^2 \delta^2}{8} \|\nabla \tilde{u}_{a_k}\|_{L^\infty(B_{c\delta/2}^+(a_n, 0))} \|\nabla \tilde{u}_{a_n}\|_{L^\infty(B_{c\delta/2}^+(a_n, 0))} \\ &\quad + \frac{\pi c^2 \delta^2}{8} \|\nabla \tilde{u}_{a_k}\|_{L^\infty(B_{c\delta/2}^+(a_n, 0))} \|\nabla u_{a_n} + \nabla \zeta_n\|_{L^\infty(B_{c\delta/2}^+(a_n, 0))} \\ &\quad + \|\nabla \tilde{u}_{a_k}\|_{L^\infty(B_{c\delta/2}^+(a_n, 0))} \|\nabla \zeta_n\|_{L^1(B_{c\delta/2}^+(a_n, 0))}. \end{aligned}$$

Using (26) and (25) and observing that

$$\|\nabla \zeta\|_{L^1(B_{c\delta/2}^+(a_n, 0))} = \frac{c\delta\pi}{2},$$

we therefore find that there exists a constant $C_6 = C_6(c)$ such that

$$\int_{\mathbb{R}_+^2} \nabla \tilde{u}_{a_k} \cdot (\nabla \tilde{u}_{a_n} - \nabla u_{a_n}) dx \leq C_6.$$

Similarly, we prove that

$$\int_{\mathbb{R}_+^2} (\nabla \tilde{u}_{a_k} - \nabla u_{a_k}) \cdot \nabla u_{a_n} dx = \int_{B_{c\delta/2}^+(a_k, 0)} (\nabla \tilde{u}_{a_k} - \nabla u_{a_k}) \cdot \nabla \tilde{u}_{a_n} dx \leq C_6,$$

because $\tilde{u}_{a_n} = u_{a_n}$ on $\mathbb{R}_+^2 \setminus B_{c\delta/2}^+(a_n, 0)$ and $|a_n - a_k| \geq 2c\delta$. By [18, Lemma 8], we obtain

$$\int_{\mathbb{R}_+^2} \nabla \tilde{u}_{a_k} \cdot \nabla \tilde{u}_{a_n} dx \leq \int_{\mathbb{R}_+^2} \nabla u_{a_k} \cdot \nabla u_{a_n} dx + 2C_6 = \pi \log \left(\frac{1 + \sqrt{1 - \varrho(a_k, a_n)^2}}{\varrho(a_k, a_n)} \right) + 2C_6. \quad (28)$$

We now set

$$\xi = \sum_{n=1}^N \gamma_n \tilde{u}_{a_n},$$

then the desired inequality for the Dirichlet energy follows from inequalities (27) and (28). The inequality for $u_{a,d}^*(x_1, 0) - \xi(x_1, 0)$ is a consequence of the construction. \square

Under assumptions similar to Proposition 13, where only neighbouring walls of different sign can approach one another, we prove the following a priori lower bound for the energy E_ϵ .

Proposition 16 (A priori lower energy bound). *Let $N \in \mathbb{N}$ and $\sigma > 0$. Suppose that $\theta_N < \alpha < \pi - \theta_N$. Then there exists C_0 with the following property. Suppose that $a \in A_N$ and $d \in \{\pm 1\}^N$ such that for $n = 1, \dots, N-1$, either $a_{n+1} - a_n \geq \sigma$ or $d_{n+1} = -d_n$. Then the inequality*

$$E_\epsilon(m) \geq \frac{\pi}{2 \log \frac{1}{\delta}} \sum_{n=1}^N \gamma_n^2 + \frac{W(a, d) - C_0}{(\log \delta)^2}$$

holds true for all $m \in M(a, d)$ and all $\epsilon \in (0, \frac{1}{4}]$.

The proof depends in part on the following uniform bound on the renormalised energy $W(a, d)$, which holds if $\rho(a) \geq c\delta$.

Lemma 17. *Let $c > 0$. There exist $\tilde{C}_0 > 0$ and $\hat{C}_0 > 0$ such that for every $a \in A_N$ with $\rho(a) \geq c\delta$ and for every $d \in \{\pm 1\}^N$, if $m \in M(a, d)$ satisfies*

$$E_\epsilon(m) \leq \frac{\pi}{2 \log \frac{1}{\delta}} \sum_{n=1}^N \gamma_n^2 + \frac{W(a, d) - \tilde{C}_0}{(\log \delta)^2},$$

then $|W(a, d)| \leq \hat{C}_0$.

Proof. We first note that E_ϵ attains its minimum in $M(a, d)$, as observed in our previous paper [18, Proposition 1]. We may therefore assume that m minimises E_ϵ in $M(a, d)$, as otherwise, we may simply replace it with a minimiser.

If $\rho(a) \geq c\delta$, then $W(a, d)$ is of order $|\log \delta|$. In particular, there exists a constant $C_1 = C_1(c, N)$ such that $E_\epsilon(m) \leq \frac{C_1}{|\log \delta|}$. Thus, writing $m = (\cos \phi, \sin \phi)$, by [18, Lemma 13] and the Cauchy-Schwarz inequality, we obtain

$$\sum_{n=1}^N \int_{a_n - c\delta}^{a_n + c\delta} |m'_1| dx_1 \leq \sum_{n=1}^N \sqrt{\int_{a_n - c\delta}^{a_n + c\delta} \sin^2 \varphi dx_1} \sqrt{\int_{a_n - c\delta}^{a_n + c\delta} (\varphi')^2 dx_1} \leq \frac{C_2}{\log \frac{1}{\delta}}$$

for a constant C_2 that depends only on c and N . Set

$$\Gamma = \sum_{n=1}^N \gamma_n^2$$

and let ξ be the function from Lemma 15. Note that $u_{a,d}^*(\cdot, 0)$ is constant in (a_{n-1}, a_n) for $n = 1, \dots, N+1$ (where $a_0 = -1$ and $a_{N+1} = 1$) with a jump of size $-\gamma_n \pi$ at a_n for $n = 1, \dots, N$. The function m_1 , on the other hand, satisfies $m_1(a_n) = d_n$. We use the fundamental theorem of calculus in each of the intervals (a_{n-1}, a_n) , obtaining

$$\begin{aligned} \pi \Gamma &= \int_{-1}^1 u_{a,d}^*(x_1, 0) m'_1(x_1) dx_1 \\ &= \int_{-1}^1 (u_{a,d}^*(x_1, 0) - \xi(x_1, 0)) m'_1(x_1) dx_1 + \int_{-1}^1 \xi(x_1, 0) m'_1(x_1) dx_1. \end{aligned}$$

Moreover,

$$\int_{-1}^1 (u_{a,d}^*(x_1, 0) - \xi(x_1, 0)) m'_1(x_1) dx_1 \leq \pi \sum_{n=1}^N \int_{a_n - c\delta}^{a_n + c\delta} |m'_1| dx_1 \leq \frac{C_2 \pi}{\log \frac{1}{\delta}},$$

while

$$\int_{-1}^1 \xi(x_1, 0) m'_1(x_1) dx_1 = \int_{\mathbb{R}_+^2} \nabla \xi \cdot \nabla u dx \leq \|\nabla \xi\|_{L^2(\mathbb{R}_+^2)} \sqrt{2E_\epsilon(m)}.$$

The inequality of Lemma 15 then gives a constant $C_3 = C_3(N, c) > 0$ such that

$$\pi^2 \left(\Gamma - \frac{C_2}{\log \frac{1}{\delta}} \right)^2 \leq \left(\pi \Gamma \log \frac{1}{\delta} - 2W(a, d) + C_3 \right) \left(\frac{\pi \Gamma}{\log \frac{1}{\delta}} + \frac{2W(a, d) - 2\tilde{C}_0}{(\log \delta)^2} \right),$$

which implies that

$$\begin{aligned} \pi^2 \Gamma^2 - \frac{2C_2 \pi^2 \Gamma}{\log \frac{1}{\delta}} + \frac{C_2^2 \pi^2}{(\log \delta)^2} \\ \leq \pi^2 \Gamma^2 + \frac{\pi \Gamma (C_3 - 2\tilde{C}_0)}{\log \frac{1}{\delta}} - \frac{4(W(a, d))^2}{(\log \delta)^2} + \frac{2W(a, d)(C_3 + 2\tilde{C}_0)}{(\log \delta)^2} - \frac{2\tilde{C}_0 C_3}{(\log \delta)^2}. \end{aligned}$$

If we choose $\tilde{C}_0 \geq C_2\pi - \frac{C_3}{2}$, then it follows that

$$4(W(a, d))^2 \leq 2W(a, d)(C_3 + 2\tilde{C}_0).$$

Therefore, the quantity $W(a, d)$ is bounded by a constant that depends only on N and c . \square

Proof of Proposition 16. We use different arguments depending on the magnitude of $\rho(a)/\delta$.

Step 1: small distances. For a fixed number $c \leq 1$, depending only on N (but to be determined later), we first show that the desired inequality holds when $\rho(a) < c\delta$.

To this end, let $h = \rho(a)$ and set $q = 1 - \cos \theta_N$. Since $m \in M(a, d)$, there exist $b_1, b_2 \in [-1, 1]$ with $b_1 < b_2$ and $b_2 - b_1 \leq h$ such that $m_1(b_1) = \pm 1$ and $m_1(b_2) = \cos \alpha$ or vice versa. Hence

$$q \leq \left| \int_{b_1}^{b_2} m'_1 dx \right| \leq \sqrt{\frac{2h}{\epsilon}} E_\epsilon(m)$$

by the fundamental theorem of calculus and the Cauchy-Schwarz inequality. It follows that

$$E_\epsilon(m) \geq \frac{q^2 \epsilon}{2h}.$$

On the other hand, by the definition of W , it is clear that there exists a constant $C_1 = C_1(N)$ such that the inequality $h < \delta$ will imply that

$$\frac{\pi}{2 \log \frac{1}{\delta}} \sum_{n=1}^N \gamma_n^2 + \frac{W(a, d)}{(\log \delta)^2} \leq \frac{C_1 \log \frac{1}{h}}{(\log \delta)^2}.$$

Thus if we choose $c \leq 1$, then it suffices to show that

$$\frac{q^2 \epsilon}{2h} \geq \frac{C_1 \log \frac{1}{h}}{(\log \delta)^2}.$$

As $\epsilon |\log \delta| = \delta - \epsilon \log |\log \epsilon|$, this inequality is equivalent to

$$q^2 \left(\delta \log \frac{1}{\delta} - \epsilon \log \frac{1}{\delta} \log \log \frac{1}{\epsilon} \right) \geq 2C_1 h \log \frac{1}{h}.$$

It is easy to see that $\epsilon \log \log \frac{1}{\epsilon} \leq \delta/2$ for all $\epsilon \in (0, 1)$. Thus it suffices to show that

$$q^2 \delta \log \frac{1}{\delta} \geq 4C_1 h \log \frac{1}{h}$$

for $h \leq c\delta$. But this is clear if $c > 0$ is chosen sufficiently small.

Before we proceed to the second step, we point out that so far, we have not used the hypothesis that $a_{n+1} - a_n \geq \sigma$ or $d_{n+1} = -d_n$ for all n . This is necessary only when $\rho(a) \geq c\delta$.

Step 2: large distances. Now we assume that $\rho(a) \geq c\delta$. If m does *not* satisfy the inequality in the hypothesis of Lemma 17, then there is nothing to prove. Thus, by Lemma 17, we may assume that $|W(a, d)|$ is bounded by a constant independent of a or d .

Due to the hypothesis that $\alpha \in (\theta_N, \pi - \theta_N)$ and that either $a_{n+1} - a_n \geq \sigma$ or $d_{n+1} = -d_n$, Proposition 13 applies. It follows that the Néel walls are uniformly separated in the sense that $\rho(a) \geq C_2$ for a constant $C_2 > 0$ that depends only on N . Now we can use the theory of our previous paper [18], and in particular Theorem 28, which gives a constant $C_3 = C_3(N, \sigma)$ such that

$$E_\epsilon(m) \geq \frac{\pi}{2 \log \frac{1}{\delta}} \sum_{n=1}^N \gamma_n^2 - \frac{C_3}{(\log \delta)^2}.$$

As we already have a bound for $|W(a, d)|$, it now suffices to choose C_0 sufficiently large. \square

2.4 Proofs of the main results

Proof of Proposition 2. Let $a \in A_N$ and set $R = \frac{1}{2}\rho(a)$. Fix $d \in \{\pm 1\}^N$. For $\epsilon \in (0, \frac{1}{4}]$, choose minimisers m_ϵ of E_ϵ in $M(a, d)$. These exist and are smooth by [18, Proposition 1].

Given $b \in A_N$ with $|b_n - a_n| \leq r \leq R/2$ for $n = 1, \dots, N$, we construct a map $G: \overline{\mathbb{R}_+^2} \rightarrow \overline{\mathbb{R}_+^2}$ as follows. For $n = 1, \dots, N$, if $x \in B_R^+(a_n, 0)$, then we set $G(x) = x + (b_n - a_n, 0)$. Furthermore, we set $G(x) = x$ for all $x \in \Omega_{2R}(a)$. Then we extend G to $\overline{\mathbb{R}_+^2}$ such that it becomes bijective and a bi-Lipschitz map, and such that $G(x_1, 0) \in \mathbb{R} \times \{0\}$ for all $x_1 \in \mathbb{R}$. We set $g(x_1) = G(x_1, 0)$ for $x_1 \in (-1, 1)$. Now consider $\tilde{m}_\epsilon = m_\epsilon \circ g^{-1} \in M(b, d)$. Using a change of variable, we compute

$$\int_{-1}^1 |\tilde{m}'_\epsilon|^2 dx_1 = \int_{-1}^1 \frac{|m'_\epsilon|^2}{|g'|} dx_1 \leq \|1/g'\|_{L^\infty(-1,1)} \int_{-1}^1 |m'_\epsilon|^2 dx_1.$$

In order to compare the stray field energies of m_ϵ and \tilde{m}_ϵ , we consider the harmonic extension $v_\epsilon \in \dot{H}^1(\mathbb{R}_+^2)$ of $m_{\epsilon 1} - \cos \alpha$ and set $\tilde{v}_\epsilon = v_\epsilon \circ G^{-1}$. Then

$$\begin{aligned} \int_{\mathbb{R}_+^2} |\nabla \tilde{v}_\epsilon|^2 dx &\leq \int_{\mathbb{R}_+^2} |\nabla v_\epsilon|^2 |(DG)^{-1}|^2 |\det DG| dx \\ &\leq \int_{\Omega_{2R}(a) \cup \bigcup_{n=1}^N B_R^+(a_n, 0)} |\nabla v_\epsilon|^2 dx + \| |(DG)^{-1}|^2 \det DG \|_{L^\infty(\mathbb{R}_+^2)} \int_{\Omega_R(a) \setminus \Omega_{2R}(a)} |\nabla v_\epsilon|^2 dx. \end{aligned}$$

Hence

$$\begin{aligned} \inf_{M(b,d)} E_\epsilon - \inf_{M(a,d)} E_\epsilon &\leq E_\epsilon(\tilde{m}_\epsilon) - E_\epsilon(m_\epsilon) \\ &\leq \frac{\epsilon}{2} (\|1/g'\|_{L^\infty(-1,1)} - 1) \int_{-1}^1 |m'_\epsilon|^2 dx_1 \\ &\quad + \left(\| |(DG)^{-1}|^2 \det DG \|_{L^\infty(\mathbb{R}_+^2)} - 1 \right) \int_{\Omega_R(a) \setminus \Omega_{2R}(a)} \frac{|\nabla v_\epsilon|^2}{2} dx. \end{aligned}$$

By Theorem 1, we know that m satisfies the assumptions of [18, Theorem 28] for a constant C_0 that is independent of ϵ . Hence, by Remark 30 and (62) in [18], together with Lemma 14, there exists another constant C , also independent of ϵ (but depending on R), such that

$$\epsilon \int_{-1}^1 |m'_\epsilon|^2 dx_1 + \int_{\Omega_R(a) \setminus \Omega_{2R}(a)} |\nabla v_\epsilon|^2 dx \leq \frac{C}{(\log \delta)^2}.$$

The quantities

$$\|1/g'\|_{L^\infty(-1,1)} - 1 \quad \text{and} \quad \| |(DG)^{-1}|^2 \det DG \|_{L^\infty(\mathbb{R}_+^2)} - 1$$

can be made arbitrarily small when r is small enough. Thus we obtain the inequality

$$\inf_{M(b,d)} E_\epsilon - \inf_{M(a,d)} E_\epsilon \leq \frac{c_0}{(\log \delta)^2}$$

for an arbitrarily small $c_0 > 0$.

The reverse inequality is proved with essentially the same arguments, but we exchange the roles of a_n and b_n . This has the consequence that instead of working with one minimiser for a given ϵ , we have a family depending on the position of b_n . We can check that all the relevant quantities in the resulting inequalities are uniformly bounded, and we obtain the desired estimate. \square

Proof of Theorem 4. The first statement of the theorem is an obvious consequence of Proposition 10. Thus only the second statement remains to be proved. Suppose, therefore, that we have $\epsilon_k \searrow 0$

and $\phi_0 \in \Phi$ such that $\phi_{\epsilon_k} \rightarrow \phi_0$ in $L^2(-1, 1)$. We may further assume that the convergence holds pointwise almost everywhere. Since $\phi_0 \in \Phi$, there exist $\tilde{N} \in \mathbb{N} \cup \{0\}$, $a \in A_{\tilde{N}}$, and $\omega \in ((2\pi\mathbb{Z} + \{0, \pm 2\alpha\}) \setminus \{0\})^{\tilde{N}}$ such that

$$\phi'_0 = \sum_{n=1}^{\tilde{N}} \omega_n \delta_{a_n}.$$

Now suppose that

$$E_{\epsilon_k}(\phi_{\epsilon_k}) \leq \frac{\eta(\phi_0)}{\log \frac{1}{\delta_k}} + \frac{C_0}{(\log \delta_k)^2} \quad (29)$$

for all $k \in \mathbb{N}$, where $\delta_k = \epsilon_k \log \frac{1}{\epsilon_k}$.

Fix $n \in \{1, \dots, \tilde{N}\}$ for the moment. Then there exists $r \in (0, \frac{\rho(a)}{2}]$ such that $\phi_{\epsilon_k}(a_n \pm r) \rightarrow \phi_0(a_n \pm r)$ as $k \rightarrow \infty$ and ϕ_0 is continuous at $a_n \pm r$. We know that $\phi_0(a_n + r) - \phi_0(a_n - r) = \omega_n$ and ϕ_{ϵ_k} is continuous. If $\omega_n > 0$, then we choose

$$L_n^- = \min \{ \ell \in \mathbb{Z} : \pi \ell > \phi_0(a_n - r) \}$$

and

$$L_n^+ = \max \{ \ell \in \mathbb{Z} : \pi \ell < \phi_0(a_n + r) \}.$$

For sufficiently large value of k , we can choose $b_{nL_n^-}^k, \dots, b_{nL_n^+}^k \in (a_n - r, a_n + r)$ such that $\phi_{\epsilon_k}(b_{n\ell}^k) = \pi \ell$ for all $\ell = L_n^-, \dots, L_n^+$ and such that $b_{nL_n^-}^k < \dots < b_{nL_n^+}^k$. If $\omega_n < 0$, we choose $b_{n\ell}^k$ similarly.

Let $N = \iota(\phi_0)$ and let $a^k \in A_N$ comprise all these points, i.e.,

$$a^k = (b_{1L_1^-}^k, \dots, b_{1L_1^+}^k, \dots, b_{\tilde{N}L_{\tilde{N}}^-}^k, \dots, b_{\tilde{N}L_{\tilde{N}}^+}^k).$$

Then $(\cos \phi_{\epsilon_k}, \sin \phi_{\epsilon_k}) \in M(a^k, d)$ for some $d \in \{\pm 1\}^N$, which will satisfy $d_{n-1} = -d_n$ or $|a_{n-1}^k - a_n^k| \geq \rho(a)$ for $n = 1, \dots, N$ whenever k is sufficiently large. If $\theta_N < \alpha < \pi - \theta_N$, then Proposition 16 applies. Thus we obtain a constant C_1 such that

$$E_{\epsilon_k}(\phi_{\epsilon_k}) \geq \frac{\pi}{2 \log \frac{1}{\delta_k}} \sum_{n=1}^N \gamma_n^2 + \frac{W(a^k, d) - C_1}{(\log \delta_k)^2} \quad (30)$$

for every sufficiently large k , where, as usual, $\gamma_n = d_n - \cos \alpha$. It is readily checked that

$$\eta(\phi_0) = \frac{\pi}{2} \sum_{n=1}^N \gamma_n^2$$

(indeed, the function η is defined with this identity in mind). Hence (29) and (30) imply that

$$\limsup_{k \rightarrow \infty} W(a^k, d) < \infty.$$

According to Proposition 13, this means that all the points of a^k remain separated from one another when $k \rightarrow \infty$. By construction, this is only possible when $N = \tilde{N}$ and $|\omega_n| = 2\alpha$ or $2\pi - 2\alpha$ for every $n = 1, \dots, \tilde{N}$. Hence ϕ_0 is simple. \square

For the Γ -convergence result of Corollary 5, we have matching lower and upper bounds of E_ϵ only in the case of a limiting magnetisation $m_0 = (\cos \phi_0, \sin \phi_0)$ for ϕ_0 simple, i.e., when all jumps come from individual domain walls of sign ± 1 . This is a common feature in Γ -convergence results for Ginzburg-Landau models where the vortex points carry winding numbers ± 1 .

Proof of Corollary 5. For the lower bound, we consider a sequence $\epsilon_k \searrow 0$ and then write $m_k = (\cos \phi_{\epsilon_k}, \sin \phi_{\epsilon_k})$. Extracting a subsequence if necessary, we may assume that $\phi_k \rightarrow \phi_0$ almost everywhere in $(-1, 1)$. We represent ϕ' as in (6). Then $\omega \in \{\pm 2\alpha, \pm 2(\pi - \alpha)\}^N$, since ϕ is simple. As in the proof of Theorem 4, we construct the points $a^k \in A_N$ such that $|a_{n-1}^k - a_n^k| \geq \rho(a)$ and $m_k \in M(a^k, d)$ for k sufficiently large. (Note that for this construction there is no need of the angle restriction $\theta_N < \alpha < \pi - \theta_N$, which is imposed in Theorem 4 for a different reason.) Since $\eta(\phi_0) = \frac{\pi}{2} \sum_{n=1}^N \gamma_n^2$ and $a^k \rightarrow a$ as $k \rightarrow \infty$, by Theorem 1 and Proposition 2, we deduce the desired lower bound:

$$\begin{aligned} E_{\epsilon_k}(\phi_{\epsilon_k}) &\geq \inf_{M(a^k, d)} E_{\epsilon_k} \geq \inf_{M(a, d)} E_{\epsilon_k} - o\left(\frac{1}{(\log \delta_k)^2}\right) \\ &= \frac{\eta(\phi_0)}{\log \frac{1}{\delta_k}} + \frac{\sum_{n=1}^N e(d_n) + W(a, d)}{(\log \delta_k)^2} - o\left(\frac{1}{(\log \delta_k)^2}\right). \end{aligned}$$

The upper bound follows from Theorem 1 as follows. Given a simple ϕ_0 with transition profile (a, d) , Theorem 1 gives a family $(m_\epsilon)_{\epsilon > 0}$ in $M(a, d)$ such that

$$E_\epsilon(m_\epsilon) \leq \frac{\eta(\phi_0)}{\log \frac{1}{\delta}} + \frac{\sum_{n=1}^N e(d_n) + W(a, d)}{(\log \frac{1}{\delta})^2} + o\left(\frac{1}{(\log \frac{1}{\delta})^2}\right).$$

Next we note that m_ϵ can always be modified, without changing the energy, such that between a_n and a_{n+1} (for $n = 1, \dots, N-1$), as well as between -1 and a_1 and between a_N and 1 , the sign of the second component $m_{\epsilon 2}$ is the same as the sign of $\sin \phi_0$. To this end, we merely replace m_ϵ by $(m_{\epsilon 1}, \pm |m_{\epsilon 2}|)$, with the sign chosen appropriately in each of these intervals. Thus we may assume that each m_ϵ has this property. Now let ϕ_ϵ denote the phase of m_ϵ (i.e., such that $m_\epsilon = (\cos \phi_\epsilon, \sin \phi_\epsilon)$) with $\phi_\epsilon(-1) = \phi_0(-1)$. Then automatically

$$\phi_\epsilon(a_n) = \frac{1}{2} \left(\lim_{x_1 \nearrow a_n} \phi_0(x_1) + \lim_{x_1 \searrow a_n} \phi_0(x_1) \right)$$

for $n = 1, \dots, N$, and $\phi_\epsilon(1) = \phi_0(1)$. Hence the only possible accumulation point for $(\phi_\epsilon)_{\epsilon > 0}$ in Φ is ϕ_0 . The compactness of Theorem 4 then yields $\phi_\epsilon \rightarrow \phi_0$ in $L^2(-1, 1)$ as $\epsilon \searrow 0$. \square

Proof of Corollary 7. For the first statement, assume that the functions $\phi_\epsilon \in H^1(-1, 1)$ satisfy $\phi_\epsilon(-1) = \alpha$ and $\phi_\epsilon(1) = 2\pi\ell + \alpha$ for every $\epsilon > 0$, where $N = 2\ell + 1$. It suffices to show that any subsequence $(\phi_{\epsilon_k})_{k \in \mathbb{N}}$ with $\epsilon_k \searrow 0$ contains another subsequence that satisfies the desired inequality. In order to keep the notation simple, we suppress the subsequences in the following. By continuity of ϕ_ϵ , we can choose $a^\epsilon \in A_N$ such that $m_\epsilon = (\cos \phi_\epsilon, \sin \phi_\epsilon) \in M(a^\epsilon, d_N^+)$. Since $\alpha \in (\theta_N, \pi - \theta_N)$, Proposition 16 implies that for some $C > 0$,

$$E_\epsilon(\phi_\epsilon) \geq \frac{\mathcal{E}_0}{\log \frac{1}{\delta}} + \frac{W(a^\epsilon, d_N^+) - C}{(\log \frac{1}{\delta})^2} \quad \text{for every } \epsilon.$$

If $\rho(a^\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$ (for some subsequence), then we use Proposition 13, which implies that $W(a^\epsilon, d_N^+) \rightarrow \infty$ as $\epsilon \rightarrow 0$. This immediately gives the desired inequality. Otherwise, the points of a^ϵ stay uniformly separated from one another and uniformly away from the boundary as $\epsilon \rightarrow 0$. By Proposition 2 and a compactness argument, we find $a \in A_N$ such that $a_\epsilon \rightarrow a$ (for a subsequence) and $E_\epsilon(\phi_\epsilon) \geq \inf_{M(a, d_N^+)} E_\epsilon - o(1/(\log \delta)^2)$ as $\epsilon \searrow 0$. Theorem 1 now gives the conclusion.

For the proof of statement 2, assume that

$$E_\epsilon(\phi_\epsilon) \leq \frac{\mathcal{E}_0}{\log \frac{1}{\delta}} + O\left(\frac{1}{(\log \delta)^2}\right). \quad (31)$$

By Theorem 4, there exist a sequence $\epsilon_k \searrow 0$ and a limit $\phi_0 \in \Phi$ such that $\phi_{\epsilon_k} \rightarrow \phi_0$ as $k \rightarrow \infty$ in $L^2(-1, 1)$. We may further assume that we have pointwise convergence almost everywhere. We claim that ϕ_0 is simple (i.e., has jumps of size 2α and $2(\pi - \alpha)$ only), as required for the statement. In order to show this, consider a jump of ϕ_0 , of any size, at a point $b \in (-1, 1)$. Then there exist $\psi_-, \psi_+ \in 2\pi\mathbb{Z} \pm \alpha$ and $r > 0$ such that $\phi_0 = \psi_-$ in $(b-r, b)$ and $\phi_0 = \psi_+$ in $(b, b+r)$. Furthermore, we may choose r such that $\phi_{\epsilon_k}(b-r) \rightarrow \psi_-$ and $\phi_{\epsilon_k}(b+r) \rightarrow \psi_+$ as $k \rightarrow \infty$. If, say, $\psi_+ - \psi_- = 2\pi j$ for some $j \in \mathbb{Z} \setminus \{0\}$, then by the continuity of ϕ_{ϵ_k} , the set

$$\{x_1 \in (b-r, b+r) : \phi_{\epsilon_k}(x_1) \in \pi\mathbb{Z}\}$$

has at least $2|j|$ points whenever k is sufficiently large. We may select $2|j|$ of them, say $\{t_1^{(k)}, \dots, t_{2|j|}^{(k)}\}$, such that $\cos \phi_{\epsilon_k}(t_i^{(k)}) = (-1)^i$ for $i = 1, \dots, 2|j|$ or $\cos \phi_{\epsilon_k}(t_i^{(k)}) = (-1)^{i+1}$ for $i = 1, \dots, 2|j|$. Similar statements hold if $\psi_+ - \psi_- = 2\pi j \pm 2\alpha$ (but now we have $2|j| \pm 1$ points). Near the boundary, the function ϕ_0 is constant, too. More precisely, there exists $r > 0$ such that $\phi_0 = \chi_-$ in $(-1, r-1)$ and $\phi_0 = \chi_+$ in $(1-r, 1)$ for two numbers $\chi_-, \chi_+ \in 2\pi\mathbb{Z} \pm \alpha$. If $\chi_- \neq -\alpha$, say $\chi_- = 2\pi j - \alpha$ for $j \in \mathbb{Z} \setminus \{0\}$, then the set

$$\{x_1 \in (-1, r-1) : \phi_{\epsilon_k}(x_1) \in \pi\mathbb{Z}\}$$

has at least $2|j|$ points for k large enough. (We may think of this situation as a jump at the boundary.) Again we may select $2|j|$ of them such that the sign of $\cos \phi_{\epsilon_k}$ oscillates between ± 1 . Similar statements hold if $\chi_- = 2\pi j + \alpha$ and for the other boundary point. The prescribed boundary conditions entail that the number of points of $\phi_{\epsilon_k}^{-1}(\pi\mathbb{Z})$ covered by the above discussion is at least N .

Suppose first that $\ell \geq 1$ (and thus $N \geq 3$). In this case, if ϕ_0 were *not* simple or did *not* match the given boundary data, then we could construct $a^{(k)} \in A_N$ (from points chosen among the above $t_i^{(k)}$ for all the jumps, including jumps at the boundary) and $\sigma > 0$ such that for $n = 1, \dots, N-1$, either $a_{n+1}^{(k)} - a_n^{(k)} \geq \sigma$ or $\cos \phi_{\epsilon_k}(a_{n+1}^{(k)}) = -\cos \phi_{\epsilon_k}(a_n^{(k)})$, and such that there are exactly $\ell + 1$ positive and ℓ negative signs, but $\rho(a^{(k)}) \rightarrow 0$ as $k \rightarrow \infty$. Proposition 16 and Proposition 13 would then give an estimate for the energy incompatible with the assumption (31). Therefore, ϕ_0 is simple and there is no jump at the boundary, which means that $\phi_0(-1) = -\alpha$, $\phi_0(1) = 2\pi\ell + \alpha$, and there must be at least N jumps (at least $\ell + 1$ of which are of the size 2α and at least ℓ of the size $2(\pi - \alpha)$). In particular, we conclude that $\eta(\phi_0) \geq \mathcal{E}_0$. But Corollary 5, together with (31), implies that $\mathcal{E}_0 \geq \eta(\phi_0)$, so we have equality. Therefore, ϕ_0 has exactly N jumps. It also follows that ϕ_0 is of the form as described in the statement.

If $\ell = 0$ and $N = 1$, then we take advantage of the fact that $\theta_2 = \theta_1 = 0$. In this case, if ϕ_0 did *not* match the given boundary data, we would be able to construct $a^{(k)} \in A_1$ with properties as above. If ϕ_0 were *not* simple, in order to achieve that $\rho(a^{(k)}) \rightarrow 0$, we could construct $a^{(k)} \in A_2$ instead. We would then find a contradiction with the same arguments.

Statement 3 is a standard consequence of the Γ -convergence result in Corollary 5, which means that minimisers of $|\log \delta|(|\log \delta|E_\epsilon - \mathcal{E}_0) - \mathcal{E}_1$ converge to minimisers of $W(\cdot, d_N^+)$. Indeed, if \tilde{a} is any point in A_N , consider the (unique) simple function $\tilde{\phi}_0 \in \Phi$ with the jump points \tilde{a} and the structure described in statement 2, and satisfying the boundary conditions $\tilde{\phi}_0(-1) = -\alpha$, $\tilde{\phi}_0(1) = 2\pi\ell + \alpha$. By Corollary 5, there exists a family $(\tilde{\phi}_\epsilon)_{\epsilon > 0}$ with $\tilde{\phi}_\epsilon(-1) = -\alpha$, $\tilde{\phi}_\epsilon(1) = 2\pi\ell + \alpha$, such that

$$E_\epsilon(\phi_\epsilon) \leq E_\epsilon(\tilde{\phi}_\epsilon) \leq \frac{\mathcal{E}_0}{\log \frac{1}{\delta}} + \frac{\mathcal{E}_1 + W(\tilde{a}, d_N^+)}{(\log \frac{1}{\delta})^2} + o\left(\frac{1}{(\log \frac{1}{\delta})^2}\right)$$

as $\epsilon \searrow 0$, since ϕ_ϵ are minimisers of E_ϵ for their boundary data. Then statement 2 applies to the given family of minimisers $(\phi_\epsilon)_{\epsilon > 0}$, so for a subsequence, we have the convergence $\phi_{\epsilon_k} \rightarrow \phi_0$ for a simple function ϕ_0 with jump points $a \in A_N$ as in statement 2. Then the lower bound in Corollary 5 combined with the above upper bound for $E_\epsilon(\phi_\epsilon)$, yield $W(a, d_N^+) \leq W(\tilde{a}, d_N^+)$ in the limit $\epsilon \searrow 0$. That is, a is a minimizer of $W(\cdot, d_N^+)$ over A_N . \square

3 Analysis for the unconfined problem

This section analyses the problem described in Subsection 1.3. We first derive some properties of the function I appearing in the renormalised energy, then we derive and analyse the limiting stray field potential for this problem, and finally we explain how the arguments for the confined case [18] need to be adapted for the proof of Theorem 8. We also prove a Γ -convergence result in Theorem 31 below, which adapts the statements of Theorem 4 and Corollary 5 to the unconfined problem.

3.1 The function I

Here we prove a few statements about the function

$$I(t) = \int_0^\infty \frac{se^{-s}}{s^2 + t^2} ds, \quad t > 0,$$

defined in the introduction. First we have an alternative representation.

Lemma 18. *For any $t > 0$,*

$$I(t) = \int_0^\infty \frac{\cos s}{s + t} ds.$$

Remark 19. *The integral in Lemma 18 does not converge in the L^1 -sense, but the Leibniz criterion for alternating series implies that it converges as an improper Riemann integral. We always use this interpretation for oscillating integrals of this type.*

Proof. Let $t, R > 0$. Note that the functions $z \mapsto \frac{e^{\pm iz}}{z+t}$ are holomorphic in $\{z \in \mathbb{C} : \operatorname{Re} z > -t\}$. Hence using contour integrals along the boundaries of the quarter disks

$$\{z \in \mathbb{C} : \operatorname{Re} z > 0, \pm \operatorname{Im} z > 0, |z| < R\},$$

we find that

$$0 = \int_0^R \frac{e^{is}}{s+t} ds - i \int_0^R \frac{e^{-s}}{is+t} ds + iR \int_0^{\pi/2} e^{i\phi} \frac{\exp(iRe^{i\phi})}{Re^{i\phi} + t} d\phi$$

and

$$0 = \int_0^R \frac{e^{-is}}{s+t} ds + i \int_0^R \frac{e^{-s}}{-is+t} ds - iR \int_0^{\pi/2} e^{-i\phi} \frac{\exp(-iRe^{-i\phi})}{Re^{-i\phi} + t} d\phi.$$

Since

$$|\exp(iRe^{i\phi})| = e^{-R \sin \phi} \quad \text{and} \quad |Re^{i\phi} + t| \geq R$$

for $\phi \in [0, \pi/2]$, and since there exists a constant $c > 0$ such that $c\phi \leq \sin \phi$ for every $\phi \in [0, \frac{\pi}{2}]$, we deduce that

$$\left| iR \int_0^{\pi/2} e^{i\phi} \frac{\exp(iRe^{i\phi})}{Re^{i\phi} + t} d\phi \right| \leq \int_0^{\pi/2} e^{-cR\phi} d\phi \leq \frac{1}{cR}.$$

Similarly,

$$\left| iR \int_0^{\pi/2} e^{-i\phi} \frac{\exp(-iRe^{-i\phi})}{Re^{-i\phi} + t} d\phi \right| \leq \frac{1}{cR}.$$

Hence, by adding the above equalities, we obtain

$$\int_0^\infty \frac{\cos s}{s+t} ds = \frac{1}{2} \lim_{R \rightarrow \infty} \int_0^R \frac{e^{is} + e^{-is}}{s+t} ds = \frac{i}{2} \lim_{R \rightarrow \infty} \int_0^R e^{-s} \left(\frac{1}{t+is} - \frac{1}{t-is} \right) ds = \int_0^\infty \frac{se^{-s}}{s^2+t^2} ds.$$

The last integral is of course identical to $I(t)$. □

The next lemma shows that $I(t)$ is logarithmic to leading order when $t \searrow 0$ and decays quadratically as $t \rightarrow \infty$. We will see later that I can also be used to describe the tail profile of a Néel wall (see (42) below). In this respect, our decay estimate is consistent with previous estimates for the decay of a tail of a Néel wall in [5] and [19, Theorem 5.2].

Lemma 20. *The function I is positive, decreasing, and convex with $I(t) \leq 1/t^2$ for all $t > 0$. If I_0 is the number defined in (10), then*

$$0 \leq I(t) + \log t - I_0 \leq \frac{\pi t}{2}$$

for all $t > 0$ as well. In particular, $I(t) = \log \frac{1}{t} + I_0 + o(1)$ as $t \searrow 0$.

Proof. By a change of variable, we write $I(t) = \int_0^\infty \frac{se^{-st}}{s^2+1} ds > 0$ for $t > 0$, and then we compute

$$I'(t) = - \int_0^\infty \frac{s^2 e^{-st}}{s^2+1} ds < 0 \quad \text{and} \quad I''(t) = \int_0^\infty \frac{s^3 e^{-st}}{s^2+1} ds > 0. \quad (32)$$

Hence I is decreasing and convex. Moreover, using $I(t) = \int_0^\infty \frac{se^{-s}}{s^2+t^2} ds$, we see that

$$I(t) \leq \frac{1}{t^2} \int_0^\infty se^{-s} ds = \frac{1}{t^2}, \quad t > 0.$$

Furthermore, integration by parts gives

$$I(t) = -\log t + \int_0^\infty e^{-s} \log \sqrt{s^2 + t^2} ds.$$

Note that

$$\frac{d}{dt} \int_0^\infty e^{-s} \log \sqrt{s^2 + t^2} ds = \int_0^\infty \frac{te^{-s}}{s^2 + t^2} ds = \int_0^\infty \frac{e^{-st}}{s^2 + 1} ds$$

and

$$0 \leq \int_0^\infty \frac{e^{-st}}{s^2 + 1} ds \leq \int_0^\infty \frac{ds}{s^2 + 1} = \frac{\pi}{2}$$

for every $t > 0$. Thus,

$$I_0 = \int_0^\infty e^{-s} \log \sqrt{s^2 + t^2} ds \Big|_{t=0} \leq \int_0^\infty e^{-s} \log \sqrt{s^2 + t^2} ds \leq I_0 + \frac{\pi t}{2}, \quad t > 0,$$

which leads to the desired conclusion. \square

3.2 The limiting stray field potential: construction

In this section we will redefine and compute the function $u_{a,d}^* : \mathbb{R}_+^2 \rightarrow \mathbb{R}$, the limiting stray field potential for Néel walls of sign d_1, \dots, d_N at the points a_1, \dots, a_N , for the problem with anisotropy term. Simultaneously, we will obtain a limiting profile $\mu_{a,d}^* : \mathbb{R} \rightarrow \mathbb{R}$ for the tails of the Néel walls.

The two functions are determined through the boundary value problem¹¹

$$\begin{aligned}\Delta u_{a,d}^* &= 0 && \text{in } \mathbb{R}_+^2, \\ \frac{\partial u_{a,d}^*}{\partial x_1} &= \mu_{a,d}^* - \pi \sum_{n=1}^N \gamma_n \delta_{a_n} && \text{on } \mathbb{R} \times \{0\}, \\ \frac{\partial u_{a,d}^*}{\partial x_2} &= -(\mu_{a,d}^*)' && \text{on } \mathbb{R} \times \{0\}.\end{aligned}$$

In order to see that this is a well-posed boundary value problem (and also in order to compute the solution later), we consider the harmonic conjugate of $u_{a,d}^*$, i.e., the function $v_{a,d}^*: \mathbb{R}_+^2 \rightarrow \mathbb{R}$, unique up to a constant, that satisfies $\nabla v_{a,d}^* = \nabla^\perp u_{a,d}^*$. The second boundary condition for $u_{a,d}^*$ implies that $v_{a,d}^*(\cdot, 0) - \mu_{a,d}^*$ is constant. Thus we may assume without loss of generality that $v_{a,d}^*(\cdot, 0) = \mu_{a,d}^*$ in \mathbb{R} ; this will then also determine $v_{a,d}^*$ completely. We finally obtain the following boundary value problem for the conjugate harmonic function:

$$\begin{aligned}\Delta v_{a,d}^* &= 0 && \text{in } \mathbb{R}_+^2, \\ \frac{\partial v_{a,d}^*}{\partial x_2} &= v_{a,d}^* - \pi \sum_{n=1}^N \gamma_n \delta_{a_n} && \text{on } \mathbb{R} \times \{0\}.\end{aligned}$$

That is, we have a harmonic function satisfying a Robin type boundary condition here. We now give some arguments depending in part on formal calculations, but they will be justified later.

As in our previous paper [18], we can solve this problem by superimposing the solutions of simpler problems. We therefore consider the following:

$$\Delta v = 0 \quad \text{in } \mathbb{R}_+^2, \quad (33)$$

$$\frac{\partial v}{\partial x_2} = v - \pi \delta_0 \quad \text{on } \mathbb{R} \times \{0\}. \quad (34)$$

Let \hat{v} denote the Fourier transform of v with respect to x_1 , i.e.,

$$\hat{v}(\xi, x_2) = \int_{-\infty}^{\infty} e^{-i\xi x_1} v(x_1, x_2) dx_1, \quad \xi \in \mathbb{R}, \quad x_2 \geq 0.$$

Then we find that

$$\begin{aligned}\frac{\partial^2 \hat{v}}{\partial x_2^2} - \xi^2 \hat{v} &= 0 && \text{for } x_2 > 0, \\ \frac{\partial \hat{v}}{\partial x_2} &= \hat{v} - \pi && \text{for } x_2 = 0.\end{aligned}$$

We want a function v with finite Dirichlet energy at $|x| \rightarrow \infty$; thus we rule out solutions with exponential growth as $x_2 \rightarrow \infty$ and find that

$$\hat{v}(\xi, x_2) = \frac{\pi e^{-|\xi| x_2}}{1 + |\xi|}. \quad (35)$$

¹¹Note that the limiting stray-field potential u corresponding to a 180° Néel wall in the confined case (defined on page 22) satisfies $\frac{\partial u}{\partial x_1} = -\pi \delta_0$ in $(-1, 1) \times \{0\}$. In the unconfined case, an additional term $\mu_{a,d}^*$ appears in the equation for $\frac{\partial u_{a,d}^*}{\partial x_1}$, due to the presence of anisotropy. In fact, this term makes the limiting stray-field energy finite in $\mathbb{R}_+^2 \setminus B_1(0)$, see Proposition 21.

Before justifying these formal calculations, we derive some properties of the function v implicitly defined here. From here on, the arguments are fully rigorous again. As $\hat{v}(\cdot, 0) \in L^2(\mathbb{R})$, we deduce that $v(\cdot, 0) \in L^2(\mathbb{R})$ and

$$\int_{\mathbb{R}} v^2 dx_1 = \frac{1}{2\pi} \int_{\mathbb{R}} \hat{v}^2 d\xi = \frac{\pi}{2} \int_{\mathbb{R}} \frac{d\xi}{(1+|\xi|)^2} = \pi \int_0^\infty \frac{d\xi}{(1+\xi)^2} = \pi. \quad (36)$$

Applying the inverse Fourier transform for every fixed $x_2 > 0$, we obtain

$$v(x_1, x_2) = \frac{1}{2} \int_{-\infty}^\infty \frac{e^{i\xi x_1 - |\xi| x_2}}{1+|\xi|} d\xi.$$

In order to find a more convenient representation, we use a contour integral in \mathbb{C} . Consider the contour consisting of the intervals $[0, R]$ and $[0, iR]$ (the second with reverse orientation) and the quarter circle Γ_R parametrised by $g(t) = Re^{it}$ for $0 \leq t \leq \frac{\pi}{2}$. Suppose first that $x_1 > 0$ and fix $x_2 > 0$ as well. The function $z \mapsto e^{izx_1 - zx_2}/(1+z)$ is holomorphic away from $z = -1$. Hence

$$\int_0^R \frac{e^{i\xi x_1 - |\xi| x_2}}{1+|\xi|} d\xi = \int_0^R \frac{e^{izx_1 - zx_2}}{1+z} dz = \int_0^{iR} \frac{e^{izx_1 - zx_2}}{1+z} dz - \int_{\Gamma_R} \frac{e^{izx_1 - zx_2}}{1+z} dz.$$

We observe that

$$\begin{aligned} \left| \int_{\Gamma_R} \frac{e^{izx_1 - zx_2}}{1+z} dz \right| &= R \left| \int_0^{\pi/2} \frac{\exp(R(ix_1 - x_2)(\cos t + i \sin t))}{1 + Re^{it}} e^{it} dt \right| \\ &\leq R \int_0^{\pi/2} \exp(R(-x_2 \cos t - x_1 \sin t)) dt. \end{aligned}$$

Since $x_2 \cos t + x_1 \sin t \geq \min\{x_1, x_2\} > 0$ for $t \in [0, \frac{\pi}{2}]$, we conclude that

$$\lim_{R \rightarrow \infty} \int_{\Gamma_R} \frac{e^{izx_1 - zx_2}}{1+z} dz = 0$$

and

$$\int_0^\infty \frac{e^{i\xi x_1 - |\xi| x_2}}{1+|\xi|} d\xi = i \int_0^\infty \frac{e^{-tx_1 - itx_2}}{1+it} dt.$$

Similarly, but integrating over the boundary of

$$\{z \in \mathbb{C} : \operatorname{Re} z > 0, \operatorname{Im} z < 0, |z| < R\},$$

we see that

$$\int_{-\infty}^0 \frac{e^{i\xi x_1 - |\xi| x_2}}{1+|\xi|} d\xi = -i \int_0^\infty \frac{e^{-tx_1 + itx_2}}{1-it} dt$$

and thus

$$v(x_1, x_2) = \frac{i}{2} \int_0^\infty e^{-tx_1} \left(\frac{e^{-itx_2}}{1+it} - \frac{e^{itx_2}}{1-it} \right) dt.$$

We further compute

$$\frac{e^{-itx_2}}{1+it} - \frac{e^{itx_2}}{1-it} = \frac{e^{-itx_2} - e^{itx_2}}{1+t^2} - it \frac{e^{-itx_2} + e^{itx_2}}{1+t^2} = -2i \frac{\sin(tx_2) + t \cos(tx_2)}{1+t^2}.$$

Hence

$$v(x_1, x_2) = \int_0^\infty e^{-tx_1} \frac{\sin(tx_2) + t \cos(tx_2)}{1+t^2} dt$$

for $x_1 > 0$. Similar computations for $x_1 < 0$ show that

$$v(x_1, x_2) = \int_0^\infty e^{-t|x_1|} \frac{\sin(tx_2) + t \cos(tx_2)}{1+t^2} dt \quad (37)$$

for all $x_1 \neq 0$ and $x_2 > 0$. (This is consistent with the expectation that v is an even function in x_1 , which comes from the symmetry of the boundary value problem (33), (34).) Furthermore, by the dominated convergence theorem, there is a continuous extension to $x_2 = 0$ (as long as $x_1 \neq 0$), given by the obvious integral. We then check that

$$\frac{\partial v}{\partial x_2}(x_1, 0) = v(x_1, 0) \quad \text{for every } x_1 \neq 0. \quad (38)$$

We now want to find the corresponding conjugate harmonic function $u: \mathbb{R}_+^2 \rightarrow \mathbb{R}$. The condition $\nabla v = \nabla^\perp u$ suggests that¹² there exist two constants $c_1, c_2 \in \mathbb{R}$ such that

$$u(x_1, x_2) = - \int_0^\infty e^{-tx_1} \frac{\cos(tx_2) - t \sin(tx_2)}{1+t^2} dt + c_1$$

for $x_1 > 0$ and

$$u(x_1, x_2) = \int_0^\infty e^{tx_1} \frac{\cos(tx_2) - t \sin(tx_2)}{1+t^2} dt + c_2$$

for $x_1 < 0$. We expect, however, that u has a limit as $|x| \rightarrow \infty$, and we may set this limit 0. Letting $x_1 \rightarrow \infty$, we see that this would imply that $c_1 = c_2 = 0$. Thus

$$u(x_1, x_2) = - \frac{x_1}{|x_1|} \int_0^\infty e^{-t|x_1|} \frac{\cos(tx_2) - t \sin(tx_2)}{1+t^2} dt. \quad (39)$$

In particular, u is odd in x_1 and $u(0, x_2) = 0$ for every $x_2 > 0$. Here again, we have a continuous extension to $x_2 = 0$ when $x_1 \neq 0$.

We now justify these formal computations. Moreover, we prove in Propositions 21 and 23 below that $u(x) = -\arctan \frac{x_1}{x_2} + o(1)$ and $v(x) = \log \frac{1}{|x|} + I_0 + o(1)$ as $x \rightarrow 0$ and $u, v \rightarrow 0$ as $|x| \rightarrow \infty$.

Proposition 21. *Formulas (37) and (39) define a pair of conjugate harmonic functions $u, v: \mathbb{R}_+^2 \rightarrow \mathbb{R}$. There exists a universal constant C with*

$$\pi \log \frac{1}{r} - C \leq \int_{\mathbb{R}_+^2 \setminus B_r(0)} |\nabla v|^2 dx \leq \pi \log \frac{1}{r} + C$$

for all $r \in (0, 1]$. Moreover, $|u| \leq \frac{\pi}{2}$ in \mathbb{R}_+^2 and $u, v \rightarrow 0$ as $|x| \rightarrow \infty$.

Proof. Step 1: harmonicity and limit of v . We have seen that for every fixed $x_2 > 0$, the Fourier transform of v with respect to x_1 is given by (35). It is thus clear that the function $x_1 \mapsto v(x_1, x_2)$ belongs to $H^k(\mathbb{R})$ for every $k \in \mathbb{N}$ for all $x_2 > 0$. Moreover, we see that $v \in L_{\text{loc}}^2((0, \infty), H^k(\mathbb{R}))$. Integrating against a test function and using Plancherel's theorem, we further see that the distributional derivative $\frac{\partial^2 v}{\partial x_2^2}$ satisfies

$$\widehat{\frac{\partial^2 v}{\partial x_2^2}} = \frac{\pi \xi^2 e^{-|\xi|x_2}}{1+|\xi|} = \xi^2 \widehat{v} = -\widehat{\frac{\partial^2 v}{\partial x_1^2}}.$$

Hence v is harmonic in \mathbb{R}_+^2 .

¹²Thus u is harmonic in \mathbb{R}_+^2 and $\frac{\partial u}{\partial x_1} = v - \pi \delta_0$ in $\mathbb{R} \times \{0\}$, so that the Fourier transform of u in x_1 is given by $\widehat{u}(\xi, x_2) = \frac{i\pi\xi}{|\xi|(1+|\xi|)} e^{-|\xi|x_2}$ for $\xi \neq 0$ and $x_2 > 0$.

Furthermore, we see that the norms $\|v(\cdot, x_2)\|_{H^k(\mathbb{R})}$ tend to 0 as $x_2 \rightarrow \infty$. In particular, by the embedding $H^1(\mathbb{R}) \subset L^\infty(\mathbb{R})$, we deduce that $v(x_1, x_2) \rightarrow 0$ as $x_2 \rightarrow \infty$ uniformly in x_1 . With the formula (37), it is easy to see that $v(x_1, x_2) \rightarrow 0$ as $x_1 \rightarrow \pm\infty$ uniformly in x_2 . Hence $\lim_{|x| \rightarrow \infty} v(x) = 0$.

Step 2: estimate of $\int_{\mathbb{R}_+^2 \setminus B_r(0)} |\nabla v|^2 dx$. We consider the Dirichlet energy of v first in $\mathbb{R} \times (s, +\infty)$ and then separately in the infinite strips $(s, +\infty) \times (0, 2s)$ for $s \in (0, 1]$.

Fix $s \in (0, 1]$. Then in $\mathbb{R} \times (s, +\infty)$, we compute

$$\begin{aligned} \int_s^\infty \int_{-\infty}^\infty \left(\xi^2 (\hat{v}(\xi, x_2))^2 + \left(\frac{\partial \hat{v}}{\partial x_2}(\xi, x_2) \right)^2 \right) d\xi dx_2 &= 2\pi^2 \int_s^\infty \int_{-\infty}^\infty \frac{\xi^2 e^{-2|\xi|x_2}}{(1+|\xi|)^2} d\xi dx_2 \\ &= 2\pi^2 \int_{-\infty}^\infty \frac{\xi^2}{(1+|\xi|)^2} \int_s^\infty e^{-2|\xi|x_2} dx_2 d\xi = \pi^2 \int_{-\infty}^\infty \frac{|\xi| e^{-2|\xi|s}}{(1+|\xi|)^2} d\xi \\ &= 2\pi^2 \int_0^\infty \frac{\xi e^{-2\xi s}}{(1+\xi)^2} d\xi = 2\pi^2 \int_0^\infty \frac{te^{-2t}}{(s+t)^2} dt. \end{aligned}$$

We note that

$$\frac{t}{(s+t)^2} = \frac{1}{s+t} - \frac{s}{(s+t)^2} = \frac{d}{dt} \left(\log(s+t) + \frac{s}{s+t} \right). \quad (40)$$

Hence an integration by parts gives

$$\int_0^\infty \frac{te^{-2t}}{(s+t)^2} dt = \log \frac{1}{s} - 1 + 2 \int_0^\infty e^{-2t} \left(\log(s+t) + \frac{s}{s+t} \right) dt.$$

Finally, the concavity of the logarithm yields the inequality $\log(s+t) \leq \log t + \frac{s}{t}$, and thus,

$$\begin{aligned} \int_0^\infty e^{-2t} \log(s+t) dt &\leq \int_0^1 e^{-2t} \log 2 dt + \int_1^\infty e^{-2t} \left(\log t + \frac{s}{t} \right) dt \\ &\leq \log 2 + \int_1^\infty e^{-2t} \log t dt + s \int_1^\infty \frac{e^{-2t}}{t} dt \end{aligned}$$

for $s \leq 1$. Noting that the right hand side is bounded by a constant, we conclude that there exists a number C_1 satisfying

$$\pi \log \frac{1}{s} - C_1 \leq \int_s^\infty \int_{-\infty}^\infty |\nabla v|^2 dx_1 dx_2 \leq \pi \log \frac{1}{s} + C_1 \quad \text{for all } s \leq 1.$$

Next we consider the strip $(s, +\infty) \times (0, 2s)$. For $x_1 > 0$, we find that

$$\begin{aligned} \left| \frac{\partial v}{\partial x_1}(x_1, x_2) \right| &= \left| \int_0^\infty te^{-tx_1} \frac{\sin(tx_2) + t \cos(tx_2)}{1+t^2} dt \right| \\ &\leq \int_0^\infty \frac{t+t^2}{1+t^2} e^{-tx_1} dt = \frac{1}{x_1} \int_0^\infty \frac{sx_1 + s^2}{x_1^2 + s^2} e^{-s} ds. \end{aligned}$$

Young's inequality gives

$$sx_1 + s^2 \leq \frac{1}{2(\sqrt{2}-1)} x_1^2 + \left(1 + \frac{\sqrt{2}-1}{2} \right) s^2 = \frac{\sqrt{2}+1}{2} (x_1^2 + s^2).$$

Hence

$$\left| \frac{\partial v}{\partial x_1}(x_1, x_2) \right| \leq \frac{\sqrt{2}+1}{2x_1}.$$

Similarly, we find that

$$\left| \frac{\partial v}{\partial x_2}(x_1, x_2) \right| \leq \frac{\sqrt{2} + 1}{2x_1}$$

when $x_1 > 0$. We therefore compute

$$\int_0^{2s} \int_s^\infty |\nabla v(x_1, x_2)|^2 dx_1 dx_2 \leq \left(\frac{3}{2} + \sqrt{2} \right) \int_0^{2s} \int_s^\infty \frac{dx_1}{x_1^2} dx_2 = 3 + 2\sqrt{2}.$$

Similar estimates hold for $x_1 < 0$.

Finally, we can conclude Step 2 and estimate the energy outside the disk $B_r(0)$ as follows. Given $r \in (0, 1]$, we now apply the above inequalities for $s = r$ and for $s = r/2$ and we conclude that

$$\pi \log \frac{1}{r} - C_1 \leq \int_{\mathbb{R}_+^2 \setminus B_r(0)} |\nabla v|^2 dx \leq \pi \log \frac{2}{r} + C_1 + 6 + 4\sqrt{2}. \quad (41)$$

This proves the inequalities for the Dirichlet energy.

Step 3: u and v are conjugate harmonic functions. Since $\text{curl } \nabla^\perp v = 0$, there exists a function $\tilde{u}: \mathbb{R}_+^2 \rightarrow \mathbb{R}$ with $\nabla v = \nabla^\perp \tilde{u}$. This function satisfies

$$\begin{aligned} \Delta \tilde{u} &= 0 \quad \text{in } \mathbb{R}_+^2, \\ \frac{\partial \tilde{u}}{\partial x_1} &= v \quad \text{on } (-\infty, 0) \text{ and } (0, \infty), \end{aligned}$$

owing to (38). Moreover, we know that

$$\int_{\mathbb{R}_+^2 \setminus B_r(0)} |\nabla \tilde{u}|^2 dx < \infty$$

for any $r > 0$. Now define $w(x) = \tilde{u}(x/|x|^2)$. Then w is again harmonic in \mathbb{R}_+^2 and belongs to $H^1(B_R^+(0))$ for any $R > 0$. We compute, for $x_1 \neq 0$:

$$\frac{\partial w}{\partial x_1}(x_1, 0) = -\frac{1}{x_1^2} \frac{\partial \tilde{u}}{\partial x_1} \left(\frac{1}{x_1}, 0 \right) = -\frac{1}{x_1^2} v \left(\frac{1}{x_1}, 0 \right).$$

Note that by Lemma 20,

$$v(x_1, 0) = \int_0^\infty \frac{te^{-t|x_1|}}{1+t^2} dt = \int_0^\infty \frac{se^{-s}}{x_1^2 + s^2} ds = I(|x_1|) \leq \frac{1}{x_1^2}. \quad (42)$$

Thus $\frac{\partial w}{\partial x_1}(x_1, 0)$ is bounded near 0. The fact that $w \in H^1(B_1^+(0))$ prevents jumps of $w(\cdot, 0)$. Hence, $w(\cdot, 0)$ is Lipschitz continuous near the origin. It follows that w (as a harmonic function in \mathbb{R}_+^2) is continuous at $x = 0$ (see [11, Lemma 2.13]), and hence $c := \lim_{|x| \rightarrow \infty} \tilde{u}(x)$ exists. It is easy to check that $\nabla u = -\nabla^\perp v = \nabla \tilde{u}$ pointwise in $(0, \infty)^2$ as well as in $(-\infty, 0) \times (0, \infty)$. Since $\lim_{x_1 \rightarrow \pm\infty} u(x_1, 0) = 0$ (by (39)), we conclude that $\tilde{u} - c$ must coincide with the function u determined in formula (39).

Step 4: Other properties of u . An immediate consequence of the above is that $u \rightarrow 0$ as $|x| \rightarrow \infty$. Also, note that (39) implies that

$$|u(x_1, 0)| \leq \int_0^\infty \frac{1}{1+t^2} dt = \frac{\pi}{2}, \quad \text{for every } x_1 \neq 0.$$

Since $u \rightarrow 0$ at infinity and u is harmonic in \mathbb{R}_+^2 , the maximum principle yields $|u| \leq \frac{\pi}{2}$ in \mathbb{R}_+^2 . \square

Having the functions u and v given by (39) and (37), we also define, for a given $b \in \mathbb{R}$, the translated functions

$$u_b(x) = u(x_1 - b, x_2) \quad \text{and} \quad v_b(x) = v(x_1 - b, x_2). \quad (43)$$

Recall that for the unconfined problem, we define

$$A_N = \{a = (a_1, \dots, a_N) \in \mathbb{R}^N : a_1 < \dots < a_N\}.$$

Then, given $a \in A_N$ and $d \in \{\pm 1\}^N$, we define

$$u_{a,d}^* = \sum_{n=1}^N \gamma_n u_{a_n} \quad \text{and} \quad v_{a,d}^* = \sum_{n=1}^N \gamma_n v_{a_n}.$$

Here $\gamma_n = d_n - \cos \alpha$, as defined in the introduction. We also write

$$\mu_{a,d}^*(x_1) = v_{a,d}^*(x_1, 0).$$

3.3 The limiting stray field potential: further properties

The above functions $u_{a,d}^*$ and $v_{a,d}^*$ have infinite Dirichlet energy due to the singularities at $(a_n, 0)$ for $n = 1, \dots, N$. It follows from the inequality of Proposition 21, however, that $u_{a,d}^*, v_{a,d}^* \in W_{\text{loc}}^{1,p}(\mathbb{R}_+^2)$ for any $p \in [1, 2)$ (by the dyadic decomposition argument of Struwe [37]). Indeed, given $p \in [1, 2)$, $R > 0$, $N \in \mathbb{N}$, and a compact set $K \subset \mathbb{R}^2$, there exists a constant C such that

$$\int_{\mathbb{R}_+^2 \cap K} |\nabla u_{a,d}^*|^p dx \leq C$$

for all $a \in A_N$ with $a_{n+1} - a_n \geq R$ for $n = 1, \dots, N-1$ and for all $d \in \{\pm 1\}^N$.

In contrast to (19), we now define

$$\rho(a) = \frac{1}{2} \min\{a_2 - a_1, \dots, a_N - a_{N-1}\} \quad (44)$$

for $a \in A_N$. Moreover, we define

$$\Omega_r(a) = \mathbb{R}_+^2 \setminus \bigcup_{n=1}^N B_r(a_n, 0)$$

again for $r > 0$ and consider

$$W_1(a, d) = \frac{1}{2} \lim_{r \searrow 0} \left(\int_{\Omega_r(a)} |\nabla u_{a,d}^*|^2 dx + \int_{-\infty}^{\infty} (\mu_{a,d}^*)^2 dx_1 - \pi \log \frac{1}{r} \sum_{n=1}^N \gamma_n^2 \right). \quad (45)$$

We will prove that this limit indeed exists and is finite as well. This is an important quantity that will appear in the renormalised energy. It is the counterpart of the contribution of the tail-tail interaction in the confined case, cf. (23). The difference consists in the additional term involving $\mu_{a,d}^*$, which comes from the anisotropy. We will prove the following.

Proposition 22. *If I_0 is defined as in (10), then*

$$W_1(a, d) = \frac{\pi}{2} I_0 \sum_{n=1}^N \gamma_n^2 + \frac{\pi}{2} \sum_{n=1}^N \sum_{k \neq n} \gamma_k \gamma_n I(|a_k - a_n|).$$

As we prove in Lemma 24 below, the first term above (independent of a) represents the (intrinsic) renormalised energy induced by the limiting stray field and anisotropy energy of every Néel wall. We observe the same phenomenon as in the confined case: $W_1(a, d) = -W(a, d)$, meaning that the contribution of the interactions between the tails of one Néel walls and the core of another is twice the size in absolute value but of opposite sign.

We decompose the proof of this statement into several lemmas. As before, we consider the functions v and u given by the formulas (37) and (39). We first need some pointwise estimates. The next proposition states that like in the confined case, u behaves analogously to the phase of a vortex of degree $+1$ close to the origin. But in contrast to the confined case, the error is logarithmic here.

Proposition 23. *Let*

$$w_0(x) = -\arctan \frac{x_1}{x_2}$$

for $x \in \mathbb{R}_+^2$. Then

$$\sup_{x \in \mathbb{R}_+^2} \frac{|\nabla u(x) - \nabla w_0(x)|}{\log\left(\frac{1}{|x|} + 2\right)} < \infty. \quad (46)$$

As a consequence,

$$\sup_{x \in \mathbb{R}_+^2} \frac{|u(x) - w_0(x)|}{|x| \log\left(\frac{1}{|x|} + 2\right)} < \infty \quad \text{and} \quad \sup_{x \in \mathbb{R}_+^2} \frac{\left|v(x) - \log\left(\frac{1}{|x|}\right) - I_0\right|}{|x| \log\left(\frac{1}{|x|} + 2\right)} < \infty.$$

Therefore, for any $a \in A_N$, there exists $C > 0$ such that

$$|\nabla u_{a,d}^*(x) - \gamma_n \nabla w_0(x_1 - a_n, x_2)| \leq C \log\left(\frac{1}{(x_1 - a_n)^2 + x_2^2} + 2\right) \quad \text{in } B_{\rho(a)}^+(a_n, 0)$$

for $n = 1, \dots, N$. Moreover,¹³

$$\int_{\Omega_\delta(a)} |\nabla u_{a,d}^*|^2 dx = \pi |\log \delta| \sum_{n=1}^N \gamma_n^2 + O(1) \quad \text{as } \delta \searrow 0. \quad (47)$$

Proof. We divide the proof in several steps.

Step 1: proof of (46). We compute, for $x \neq 0$:¹⁴

$$\begin{aligned} \frac{\partial u}{\partial x_1}(x) &= \int_0^\infty e^{-t|x_1|} \frac{t \cos(tx_2) - t^2 \sin(tx_2)}{1+t^2} dt \\ &= \int_0^\infty e^{-t|x_1|} \left(\frac{t \cos(tx_2) + \sin(tx_2)}{1+t^2} - \sin(tx_2) \right) dt \\ &= \int_0^\infty e^{-t|x_1|} \frac{t \cos(tx_2) + \sin(tx_2)}{1+t^2} dt - \frac{x_2}{x_1^2 + x_2^2} \end{aligned}$$

and

$$\begin{aligned} \frac{\partial u}{\partial x_2}(x) &= \frac{x_1}{|x_1|} \int_0^\infty e^{-t|x_1|} \frac{t \sin(tx_2) + t^2 \cos(tx_2)}{1+t^2} dt \\ &= \frac{x_1}{|x_1|} \int_0^\infty e^{-t|x_1|} \left(\frac{t \sin(tx_2) - \cos(tx_2)}{1+t^2} + \cos(tx_2) \right) dt \\ &= \frac{x_1}{|x_1|} \int_0^\infty e^{-t|x_1|} \frac{t \sin(tx_2) - \cos(tx_2)}{1+t^2} dt + \frac{x_1}{x_1^2 + x_2^2}. \end{aligned}$$

¹³The term $O(1)$ depends on a .

¹⁴Recall that for $z = x_1 + ix_2 \in \mathbb{C}$ with $x_1 > 0$, we have $\int_0^\infty e^{-tz} dt = \frac{1}{z}$.

Hence

$$\frac{\partial u}{\partial x_1}(x) - \frac{\partial w_0}{\partial x_1}(x) = \int_0^\infty e^{-t|x_1|} \frac{t \cos(tx_2) + \sin(tx_2)}{1+t^2} dt$$

and

$$\frac{\partial u}{\partial x_2}(x) - \frac{\partial w_0}{\partial x_2}(x) = \frac{x_1}{|x_1|} \int_0^\infty e^{-t|x_1|} \frac{t \sin(tx_2) - \cos(tx_2)}{1+t^2} dt.$$

Clearly

$$\left| \int_0^\infty e^{-t|x_1|} \frac{\sin(tx_2)}{1+t^2} dt \right| \leq \int_0^\infty \frac{dt}{1+t^2} = \frac{\pi}{2},$$

and similarly,

$$\left| \int_0^\infty e^{-t|x_1|} \frac{\cos(tx_2)}{1+t^2} dt \right| \leq \frac{\pi}{2}.$$

For the estimate of

$$\int_0^\infty e^{-t|x_1|} \frac{t \cos(tx_2)}{1+t^2} dt \quad \text{and} \quad \int_0^\infty e^{-t|x_1|} \frac{t \sin(tx_2)}{1+t^2} dt,$$

it suffices to consider $x_1 > 0$ by the symmetry. We distinguish the following three cases.

Case A: $x_1 \geq x_2$. Note that

$$\left| \int_0^\infty e^{-t|x_1|} \frac{t \cos(tx_2)}{1+t^2} dt \right| \leq \int_0^\infty \frac{te^{-t|x_1|}}{1+t^2} dt = \int_0^\infty \frac{se^{-s}}{x_1^2 + s^2} ds = I(|x_1|),$$

and similarly

$$\left| \int_0^\infty e^{-t|x_1|} \frac{t \sin(tx_2)}{1+t^2} dt \right| \leq I(|x_1|).$$

By Lemma 20, there exists a universal constant C such that

$$I(x_1) \leq \min \left\{ \log \left(\frac{1}{x_1} \right) + I_0 + \frac{\pi x_1}{2}, \frac{1}{x_1^2} \right\} \leq C \log \left(\frac{1}{x_1} + 2 \right) \quad \text{for all } x_1 > 0.$$

Together with the previous inequalities, this implies (46) for all points $(x_1, x_2) \in \mathbb{R}_+^2$ with $x_1 \geq x_2$.

Case B: $0 < x_1 < x_2 \leq 1$. A substitution gives

$$\int_0^\infty e^{-t|x_1|} \frac{t \cos(tx_2)}{1+t^2} dt = \int_0^\infty e^{-s|x_1|/x_2} \frac{s \cos s}{x_2^2 + s^2} ds.$$

For $k \in \mathbb{N}$, let

$$s_k = \int_{\pi k - \pi/2}^{\pi k + \pi/2} e^{-s|x_1|/x_2} \frac{s \cos s}{x_2^2 + s^2} ds.$$

Then $s_k > 0 > s_{k+1}$ whenever k is even. Moreover, since the function $s \mapsto e^{-s|x_1|/x_2} s/(x_2^2 + s^2)$ is decreasing for $s \geq x_2$, we also conclude that $|s_{k+1}| < |s_k|$ for all $k \in \mathbb{N}$ (because $x_2 \leq 1$). Therefore,

$$\sum_{k=1}^{\infty} s_k = \sum_{k=1}^{\infty} (s_{2k-1} + s_{2k}) < 0$$

and

$$\sum_{k=2}^{\infty} s_k = \sum_{k=1}^{\infty} (s_{2k} + s_{2k+1}) > 0.$$

These series converge by the Leibniz criterium for alternating series. Moreover, they correspond to certain integrals, and therefore, we obtain

$$\begin{aligned} \int_0^\infty e^{-s|x_1|/x_2} \frac{s \cos s}{x_2^2 + s^2} ds &= \int_0^{\pi/2} e^{-s|x_1|/x_2} \frac{s \cos s}{x_2^2 + s^2} ds + \sum_{k=1}^\infty s_k \\ &< \int_0^{\pi/2} e^{-s|x_1|/x_2} \frac{s \cos s}{x_2^2 + s^2} ds \leq \int_0^{\pi/2} \frac{s}{x_2^2 + s^2} ds = \log \sqrt{\frac{\pi^2}{4x_2^2} + 1} \end{aligned}$$

and

$$\begin{aligned} \int_0^\infty e^{-s|x_1|/x_2} \frac{s \cos s}{x_2^2 + s^2} ds &= \int_0^{3\pi/2} e^{-s|x_1|/x_2} \frac{s \cos s}{x_2^2 + s^2} ds + \sum_{k=2}^\infty s_k \\ &> \int_0^{3\pi/2} e^{-s|x_1|/x_2} \frac{s \cos s}{x_2^2 + s^2} ds \geq - \int_0^{3\pi/2} \frac{s}{x_2^2 + s^2} ds = - \log \sqrt{\frac{9\pi^2}{4x_2^2} + 1}. \end{aligned}$$

The same kind of estimate holds for $\int_0^\infty e^{-s|x_1|/x_2} \frac{s \sin s}{x_2^2 + s^2} ds$. Again we have suitable estimates for all the above integrals, which means that (46) is proved in this case.

Case C: $0 < x_1 < x_2$ and $x_2 > 1$. Here we proceed differently. We apply the mean value formula for the harmonic functions $\frac{\partial u}{\partial x_j}$, $j = 1, 2$, in the ball $B_{1/2}(x)$. We combine the resulting formula with the estimate for the Dirichlet energy in Proposition 21, which yields a uniform bound for $|\nabla u|$ in $\mathbb{R} \times [1, \infty)$. Thus (46) is proved in this case as well.

Step 2: behaviour of u and v near the origin. By (39), the dominated convergence theorem implies that $u(x_1, 0) \rightarrow -\text{sign}(x_1) \int_0^\infty \frac{dt}{1+t^2} = w_0(0\pm)$ as $x_1 \searrow 0$ or $x_1 \nearrow 0$. By Lemma 20 and (42), we also know that $v(x_1, 0) - \log \frac{1}{|x_1|} - I_0 \rightarrow 0$ as $x_1 \rightarrow 0$. Then the fundamental theorem of calculus, combined with (46) and $\nabla v = \nabla^\perp u$, gives the second statement of the proposition.

Step 3: estimates on $\nabla u_{a,d}^$.* The desired pointwise estimate for $\nabla u_{a,d}^*$ is an obvious consequence of (46), while (47) follows from Proposition 21. \square

The contribution of the tail of a single Néel wall to the renormalised energy is computed in the next lemma. As mentioned previously, in the unconfined model, this contribution depends both on the stray-field energy and the anisotropy energy. This also improves the estimate of Proposition 21.

Lemma 24. *The following limit exists and is given by*

$$\lim_{r \searrow 0} \left(\int_{\mathbb{R}_+^2 \setminus B_r(0)} |\nabla v|^2 dx + \int_{-\infty}^\infty v^2(x_1, 0) dx_1 - \pi \log \frac{1}{r} \right) = \pi I_0,$$

where I_0 is defined by (10).

Proof. Let $r \in (0, 1)$. Denote $\partial^+ B_r(0) = \partial B_r(0) \cap \mathbb{R}_+^2$ and write \mathcal{H}^1 for the 1-dimensional Hausdorff measure. Furthermore, we write $\frac{\partial v}{\partial r} = \frac{x}{|x|} \cdot \nabla v$. Since v is the harmonic conjugate of u , we have

$$\int_{\mathbb{R}_+^2 \setminus B_r(0)} |\nabla u|^2 dx = \int_{\mathbb{R}_+^2 \setminus B_r(0)} |\nabla v|^2 dx = - \int_{\partial^+ B_r(0)} v \frac{\partial v}{\partial r} d\mathcal{H}^1 - \int_{\mathbb{R} \setminus (-r, r)} v \frac{\partial v}{\partial x_2} dx_1.$$

For the last integral, since $v \in L^2(\mathbb{R})$ (see (36)), we know by (34) that

$$\int_{\mathbb{R} \setminus (-r, r)} v \frac{\partial v}{\partial x_2} dx_1 = \int_{\mathbb{R} \setminus (-r, r)} v^2 dx_1 \rightarrow \int_{-\infty}^\infty v^2(x_1, 0) dx_1$$

as $r \rightarrow 0$. For the remaining integral, denoting

$$v_0(x) = -\log|x|, \quad x \in \mathbb{R}^2 \setminus \{0\},$$

we compute:

$$\int_{\partial^+ B_r(0)} v \frac{\partial v}{\partial r} d\mathcal{H}^1 = \int_{\partial^+ B_r(0)} v \left(\frac{\partial v}{\partial r} - \frac{\partial v_0}{\partial r} \right) d\mathcal{H}^1 + \int_{\partial^+ B_r(0)} (v - v_0) \frac{\partial v_0}{\partial r} d\mathcal{H}^1 + \int_{\partial^+ B_r(0)} v_0 \frac{\partial v_0}{\partial r} d\mathcal{H}^1.$$

Note that

$$\int_{\partial^+ B_r(0)} v_0 \frac{\partial v_0}{\partial r} d\mathcal{H}^1 = \pi \log r.$$

By Proposition 23, we know that

$$\lim_{x \rightarrow 0} (v(x) - v_0(x)) = I_0.$$

Consequently,

$$\int_{\partial^+ B_r(0)} (v - v_0) \frac{\partial v_0}{\partial r} d\mathcal{H}^1 = -\frac{1}{r} \int_{\partial^+ B_r(0)} (v - v_0) d\mathcal{H}^1 \rightarrow -\pi I_0$$

as $r \rightarrow 0$. Finally, by Proposition 23 again,

$$\begin{aligned} \left| \int_{\partial^+ B_r(0)} v \left(\frac{\partial v}{\partial r} - \frac{\partial v_0}{\partial r} \right) d\mathcal{H}^1 \right| &\leq C \int_{\partial^+ B_r(0)} |v| \log \left(\frac{1}{r} + 2 \right) d\mathcal{H}^1 \\ &\leq C \int_{\partial^+ B_r(0)} \left(\log \frac{1}{r} + C' \right) \log \left(\frac{1}{r} + 2 \right) d\mathcal{H}^1 \rightarrow 0, \end{aligned}$$

for some universal constants C and C' , as $r \rightarrow 0$. The claim now follows. \square

The contribution of the tail-tail interaction between two Néel walls is computed as follows.

Lemma 25. *Let $b, c \in \mathbb{R}$ with $b \neq c$. If v_b and v_c are defined as in (43), then*

$$\int_{\mathbb{R}_+^2} \nabla v_b \cdot \nabla v_c dx + \int_{-\infty}^{\infty} v_b(x_1, 0) v_c(x_1, 0) dx_1 = \pi I(|b - c|).$$

Proof. Clearly it suffices to prove this identity for $c = 0$ and $b > 0$.

Recalling (35), we obtain a similar formula for the Fourier transform of v_b :

$$\hat{v}_b(\xi, x_2) = \frac{\pi e^{-ib\xi - |\xi|x_2}}{1 + |\xi|}.$$

Thus for $r > 0$, Plancherel's theorem and Fubini's theorem yield

$$\begin{aligned} \int_{\mathbb{R} \times (r, \infty)} \frac{\partial v_b}{\partial x_1} \frac{\partial v}{\partial x_1} dx &= \frac{\pi}{2} \int_r^\infty \int_{-\infty}^\infty \frac{\xi^2 e^{-ib\xi - 2|\xi|x_2}}{(1 + |\xi|)^2} d\xi dx_2 \\ &= \frac{\pi}{2} \int_{-\infty}^\infty \frac{\xi^2 e^{-ib\xi}}{(1 + |\xi|)^2} \int_r^\infty e^{-2|\xi|x_2} dx_2 d\xi = \frac{\pi}{4} \int_{-\infty}^\infty \frac{|\xi| e^{-ib\xi - 2|\xi|r}}{(1 + |\xi|)^2} d\xi \\ &= \frac{\pi}{4} \left(\int_0^\infty \frac{\xi e^{-ib\xi - 2\xi r}}{(1 + \xi)^2} d\xi + \int_0^\infty \frac{\xi e^{ib\xi - 2\xi r}}{(1 + \xi)^2} d\xi \right) = \frac{\pi}{2} \int_0^\infty \frac{\xi \cos(b\xi)}{(1 + \xi)^2} e^{-2\xi r} d\xi \\ &= \frac{\pi}{2} \int_0^\infty \frac{t \cos t}{(b + t)^2} e^{-2tr/b} dt. \end{aligned}$$

We wish to consider the limit $r \searrow 0$, but since the function $t \mapsto \frac{t \cos t}{(b+t)^2}$ does not belong to $L^1(0, \infty)$, some care is required here.

Claim. For any $b > 0$,

$$\lim_{r \searrow 0} \int_0^\infty \frac{t \cos t}{(b+t)^2} e^{-2tr/b} dt = \int_0^\infty \frac{t \cos t}{(b+t)^2} dt.$$

Proof of the claim. For $k \in \mathbb{N}$ and $r \geq 0$, define $s_k(r) = s_k^1(r) + s_k^2(r)$, where

$$s_k^1(r) = \int_{2k\pi-\pi/2}^{2k\pi+\pi/2} \frac{t \cos t}{(b+t)^2} e^{-2tr/b} dt, \quad s_k^2(r) = \int_{2k\pi+\pi/2}^{2k\pi+3\pi/2} \frac{t \cos t}{(b+t)^2} e^{-2tr/b} dt,$$

and define $\sigma_k = \sigma_k^1 + \sigma_k^2$, where

$$\sigma_k^1 = \int_{2k\pi-\pi/2}^{2k\pi+\pi/2} \frac{\cos t}{b+t} dt, \quad \sigma_k^2 = \int_{2k\pi+\pi/2}^{2k\pi+3\pi/2} \frac{\cos t}{b+t} dt.$$

Since the functions $t \mapsto te^{-2tr/b}/(b+t)^2$ are decreasing for $t > b$, there exists $k_0 \in \mathbb{N}$ such that $s_k^1(r) > |s_k^2(r)|$ for $k \geq k_0$. Hence $s_k(r) > 0$ for every $k \geq k_0$ and every $r \geq 0$. Similarly, $\sigma_k > 0$ for all $k \in \mathbb{N}$. We have the formulas

$$\int_{3\pi/2}^\infty \frac{t \cos t}{(b+t)^2} e^{-2tr/b} dt = \sum_{k=1}^\infty s_k(r) < \infty \quad \text{for } r > 0,$$

and¹⁵

$$\int_{3\pi/2}^\infty \frac{\cos t}{b+t} dt = \sum_{k=1}^\infty \sigma_k < \infty.$$

The convergence of these series follows from the Leibniz criterion for alternating series, since $s_k^1(r) > 0 > s_k^2(r)$ and $|s_k^i(r)| \searrow 0$ as $k \rightarrow \infty$ for $i = 1, 2$, and similarly, $\sigma_k^1 > 0 > \sigma_k^2$ and $|\sigma_k^i| \searrow 0$ as $k \rightarrow \infty$ for $i = 1, 2$. We now consider, for $r > 0$, the function

$$f_r(t) = \frac{1}{b+t} - \frac{te^{-2rt/b}}{(b+t)^2} \quad \text{for } t > 0$$

and claim that f_r is decreasing in t . Indeed, we compute

$$f_r'(t) = \frac{t}{(b+t)^3} \left(\frac{2rt}{b} e^{-2rt/b} - 1 + e^{-2rt/b} \right) + \frac{b}{(b+t)^3} \left(\frac{2rt}{b} e^{-2rt/b} - 1 - e^{-2rt/b} \right).$$

For any $R \geq 0$, the inequality $Re^{-R} \leq 1 - e^{-R}$ is easy to see. Using it for $R = 2rt/b$, we find that $f_r'(t) < 0$. In particular, we have $f_r \geq \lim_{t \rightarrow \infty} f_r(t) = 0$.

Now we observe that

$$\sigma_k - s_k(r) = \int_{2k\pi-\pi/2}^{2k\pi+3\pi/2} f_r(t) \cos t dt.$$

By the monotonicity of f_r , as we integrate over a full period of \cos beginning with its positive part, we conclude that $\sigma_k \geq s_k(r)$ for all $r > 0$ and all $k \in \mathbb{N}$. Also, we clearly have the convergence

$$s_k(0) = \lim_{r \searrow 0} s_k(r).$$

As $\sum_k \sigma_k$ converges, Lebesgue's dominated convergence theorem, applied to the counting measure, now implies that

$$\sum_{k=1}^\infty s_k(0) = \lim_{r \searrow 0} \sum_{k=1}^\infty s_k(r).$$

¹⁵The convergence of the integral was also proved via the contour integral in Lemma 18.

This completes the proof of the claim.

Finally, Propositions 21 and 23 imply that $\frac{\partial v_b}{\partial x_1} \frac{\partial v}{\partial x_1} \in L^1(\mathbb{R}_+^2)$. Therefore, the above claim implies

$$\int_{\mathbb{R}_+^2} \frac{\partial v_b}{\partial x_1} \frac{\partial v}{\partial x_1} dx = \lim_{r \searrow 0} \int_{\mathbb{R} \times (r, \infty)} \frac{\partial v_b}{\partial x_1} \frac{\partial v}{\partial x_1} dx = \frac{\pi}{2} \int_0^\infty \frac{t \cos t}{(b+t)^2} dt.$$

Moreover, the same computations give

$$\int_{\mathbb{R}_+^2} \frac{\partial v_b}{\partial x_2} \frac{\partial v}{\partial x_2} dx = \frac{\pi}{2} \int_0^\infty \frac{t \cos t}{(b+t)^2} dt$$

as well. Thus

$$\int_{\mathbb{R}_+^2} \nabla v_b \cdot \nabla v dx = \pi \int_0^\infty \frac{t \cos t}{(b+t)^2} dt.$$

Set $\mu(x_1) = v(x_1, 0)$ and $\mu_b(x_1) = v(x_1 - b, 0)$. Then

$$\hat{\mu}(\xi) = \frac{\pi}{1 + |\xi|} \quad \text{and} \quad \hat{\mu}_b(\xi) = \frac{\pi e^{-ib\xi}}{1 + |\xi|}.$$

Hence

$$\int_{-\infty}^\infty \mu_b \mu dx_1 = \frac{\pi}{2} \int_{-\infty}^\infty \frac{e^{-ib\xi}}{(1 + |\xi|)^2} d\xi = \pi \int_0^\infty \frac{\cos(b\xi)}{(1 + \xi)^2} d\xi = b\pi \int_0^\infty \frac{\cos t}{(b+t)^2} dt.$$

If we combine this with the previous identity, we obtain, by Lemma 18,

$$\int_{\mathbb{R}_+^2} \nabla v_b \cdot \nabla v dx + \int_{-\infty}^\infty \mu_b \mu dx_1 = \pi \int_0^\infty \frac{\cos t}{b+t} dt = \pi I(b),$$

which is the desired identity. \square

Proof of Proposition 22. This is now an easy consequence of Lemmas 24 and 25. \square

3.4 Preliminary estimates

The rest of the arguments for the proof of Theorem 8 follow the strategy of the proof of Theorem 1, given in our previous paper [18]. Because these arguments are lengthy, we do not repeat the details and merely highlight the modifications where necessary.

In the first step, we prove the following result. This has a direct counterpart for the confined problem from Section 2, see [18, Theorem 28].

Theorem 26 (Stray field energy estimate). *Let $N \in \mathbb{N}$, $R > 0$, and $C_0 > 0$. Then there exists $C_1 > 0$ such that for any $a \in A_N$ with $\rho(a) \geq R$ and any $d \in \{\pm 1\}^N$, the following holds true. Let*

$$\Gamma = \sum_{n=1}^N (d_n - \cos \alpha)^2. \tag{48}$$

Suppose that $\epsilon \in (0, \frac{1}{2}]$ with $\delta \leq R$ and $m \in M(a, d)$ with

$$E_\epsilon(m) \leq \frac{\pi\Gamma}{2 \log \frac{1}{\delta}} + \frac{C_0}{(\log \delta)^2}.$$

Let $u \in \dot{H}^1(\mathbb{R}_+^2)$ be the solution of (14) and (15). Then

$$\epsilon \int_{\mathbb{R}} |m'|^2 dx_1 + \int_{\mathbb{R}} (m_1 - \cos \alpha)^2 dx_1 + \int_{\Omega_\delta(a)} \left| \nabla u - \frac{\nabla u_{a,d}^*}{\log \frac{1}{\delta}} \right|^2 dx \leq \frac{C_1}{(\log \delta)^2}$$

and

$$\int_{\Omega_\delta(a)} |\nabla u|^2 dx \geq \frac{\pi \Gamma}{\log \frac{1}{\delta}} - \frac{C_1}{(\log \delta)^2}.$$

Proof. We first note that it suffices to prove these inequalities for ϵ small. Consider the function $u_{a,d}^*/|\log \delta|$ and modify it near the points $(a_n, 0)$, $n = 1, \dots, N$, similarly to the proof of [18, Theorem 28]. To this end, note first that by Proposition 23, we know that $u_{a,d}^*(\cdot, 0)$ is continuous on (a_{n-1}, a_n) for $n = 2, \dots, N$ and on $(-\infty, a_1)$ and (a_N, ∞) , and the one-sided limits

$$\tau_n^+ = \lim_{x_1 \searrow a_n} u_{a,d}^*(x_1, 0) \quad \text{and} \quad \tau_n^- = \lim_{x_1 \nearrow a_n} u_{a,d}^*(x_1, 0)$$

exist. Then $\tau_n^+ - \tau_n^- = -\gamma_n \pi$ for $n = 1, \dots, N$ (by Proposition 23 again). For a given number $s \in (0, R]$, we now define $\xi_s \in \dot{H}^1(\mathbb{R}_+^2)$ such that $\xi_s = u_{a,d}^*/|\log \delta|$ in $\Omega_s(a)$ and

$$\xi_s(a_n + r \cos \theta, r \sin \theta) = \frac{r u_{a,d}^*(a_n + s \cos \theta, s \sin \theta)}{s \log \frac{1}{\delta}} + \left(1 - \frac{r}{s}\right) \frac{\tau_n^+ + \tau_n^-}{2 \log \frac{1}{\delta}} \quad (49)$$

for $0 < r \leq s$, $0 < \theta < \pi$, and $n = 1, \dots, N$. Then the inequalities of Propositions 21 and 23 imply

$$\int_{\bigcup_{n=1}^N B_s(a_n, 0)} |\nabla \xi_s|^2 dx \leq \frac{C_2}{(\log \delta)^2} \quad \text{and} \quad \int_{\mathbb{R}_+^2} |\nabla \xi_s|^2 dx \leq \frac{\pi \Gamma \log \frac{1}{s} + C_2}{(\log \delta)^2} \quad (50)$$

for some constant $C_2 = C_2(N, R)$. Also define

$$\zeta(x_1) = \frac{v_{a,d}^*(x_1, 0)}{\log \frac{1}{\delta}}.$$

Using (36), we easily find a constant $C_3 = C_3(N)$ such that

$$\|\zeta\|_{L^2(\mathbb{R})} \leq \frac{C_3}{\log \frac{1}{\delta}}. \quad (51)$$

Since $m_1(a_n) = d_n$ for $n = 1, \dots, N$ and $\lim_{x_1 \rightarrow \pm\infty} m_1(x_1) = \cos \alpha$, the fundamental theorem of calculus gives the identity

$$\int_{a_{n-1}}^{a_n} \frac{d}{dx_1} ((m_1(x_1) - \cos \alpha) u_{a,d}^*(x_1, 0)) dx_1 = \gamma_n \tau_n^- - \gamma_{n-1} \tau_{n-1}^+$$

for $n = 2, \dots, N$, while

$$\int_{-\infty}^{a_1} \frac{d}{dx_1} ((m_1(x_1) - \cos \alpha) u_{a,d}^*(x_1, 0)) dx_1 = \gamma_1 \tau_1^-$$

and

$$\int_{a_N}^{\infty} \frac{d}{dx_1} ((m_1(x_1) - \cos \alpha) u_{a,d}^*(x_1, 0)) dx_1 = -\gamma_N \tau_N^+.$$

Observing that $\frac{\partial}{\partial x_1} u_{a,d}^* = \frac{\partial}{\partial x_2} v_{a,d}^* = v_{a,d}^*$ on $(\mathbb{R} \setminus \{a_1, \dots, a_N\}) \times \{0\}$, we find that

$$\frac{d}{dx_1} ((m_1(x_1) - \cos \alpha) u_{a,d}^*(x_1, 0)) = m_1'(x_1) u_{a,d}^*(x_1, 0) + (m_1(x_1) - \cos \alpha) v_{a,d}^*(x_1)$$

away from a_1, \dots, a_N . Therefore,

$$\int_{-\infty}^{\infty} (m'_1(x_1)u_{a,d}^*(x_1, 0) + (m_1(x_1) - \cos \alpha)v_{a,d}^*(x_1)) dx_1 = \sum_{n=1}^N \gamma_n(\tau_n^- - \tau_n^+) = \pi\Gamma.$$

It follows from (14), (15), the definition of ξ_s , and [18, Lemma 29] that

$$\begin{aligned} \frac{\pi\Gamma}{\log \frac{1}{\delta}} &= \int_{-\infty}^{\infty} \xi_s(x_1, 0)m'_1(x_1) dx_1 + \int_{\mathbb{R}} (m_1 - \cos \alpha)\zeta dx_1 - \int_{\mathbb{R}} \left(\xi_s(x_1, 0) - \frac{u_{a,d}^*(x_1, 0)}{\log \frac{1}{\delta}} \right) m'_1(x_1) dx_1 \\ &\leq \int_{\mathbb{R}_+^2} \nabla \xi_s \cdot \nabla u dx + \int_{-\infty}^{\infty} (m_1 - \cos \alpha)\zeta dx_1 + \frac{C_4 s}{\log \frac{1}{\delta}} \int_{-\infty}^{\infty} |m'|^2 dx_1 \end{aligned}$$

for a constant $C_4 = C_4(\alpha, N, R) > 0$. Hence

$$\begin{aligned} \frac{\pi\Gamma}{\log \frac{1}{\delta}} &\leq \frac{C_4 s}{\epsilon \log \frac{1}{\delta}} \left(2E_\epsilon(m) - \|\nabla u\|_{L^2(\mathbb{R}_+^2)}^2 - \|m_1 - \cos \alpha\|_{L^2(\mathbb{R})}^2 \right) \\ &\quad + \int_{\mathbb{R}_+^2} \nabla \xi_s \cdot \nabla u dx + \int_{-\infty}^{\infty} (m_1 - \cos \alpha)\zeta dx_1 \\ &\leq \frac{C_4 s}{\epsilon \log \frac{1}{\delta}} \left(2E_\epsilon(m) - \|\nabla u\|_{L^2(\mathbb{R}_+^2)}^2 - \|m_1 - \cos \alpha\|_{L^2(\mathbb{R})}^2 \right) \\ &\quad + \sqrt{\|\nabla \xi_s\|_{L^2(\mathbb{R}_+^2)}^2 + \|\zeta\|_{L^2(\mathbb{R})}^2} \sqrt{\|\nabla u\|_{L^2(\mathbb{R}_+^2)}^2 + \|m_1 - \cos \alpha\|_{L^2(\mathbb{R})}^2} \\ &\leq \frac{C_4 s}{\epsilon \log \frac{1}{\delta}} \left(2E_\epsilon(m) - \|\nabla u\|_{L^2(\mathbb{R}_+^2)}^2 - \|m_1 - \cos \alpha\|_{L^2(\mathbb{R})}^2 \right) \\ &\quad + \frac{\sqrt{\pi\Gamma \log \frac{1}{s} + C_5}}{\log \frac{1}{\delta}} \sqrt{\|\nabla u\|_{L^2(\mathbb{R}_+^2)}^2 + \|m_1 - \cos \alpha\|_{L^2(\mathbb{R})}^2}, \end{aligned} \tag{52}$$

where $C_5 = C_2 + C_3^2$. If we choose s such that

$$\|\nabla u\|_{L^2(\mathbb{R}_+^2)}^2 + \|m_1 - \cos \alpha\|_{L^2(\mathbb{R})}^2 = \frac{\pi\Gamma}{\log \frac{1}{s}} - \frac{2C_5}{(\log s)^2},$$

then we obtain essentially the same inequality as in the proof of [18, Theorem 28], except that we must now consider the quantities ∇u and $m_1 - \cos \alpha$ jointly. The argument in the proof of [18, Theorem 28] still applies and yields $s \geq \tilde{C}\delta$ for some $\tilde{C} = \tilde{C}(\alpha, R, N, C_0)$. We keep following the reasoning of [18, Theorem 28] and obtain a constant $C_6 = C_6(\alpha, N, R)$ such that

$$\int_{\mathbb{R}_+^2} |\nabla u|^2 dx + \int_{-\infty}^{\infty} (m_1 - \cos \alpha)^2 dx_1 \geq \frac{\pi\Gamma}{\log \frac{1}{\delta}} - \frac{C_6}{(\log \delta)^2}. \tag{53}$$

Next we use the first inequality in (52) again, but with $s = \delta$. Combining it with (53), we obtain a constant $C_7 = C_7(\alpha, N, R)$ such that

$$\int_{\mathbb{R}_+^2} \nabla \xi_\delta \cdot \nabla u dx + \int_{-\infty}^{\infty} (m_1 - \cos \alpha)\zeta dx_1 \geq \frac{\pi\Gamma}{\log \frac{1}{\delta}} - \frac{C_7}{(\log \delta)^2}.$$

Furthermore, the arguments in the proof of [18, Theorem 28] give

$$\int_{\mathbb{R}_+^2} |\nabla u - \nabla \xi_\delta|^2 dx + \int_{-\infty}^{\infty} (m_1 - \cos \alpha - \zeta)^2 dx_1 \leq \frac{C_8}{(\log \delta)^2} \tag{54}$$

for some constant $C_8 = C_8(\alpha, N, R)$. Since $\xi_\delta = u_{a,d}^*/|\log \delta|$ in $\Omega_\delta(a)$, we deduce that

$$\int_{\Omega_\delta(a)} \left| \nabla u - \frac{\nabla u_{a,d}^*}{\log \frac{1}{\delta}} \right|^2 dx + \int_{-\infty}^{\infty} (m_1 - \cos \alpha - \zeta)^2 dx_1 \leq \frac{C_8}{(\log \delta)^2}.$$

Now inequalities (50), (51), and (54) imply that

$$\int_{\bigcup_{n=1}^N B_\delta(a_n, 0)} |\nabla u|^2 dx = O\left(\frac{1}{(\log \delta)^2}\right) = \int_{-\infty}^{\infty} (m_1 - \cos \alpha)^2 dx_1.$$

We may combine this with (53). We conclude that

$$\int_{\Omega_\delta(a)} |\nabla u|^2 dx \geq \frac{\pi\Gamma}{\log \frac{1}{\delta}} - O\left(\frac{1}{(\log \delta)^2}\right).$$

Finally, this estimate, combined with the bound on $E_\epsilon(m)$, gives rise to the remaining inequality. \square

We will also need some estimates for higher derivatives of critical points of E_ϵ . These will satisfy an Euler-Lagrange equation that is most easily stated in terms of the lifting ϕ of a map $m: \mathbb{R} \rightarrow \mathbb{S}^1$. Suppose that $m = (\cos \phi, \sin \phi)$ is a critical point of E_ϵ and let $u \in \dot{H}^1(\mathbb{R}_+^2)$ be a solution of (14), (15). Then

$$\epsilon \phi'' = (\cos \alpha - \cos \phi + u') \sin \phi \quad \text{in } \mathbb{R}. \quad (55)$$

Here we use the shorthand notation u' for $\frac{\partial u}{\partial x_1}(\cdot, 0)$. We refer to our previous paper [19] for a derivation. Equation (15) can be expressed in terms of ϕ , too, yielding

$$\frac{\partial u}{\partial x_2} = \phi' \sin \phi.$$

For the functional without anisotropy, estimates for higher derivatives were obtained in [18, Lemma 11]. For the case $\epsilon = 1$ (but with anisotropy term), the same arguments were used in [19, Lemma 3.3]. Examining both proofs, it is easy to see that the following statement is true. (We do not repeat the arguments here.)

Lemma 27. *Let $0 \leq r < r' < R' < R$. Then there exists a constant C (depending only on $r' - r$ and $R - R'$) such that the following holds true. Let $\epsilon > 0$ and $J = (-R, -r) \cup (r, R)$. Suppose that $\phi \in H^1(J)$ and $u \in H^1(B_R^+(0) \setminus B_r(0))$ solve the system*

$$\begin{aligned} \Delta u &= 0 && \text{in } B_R^+(0) \setminus B_r(0), \\ \frac{\partial u}{\partial x_2} &= \phi' \sin \phi && \text{on } J \times \{0\}, \\ \epsilon \phi'' &= (\cos \alpha - \cos \phi + u') \sin \phi && \text{in } J. \end{aligned}$$

Further suppose that $\sin \phi \neq 0$ in J . Then

$$\begin{aligned} \int_{B_{R'}^+(0) \setminus B_{r'}(0)} |\nabla^2 u|^2 dx + \int_{(-R', -r') \cup (r', R')} (\epsilon (\phi'')^2 + \epsilon (\phi')^4 (1 + \cot^2 \phi) + (\phi')^2 \sin^2 \phi) dx_1 \\ \leq C \epsilon \int_J (\phi')^2 dx_1 + C \int_{B_R^+(0) \setminus B_r(0)} |\nabla u|^2 dx. \end{aligned}$$

The following statement relies on the Euler-Lagrange equation as well, but applies to minimisers of the energy in $M(a, d)$. The arguments here are similar to [19, Lemma 3.1]. The result is useful above all in view of the condition $\sin \phi \neq 0$ in the preceding lemma.

Lemma 28. *Suppose that $m \in M(a, d)$ satisfies*

$$E_\epsilon(m) = \inf_{M(a,d)} E_\epsilon.$$

Then $m_1(x_1) \neq \pm 1$ for all $x_1 \in \mathbb{R} \setminus \{a_1, \dots, a_N\}$.

Proof. Choose $\phi: \mathbb{R} \rightarrow \mathbb{R}$ such that $m = (\cos \phi, \sin \phi)$. Then ϕ satisfies the Euler-Lagrange equation (55) away from a_1, \dots, a_N . Assume, by way of contradiction, that we have $b \in \mathbb{R} \setminus \{a_1, \dots, a_N\}$ with $\sin \phi(b) = 0$. Consider the initial value problem

$$\epsilon \psi'' = (\cos \alpha - \cos \phi + u') \sin \psi, \quad \psi(b) = \phi(b), \quad \psi'(b) = 0.$$

Then $\psi \equiv \phi(b)$ is the unique solution. The function ϕ also satisfies the differential equation and the first initial condition. But clearly it cannot be constant, so we conclude that $\phi'(b) \neq 0$.¹⁶ Adding a multiple of 2π if necessary, we may assume that $\phi(b) = 0$ or $\phi(b) = \pi$. In the first case, we define $\tilde{\phi}: \mathbb{R} \rightarrow \mathbb{R}$ by

$$\tilde{\phi}(x_1) = \begin{cases} \phi(x_1) & \text{if } x_1 \leq b, \\ -\phi(x_1) & \text{if } x_1 > b. \end{cases}$$

Then $\tilde{m} = (\cos \tilde{\phi}, \sin \tilde{\phi})$ belongs to $M(a, d)$ as well, and $E_\epsilon(\tilde{\phi}) = E_\epsilon(\phi)$. Hence \tilde{m} minimises the energy in $M(a, d)$ and $\tilde{\phi}$ is a solution of the Euler-Lagrange equation away from a_1, \dots, a_N . We can show, however, that solutions of this equation are necessarily smooth (see e.g. [16, Theorem 1.1]), whereas $\tilde{\phi}$ is clearly not smooth at b . In the case $\phi(b) = \pi$, we can use a similar construction and obtain the same contradiction. Hence there is no point $b \in \mathbb{R} \setminus \{a_1, \dots, a_N\}$ with $\sin \phi(b) = 0$. \square

3.5 Proof of Theorem 8

As mentioned previously, we follow the arguments from [18, Section 6] in the proof of Theorem 8 without repeating all of the details. In order to help the reader follow these arguments, we mimic the presentation of the proof as well.

We fix $a \in A_N$ and $d \in \{\pm 1\}^N$. Set $\gamma_n = d_n - \cos \alpha$ for $n = 1, \dots, N$ and recall the definition $\Gamma = \sum_{n=1}^N \gamma_n^2$ in (48). Throughout the following arguments, we use the symbol C to indiscriminately denote various constants depending only on α , N , and a , and occasionally on the exponent of an L^p -space used in the context.

Part of the proof requires a construction where we glue a ‘tail’ profile together with a number of ‘core’ profiles (one for every $n = 1, \dots, N$). While the tail profile comes from the previously constructed function $u_{a,d}^*$ (and is different from [18]), we use exactly the same core profile as in the previous paper. This may seem somewhat inconsistent, as we neglect the anisotropy there, but in the limit the difference will be invisible. The core profiles are minimisers of an auxiliary functional $E_\epsilon^{\gamma_\pm}$, see [18, Section 4]. Hence the quantities γ_\pm and $E_\epsilon^{\gamma_\pm}$ are the same as in the other paper. Furthermore, we now use the symbol e for the function $e: \{\pm 1\} \rightarrow \mathbb{R}$ defined in our previous paper [18, Definition 26] (and also mentioned in Theorem 1) and called the ‘core energy’. To avoid confusion, we do not use the exponential function any more.

3.5.1 Preparation

Recall the function

$$w_0(x) = -\arctan \frac{x_1}{x_2}$$

¹⁶This kind of argument was also used by Capella-Melcher-Otto [4].

defined in Proposition 23. (The same function is defined by a different formula in [18].) Recall that

$$u_{a,d}^* = \sum_{n=1}^N \gamma_n u_{a_n} \quad \text{and} \quad v_{a,d}^* = \sum_{n=1}^N \gamma_n v_{a_n}$$

for the functions constructed in Section 3.2 (see (43) and the subsequent formulas). We also set $\mu_{a,d}^*(x_1) = v_{a,d}^*(x_1, 0)$. By (42),

$$\mu_{a,d}^*(x_1) = \sum_{n=1}^N \gamma_n I(|x_1 - a_n|).$$

Let $r \in (0, 1/2]$ with $r \leq \rho(a)$ (where $\rho(a)$ is defined as in (44)). For $n = 1, \dots, N$, let

$$\lambda_n = \sum_{k \neq n} \gamma_k I(|a_k - a_n|).$$

As I is locally Lipschitz continuous in $(0, \infty)$, we find that

$$|\mu_{a,d}^*(x_1) - \lambda_n - \gamma_n I(|x_1 - a_n|)| \leq Cr \tag{56}$$

for $x_1 \in [a_n - r, a_n + r]$. Also define

$$\omega_n = \sum_{k \neq n} \gamma_k u_{a_k}(a_n, 0).$$

Then by Proposition 23,

$$|u_{a,d}^*(x) - \omega_n - \gamma_n w_0(x_1 - a_n, x_2)| \leq Cr \log \frac{1}{r} \quad \text{in } B_r^+(a_n, 0) \tag{57}$$

and

$$|\nabla u_{a,d}^* - \gamma_n \nabla w_0(x_1 - a_n, x_2)| \leq C \log \frac{1}{r} \quad \text{in } B_r^+(a_n, 0). \tag{58}$$

We define $W_1(a, d)$ as in Section 3.3 and

$$W_2(a, d) = -\pi \sum_{n=1}^N \gamma_n \lambda_n = -\pi \sum_{n=1}^N \sum_{k \neq n} \gamma_k \gamma_n I(|a_k - a_n|).$$

If W is the function defined in Section 1.3, then by Proposition 22:

$$W(a, d) = W_1(a, d) + W_2(a, d) - \pi I_0 \sum_{n=1}^N \gamma_n^2 = -W_1(a, d). \tag{59}$$

3.5.2 A lower bound for the interaction energy

We first want to prove that

$$\liminf_{\epsilon \searrow 0} \left((\log \delta)^2 \inf_{M(a,d)} E_\epsilon - \frac{\pi \Gamma}{2} \log \frac{1}{\delta} \right) \geq \sum_{n=1}^N e(d_n) + W(a, d).$$

First step: use minimisers Similarly to [20, Proposition 1], we conclude that E_ϵ has minimisers m_ϵ in $M(a, d)$. Clearly it suffices to consider these minimisers. We claim that

$$\limsup_{\epsilon \searrow 0} \left((\log \delta)^2 E_\epsilon(m_\epsilon) - \frac{\pi\Gamma}{2} \log \frac{1}{\delta} \right) < \infty.$$

In order to prove this, we first consider the case where $a_n \in (-1, 1)$ for $n = 1, \dots, N$. Then we can apply the results from our previous paper [18], in particular Proposition 27, which states that there exist a constant C_0 and $\hat{m}_\epsilon \in M(a, d)$ with $\hat{m}_{\epsilon 1} \equiv \cos \alpha$ outside of $(-1, 1)$ such that

$$\frac{\epsilon}{2} \int_{-1}^1 |\hat{m}'_\epsilon|^2 dx_1 + \frac{1}{2} \int_{\mathbb{R}_+^2} |\nabla \hat{u}_\epsilon|^2 dx \leq \frac{\pi\Gamma}{2 \log \frac{1}{\delta}} + \frac{C_0}{(\log \delta)^2},$$

where $\hat{u}_\epsilon \in \dot{H}^1(\mathbb{R}_+^2)$ solves (14), (15) for \hat{m}_ϵ instead of m . Let $\hat{v}_\epsilon \in \dot{H}^1(\mathbb{R}_+^2)$ be the harmonic extension of $\hat{m}_{\epsilon 1} - \cos \alpha$ to \mathbb{R}_+^2 (so that \hat{u}_ϵ and \hat{v}_ϵ are conjugate harmonic functions). Theorem 28 and Remark 31 in [18] imply (by the dyadic decomposition argument of Struwe [37]) that for any $p \in [1, 2)$, the following inequality holds true:

$$\frac{\epsilon}{2} \int_{-1}^1 |\hat{m}'_\epsilon|^2 dx_1 + \|\nabla \hat{v}_\epsilon\|_{L^p(B_2^+(0))}^2 \leq \frac{C}{(\log \delta)^2}.$$

Standard trace theorems for Sobolev spaces then imply that

$$\int_{-1}^1 (\hat{m}_{\epsilon 1} - \cos \alpha)^2 dx_1 \leq \frac{C}{(\log \delta)^2}.$$

Hence

$$E_\epsilon(m_\epsilon) \leq E_\epsilon(\hat{m}_\epsilon) \leq \frac{\pi\Gamma}{2 \log \frac{1}{\delta}} + \frac{C}{(\log \delta)^2}.$$

For any other $a \in A_N$, we may scale the domain such that we are in the above situation and observe that the stray field energy does not change under such scaling. The exchange energy and the anisotropy energy will change (one of them will decrease and the other increase), but only by a factor depending on a . As both of them are of order $1/(\log \delta)^2$, we still obtain an inequality of the same form, albeit for a different constant C .

Second step: prove convergence away from the walls By Theorem 26 and Proposition 21, we have a sequence $\epsilon_k \searrow 0$ such that the functions $w_k = u_{\epsilon_k} \log \frac{1}{\delta_k}$ satisfy $w_k \rightharpoonup w$ weakly in $\dot{H}^1(\Omega_s(a)) \cap W_{\text{loc}}^{1,p}(\mathbb{R}_+^2)$ for all $s > 0$ and $p \in [1, 2)$, where $w \in u_{a,d}^* + \dot{H}^1(\mathbb{R}_+^2)$ and $\delta_k = \epsilon_k \log \frac{1}{\epsilon_k}$. Moreover, by Theorem 26, we may choose this subsequence such that $\mu_k = (m_{\epsilon_k 1} - \cos \alpha) \log \frac{1}{\delta_k} \rightharpoonup \mu$ weakly in $L^2(\mathbb{R})$ for some $\mu \in L^2(\mathbb{R})$.

But we have in fact better convergence: Lemma 27 and Theorem 26 imply that $w_k \rightharpoonup w$ weakly in $H^2(B_R(0) \cap \Omega_s(a))$ for any $R, s > 0$ and $\mu_k \rightarrow \mu$ in the strong L^∞ and weak H^1 sense in $(-R, R) \setminus \bigcup_{n=1}^N (a_n - s, a_n + s)$ for any $s > 0$. (We can use Lemma 27 here because of Lemma 28.) Furthermore,

$$\limsup_{k \rightarrow \infty} \left((\log \delta_k)^2 \epsilon_k \int_{(-R, R) \setminus \bigcup_{n=1}^N (a_n - s, a_n + s)} (\phi''_{\epsilon_k})^2 dx_1 \right) < \infty.$$

By the Euler-Lagrange equation,

$$u'_\epsilon + \cos \alpha - \cos \phi_\epsilon = \frac{\epsilon \phi''_\epsilon}{\sin \phi_\epsilon} \quad \text{in } \mathbb{R} \setminus \{a_1, \dots, a_N\}.$$

It follows that $w'_k(\cdot, 0) - \mu_k \rightharpoonup 0$ in the distribution sense in $\mathbb{R} \setminus \{a_1, \dots, a_N\}$. Since by the trace theorem, $w'_k(\cdot, 0) \rightharpoonup w'(\cdot, 0)$ as distributions in \mathbb{R} , and since $\mu_k \rightharpoonup \mu$ weakly in $L^2(\mathbb{R})$, we conclude that $w'(\cdot, 0) = \mu$ in $\mathbb{R} \setminus \{a_1, \dots, a_N\}$. Clearly $\Delta w = 0$ in \mathbb{R}_+^2 . If we write $\frac{\partial w}{\partial x_2}$ for the distribution on \mathbb{R} such that for any $\eta \in C_0^\infty(\overline{\mathbb{R}_+^2})$,

$$\int_{-\infty}^{\infty} \frac{\partial w}{\partial x_2} \eta dx_1 = - \int_{\mathbb{R}_+^2} \nabla w \cdot \nabla \eta dx,$$

then the equations $\frac{\partial w_k}{\partial x_2} = -\mu'_k$ and the weak $W_{\text{loc}}^{1,p}$ -convergence for $p < 2$ imply that $\frac{\partial w}{\partial x_2} = -\mu'$ in the sense of distributions.

We claim that only the function $u_{a,d}^*$ has these properties. In order to prove this claim, set $f = w - u_{a,d}^*$ and $h = \mu - \mu_{a,d}^*$. Then $f \in \dot{H}^1(\mathbb{R}_+^2)$, $h \in L^2(\mathbb{R})$, and $\Delta f = 0$ in \mathbb{R}_+^2 . Moreover, $f' = h$ in $\mathbb{R} \setminus \{a_1, \dots, a_N\}$. But since $f(\cdot, 0) \in \dot{H}^{1/2}(\mathbb{R})$ cannot have any jumps, this implies that $f' = h$ in all of \mathbb{R} . We also know that $\frac{\partial f}{\partial x_2} = -h'$ on $\mathbb{R} \times \{0\}$ in the sense that

$$\int_{-\infty}^{\infty} h \eta' dx_1 = - \int_{\mathbb{R}_+^2} \nabla f \cdot \nabla \eta dx$$

for any $\eta \in C_0^\infty(\overline{\mathbb{R}_+^2})$. If g denotes the conjugate harmonic function to f such that $\nabla g = \nabla^\perp f$, then

$$\begin{aligned} \int_{\mathbb{R} \times \{0\}} h \eta' dx_1 &= \int_{\mathbb{R}_+^2} \nabla^\perp g \cdot \nabla \eta dx = \lim_{s \searrow 0} \int_{\mathbb{R} \times (s, \infty)} \nabla^\perp g \cdot \nabla \eta dx = - \lim_{s \searrow 0} \int_{-\infty}^{\infty} \frac{\partial g}{\partial x_1}(x_1, s) \eta(x_1, s) dx_1 \\ &= \lim_{s \searrow 0} \int_{-\infty}^{\infty} g(x_1, s) \frac{\partial \eta}{\partial x_1}(x_1, s) dx_1 = \int_{\mathbb{R} \times \{0\}} g \eta' dx_1. \end{aligned}$$

Thus after adding a suitable constant, we find that $g = h$ and $\frac{\partial}{\partial x_2} g = g$ on $\mathbb{R} \times \{0\}$. As g is harmonic, the last identity, combined with an integration by parts, implies that $\int_{\mathbb{R}_+^2} |\nabla g|^2 dx = - \int_{\mathbb{R}} g^2 dx_1$. That is, $g = 0$ in \mathbb{R}_+^2 , and hence $w = u_{a,d}^*$ and $\mu = \mu_{a,d}^*$. As we have thus identified unique limits, the above convergence holds in fact not just for the sequences w_k and μ_k , but for the full families $u_\epsilon |\log \delta|$ and $(m_{\epsilon_1} - \cos \alpha) |\log \delta|$, in the same sense, as $\epsilon \searrow 0$.

We conclude in particular that

$$\int_{\Omega_r(a)} |\nabla u_{a,d}^*|^2 dx \leq \liminf_{\epsilon \searrow 0} \left((\log \delta)^2 \int_{\Omega_r(a)} |\nabla u_\epsilon|^2 dx \right)$$

and

$$\int_{-\infty}^{\infty} (\mu_{a,d}^*)^2 dx_1 \leq \liminf_{\epsilon \searrow 0} \left((\log \delta)^2 \int_{-\infty}^{\infty} (m_{\epsilon_1} - \cos \alpha)^2 dx_1 \right).$$

Note also that for every $r \in (0, 1/2]$ with $r \leq \rho(a)$,

$$\left| m_{\epsilon_1}(a_n \pm r) - \cos \alpha - \frac{\gamma_n I(r) + \lambda_n}{\log \frac{1}{\delta}} \right| \leq \frac{Cr + o(1)}{\log \frac{1}{\delta}}$$

by (56) and the above convergence.

Third step: rescale the cores Fix $n \in \{1, \dots, N\}$, $r \in (0, \rho(a))$ and consider the rescaled functions $\tilde{m}_\epsilon : \mathbb{R} \rightarrow \mathbb{S}^1$ and $\tilde{u}_\epsilon : \mathbb{R}_+^2 \rightarrow \mathbb{R}$ defined by

$$\tilde{m}_\epsilon(x_1) = m_\epsilon(rx_1 + a_n), \quad \tilde{u}_\epsilon(x) = u_\epsilon(rx_1 + a_n, rx_2), \quad \tilde{\epsilon} = \frac{\epsilon}{r}.$$

Then the arguments from the first half of p. 475 in [18] apply, the only difference being that some of the right-hand sides of the estimates need to be multiplied by $\log \frac{1}{r}$, due to the appearance of this factor in (58). We then apply [18, Corollary 21] to $\mu = d_n \tilde{m}_\epsilon$ with

$$\gamma = d_n \gamma_n, \quad \eta = Cr \log \frac{1}{r} + o(1), \quad \text{and} \quad \zeta = d_n(\lambda_n + \gamma_n I(r)) + Cr + o(1).$$

This eventually gives the inequality

$$\begin{aligned} (\log \delta)^2 & \left(\epsilon \int_{a_n-r}^{a_n+r} |m'_\epsilon|^2 dx_1 + \int_{B_r^+(a_n,0)} |\nabla u_\epsilon|^2 dx \right) - \pi \gamma_n^2 \left(\log \frac{1}{\delta} - \log \frac{1}{r} \right) \\ & \geq 2e(d_n) - 2\pi \gamma_n \lambda_n - 2\pi \gamma_n^2 I(r) + 2\pi \gamma_n^2 \log \frac{1}{r} - Cr \log \frac{1}{r} - o(1) \\ & \geq 2e(d_n) - 2\pi \gamma_n \lambda_n - 2\pi \gamma_n^2 I_0 - Cr \log \frac{1}{r} - o(1). \end{aligned}$$

(The last inequality comes from Lemma 20.)

Fourth step: combine the estimates We can now adapt the arguments on p. 476 in [18] to our unconfined model, recalling identity (59). The desired lower bound follows.

3.5.3 An upper bound for the interaction energy

We now want to prove the inequality

$$\limsup_{\epsilon \searrow 0} \left((\log \delta)^2 \inf_{M(a,d)} E_\epsilon - \frac{\pi \Gamma}{2} \log \frac{1}{\delta} \right) \leq \sum_{n=1}^N e(d_n) + W(a,d).$$

First step: glue energy minimising cores into the tail profile Define

$$\kappa_n^r = \frac{\lambda_n + \gamma_n I(r)}{\gamma_n \log \frac{1}{\delta}}.$$

Then we define certain profiles m_ϵ and approximate stray field potentials \tilde{u}_ϵ with the same formulas as on p. 477 of [18]. The functions $\mu_{a,d}^*$ and $u_{a,d}^*$ appearing in this construction are as in Section 3.5.1, while $\mu_{\epsilon/r}^n$ and $u_{\epsilon/r}^n$ are as in the other paper [18]. Furthermore, the function $u_\epsilon: \mathbb{R}_+^2 \rightarrow \mathbb{R}$ is the solution of (14), (15) for m_ϵ instead of m . In the rest of the proof, we estimate the energy of m_ϵ , showing that it provides the desired bounds.

Second step: estimate the magnetostatic energy in terms of \tilde{u}_ϵ The Laplacian $\Delta \tilde{u}_\epsilon$ satisfies the same formula as in our previous paper [18]. But in the subsequent estimates, the inequalities (72) and (73) of [18] have to be replaced by (57) and (58). This means that the additional factor $\log \frac{1}{r}$ appears in some of the estimates. This eventually leads to the conclusion that

$$\|\Delta \tilde{u}_\epsilon\|_{L^p(\mathbb{R}_+^2)} \leq \frac{Cr^{2/p-1} \log \frac{1}{r} + o(1)}{\log \frac{1}{\delta}} \quad (60)$$

for any fixed $p < \infty$. We can still use [18, Theorem 22] to conclude that

$$\left| \mu_\epsilon^n(x_1) - \cos \alpha + \frac{\gamma_n \log |x_1|}{\log \frac{1}{\delta}} \right| \leq \frac{o(1)}{\log \frac{1}{\delta}}$$

for any $x_1 \in [-\frac{3}{4}, -\frac{1}{2}] \cup [\frac{1}{2}, \frac{3}{4}]$. We now combine this estimate with (56) and Lemma 20. For $x_1 \in [a_n - \frac{3r}{4}, a_n - \frac{r}{2}] \cup [a_n + \frac{r}{2}, a_n + \frac{3r}{4}]$, this gives rise to:

$$\begin{aligned} & \left| (1 - \kappa_n^r) \mu_{\epsilon/r}^n \left(\frac{x_1 - a_n}{r} \right) + d_n \kappa_n^r - \cos \alpha - \frac{\mu_{a,d}^*(x_1)}{\log \frac{1}{\delta}} \right| \\ &= \left| (1 - \kappa_n^r) \left(\cos \alpha - \frac{\gamma_n \log \frac{|x_1 - a_n|}{r}}{\log \frac{1}{\delta}} \right) + d_n \kappa_n^r - \cos \alpha - \frac{\lambda_n + \gamma_n I(|x_1 - a_n|)}{\log \frac{1}{\delta}} \right| + \frac{Cr + o(1)}{\log \frac{1}{\delta}} \quad (61) \\ &= \left| \kappa_n^r \gamma_n - \frac{\gamma_n \log \frac{1}{r}}{\log \frac{1}{\delta}} - \frac{\lambda_n + \gamma_n I_0}{\log \frac{1}{\delta}} \right| + \frac{Cr + o(1)}{\log \frac{1}{\delta}} = \frac{Cr + o(1)}{\log \frac{1}{\delta}}. \end{aligned}$$

The next arguments are similar to our previous paper [18] again, but with two adaptations: first, we will need to multiply some of the terms in the estimates with $\log \frac{1}{r}$, because we use (57) and (58) instead of (72) and (73) in [18]. Second, when we restrict the functions u_ϵ and \tilde{u}_ϵ to a half-disk in \mathbb{R}_+^2 , we need to make sure that all the Néel walls are included.

Therefore, we fix $R \geq 1$ such that $a_1, \dots, a_N \in B_{R/2}(0)$. We now have the inequality

$$\|\Delta(u_\epsilon - \tilde{u}_\epsilon)\|_{L^p(\mathbb{R}_+^2)} \leq \frac{Cr^{2/p-1} \log \frac{1}{r} + o(1)}{\log \frac{1}{\delta}} \quad (62)$$

for an arbitrary fixed $p \in (1, 2)$, and still

$$\left\| \frac{\partial}{\partial x_2} (u_\epsilon - \tilde{u}_\epsilon) \right\|_{L^\infty(\mathbb{R})} \leq \frac{C + o(1)}{\log \frac{1}{\delta}}. \quad (63)$$

The support of $\Delta(u_\epsilon - \tilde{u}_\epsilon)$ is contained in $\bigcup_{n=1}^N B_r^+(a_n, 0)$ and the support of $\frac{\partial}{\partial x_2} (u_\epsilon - \tilde{u}_\epsilon)$ is contained in $\bigcup_{n=1}^N (a_n - r, a_n + r)$. Thus we conclude that

$$\|\Delta(u_\epsilon - \tilde{u}_\epsilon)\|_{\mathcal{M}(\mathbb{R}^2)} \leq \frac{Cr \log \frac{1}{r} + o(1)}{\log \frac{1}{\delta}},$$

where $\mathcal{M}(\mathbb{R}^2)$ is the space of Radon measures on \mathbb{R}^2 . Hence by well-known estimates for the Poisson equation with source term in $\mathcal{M}(\mathbb{R}^2)$, we find the estimate

$$\|\nabla(u_\epsilon - \tilde{u}_\epsilon)\|_{L^q(B_R^+(0))} \leq R^{2/q-1} \frac{C \log \frac{1}{r} + o(1)}{\log \frac{1}{\delta}}$$

for an arbitrary fixed $q \in [1, 2)$. By the arguments in [18], we have the inequality

$$\|\nabla \tilde{u}_\epsilon\|_{L^q(B_R^+(0))} \leq R^{2/q-1} \frac{C \log \frac{1}{r} + o(1)}{\log \frac{1}{\delta}}.$$

Hence

$$\|\nabla u_\epsilon\|_{L^q(B_R^+(0))} \leq R^{2/q-1} \frac{C \log \frac{1}{r} + o(1)}{\log \frac{1}{\delta}}. \quad (64)$$

Setting

$$\bar{u}_\epsilon = \int_{B_R^+(0)} u_\epsilon \, dx,$$

we now compute

$$\int_{\mathbb{R}_+^2} |\nabla u_\epsilon|^2 \, dx = \int_{\mathbb{R}_+^2} \nabla \tilde{u}_\epsilon \cdot \nabla u_\epsilon \, dx - \int_{(-R, R) \times \{0\}} (u_\epsilon - \bar{u}_\epsilon) \frac{\partial}{\partial x_2} (u_\epsilon - \tilde{u}_\epsilon) \, dx_1 + \int_{\mathbb{R}_+^2} (u_\epsilon - \bar{u}_\epsilon) \Delta \tilde{u}_\epsilon \, dx. \quad (65)$$

By the Hölder inequality, the trace theorem, and (64), we obtain

$$\begin{aligned} \|u_\epsilon - \bar{u}_\epsilon\|_{L^1(\bigcup_{n=1}^N (a_n - r, a_n + r))} &\leq Cr^{2-2/q} \|u_\epsilon - \bar{u}_\epsilon\|_{L^{q/(2-q)}(-R, R)} \\ &\leq Cr^{2-2/q} \|\nabla u_\epsilon\|_{L^q(B_R^+)} \leq Cr^{2-2/q} R^{2/q-1} \frac{C \log \frac{1}{r} + o(1)}{\log \frac{1}{\delta}}, \end{aligned}$$

and since we have inequality (63), we obtain

$$- \int_{(-R, R) \times \{0\}} (u_\epsilon - \bar{u}_\epsilon) \frac{\partial}{\partial x_2} (u_\epsilon - \bar{u}_\epsilon) dx_1 \leq R^{2/q-1} \frac{Cr^{2-2/q} \log \frac{1}{r} + o(1)}{(\log \delta)^2}.$$

Moreover, as the support of $\Delta \tilde{u}_\epsilon$ is included in $\bigcup_{n=1}^N B_r^+(a_n, 0) \subset B_R(0)$, the Hölder inequality and Sobolev embedding theorem, combined with (60) and (64), yield

$$\int_{\mathbb{R}_+^2} (u_\epsilon - \bar{u}_\epsilon) \Delta \tilde{u}_\epsilon dx \leq R^{2-2/p} \frac{Cr^{2/p-1} (\log r)^2 + o(1)}{(\log \delta)^2}.$$

For $p = \frac{4}{3}$ and $q = \frac{2p}{3p-2} = \frac{4}{3}$, identity (65) and the above inequalities imply that

$$\int_{\mathbb{R}_+^2} |\nabla u_\epsilon|^2 dx \leq \int_{\mathbb{R}_+^2} |\nabla \tilde{u}_\epsilon|^2 dx + \sqrt{R} \frac{C\sqrt{r}(\log r)^2 + o(1)}{(\log \delta)^2}.$$

Hence

$$E_\epsilon(m_\epsilon) \leq \frac{1}{2} \int_{-\infty}^{\infty} (\epsilon |m'_\epsilon|^2 + (m_{\epsilon 1} - \cos \alpha)^2) dx_1 + \frac{1}{2} \int_{\mathbb{R}_+^2} |\nabla \tilde{u}_\epsilon|^2 dx + \sqrt{R} \frac{C\sqrt{r}(\log r)^2 + o(1)}{(\log \delta)^2}. \quad (66)$$

Third step: estimate $\|\nabla \tilde{u}_\epsilon\|_{L^2(\mathbb{R}_+^2)}$ Identity [18, (91)] remains true here. The subsequent calculations in [18] still work, but instead of [18, (92)] we now obtain

$$\|\nabla \tilde{u}_\epsilon\|_{L^2(B_r^+(a_n, 0))}^2 \leq (1 - \kappa_n^r)^2 \|\nabla u_{\epsilon/r}^n\|_{L^2(B_1^+(0))}^2 + \frac{Cr \log \frac{1}{r} + o(1)}{(\log \delta)^2}.$$

By the above definition of κ_n^r , the same calculations as on p. 481 of [18] now yield the estimate

$$\begin{aligned} \int_{\mathbb{R}_+^2} |\nabla \tilde{u}_\epsilon|^2 dx &\leq \frac{1}{(\log \delta)^2} \int_{\Omega_r(a)} |\nabla u_{a,d}^*|^2 dx + \sum_{n=1}^N \int_{B_1^+(0)} |\nabla u_{\epsilon/r}^n|^2 dx \\ &\quad - \sum_{n=1}^N \frac{2\pi\gamma_n(\lambda_n + \gamma_n I(r))}{(\log \delta)^2} + \frac{Cr \log \frac{1}{r} + o(1)}{(\log \delta)^2}. \quad (67) \end{aligned}$$

Fourth step: estimate the exchange energy The calculations here are the same as in [18], but as inequality (73) there is replaced by (58) in the current article, the first inequality on p. 482 in [18] becomes

$$\left\| \frac{1 - \kappa_n^r}{r} \frac{d\mu_{\epsilon/r}^n}{dx_1} \left(\frac{x_1 - a_n}{r} \right) - \frac{\frac{d}{dx_1} \mu_{a,d}^*(x_1)}{\log \frac{1}{\delta}} \right\|_{L^2(T_n^r)} \leq \frac{C\sqrt{r} \log \frac{1}{r} + o(1)}{\log \frac{1}{\delta}}.$$

(Incidentally, the set T_n^r should be defined as $T_n^r = (a_n - \frac{3r}{4}, a_n - \frac{r}{2}) \cup (a_n + \frac{r}{2}, a_n + \frac{3r}{4})$, not as stated in [18].) Since the core profiles $\mu_{\epsilon/r}^n$ used here are exactly the same as in [18], we can use [18, Theorem 17] to estimate them. Furthermore, we have inequality (61). Thus we find that

$$\frac{\epsilon}{2} \int_{-\infty}^{\infty} |m'_\epsilon|^2 dx_1 = \frac{\epsilon}{2r} \sum_{n=1}^N \int_{-1}^1 \frac{\left(\frac{d}{dx_1} \mu_{\epsilon/r}^n \right)^2}{1 - (\mu_{\epsilon/r}^n)^2} dx_1 + \frac{o(1)}{(\log \delta)^2}. \quad (68)$$

Extra step: estimate the anisotropy energy For the problem studied here, we need to estimate the anisotropy energy as well, of course. Fortunately, this is rather straightforward. Owing to inequality (61), we can immediately conclude that

$$\int_{-\infty}^{\infty} (m_{\epsilon 1} - \cos \alpha)^2 dx_1 = \frac{1}{(\log \delta)^2} \int_{-\infty}^{\infty} (\mu_{a,d}^*)^2 dx_1 + \frac{Cr + o(1)}{(\log \delta)^2}. \quad (69)$$

Here we also use the fact that φ_n (in the definition of $m_{\epsilon 1}$ in [18]) is supported in the set $\bigcup_{n=1}^N T_n^r$ of measure $Nr/2$ and $\|\mu_{a,d}^*\|_{L^1(\mathbb{R})} \leq C$ by Lemma 20.

Combining (66), (67), (68), and (69), we obtain (using the notation of [18]):

$$\begin{aligned} E_{\epsilon}(m_{\epsilon}) &\leq \frac{1}{2(\log \delta)^2} \int_{\Omega_r(a)} |\nabla u_{a,d}^*|^2 dx + \sum_{n=1}^N \inf_{M|\gamma_n|} E_{\epsilon/r}^{|\gamma_n|} \\ &\quad - \sum_{n=1}^N \frac{\pi\gamma_n(\lambda_n + \gamma_n I(r))}{(\log \delta)^2} + \frac{1}{2(\log \delta)^2} \int_{-\infty}^{\infty} (\mu_{a,d}^*)^2 dx_1 + \sqrt{R} \frac{C\sqrt{r}(\log r)^2 + o(1)}{(\log \delta)^2}. \end{aligned}$$

Fifth step: estimate the core energy This is exactly the same as on p. 483 of [18].

Sixth step: combine the estimates It follows that

$$\begin{aligned} (\log \delta)^2 E_{\epsilon}(m_{\epsilon}) &\leq \frac{1}{2} \int_{\Omega_r(a)} |\nabla u_{a,d}^*|^2 dx + \frac{1}{2} \int_{-\infty}^{\infty} (\mu_{a,d}^*)^2 dx_1 \\ &\quad + \frac{\pi\Gamma}{2} \log \frac{1}{\delta} - \frac{\pi\Gamma}{2} \log \frac{1}{r} + \sum_{n=1}^N e(d_n) + W_2(a, d) - \pi I_0 \sum_{n=1}^N \gamma_n^2 \\ &\quad + C\sqrt{Rr}(\log r)^2 + o(1). \end{aligned}$$

Thus by (45) and (59),

$$(\log \delta)^2 E_{\epsilon}(m_{\epsilon}) \leq W(a, d) + \frac{\pi\Gamma}{2} \log \frac{1}{\delta} + \sum_{n=1}^N e(d_n) + C\sqrt{Rr}(\log r)^2 + o(1).$$

When we pass to the limit $r \rightarrow 0$, this proves the upper bound and completes the proof of Theorem 8.

3.6 Renormalised energy, separation and Γ -convergence

Now that the proof of Theorem 8 is complete, the theory for the unconfined model is at the same stage as it has been developed for the confined model in our previous paper [18]. As discussed in the introduction, the model also permits counterparts to the further results from Subsection 1.2. Fortunately, the proofs require few fundamentally new ideas, and therefore, we can keep this discussion relatively short.

We also prove Proposition 9 here. For this purpose, we analyse the renormalised energy in the unconfined case. First, we need the following result, which is similar to Lemma 11. For $a \in A_N$, recall the quantity $\rho(a)$, defined in (44), as it will appear in this statement.

Lemma 29. *Let $N \geq 2$ and $A_{k\ell} \in \mathbb{R}$ for $1 \leq k < \ell \leq N$, and let $f: A_N \rightarrow \mathbb{R}$ be defined by*

$$f(a) = - \sum_{1 \leq k < \ell \leq N} A_{k\ell} I(a_{\ell} - a_k), \quad a \in A_N.$$

1. If $\sum_{K \leq k < \ell \leq L} A_{k\ell} < 0$ for all $K, L \in \{1, \dots, N\}$ with $K < L$, then $\inf_{A_N} f > -\infty$ and $f(a) \rightarrow \infty$ as $\rho(a) \rightarrow 0$.

2. If there exists $K < L$ such that $\sum_{K \leq k < \ell \leq L} A_{k\ell} > 0$, then there exists a sequence $(a^{(i)})_{i \in \mathbb{N}}$ in A_N such that $\rho(a^{(i)}) \rightarrow 0$ and $f(a^{(i)}) \rightarrow -\infty$ as $i \rightarrow \infty$.

Proof. We follow the arguments in the proof of Lemma 11, pointing out only the differences (which are concerned with the possibility that $\rho(a)$ may be unbounded).

For statement 1, we proceed, as before, by induction. For $N = 2$, the statement follows from the properties of I proved in Lemma 20. In the induction step, we consider an arbitrary sequence $(a^{(i)})_{i \in \mathbb{N}}$ in A_N (not necessarily satisfying $\rho(a^{(i)}) \rightarrow 0$, because we also make a statement about the infimum of f over A_N) and we distinguish two cases.

Case A: $\limsup_{i \rightarrow \infty} (a_N^{(i)} - a_1^{(i)}) = \infty$. We partition $\{1, \dots, N\}$ into $\Lambda_1^{(i)}$ and $\Lambda_2^{(i)}$ with $\Lambda_1^{(i)} = \{1, \dots, n^{(i)}\}$, $\Lambda_2^{(i)} = \{n^{(i)} + 1, \dots, N\}$ such that $a_{n^{(i)}+1}^{(i)} - a_{n^{(i)}}^{(i)} \rightarrow \infty$ as $i \rightarrow \infty$. We may assume that the inequalities $2 \leq n^{(i)} \leq N - 2$ hold for all values of i or for none; if necessary, we pass to a subsequence with this property. In the first case, we conclude that $|\Lambda_1^{(i)}| \geq 2$ and $|\Lambda_2^{(i)}| \geq 2$. Otherwise, they will satisfy $|\Lambda_1^{(i)}| = 1$ or $|\Lambda_2^{(i)}| = 1$.

Case A.1: $|\Lambda_1^{(i)}|, |\Lambda_2^{(i)}| \geq 2$. Then

$$f(a^{(i)}) = - \sum_{1 \leq k < \ell \leq n^{(i)}} A_{k\ell} I(a_\ell^{(i)} - a_k^{(i)}) - \sum_{n^{(i)} < k < \ell \leq N} A_{k\ell} I(a_\ell^{(i)} - a_k^{(i)}) - \sum_{k \in \Lambda_1^{(i)}, \ell \in \Lambda_2^{(i)}} A_{k\ell} I(a_\ell^{(i)} - a_k^{(i)}).$$

Since $2 \leq |\Lambda_1^{(i)}|, |\Lambda_2^{(i)}| < N$, by induction, we know that the first two terms are uniformly bounded below. The last term converges to 0 by Lemma 20. Moreover, if $\rho(a^{(i)}) \rightarrow 0$, then at least one of $\Lambda_1^{(i)}$ or $\Lambda_2^{(i)}$ has points that collide as $i \rightarrow \infty$. Thus, by induction, $f(a^{(i)}) \rightarrow \infty$ as $i \rightarrow \infty$.

Case A.2: $|\Lambda_1^{(i)}| = 1$ or $|\Lambda_2^{(i)}| = 1$. Without loss of generality, we may assume that $|\Lambda_1^{(i)}| = 1$ and $N > |\Lambda_2^{(i)}| \geq 2$. Then

$$f(a^{(i)}) = - \sum_{2 \leq k < \ell \leq N} A_{k\ell} I(a_\ell^{(i)} - a_k^{(i)}) - \sum_{\ell=2}^N A_{k\ell} I(a_\ell^{(i)} - a_1^{(i)}).$$

The conclusion follows as in Case A.1.

Case B: $\limsup_{i \rightarrow \infty} (a_N^{(i)} - a_1^{(i)}) < \infty$. As f is translation invariant, we may assume that $a_1^{(i)} = 0$ for all $i \in \mathbb{N}$. Then we use the arguments in Lemma 11, since $I(t)$ behaves like $\log \frac{1}{t}$ as $t \rightarrow 0$.

For statement 2, we use the same arguments as in Lemma 11. \square

For the proof of Proposition 9, we require the following lemma.

Lemma 30. *The function $(0, \infty) \rightarrow (\frac{1}{8}, \frac{1}{2})$, $t \mapsto \frac{I'(2t)}{I'(t)}$, is well-defined and bijective.*

Proof. By (32), a change of variables implies

$$\begin{aligned} 2I'(2t) - I'(t) &= \int_0^\infty \frac{s^2}{s^2 + 1} (e^{-st} - 2e^{-2st}) ds = \int_0^\infty \left(\frac{s^2}{s^2 + 1} - \frac{s^2}{s^2 + 4} \right) e^{-st} ds > 0, \\ 8I'(2t) - I'(t) &= \int_0^\infty \frac{s^2}{s^2 + 1} (e^{-st} - 8e^{-2st}) ds = \int_0^\infty \left(\frac{s^2}{s^2 + 1} - \frac{4s^2}{s^2 + 4} \right) e^{-st} ds < 0. \end{aligned}$$

Since $I' < 0$ in $(0, \infty)$, then $\frac{1}{8} < \frac{I'(2t)}{I'(t)} < \frac{1}{2}$ in $(0, \infty)$. By (32), another change of variables gives

$$\begin{aligned} t^3 I'(t) &= -t^3 \int_0^\infty \frac{s^2}{s^2+1} e^{-st} ds = - \int_0^\infty \frac{t^2 s^2}{s^2+t^2} e^{-s} ds \rightarrow - \int_0^\infty s^2 e^{-s} ds = -2 \quad \text{as } t \rightarrow \infty, \\ t I'(t) &= -t \int_0^\infty \frac{s^2}{s^2+1} e^{-st} ds = - \int_0^\infty \frac{s^2}{s^2+t^2} e^{-s} ds \rightarrow - \int_0^\infty e^{-s} ds = -1 \quad \text{as } t \rightarrow 0, \end{aligned}$$

by the dominated convergence theorem. In particular, we have $\lim_{t \rightarrow 0} \frac{I'(2t)}{I'(t)} = \frac{1}{2}$ and $\lim_{t \rightarrow \infty} \frac{I'(2t)}{I'(t)} = \frac{1}{8}$. By the intermediate value theorem, the function $t \mapsto I'(2t)/I'(t)$ attains every value in $(\frac{1}{8}, \frac{1}{2})$.

In order to prove injectivity, we consider the function $f(t) = I'(2t) - qI'(t)$ for a fixed $q \in (\frac{1}{8}, \frac{1}{2})$. We also consider $t_0 > 0$ with $f(t_0) = 0$. By computations similar to the above, we may write

$$f(t_0) = \int_0^\infty g(s) e^{-st_0} ds,$$

where

$$g(s) = \frac{qs^2}{s^2+1} - \frac{s^2}{2(s^2+4)} = \frac{s^2}{2(s^2+1)(s^2+4)} ((8q-1) - (1-2q)s^2).$$

We note that $8q-1 > 0$ and $1-2q > 0$. Hence there exists $\sigma > 0$ such that $g(s) > 0$ for $s < \sigma$ and $g(s) < 0$ for $s > \sigma$. Therefore $(\sigma-s)g(s) > 0$ for all $s \neq \sigma$. Next we compute

$$f'(t_0) = - \int_0^\infty sg(s) e^{-st_0} ds > -\sigma \int_0^\infty g(s) e^{-st_0} ds = -\sigma f(t_0) = 0.$$

To summarise, if $f(t_0) = 0$, then $f'(t_0) > 0$. Of course it follows that f can have at most one zero. Thus for any $q \in (\frac{1}{8}, \frac{1}{2})$, there exists exactly one $t_0 > 0$ such that $I'(2t_0)/I'(t_0) = q$. \square

Proof of Proposition 9. The first two statements are consequences of Lemmas 12 and 29.

For statement 3, if $N = 3$, assume that $a \in A_3$ is a critical point of $W(\cdot, d)$. Then a satisfies two equations: $I'(a_2 - a_1) = I'(a_3 - a_2)$ and $I'(a_3 - a_1) = -\frac{\gamma_2}{\gamma_1} I'(a_3 - a_2)$. As I' is increasing (see Lemma 20), we deduce that $a_2 - a_1 = a_3 - a_2 =: t$ (i.e., the points a are equidistributed). Thus, the existence of a is equivalent to the existence of a solution of $I'(2t) = cI'(t)$ with $c = -\frac{\gamma_2}{\gamma_1}$.

Suppose that $d_1 = 1$. By Lemma 30, the equation $I'(2t) = cI'(t)$ has a solution if, and only if, $\frac{1}{8} < c < \frac{1}{2}$. Moreover, this solution is unique. Hence $W(\cdot, d)$ has a critical point exactly under these conditions, and the critical point is unique. Since $c = -\frac{\gamma_2}{\gamma_1} = \frac{1+\cos\alpha}{1-\cos\alpha}$, the above inequalities hold true if, and only if, $-\frac{7}{9} < \cos\alpha < -\frac{1}{3}$, which corresponds to the range of angles α given in Proposition 9. For $d_1 = -1$, the arguments are analogous. Moreover, these critical points are *not* minimisers. Indeed, $W(\cdot, d)$ does not have any minimiser, since statement 2 implies that $\inf_{A_3} W(\cdot, d) = -\infty$ for $d_1 \cos\alpha < -\frac{1}{3}$.

For statement 4, we first consider the case $N = 2$ and $\alpha \in (0, \pi)$. By Lemma 20, we know that $I' < 0$, so I does not have any critical point and neither does $W(\cdot, d)$. In the general case, we only consider the case where $d_1 \cos\alpha \geq 0$, as the arguments are the same in the other case. This is equivalent to $0 < |\gamma_1| \leq |\gamma_2|$.

Case A: N is odd. Then, given $a \in A_N$, we compute

$$\frac{1}{\pi\gamma_1} \frac{\partial W(\cdot, d)}{\partial a_1}(a) = (\gamma_2 I'(a_2 - a_1) + \gamma_1 I'(a_3 - a_1)) + \cdots + (\gamma_2 I'(a_{N-1} - a_1) + \gamma_1 I'(a_N - a_1)).$$

According to Lemma 20, we know that $I'(a_n - a_1) < I'(a_{n+1} - a_1) < 0$ for $2 \leq n \leq N-1$. The assumption $|\gamma_1| \leq |\gamma_2|$ then implies that either all terms of the form $\gamma_2 I'(a_n - a_1) + \gamma_1 I'(a_{n+1} - a_1)$ (for n even) give a positive contribution to the above sum, or all give a negative contribution. In particular, $\frac{\partial W(a, d)}{\partial a_1} \neq 0$ for every $a \in A_N$, and $W(\cdot, d)$ can have no critical points.

Case B: N is even. In this case, we have one additional term at the end of the above sum, which is $\gamma_2 I'(a_N - a_1)$. It has the same sign as the sum of the other terms. Thus, the conclusion holds. \square

We now turn to the separation and Γ -convergence result in the unconfined case (the counterparts of Theorem 4 and Corollary 5). Writing $m = (\cos \phi, \sin \phi) : \mathbb{R} \rightarrow \mathbb{S}^1$, we recall that we sometimes write $E_\epsilon(\phi)$ instead of $E_\epsilon(m)$. Note that a finite energy configuration $E_\epsilon(m)$ has the property that $m_1 - \cos \alpha \in H^1(\mathbb{R})$. In particular, $m_1 \rightarrow \cos \alpha$ as $x_1 \rightarrow \pm\infty$. With some abuse of notation, we express this fact as $m_1(\pm\infty) = \cos \alpha$. This implies that m_2 also has a limit at $\pm\infty$, which belongs to $\{\pm \sin \alpha\}$. Therefore, any continuous lifting ϕ of m has limits at $\pm\infty$, belonging to $2\pi\mathbb{Z} \pm \alpha$.

When $\epsilon \searrow 0$, we expect to find limit configurations given by piecewise constant functions $\phi_0 : \mathbb{R} \rightarrow 2\pi\mathbb{Z} \pm \alpha$ with finite total variation. A lot of the terminology from Subsection 1.2 is adapted to this situation in an obvious manner. We therefore use it here as well, including in particular the notation $\iota(\phi_0)$ and $\eta(\phi_0)$ for a function ϕ_0 as above.

Theorem 31. *The following holds true.*

1. **Compactness and Separation.** *Suppose that $(\phi_\epsilon)_{\epsilon>0}$ is a family of continuous functions in \mathbb{R} such that $\phi_\epsilon(-\infty) = \pm\alpha$ and $\phi_\epsilon(+\infty) \in 2\pi\mathbb{Z} \pm \alpha$ for all $\epsilon > 0$. Suppose that*

$$\limsup_{\epsilon \searrow 0} |\log \epsilon| E_\epsilon(\phi_\epsilon) < \infty.$$

Then there exist a sequence of numbers $\epsilon_k \searrow 0$ and a function $\phi_0 : \mathbb{R} \rightarrow 2\pi\mathbb{Z} \pm \alpha$ of finite total variation such that $\phi_{\epsilon_k} \rightarrow \phi_0$ in $L^2_{\text{loc}}(\mathbb{R})$. If $N = \iota(\phi_0)$ and $\theta_N < \alpha < \pi - \theta_N$, and if inequality (8) is satisfied, then ϕ_0 is simple.

2. **Lower bound.** *Let $(\phi_\epsilon)_{\epsilon>0}$ be a family of continuous functions in \mathbb{R} such that $\phi_\epsilon(-\infty) = \pm\alpha$, $\phi_\epsilon(+\infty) \in 2\pi\mathbb{Z} \pm \alpha$ for all $\epsilon > 0$ and $\phi_\epsilon \rightarrow \phi_0$ in $L^2_{\text{loc}}(\mathbb{R})$ for some simple, piecewise constant limit $\phi_0 : \mathbb{R} \rightarrow 2\pi\mathbb{Z} \pm \alpha$. Suppose that (a, d) is the transition profile of ϕ_0 and $N = \iota(\phi_0)$. Then*

$$E_\epsilon(\phi_\epsilon) \geq \frac{\eta(\phi_0)}{\log \frac{1}{\delta}} + \frac{\sum_{n=1}^N e(d_n) + W(a, d)}{(\log \frac{1}{\delta})^2} - o\left(\frac{1}{(\log \frac{1}{\delta})^2}\right).$$

3. **Upper bound.** *If $\phi_0 : \mathbb{R} \rightarrow 2\pi\mathbb{Z} \pm \alpha$ is simple with transition profile (a, d) , and if $N = \iota(\phi_0)$, then there exists a family $(\phi_\epsilon)_{\epsilon>0}$ of continuous functions in \mathbb{R} such that $\phi_\epsilon(-\infty) = \pm\alpha$, $\phi_\epsilon(+\infty) \in 2\pi\mathbb{Z} \pm \alpha$, $\phi_\epsilon \rightarrow \phi_0$ in $L^2(\mathbb{R})$ and*

$$E_\epsilon(\phi_\epsilon) \leq \frac{\eta(\phi_0)}{\log \frac{1}{\delta}} + \frac{\sum_{n=1}^N e(d_n) + W(a, d)}{(\log \frac{1}{\delta})^2} + o\left(\frac{1}{(\log \frac{1}{\delta})^2}\right).$$

The proof follows the same arguments as the proofs of Theorem 4 and Corollary 5. Since these arguments are based on Propositions 13 and 16, we need to discuss why they are, *mutatis mutandis*, still valid in the unconfined case.

- A version of Proposition 13 for the unconfined model can be proved with the same arguments. The proof is now based on Lemmas 12 and 29 and the fact that $I(|a_\ell - a_k|) \leq 1$ whenever $|a_\ell - a_k| \geq 1$.
- The proof of Proposition 16 relies mostly on theory from [18], the counterpart of which for the unconfined model has been developed above. The additional ingredient is the energy estimate of Lemma 14 (which is used in Lemma 15, which in turn feeds into Lemma 17 and finally Proposition 16). The inequality of Lemma 14, however, is easier in the unconfined case and follows immediately from Proposition 21.

There is one result in [18] that has not explicitly been adapted to the unconfined case in this paper, but is used in the proof of Proposition 16 (via Lemma 17) as well. This is [18, Lemma 13]. It is readily checked, however, that the arguments for this result carry over to the unconfined model.

Corollary 7, on the other hand, cannot be generalised directly to the unconfined model. This becomes evident when we compare statement 3 with the results of Proposition 9. Even if $\alpha \in \Theta$, the function $W(\cdot, d_N^\pm)$ does not have a minimiser in general (or perhaps it never does). Hence a result such as statement 3 in Corollary 7 is inconceivable in the unconfined model.

Of course, Theorem 31 still gives some information about topological Néel walls in the unconfined model, but this information will be less useful than in the confined case, because we have to expect that part of the topology will disappear at $\pm\infty$ in the limit.

References

- [1] R. Alicandro and M. Ponsiglione. Ginzburg-Landau functionals and renormalized energy: a revised Γ -convergence approach. *J. Funct. Anal.*, 266(8):4890–4907, 2014.
- [2] F. Bethuel, H. Brezis, and F. Hélein. *Ginzburg-Landau vortices*. Birkhäuser, Boston, 1994.
- [3] A. Braides and L. Truskinovsky. Asymptotic expansions by Γ -convergence. *Contin. Mech. Thermodyn.*, 20(1):21–62, 2008.
- [4] A. Capella, C. Melcher, and F. Otto. Wave-type dynamics in ferromagnetic thin films and the motion of Néel walls. *Nonlinearity*, 20(11):2519–2537, 2007.
- [5] M. Chermisi and C. B. Muratov. One-dimensional Néel walls under applied external fields. *Nonlinearity*, 26:2935–2950, 2013.
- [6] J. E. Colliander and R. L. Jerrard. Ginzburg-Landau vortices: weak stability and Schrödinger equation dynamics. *J. Anal. Math.*, 77:129–205, 1999.
- [7] A. DeSimone, H. Knüpfer, and F. Otto. 2-d stability of the Néel wall. *Calc. Var. Partial Differential Equations*, 27:233–253, 2006.
- [8] A. DeSimone, R. V. Kohn, S. Müller, and F. Otto. Magnetic microstructures—a paradigm of multiscale problems. In *ICIAM 99 (Edinburgh)*, pages 175–190. Oxford Univ. Press, Oxford, 2000.
- [9] A. DeSimone, R. V. Kohn, S. Müller, and F. Otto. Repulsive interaction of Néel walls, and the internal length scale of the cross-tie wall. *Multiscale Model. Simul.*, 1:57–104, 2003.
- [10] A. DeSimone, R.V. Kohn, S. Müller, and F. Otto. Recent analytical developments in micromagnetics. In *The Science of Hysteresis, Vol. 2*, pages 269–381. Elsevier Academic Press, 2005.
- [11] D. Gilbarg and N.S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [12] A. Hubert and R. Schäfer. *Magnetic domains*. Springer, Berlin–Heidelberg–New York, 1998.
- [13] R. Ignat. A Γ -convergence result for Néel walls in micromagnetics. *Calc. Var. Partial Differential Equations*, 36:285–316, 2009.

- [14] R. Ignat and R.L. Jerrard. Interaction energy between vortices of vector fields on Riemannian surfaces. *C. R. Math. Acad. Sci. Paris*, 355(5):515–521, 2017.
- [15] R. Ignat and R.L. Jerrard. Renormalized energy between vortices in some Ginzburg-Landau models on 2-dimensional Riemannian manifolds,. *arXiv:1910.02921*, 2019.
- [16] R. Ignat and H. Knüpfner. Vortex energy and 360° Néel walls in thin-film micromagnetics. *Comm. Pure Appl. Math.*, 63:1677–1724, 2010.
- [17] R. Ignat and M. Kurzke. Global Jacobian and Gamma-convergence in a two-dimensional Ginzburg-Landau model for boundary vortices. *arXiv:1910.06039*.
- [18] R. Ignat and R. Moser. Interaction energy of domain walls in a nonlocal Ginzburg-Landau type model from micromagnetics. *Arch. Ration. Mech. Anal.*, 221:419–485, 2016.
- [19] R. Ignat and R. Moser. Néel walls with prescribed winding number and how a nonlocal term can change the energy landscape. *J. Differential Equations*, 263:5846–5901, 2017.
- [20] R. Ignat and R. Moser. Energy minimisers of prescribed winding number in an \mathbb{S}^1 -valued nonlocal Allen-Cahn type model. *Adv. Math.*, 357:106819, 45, 2019.
- [21] R. Ignat and F. Otto. A compactness result in thin-film micromagnetics and the optimality of the Néel wall. *J. Eur. Math. Soc. (JEMS)*, 10:909–956, 2008.
- [22] R. Jerrard and D. Spirn. Refined Jacobian estimates for Ginzburg-Landau functionals. *Indiana Univ. Math. J.*, 56(1):135–186, 2007.
- [23] R.L. Jerrard. Lower bounds for generalized Ginzburg-Landau functionals. *SIAM J. Math. Anal.*, 30(4):721–746, 1999.
- [24] R.L. Jerrard and H.M. Soner. The Jacobian and the Ginzburg-Landau energy. *Calc. Var. Partial Differential Equations*, 14(2):151–191, 2002.
- [25] M. Kurzke. A nonlocal singular perturbation problem with periodic well potential. *ESAIM Control Optim. Calc. Var.*, 12(1):52–63, 2006.
- [26] J.C. Lagarias. Euler’s constant: Euler’s work and modern developments. *Bull. Amer. Math. Soc. (N.S.)*, 50:527–628, 2013.
- [27] E. H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001.
- [28] F.H. Lin. Vortex dynamics for the nonlinear wave equation. *Comm. Pure Appl. Math.*, 52(6):737–761, 1999.
- [29] R. G. Lund and C. B. Muratov. One-dimensional domain walls in thin ferromagnetic films with fourfold anisotropy. *Nonlinearity*, 29:1716–1734, 2016.
- [30] C. Melcher. The logarithmic tail of Néel walls. *Arch. Ration. Mech. Anal.*, 168:83–113, 2003.
- [31] C. Melcher. Logarithmic lower bounds for Néel walls. *Calc. Var. Partial Differential Equations*, 21:209–219, 2004.
- [32] C. B. Muratov and X. Yan. Uniqueness of one-dimensional Néel wall profiles. *Proc. A.*, 472:20150762, 15, 2016.
- [33] T. Rivière and S. Serfaty. Limiting domain wall energy for a problem related to micromagnetics. *Comm. Pure Appl. Math.*, 54(3):294–338, 2001.

- [34] E. Sandier. Lower bounds for the energy of unit vector fields and applications. *J. Funct. Anal.*, 152(2):379–403, 1998.
- [35] E. Sandier and S. Serfaty. Gamma-convergence of gradient flows with applications to Ginzburg-Landau. *Comm. Pure Appl. Math.*, 57(12):1627–1672, 2004.
- [36] E. Sandier and S. Serfaty. *Vortices in the magnetic Ginzburg-Landau model*, volume 70 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Boston, Inc., Boston, MA, 2007.
- [37] M. Struwe. On the asymptotic behavior of minimizers of the Ginzburg-Landau model in 2 dimensions. *Differential Integral Equations*, 7(5-6):1613–1624, 1994.