



Citation for published version:

Mccluskey, A, Squires, A, Coles, S & Morgan, B 2023, 'kinisi: Bayesian analysis of mass transport from molecular dynamics simulations', *The Journal of Open Source Software*.

Publication date:
2023

Document Version
Early version, also known as pre-print

[Link to publication](#)

Publisher Rights
CC BY

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

kinisi: Bayesian analysis of mass transport from molecular dynamics simulations

Andrew R. McCluskey^{1,2}, Alexander G. Squires^{3,4}, Samuel W. Coles^{5,4}, and Benjamin J. Morgan^{5,4}

¹ School of Chemistry, University of Bristol, Cantock's Close, Bristol, BS8 1TS, UK ² European Spallation Source ERIC, Ole Maaløes vej 3, 2200 København N, DK ³ Department of Chemistry, University College London, 20 Gordon Street, London WC1H 0AJ, UK ⁴ The Faraday Institution, Quad One, Harwell Science and Innovation Campus, Didcot, OX11 0RA, UK ⁵ Department of Chemistry, University of Bath, Claverton Down, Bath, BA2 7AY, UK

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

kinisi is a Python package for estimating transport coefficients—e.g., self-diffusion coefficients, D^* —and their corresponding uncertainties from molecular dynamics simulation data: it includes an implementation of the approximate Bayesian regression scheme described in McCluskey et al. (2023), wherein the mean-squared displacement (MSD) of mobile atoms is modelled as a multivariate normal distribution that is parametrised from the input simulation data. kinisi uses Markov-chain Monte Carlo (Foreman-Mackey et al., 2019; Goodman & Weare, 2010) to sample this model multivariate normal distribution to give a posterior distribution of linear model ensemble MSDs that are compatible with the observed simulation data. For each linear ensemble MSD, $x(t)$, a corresponding estimate of the diffusion coefficient, \hat{D}^* is given via the Einstein relation,

$$\hat{D}^* = \frac{1}{6} \frac{dx(t)}{dt},$$

where t is time. The posterior distribution of compatible model ensemble MSDs calculated by kinisi gives a point estimate for the most probable value of D^* , given the observed simulation data, and an estimate of the corresponding uncertainty in \hat{D}^* . kinisi also provides equivalent functionality for estimating collective transport coefficients, i.e., jump-diffusion coefficients and ionic conductivities.

Statement of Need

Molecular dynamics simulations are widely used to calculate transport coefficients such as self-diffusion coefficients and ionic conductivities (Klepeis et al., 2009; B. J. Morgan, 2021; Benjamin J. Morgan & Madden, 2014; Poletayev et al., 2022; Sendner et al., 2009; Shimizu et al., 2015; Wang & Hou, 2011; Zelovich et al., 2019). Because molecular dynamics simulations are limited in size and timescale, ensemble parameters, such as transport coefficients, that are calculated from simulation trajectories are estimates of the corresponding true (unknown) parameter of interest and suffer from statistical uncertainty. The statistical properties of any calculated ensemble parameters depend on the details of the input molecular dynamics simulation—e.g., the choice of interaction potential, system size, and simulation timescale—and the choice of estimator for the target parameter to be calculated. An optimal estimation method should minimise the statistical uncertainty in the derived parameter of interest—the method should be statistically efficient—and should provide an accurate estimate of this uncertainty, so that calculated values can be used in downstream statistical analyses.

40 One widely-used approach to estimating the self-diffusion coefficient, D^* , from molecular
41 dynamics simulation is to fit a linear model to the observed mean-square displacement, xt (Allen
42 & Tildesley, 2017), where the slope of this “best fit” linear relationship gives a point-estimate
43 for D^* via the corresponding Einstein relation. The simplest approach to fitting a linear
44 model to observed MSD data is ordinary least squares (OLS). OLS, however, is statistically
45 inefficient and gives a large uncertainty in the resulting estimate of D^* , while also significantly
46 underestimating this uncertainty (McCluskey et al., 2023). `kinisi` implements the alternative
47 approximate Bayesian regression scheme described in McCluskey et al. (2023), which gives a
48 statistically efficient estimate for D^* and an accurate estimate for the associated uncertainty
49 $\sigma^2[\widehat{D}^*]$. This approach gives more accurate estimates of D^* from a given size of simulation
50 data (number of atoms and simulation timescale) than ordinary least-squares or weighted
51 least-squares, while the calculated uncertainties allow robust downstream analysis, such as
52 estimating activation energies by fitting an Arrhenius model to $D^*(T)$.

53 `kinisi` supports simulation output from a variety of common simulation software packages,
54 including VASP (Kresse & Furthmüller, 1996a, 1996b; Kresse & Hafner, 1993, 1994) and
55 those compatible with Pymatgen (Ong et al., 2013), atomic simulation environment (ASE)
56 (Larsen et al., 2017), and MDAnalysis (Gowers et al., 2016; Michaud-Agrawal et al., 2011).
57 Tutorials and API-level documentation are provided online at kinisi.rtdf.io.

58 Full details of the approximate Bayesian regression method implemented in `kinisi` are provided
59 in McCluskey et al. (2023). A list of publications where `kinisi` has been used in the analysis
60 of simulation data is available at kinisi.readthedocs.io/en/latest/papers.html.

61 Acknowledgements

62 The authors thank all of the users of `kinisi` for contributing feedback, suggesting new features
63 and filing bug reports. S.W.C., A.G.S. and B.J.M. acknowledge the support of the Faraday
64 Institution (grant numbers FIRG016, FIG017). B.J.M. acknowledges support from the Royal
65 Society (UF130329 and URF\R\191006).

66 Allen, M. P., & Tildesley, D. K. (2017). *Computer simulation of liquids* (2nd ed.). Oxford
67 University Press.

68 Foreman-Mackey, D., Farr, W. M., Sinha, M., Archibald, A. M., Hogg, D. W., Sanders, J. S.,
69 Zuntz, J., Williams, P. K. G., Nelson, A. R. J., de Val-Borro, M., Erhardt, T., Pashchenko,
70 I., & Pla, O. A. (2019). Emcee v3: A python ensemble sampling toolkit for affine-invariant
71 MCMC. *J. Open Source Softw.*, 4(43), 1864. <https://doi.org/10.21105/joss.01864>

72 Goodman, J., & Weare, J. (2010). Ensemble samplers with affine invariance. *Communications*
73 *in Applied Mathematics and Computational Science*, 5(1), 65–80. <https://doi.org/10.2140/camcos.2010.5.65>

74
75 Gowers, R. J., Linke, M., Barnoud, J., Reddy, T. J. E., Melo, M. N., Seyler, S. L., Dotson, D.
76 L., Domanski, J., Buchoux, S., Kenney, I. M., & Beckstein, O. (2016). MDAnalysis: A
77 python package for the rapid analysis of molecular dynamics simulations. In S. Benthall &
78 S. Rostrup (Eds.), *Proceedings of the 15th python in science conference* (pp. 98–105).
79 SciPy. <https://doi.org/10.25080/majora-629e541a-00e>

80 Klepeis, J. L., Lindorff-Larsen, K., Dror, R. O., & Shaw, D. E. (2009). Long-timescale molecular
81 dynamics simulations of protein structure and function. *Curr. Op. Struct. Biol.*, 19(2),
82 120–127. <https://doi.org/10.1016/j.sbi.2009.03.004>

83 Kresse, G., & Furthmüller, J. (1996a). Efficiency of ab-initio total energy calculations for
84 metals and semiconductors using a plane-wave basis set. *Comput. Mat. Sci.*, 6(1), 15–50.
85 [https://doi.org/10.1016/0927-0256\(96\)00008-0](https://doi.org/10.1016/0927-0256(96)00008-0)

- 86 Kresse, G., & Furthmüller, J. (1996b). Efficient iterative schemes for ab initio total-energy
87 calculations using a plane-wave basis set. *Phys. Rev. B*, *54*(16), 11169. [https://doi.org/](https://doi.org/10.1103/PhysRevB.54.11169)
88 [10.1103/PhysRevB.54.11169](https://doi.org/10.1103/PhysRevB.54.11169)
- 89 Kresse, G., & Hafner, J. (1993). Ab initio molecular dynamics for liquid metals. *Phys. Rev. B*,
90 *47*(1), 558(R). <https://doi.org/10.1103/PhysRevB.47.558>
- 91 Kresse, G., & Hafner, J. (1994). Ab initio molecular-dynamics simulation of the liquid-
92 metal-amorphous-semiconductor transition in germanium. *Phys. Rev. B*, *49*(20), 14251.
93 <https://doi.org/10.1103/PhysRevB.49.14251>
- 94 Larsen, A. H., Mortensen, J. J., Blomqvist, J., Castelli, I. E., Christensen, R., Duřak, M.,
95 Friis, J., Groves, M. N., Hammer, B., Hargus, C., Hermes, E. D., Jennings, P. C.,
96 Jensen, P. B., Kermode, J., Kitchin, J. R., Kolsbjerg, E. L., Kubal, J., Kaasbjerg, K.,
97 Lysgaard, S., ... Jacobsen, K. W. (2017). The atomic simulation environment—a python
98 library for working with atoms. *J. Phys.: Condens. Matter*, *29*(27), 273002. [https://doi.org/](https://doi.org/10.1088/1361-648x/aa680e)
99 [10.1088/1361-648x/aa680e](https://doi.org/10.1088/1361-648x/aa680e)
- 100 McCluskey, A. R., Coles, S. W., & Morgan, B. J. (2023). *Accurate estimation of diffusion*
101 *coefficients and their uncertainties from computer simulation*. [https://arxiv.org/abs/2305.](https://arxiv.org/abs/2305.18244)
102 [18244](https://arxiv.org/abs/2305.18244)
- 103 Michaud-Agrawal, N., Denning, E. J., Woolf, T. B., & Beckstein, O. (2011). MDAAnalysis:
104 A toolkit for the analysis of molecular dynamics simulations. *J. Comput. Chem.*, *32*(10),
105 2319–2327. <https://doi.org/10.1002/jcc.21787>
- 106 Morgan, B. J. (2021). Mechanistic origin of superionic lithium diffusion in anion-disordered
107 $\text{Li}_6\text{PS}_5\text{X}$ argyrodites. *Chem. Mater.*, *33*(6), 2004–2018. [https://doi.org/10.1021/acs.](https://doi.org/10.1021/acs.chemmater.0c03738)
108 [chemmater.0c03738](https://doi.org/10.1021/acs.chemmater.0c03738)
- 109 Morgan, Benjamin J., & Madden, P. A. (2014). Relationships between atomic diffusion
110 mechanisms and ensemble transport coefficients in crystalline polymorphs. *Phys. Rev.*
111 *Lett.*, *112*(14). <https://doi.org/10.1103/physrevlett.112.145901>
- 112 Ong, S. P., Richards, W. D., Jain, A., Hautier, G., Kocher, M., Cholia, S., Gunter, D., Chevrier,
113 V., Persson, K. A., & Ceder, G. (2013). Python materials genomics (pymatgen) : A
114 robust, open-source python library for materials analysis. *Comput. Mat. Sci.*, *68*, 314–319.
115 <https://doi.org/10.1016/j.commatsci.2012.10.028>
- 116 Poletayev, A. D., Dawson, J. A., Islam, M. S., & Lindenberg, A. M. (2022). Defect-driven
117 anomalous transport in fast-ion conducting solid electrolytes. *Nat. Mater.*, *21*, 1066–1073.
118 <https://doi.org/10.1038/s41563-022-01316-z>
- 119 Sendner, C., Horinek, D., Bocquet, L., & Netz, R. R. (2009). Interfacial water at hydrophobic
120 and hydrophilic surfaces: Slip, viscosity, and diffusion. *Langmuir*, *25*(18), 10768–10781.
121 <https://doi.org/10.1021/la901314b>
- 122 Shimizu, K., Freitas, A. A., Atkin, R., Warr, G. G., FitzGerald, P. A., Doi, H., Saito, S., Ueno,
123 K., Umabayashi, Y., Watanabe, M., & Lopes, J. N. C. (2015). Structural and aggregate
124 analyses of (Li salt + glyme) mixtures: The complex nature of solvate ionic liquids. *Phys.*
125 *Chem. Chem. Phys.*, *17*(34), 22321–22335. <https://doi.org/10.1039/c5cp03414k>
- 126 Wang, J., & Hou, T. (2011). Application of molecular dynamics simulations in molecular
127 property prediction II: Diffusion coefficient. *J. Comput. Chem.*, *32*(16), 3505–3519.
128 <https://doi.org/10.1002/jcc.21939>
- 129 Zelovich, T., Vogt-Maranto, L., Hickner, M. A., Paddison, S. J., Bae, C., Dekel, D. R., &
130 Tuckerman, M. E. (2019). Hydroxide Ion Diffusion in Anion-Exchange Membranes at Low
131 Hydration: Insights from Ab Initio Molecular Dynamics. *Chem. Mater.*, *31*(15), 5778–5787.
132 <https://doi.org/10.1021/acs.chemmater.9b01824>