



Citation for published version:

Barker, T, Prange, C & Tan, J 2024, 'On Symmetry Breaking for the Navier–Stokes Equations', *Communications in Mathematical Physics*, vol. 405, no. 2, 25. <https://doi.org/10.1007/s00220-023-04897-1>

DOI:

[10.1007/s00220-023-04897-1](https://doi.org/10.1007/s00220-023-04897-1)

Publication date:

2024

Document Version

Peer reviewed version

[Link to publication](#)

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: <http://dx.doi.org/10.1007/s00220-023-04897-1>

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Citation for published version:

Barker, T, Prange, C & Tan, J 2024 'On symmetry breaking for the Navier-Stokes equations' Communications in Mathematical Physics. <<https://arxiv.org/abs/2302.02836>>

Publication date:
2024

Document Version
Peer reviewed version

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ON SYMMETRY BREAKING FOR THE NAVIER-STOKES EQUATIONS

TOBIAS BARKER, CHRISTOPHE PRANGE, AND JIN TAN

ABSTRACT. Inspired by an open question by Chemin, Zhang and Zhang about the regularity of the 3D Navier-Stokes equations with one initially small component, we investigate symmetry breaking and symmetry preservation. Our results fall in three classes. First we prove strong symmetry breaking. Specifically, we demonstrate third component norm inflation (3rdNI) and Isotropic Norm Inflation (INI) starting from zero third component. Second we prove symmetry breaking for initially zero third component, even in the presence of a favorable initial pressure gradient. Third we study certain symmetry preserving solutions with a shear flow structure. Specifically, we give applications to the inviscid limit and exhibit explicit solutions that inviscidly damp to the Kolmogorov flow.

1. INTRODUCTION

Symmetries preserved by evolution play an important role in the mathematical theory of the Navier-Stokes equations and Euler equations:

$$\partial_t u^\nu + u^\nu \cdot \nabla u^\nu + \nabla P = \nu \Delta u^\nu \quad \text{in } \mathbb{R}_+ \times \mathbb{T}^3, \quad \operatorname{div} u^\nu = 0, \quad \nu \geq 0. \quad (1.1)$$

On the one hand, certain preserved symmetries lead to the preservation of certain structures that grant smoothness of solutions [33], [41]. On the other hand, preserved symmetries reduce the number of degrees of freedom of the Navier-Stokes and Euler equations, which can make it possible to prove or numerically investigate the existence of singularities [22], [14], [26, 27].

In this vein, in recent years there has been a substantial amount of activity aimed at showing that additional assumptions of one component of the velocity field (solving the Navier-Stokes equations) imply that the solution is regular. On the other side of coin, this corresponds to showing that solutions of the Navier-Stokes equations that become singular must do so in an isotropic manner. Research in this direction was initiated in the seminal paper of Neustupa and Penel [37]. Since then there have been many contributions to one-component regularity for the Navier-Stokes equations, with recent contributions showing regularity provided that one-component of the velocity field has a finite norm either almost preserved [10, 11] or preserved¹ with respect to the Navier-Stokes scaling symmetry [12, 13, 43].

The purpose of this paper is to understand the dynamics of the Navier-Stokes equations when one-component of the initial data is zero. Throughout we will set the third component of the initial data to be zero, without loss of generality. Our main motivation is an open question raised by Chemin, Zhang and Zhang [13] when

Date: November 22, 2023.

2010 Mathematics Subject Classification. 35Q30, 35Q31, 35N20, 76F10.

Key words and phrases. Navier-Stokes equations; Euler equations; global solutions; shear flows; ill-posedness; norm inflation; inviscid limit; inviscid damping.

¹Currently one component regularity criteria in terms of norms preserved with respect to the Navier-Stokes scaling symmetry, involve spatial norms with some differentiability or Lorentz time norms. It remains a long standing open problem if a solution to the Navier-Stokes equations u , with third component $u_3 \in L^q(0, T; L^p(\mathbb{R}^3))$ ($\frac{3}{p} + \frac{2}{q} = 1$, $p \in [3, \infty)$) is smooth on $(0, T) \times \mathbb{R}^3$.

discussing endpoint one-component regularity criteria. Specially, in [13, page 873], Chemin, Zhang and Zhang formulate the following open question:

(Q) [If] for some unit vector e of \mathbb{R}^3 , [the component of the initial data] $\|u_{\text{in}} \cdot e\|_{H^{\frac{1}{2}}}$ is small with respect to some universal constant, is it implied that there is no blow up for the Fujita-Kato solution of (NS)?

1.1. Main results of the paper. In relation to the aforementioned open problem **(Q)**, our first two results show that initial data with zero third component can exhibit third component norm inflation (3rdNI, Theorem A) and Isotropic Norm Inflation (INI, Theorem A') with respect to critical norms specified in [7].

Theorem A (strong symmetry breaking). *For any $0 < \delta < 1$, there exists mean-free $C^\infty(\mathbb{T}^3)$ solenoidal initial data u_{in} and \bar{u}_{in}^2 with vanishing third component,*

$$\|u_{\text{in}} - \bar{u}_{\text{in}}\|_{\dot{B}_{\infty,\infty}^{-1}} = \|u_{\text{in}}^h - \bar{u}_{\text{in}}^h\|_{\dot{B}_{\infty,\infty}^{-1}} < \delta,$$

and such that the following holds true.

There exists a unique solution u (resp. \bar{u}) of the Cauchy problem (1.1) subject to initial data u_{in} (resp. \bar{u}_{in}) belonging to $C^\infty((0, T] \times \mathbb{T}^3)$ for some time $0 < T < \delta$ with $\bar{u}^3 \equiv 0$ on $[0, T] \times \mathbb{R}^3$ and

$$\|u^3(T, \cdot) - \bar{u}^3(T, \cdot)\|_{\dot{B}_{\infty,\infty}^{-1}} = \|u^3(T, \cdot)\|_{\dot{B}_{\infty,\infty}^{-1}} > \frac{1}{\delta}.$$

Moreover,

$$\min(\|u^1(T, \cdot)\|_{L^3}, \|u^2(T, \cdot)\|_{L^3}, \|u^3(T, \cdot)\|_{L^3}) > \frac{1}{\delta}. \quad (1.2)$$

Theorem A does not provide negative evidence towards **(Q)**, but demonstrates that regularity in that case can only be granted by a yet to be discovered mechanism unrelated to the preservation of smallness of the third component of the corresponding solution. However, the solution in Theorem A remains small in $\dot{B}_{\infty,\infty}^{-1}$ at T in certain directions (see the discussion in subsection 1.3). Thus, the construction in Theorem A does not rule out the possibility that solutions, with initial third component equal to zero, remain small along some time-varying direction. Such a possibility is in fact ruled out by our second result below.

Theorem A' (strong isotropic symmetry breaking). *For any $0 < \delta < 1$, there exists mean-free $C^\infty(\mathbb{T}^3)$ solenoidal initial data u_{in}^3 and \bar{u}_{in} with vanishing third component,*

$$\|u_{\text{in}} - \bar{u}_{\text{in}}\|_{\dot{B}_{\infty,\infty}^{-1}} = \|u_{\text{in}}^h - \bar{u}_{\text{in}}^h\|_{\dot{B}_{\infty,\infty}^{-1}} < \delta,$$

and such that the following holds true.

There exists a unique solution u of the Cauchy problem (1.1) subject to initial data u_{in} belonging to $C^\infty((0, T] \times \mathbb{T}^3)$ for some time $0 < T < \delta$ and such that

$$\inf_{\mathbf{e} \in \mathbb{R}^3: |\mathbf{e}|=1} \|u(T, \cdot) \cdot \mathbf{e}\|_{\dot{B}_{\infty,\infty}^{-1}} > \frac{1}{\delta}. \quad (1.3)$$

We dub the norm inflation in all directions in (1.3) ‘Isotropic Norm Inflation’ (INI).

Now define the initial pressure P_{in} associated to a given solenoidal initial data u_{in} , which satisfies

$$-\Delta P_{\text{in}} := \nabla u_{\text{in}} : (\nabla u_{\text{in}})^T. \quad (1.4)$$

²This data has the structure given by (1.8). Our result shows that the solution map is not continuous at \bar{u}_{in} in the critical space $\dot{B}_{\infty,\infty}^{-1}$.

³This data has the structure given by (2.21).

Note that the initial data used to prove Theorems A and A', which will heuristically be described in subsections 1.2-1.3, both necessarily generate an initial pressure P_{in} that satisfies $\partial_3 P_{\text{in}} \neq 0$. From the equation for the third component of the associated solution (1.1), it is qualitatively clear that such an initial pressure will always produce a solution that breaks the symmetry of the third component zero. In this regard, we call pressure of this type *unfavorable*.⁴ Notice that there are other examples of plane-wave initial data that demonstrate symmetry breaking. We refer for instance to Figure 1 that shows breaking for the Taylor-Green vortex

$$u_{\text{in}}(x_1, x_2, x_3) = (\sin x_1 \cos x_2 \cos x_3, -\cos x_1 \sin x_2 \cos x_3, 0).$$

Notice that

$$\Delta P_{\text{in}} = 2(\cos x_3)^2((\cos x_1 \cos x_2)^2 - (\sin x_1 \sin x_2)^2)$$

so that the pressure for the Taylor-Green vortex is also unfavorable.




FIGURE 1. Taylor-Green vortex solution of Navier-Stokes with viscosity $\nu = 10^{-1}$. From top to bottom: $\|u^1(\cdot, t)\|_{L^1(\mathbb{T}^3)}$, $\|u^2(\cdot, t)\|_{L^1(\mathbb{T}^3)}$ and $\|u^3(\cdot, t)\|_{L^1(\mathbb{T}^3)}$. This simulation shows breaking with initial pressure unfavorable (see Footnote 4 for a definition) to symmetry preservation. Choice of parameters: total time $T = 10$ and time step $dt = 10^{-2}$; spectral code by Mikael Mortensen taken from <https://github.com/spectralDNS/spectralDNS> with $(2^5)^3$ mesh points.

⁴The terminology *unfavorable* refers here to the fact that the pressure is unfavorable to symmetry preservation.

As a consequence of the above, for initial data u_{in} with zero third component, we say that an initial pressure P_{in} is *favorable*⁵ if

$$\operatorname{div}_{\text{h}} u_{\text{in}}^{\text{h}} = 0 \quad \text{with} \quad \partial_3 P_{\text{in}} = 0, \quad \text{where} \quad -\Delta P_{\text{in}} := \nabla_{\text{h}} u_{\text{in}}^{\text{h}} : (\nabla_{\text{h}} u_{\text{in}}^{\text{h}})^{\text{T}}. \quad (1.5)$$

For a *favorable* initial pressure, the equation for the third component of (1.1) does not immediately imply that the third component of the solution breaks symmetry and becomes non-zero. In the Theorem below we are able to demonstrate an initial data, which has zero third component and favorable initial pressure, yet the corresponding solution breaks the symmetry and has non-zero third component on some time interval.

Theorem B (symmetry breaking despite favorable pressure gradient). *We consider the initial data*

$$u_{\text{in}} = \left(\cos x_2 \frac{N}{N + \sin x_3}, \cos x_1 \frac{N + \sin x_3}{N}, 0 \right) \quad (1.6)$$

which has favorable initial pressure gradient in the sense that $\partial_3 P_{\text{in}} = 0$, see (1.5). Then there exists a positive constant N_0 such that for any $N > N_0$, the initial data given by (1.6) generates a unique solution u to the Navier-Stokes equations (1.1) on $[0, 1] \times \mathbb{T}^3$ that satisfies

$$\|u^3(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \sim \frac{t^2}{N^2}, \quad (1.7)$$

for any $0 \leq t \leq \frac{1}{N^2} \ll 1$.

Remark 1 (comparison to other symmetry breaking results in the context of non-uniqueness). The non-uniqueness numerical results of Guillod and Šverák [24] concern Leray-Hopf solutions of the Navier-Stokes equations that break a symmetry class. In the context of the Euler equations and convex integration, symmetry breaking and restoration mechanisms were explicitly investigated in [2]. The non-uniqueness results for dissipative solutions of Euler by Scheffer [38], Shnirelman [40], De Lellis and Székelyhidi [21], Isett [29] and for weak solutions of Navier-Stokes by Buckmaster and Vicol [8] can also be seen as symmetry breaking results. Our results are in a different vein though. We show breaking of symmetry on some time interval where the solution is unique and smooth.

Remark 2 (comparison to other symmetry breaking results in the context of singularity formation). Let us mention that symmetry breaking of the third component plays a role in Hou's recent numerical investigation of potentially singular axisymmetric Navier-Stokes solutions [27]. There the initial data has third component zero, yet it is subsequently observed in the numerics that 'near the symmetry axis, the upward axial velocity dominates the angular and radial velocities' [27, page 2].

Remark 3 (isotropic motion with initial third component zero). For the construction in Theorem A', one can also show that for all $\mathbf{e} \in \mathbb{R}^3$ with $|\mathbf{e}| = 1$:

$$\|u(T, \cdot) \cdot \mathbf{e}\|_{\dot{B}_{\infty, \infty}^{-1}} \sim \frac{1}{\delta}.$$

Thus at time T , the velocity field is comparable in all unit directions with respect to the $\dot{B}_{\infty, \infty}^{-1}$ norm, despite evolving from initial data with zero third component.

⁵The terminology *favorable* refers here to the fact that the pressure is favorable to symmetry preservation. We demonstrate, see Theorem B that favorable pressure is still not enough to preserve the vanishing of the third component of the velocity, if it is zero initially.




FIGURE 2. Simulation of the Navier-Stokes equations with viscosity $\nu = 10^{-1}$ and initial data from Theorem B taking $N = 1.1$. From top to bottom: $\|u^1(\cdot, t)\|_{L^1(\mathbb{T}^3)}$, $\|u^2(\cdot, t)\|_{L^1(\mathbb{T}^3)}$ and $\|u^3(\cdot, t)\|_{L^1(\mathbb{T}^3)}$. This simulation shows breaking despite initial pressure favorable (see Footnote 5 for a definition) to symmetry preservation. Choice of parameters: total time $T = 10$ and time step $dt = 10^{-2}$; spectral code by Mikael Mortensen taken from <https://github.com/spectralDNS/spectralDNS> with $(2^5)^3$ mesh points.

Further results. Our interest here is on flows that preserve the symmetry $u^3 = 0$ and on applications of such flows.

There are indeed flows solving Navier-Stokes and Euler that have a shear-flow structure and keep the third component identically zero, as for instance the plane parallel channel flows $(u^1(x_3, t), u^2(x_1, x_3, t), 0)$ introduced by Wang [44]. Rotating these flows gives rise to a whole family of symmetry preserving pressureless shear flows that we dub ‘2.75D shear flows’, which are defined as follows. Let $\lambda \in \mathbb{Z}$ be a constant. Consider the initial data

$$u_{\text{in}} = (f(\lambda x_1 + x_2, x_3), -\lambda f(\lambda x_1 + x_2, x_3) - g(x_3), 0), \quad x \in \mathbb{T}^3. \quad (1.8)$$

In fact, in the case that u_{in} merely belongs to $L^2(\mathbb{T}^3)$ (that corresponds to f or g being rough), one can obtain a Leray-Hopf weak solution to the problem (1.1) with $\nu > 0$, and initial data u_{in} , given by

$$u^\nu = (F_\nu(t, \lambda x_1 + x_2, x_3), -\lambda F_\nu(t, \lambda x_1 + x_2, x_3) - e^{\nu t \partial_3^2} g(x_3), 0) \quad (1.9)$$

where $F_\nu : \mathbb{R}_+ \times \mathbb{T}^2 \rightarrow \mathbb{R}$ is the unique global-in-time solution to

$$\begin{cases} \partial_t F_\nu - e^{\nu t \partial_2^2} g(y_2) \partial_1 F_\nu = \nu((\lambda^2 + 1)\partial_1^2 + \partial_2^2) F_\nu & \text{in } \mathbb{T}^2 \times \mathbb{R}_+ \\ F_\nu(0, y_1, y_2) = f(y_1, y_2). \end{cases} \quad (1.10)$$

For more insights about the derivation of these flows, we refer to Appendix A.

Remark 4 (2.75D shear flows for Euler). One also has ‘2.75D shear flows’ that solve the Euler equations in a distributional sense:

$$u^E(t, x) = (f(\lambda x_1 + x_2 + tg(x_3), x_3), -\lambda f(\lambda x_1 + x_2 + tg(x_3), x_3) - g(x_3), 0), \quad (1.11)$$

where $f \in L^2(\mathbb{T}^3)$ and $g \in C^\infty(\mathbb{T}^2)$.

Inviscid damping. In Section 4.1 we show that the 2.75D shear flows for Euler, see Remark 4, inviscidly damp to the *Kolmogorov flow*. The Kolmogorov flow $u^K = (0, \sin x_3, 0)$ is a stationary solution of the 3D Euler equations in \mathbb{T}^3 . In [20] Coti-Zelati, Elgindi and Widmayer exhibit 2D stationary solutions to the Euler equations near u^K , thus demonstrating a lack of inviscid damping near u^K . On the other hand, 2.75D shear Euler flows (1.11) can be used to produce explicit solutions that inviscidly damp⁶ to u^K for large times. This and [20] show that dynamics near the Kolmogorov flow in \mathbb{T}^3 are rich and no generic behavior can be expected.

Vanishing viscosity. In Section 4.2 we investigate the vanishing viscosity limit for 2.75D shear flow solutions of Navier-Stokes that are Onsager supercritical. Turbulence theory from [30, 31, 32] predicts that if u^ν is a weak Leray-Hopf solution in $\mathbb{T}^3 \times (0, \infty)$, with viscosity ν and initial data u_{in} , then generically one has *anomalous dissipation*:

$$\liminf_{\nu \rightarrow 0} \nu \int_0^T \int_{\mathbb{T}^3} |\nabla u^\nu|^2 dx ds > 0. \quad (1.12)$$

It is known that if (1.12) and the vanishing viscosity limit holds in suitable topology, then the corresponding Euler flow u^E must belong to *Onsager supercritical spaces* such as $C^{\frac{1}{3}-}$ or $\dot{H}^{\frac{5}{6}-}$. See [19] and [15], for example.

Using 2.75D shear flows for the Navier-Stokes and Euler equations, we show in Propositions 4.1 and 4.2 that the vanishing viscosity limit and the corresponding Euler flow belonging to *Onsager supercritical spaces* are not sufficient conditions for *anomalous dissipation*. Moreover in Proposition 4.1, we build upon the work of [4] to give an example of a rough solution to the 3D Euler equations that satisfies the local energy balance.

1.2. Heuristics for strong symmetry breaking. In this subsection, we give some heuristics for Theorem A.

The mechanism to get norm inflation in the critical $\dot{B}_{\infty, \infty}^{-1}$ space is well understood thanks to the work of Bourgain and Pavlović [7], and later Yoneda [45] and Cheskidov and Dai [16]. We mention here also the work of Wang [42], which demonstrates norm inflation phenomena in the spaces $\dot{B}_{\infty, q}^{-1}$ for $1 \leq q \leq 2$, but the construction is different from the one considered here.

Our point here is to explain how to get norm inflation on the third component (3rdNI), starting from data with third component equal to zero as in the case of Theorem A. Such norm inflation on the third component cannot be obtained from the previously known constructions.

As a starting point, let us consider the general plane-wave initial data

$$\kappa(r) \sum_{j=1}^r A_j (\mathbf{v} \cos(\mathbf{k}_j \cdot x) + \mathbf{v}' \cos(\mathbf{k}'_j \cdot x)). \quad (1.13)$$

Here $\mathbf{v}, \mathbf{v}' \in \mathbb{R}^3$ are fixed constant vectors, $\kappa(r)$ is some function such that $\kappa(r) \rightarrow 0$ when $r \rightarrow \infty$, $A_j \in [0, \infty)$ is a sequence of amplitudes growing geometrically, and $\mathbf{k}_j, \mathbf{k}'_j \in \mathbb{R}^3$ are two sequences of vectors whose magnitudes (approximately) grow at a geometric rate. Hence the initial data given by (1.13) is a superposition of highly

⁶We thank Hao Jia for this observation.

oscillating plane waves. This data covers the situations studied in [7, 45, 16]. In all these studies, $\kappa(r)$ and the sequence A_j need to be finely tuned in order to produce a small $\dot{B}_{\infty,\infty}^{-1}$ norm at the initial time but then a large $\dot{B}_{\infty,\infty}^{-1}$ norm after a short time has passed.

We now describe the geometric constraints that we put on the vectors \mathbf{v} , \mathbf{v}' , \mathbf{k}_j and \mathbf{k}'_j . There are two obvious conditions. First, in order to satisfy the divergence-free condition, we impose

$$\mathbf{v} \cdot \mathbf{k}_j = 0 = \mathbf{v}' \cdot \mathbf{k}'_j. \quad (3\text{rdNI Condition 1})$$

Second, in order to have vertical velocity zero initially, we impose

$$\mathbf{v} \cdot \mathbf{e}_3 = \mathbf{v}' \cdot \mathbf{e}_3 = 0 \quad (3\text{rdNI Condition 2})$$

where \mathbf{e}_3 is the third vector of the canonical basis of \mathbb{R}^3 . In order to produce norm inflation in $\dot{B}_{\infty,\infty}^{-1}$ from this superposition of highly oscillating plane waves, one needs to produce a non-oscillating function from the interaction of the term oscillating with wavenumber \mathbf{k}_j and the term oscillating with wavenumber \mathbf{k}'_j . Hence, following [7, 45, 16], we impose that there exists a fixed constant vector $\boldsymbol{\eta} \in \mathbb{R}^3$ such that

$$\mathbf{k}_j - \mathbf{k}'_j = \boldsymbol{\eta}. \quad (3\text{rdNI Condition 3})$$

The norm inflation mechanism can be seen as a backward frequency cascade, producing large-scale, non-oscillating, structures from small-scale, highly oscillating, structures.

We now investigate the conditions needed to get norm inflation of the third component. A computation of the second Duhamel iterate leads to the following inflation term

$$\kappa(r)^2 \sum_{j=1}^r \left(\int_0^t e^{-(|\mathbf{k}_j|^2 + |\mathbf{k}'_j|^2)(t-s)} ds \right) \mathcal{P}(\sin(\boldsymbol{\eta} \cdot x) ((\mathbf{v} \cdot \mathbf{k}'_j) \mathbf{v}' - (\mathbf{v}' \cdot \mathbf{k}_j) \mathbf{v})). \quad (1.14)$$

Notice that the third component of

$$\sin(\boldsymbol{\eta} \cdot x) ((\mathbf{v} \cdot \mathbf{k}'_j) \mathbf{v}' - (\mathbf{v}' \cdot \mathbf{k}_j) \mathbf{v}) \quad (1.15)$$

is zero. Hence, in order to get norm inflation on the third component, one needs the quantity in (1.15) to have a non-zero divergence, which will impose further constraints on \mathbf{k}_j , \mathbf{k}'_j , \mathbf{v} , \mathbf{v}' and $\boldsymbol{\eta}$. This is in stark contrast with previous studies [7, 45, 16], where the quantity in (1.15) is divergence-free and hence the norm inflation term remains in the span of \mathbf{v} and \mathbf{v}' .

Computing the Helmholtz-Leray projection in the norm inflation term (1.14) we get

$$\begin{aligned} & \mathcal{P}(\sin(\boldsymbol{\eta} \cdot x) ((\mathbf{v} \cdot \mathbf{k}'_j) \mathbf{v}' - (\mathbf{v}' \cdot \mathbf{k}_j) \mathbf{v})) \\ &= \sin(\boldsymbol{\eta} \cdot x) \left((\mathbf{v} \cdot \mathbf{k}'_j) \mathbf{v}' - (\mathbf{v}' \cdot \mathbf{k}_j) \mathbf{v} - \frac{\boldsymbol{\eta}}{|\boldsymbol{\eta}|^2} ((\mathbf{v} \cdot \mathbf{k}'_j)(\mathbf{v}' \cdot \boldsymbol{\eta}) - (\mathbf{v}' \cdot \mathbf{k}_j)(\mathbf{v} \cdot \boldsymbol{\eta})) \right). \end{aligned} \quad (1.16)$$

Therefore, we need

$$\boldsymbol{\eta} \cdot \mathbf{e}_3 \neq 0 \quad (3\text{rdNI Condition 4})$$

and

$$(\mathbf{v} \cdot \mathbf{k}'_j)(\mathbf{v}' \cdot \boldsymbol{\eta}) - (\mathbf{v}' \cdot \mathbf{k}_j)(\mathbf{v} \cdot \boldsymbol{\eta}) \neq 0$$

in order to have norm inflation on the third component of the velocity. Using (3rdNI Condition 1) we can rewrite the last condition as

$$(\mathbf{v} \cdot \mathbf{k}'_j)(\mathbf{v}' \cdot \mathbf{k}_j) \neq 0$$

i.e.

$$\mathbf{v} \cdot \mathbf{k}'_j \neq 0 \quad \text{and} \quad \mathbf{v}' \cdot \mathbf{k}_j \neq 0. \quad (3\text{rdNI Condition 5})$$

Notice that conditions (3rdNI Condition 1)-(3rdNI Condition 5) are necessary but also sufficient to have norm inflation on the third component. There are many possible choices within the constraints (3rdNI Condition 1)-(3rdNI Condition 5). In particular, taking

$$\mathbf{v} = (1, -\lambda, 0) \quad \text{and} \quad \mathbf{k}_{\mathbf{j}} = (\lambda, 1, 2^{j-1}K)$$

for $\lambda, K \in \mathbb{Z}$, one has a whole family of initial data u_{in} with $u_{\text{in}}^3 = 0$ that produces norm inflation on the third component in $\dot{B}_{\infty, \infty}^{-1}$ and such that u_{in} is close in $\dot{B}_{\infty, \infty}^{-1}$ to a 2.75D shear flow initial data defined in (1.8).

1.3. Heuristics for strong isotropic symmetry breaking. From the previous subsection, notice that for initial data of the form

$$\kappa(r) \sum_{j=1}^r A_j (\mathbf{v} \cos(\mathbf{k}_{\mathbf{j}} \cdot x) + \mathbf{v}' \cos(\mathbf{k}'_{\mathbf{j}} \cdot x))$$

that satisfies (3rdNI Condition 1)-(3rdNI Condition 5), the associated inflation term (1.16) vanishes in the direction $\boldsymbol{\eta}$. Here

$$\mathbf{k}_{\mathbf{j}} - \mathbf{k}'_{\mathbf{j}} = \boldsymbol{\eta} \quad \text{for all } \mathbf{j}. \quad (1.17)$$

This represents a block in using such initial data for obtaining the isotropic norm inflation (1.3) in Theorem A'.

To overcome this, we instead take initial data of the form

$$\kappa(r) \sum_{j=1}^{r^3} A_j (\mathbf{v}_{\mathbf{j}} \cos(\mathbf{k}_{\mathbf{j}} \cdot x) + \mathbf{v}'_{\mathbf{j}} \cos(\mathbf{k}'_{\mathbf{j}} \cdot x))$$

with $\mathbf{k}_{\mathbf{j}} - \mathbf{k}'_{\mathbf{j}} = \boldsymbol{\eta}_{\mathbf{j}}.$ (1.18)

Here, $\mathbf{v}_{\mathbf{j}}$ and $\mathbf{v}'_{\mathbf{j}}$ vary with \mathbf{j} and crucially the low frequency vector $\boldsymbol{\eta}_{\mathbf{j}}$ points in different directions depending on the index \mathbf{j} . Specifically, we glue higher frequency terms to the initial data in Theorem A, such that the added terms $\mathbf{v}_{\mathbf{j}}, \mathbf{v}'_{\mathbf{j}}$ and $\boldsymbol{\eta}_{\mathbf{j}}$ point in different directions depending on \mathbf{j} .

The initial data we design in Theorem A' can be decomposed into three pieces $u_{\text{in}} = u_{\text{in}}^{(1)} + u_{\text{in}}^{(2)} + u_{\text{in}}^{(3)}$ such that

- each $u_{\text{in}}^{(1)} - u_{\text{in}}^{(3)}$ separately generate an associated Navier-Stokes solution with a norm inflation term, with each of these norm inflation terms being of comparable size in $\dot{B}_{\infty, \infty}^{-1}$,
- the norm inflation term associated to u_{in} is the sum of the norm inflation terms associated to $u_{\text{in}}^{(1)} - u_{\text{in}}^{(3)}$.

Careful choices of $\mathbf{v}_{\mathbf{j}}, \mathbf{v}'_{\mathbf{j}}, \mathbf{k}_{\mathbf{j}}$ and $\mathbf{k}'_{\mathbf{j}}$ then give that the norm inflation terms associated to $u_{\text{in}}^{(1)} - u_{\text{in}}^{(3)}$ point in linearly independent directions. This, together with our choices of $\mathbf{v}_{\mathbf{j}}, \mathbf{v}'_{\mathbf{j}}, \mathbf{k}_{\mathbf{j}}$ and $\mathbf{k}'_{\mathbf{j}}$ and the fact that $\dot{B}_{\infty, \infty}^{-1}$ is an L^∞ -based space, enable us to show that for any fixed unit vector \mathbf{e}

- (i) the dot product of \mathbf{e} with at least one of the norm inflation terms associated to $u_{\text{in}}^{(1)} - u_{\text{in}}^{(3)}$ has a $\dot{B}_{\infty, \infty}^{-1}$ norm with a large lower bound,
- (ii) the lower bound in (i) also serves as a lower bound for the $\dot{B}_{\infty, \infty}^{-1}$ norm of the dot product of \mathbf{e} with the norm inflation term associated to u_{in} .

These features then imply that u_{in} generates a norm inflation term that has large $\dot{B}_{\infty, \infty}^{-1}$ norm in all unit directions. This in turn leads to the results described in Theorem A'.

1.4. Heuristics for symmetry breaking despite favorable pressure gradient. In this subsection, we give some heuristics for Theorem B.

Let us first explain how we design the initial data. Our objective is to find an initial data that will generate symmetry breaking despite favorable initial pressure (see Footnote 5). The two fractions in u_{in} are there to fulfill the condition $\partial_3 P_{\text{in}} \equiv 0$, where P_{in} is defined by (1.4). We also remark that the order in t in (1.7) is expected because $\partial_3 P_{\text{in}} = 0$ at $t = 0$ formally implies that $\partial_t u^3 = 0$ at $t = 0$. The breaking is not driven by the vertical derivative of the pressure at the initial time, as is the case in Theorem A and for the Taylor-Green vortex, see Figure 1.

In our proof, the condition $N > N_0$ appears for technical reasons in order to identify the leading order term. Indeed, for N large, the term involving S_1 is the dominant term in the right hand side of (3.11). Notice also that the larger the N , the closer our data is from the two-dimensional data $(\cos x_2, \cos x_1, 0)$ that generates a unique global two-dimensional solution to 3D Navier-Stokes. This has two implications. First, one sees, that u_{in} generates a unique solution to the 3D Navier-Stokes equations on $[0, 1] \times \mathbb{T}^3$ for N large. Second, the larger the N , the weaker the symmetry breaking effect should be. This observation is consistent with the bound $O(t^2/N^2)$ in (1.7).

Remark 5 (on the condition $N \gg 1$ in Theorem B). Figure 2 shows a simulation of the Navier-Stokes equations with initial data from Theorem B taking $N = 1.1$. The graph shows that symmetry breaking happens in spite of the fact that N is taken small. Therefore we expect that the result of Theorem B remains true for $1 < N \leq N_0$.

1.5. Outline of the paper. Section 2 is devoted to the proof of strong symmetry breaking, namely Theorem A (see Subsection 2.1) and to the proof of strong isotropic symmetry breaking, namely Theorem A' (see Subsection 2.2). Section 3 addresses the proof of Theorem B, i.e. symmetry breaking despite pressure favorable to symmetry preservation. The last part of the paper, Section 4 is concerned with some applications of the 2.75D shear flows, which are symmetry preserving shear flows. This section contains two types of results. First, in Section 4.1 we investigate inviscid damping effects for 2.75D shear flow solutions of Euler. Second, in Section 4.2 we study Onsager supercritical inviscid limits of 2.75D shear flows. Finally in Appendix A, we give another perspective on the derivation of 2.75D shear flows.

1.6. Notations and preliminary results. We begin this section by introducing relevant notation. We denote by C positive numerical constants that may change from one line to the other, and we sometimes write $A \lesssim B$ (resp. $A \lesssim_a B$) instead of $A \leq CB$ (resp. $A \leq C_a B$ with the constant C depending on a). Likewise, $A \sim B$ means that $C_1 B \leq A \leq C_2 B$ with absolute constants C_1, C_2 . Throughout the paper, i -th coordinate ($i = 1, 2, 3$) of a vector v will be denoted by v^i , and horizontal component of v will be denoted by v^{h} . For a real-valued matrix \mathcal{M} , \mathcal{M}^{T} represents its transpose, while for two multidimensional real-valued matrices $\mathcal{M}_1, \mathcal{M}_2$, $\mathcal{M}_1 : \mathcal{M}_2$ denotes their standard inner product. For X a Banach space, $p \in [1, \infty]$ and $T \in (0, \infty]$, the notation $L^p(0, T; X)$ or $L_T^p(X)$ stands for the set of measurable functions $f : [0, T] \rightarrow X$ with $t \mapsto \|f(t)\|_X$ in $L^p(0, T)$, endowed with the norm $\|\cdot\|_{L_T^p(X)} := \|\|\cdot\|_X\|_{L^p(0, T)}$. We keep the same notation for functions with several components.

We recall that the Besov spaces $\dot{B}_{\infty, \infty}^{-2\sigma}$ (with $\sigma > 0$) is equipped with the norm

$$\|v(\cdot)\|_{\dot{B}_{\infty, \infty}^{-2\sigma}} = \|\|s^\sigma e^{s\Delta} v(\cdot)\|_{L^\infty}\|_{L^\infty(\mathbb{R}_+)}.$$

Note also that one has the embedding

$$L^3(\mathbb{T}^3) \hookrightarrow \dot{B}_{\infty, \infty}^{-1}(\mathbb{T}^3) \quad (1.19)$$

for mean-free functions on the torus.⁷ As is usual, we define the bilinear operator

$$\mathcal{B}(u, v)(t, x) := - \int_0^t e^{(t-\tau)\Delta} \mathcal{P}(u \cdot \nabla v)(\tau, x) d\tau$$

with \mathcal{P} the projection on divergence-free vector fields (the so-called Helmholtz-Leray projection).

We need the following obvious estimates for the one-dimensional heat kernel

$$\mathcal{K}(t, x_3) := \frac{1}{\sqrt{4\pi t}} e^{-\frac{|x_3|^2}{4t}}, \quad \forall (t, x_3) \in \mathbb{R}_+ \times \mathbb{T}. \quad (1.20)$$

Lemma 1.1. *Let $g \in C^\infty(\mathbb{T})$, then for any $s \in \mathbb{R}_+$, one has*

$$\|(\mathcal{K} \star g)(s, \cdot)\|_{L^\infty(\mathbb{T})} \leq \|g\|_{L^\infty(\mathbb{T})}$$

and

$$\|(\mathcal{K} \star g)(s, \cdot) - g(\cdot)\|_{L^\infty(\mathbb{T})} \leq s \|g''\|_{L^\infty(\mathbb{T})}.$$

Finally, we state a standard absorbing lemma which is useful for our proofs.

Lemma 1.2. *Suppose that $y : [0, T] \rightarrow [0, \infty)$ is continuous and satisfies $y(0) = 0$. Furthermore suppose that for all $t \in [0, T]$, y satisfies the following inequality:*

$$\sup_{s \in [0, t]} y(s) \leq a \left(\sup_{s \in [0, t]} y(s) \right)^2 + b \sup_{s \in [0, t]} y(s) + c,$$

with $a, b, c > 0$ and $b + 2ac < \frac{1}{4}$. Then we conclude that

$$\sup_{s \in [0, T]} y(s) < 2c.$$

2. STRONG SYMMETRY BREAKING

2.1. Proof of Theorem A. In this section, our objective is to prove Theorem A. We investigate the growth of the vertical velocity for the three-dimensional Navier-Stokes problem (1.1) supplemented with initial data u_{in} that is close in the critical Besov spaces $\dot{B}_{\infty, \infty}^{-1}$ to initial data considered in (1.8). For a heuristic description of the growth mechanism with a focus on how to produce third component norm inflation from anisotropic initial data, we refer to Subsection 1.2. We proceed in three steps.

⁷Let us give a short proof of this embedding. For a mean-free function v in $L^3(\mathbb{T}^3)$, for $s > 1$,

$$\begin{aligned} \left\| \sum_{\xi \in \mathbb{Z}^3 \setminus \{0\}} e^{-s|\xi|^2} e^{ix \cdot \xi} \hat{v}(\xi) \right\|_{L^\infty(\mathbb{T}^3)} &= \left\| \sum_{\xi \in \mathbb{Z}^3 \setminus \{0\}} s|\xi|^2 e^{-s|\xi|^2} e^{ix \cdot \xi} \frac{\hat{v}(\xi)}{s|\xi|^2} \right\|_{L^\infty(\mathbb{T}^3)} \\ &\leq \frac{1}{s} \left(\sum_{\xi \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|\xi|^4} \right)^{\frac{1}{2}} \left(\sum_{\xi \in \mathbb{Z}^3 \setminus \{0\}} |\hat{v}(\xi)|^2 \right)^{\frac{1}{2}} \\ &\leq \frac{C}{s} \|v\|_{L^2(\mathbb{T}^3)} \leq \frac{C}{s} \|v\|_{L^3(\mathbb{T}^3)}, \end{aligned}$$

where $C \in (0, \infty)$ is a universal constant. Notice that we used that the function xe^{-x} is bounded on \mathbb{R} . For $s \in (0, 1]$, we rely on the result of Maekawa and Terasawa [35] for instance.

Step 1: choice of the initial data. Let $\Gamma_1, \Gamma_2 : \mathbb{N} \mapsto \mathbb{R}$ be such that

$$\Gamma_1(m) := \sum_{j=1}^m \frac{1}{j} \quad \text{and} \quad \Gamma_2(m) := \Gamma_1^{\frac{1}{3}}(m) \quad \text{for } m \in \mathbb{N}. \quad (2.1)$$

Let r be a large integer (to be specified later). We set initial data u_{in} and \bar{u}_{in} as follows:⁸

$$u_{\text{in}} = \frac{1}{\Gamma_2(r)} \sum_{j=1}^r \frac{k_j}{\sqrt{j}} \left(\mathbf{v} \cos(x_1 + x_2 + k_j x_3) + \mathbf{v}' \cos(-x_2 + (k_j + 1)x_3) \right),$$

$$\bar{u}_{\text{in}} = \frac{1}{\Gamma_2(r)} \sum_{j=1}^r \frac{k_j}{\sqrt{j}} \mathbf{v} \cos(x_1 + x_2 + k_j x_3),$$

where $\mathbf{v} = (1, -1, 0)$, $\mathbf{v}' = (1, 0, 0)$ are vectors and we define the sequence $k_j = 2^{3j} T^{-\frac{1}{2}}$ ($j = 1, \dots, r$). The existence time $0 < T < 1$ is to be determined in terms of r .

Obviously, \bar{u}_{in} has the structure (1.8) by taking

$$\lambda = 1, \quad f(y_1, y_2) = \frac{1}{\Gamma_2(r)} \sum_{j=1}^r \frac{k_j}{\sqrt{j}} \mathbf{v} \cos(y_1 + k_j y_2), \quad g = 0.$$

Thus the vertical velocity of the corresponding 2.75D shear flow remains identically zero for all positive time.

Notice that

$$u_{\text{in}} - \bar{u}_{\text{in}} = \frac{1}{\Gamma_2(r)} \sum_{j=1}^r \frac{k_j}{\sqrt{j}} \mathbf{v}' \cos(-x_2 + (k_j + 1)x_3),$$

$$e^{t\Delta}(u_{\text{in}} - \bar{u}_{\text{in}})(x) = \frac{1}{\Gamma_2(r)} \sum_{j=1}^r \frac{k_j}{\sqrt{j}} \mathbf{v}' \cos(-x_2 + (k_j + 1)x_3) e^{-((k_j+1)^2+1)t}$$

and for appropriate r , we have

$$\|u_{\text{in}} - \bar{u}_{\text{in}}\|_{\dot{B}_{\infty, \infty}^{-1}} \leq \frac{1}{\Gamma_2(r)} \sup_{s>0} \left(\sum_{j=1}^r \frac{k_j}{\sqrt{j}} s^{\frac{1}{2}} e^{-((k_j+1)^2+1)s} \right)$$

$$\lesssim \frac{1}{\Gamma_2(r)} \sup_{s>0} \left(\sum_{j=1}^r k_j s^{\frac{1}{2}} e^{-k_j^2 s} \right) \lesssim \frac{1}{\Gamma_2(r)} = \Gamma_1^{-\frac{1}{3}}(r).$$

In the above and in what follows, we use that series of the type $\sum_{j \in \mathbb{N}} k_j s^{\frac{1}{2}} e^{-k_j^2 s}$ and $\sum_{j \in \mathbb{N}} k_j^2 s e^{-k_j^2 s}$ are uniformly bounded in s . This can be easily seen by splitting the sum into $\{j : 16^j \frac{s}{T} < 1\}$ and its complement.

Step 2: analysis of the second approximation. Now, we analyze the second approximate solution associated with initial data u_{in} . In order to do that, let us first recall $u_1(t, x) = e^{t\Delta} u_{\text{in}}$ with

$$u_1(t, x) = \frac{1}{\Gamma_2(r)} \sum_{j=1}^r \frac{k_j}{\sqrt{j}} \left(\mathbf{v} \cos(x_1 + x_2 + k_j x_3) e^{-(k_j^2+2)t} \right. \\ \left. + \mathbf{v}' \cos(-x_2 + (k_j + 1)x_3) e^{-((k_j+1)^2+1)t} \right), \quad (2.2)$$

⁸We emphasize that k_j is a scalar. Comparing the data to (1.13), we see that here $\kappa(r) := \frac{1}{\Gamma_2(r)}$, $A_j = \frac{k_j}{\sqrt{j}}$, $\mathbf{k}_j = (1, 1, k_j)$ and $\mathbf{k}'_j = (0, -1, k_j + 1)$.

and $v_2 := \mathcal{B}(u_1, u_1)$ with

$$v_2(t, x) = \frac{1}{\Gamma_2^2(r)} \sum_{i=1}^r \sum_{j=1}^r \int_0^t e^{(t-\tau)\Delta} \mathcal{P}U_{i,j}(\tau, x) d\tau,$$

where

$$U_{i,j}(\tau, x) := \frac{k_i k_j}{\sqrt{i j}} \left(\mathbf{v} G_{i,j}^+(\tau, x) + \mathbf{v}' G_{i,j}^-(\tau, x) \right)$$

and

$$\begin{aligned} G_{i,j}^+(\tau, x) &:= -\frac{1}{2} \left(\sin(x_1 + 2x_2 + (k_j - k_i - 1)x_3) + \sin(x_1 + (k_j + k_i + 1)x_3) \right) \\ &\quad \times e^{-((k_j^2 + (k_i + 1)^2 + 3)\tau)}, \\ G_{i,j}^-(\tau, x) &:= -\frac{1}{2} \left(\sin(-x_1 - 2x_2 + (k_j - k_i + 1)x_3) + \sin(x_1 + (k_j + k_i + 1)x_3) \right) \\ &\quad \times e^{-((k_j + 1)^2 + k_i^2 + 3)\tau}. \end{aligned}$$

We see that

$$\begin{aligned} U_{j,j}(\tau, x) &= \frac{1}{2} \frac{k_j^2}{j} (\mathbf{v}' - \mathbf{v}) \sin(x_1 + 2x_2 - x_3) e^{-(2k_j^2 + 2k_j + 4)\tau} \\ &\quad - \frac{1}{2} \frac{k_j^2}{j} (\mathbf{v} + \mathbf{v}') \sin(x_1 + (2k_j + 1)x_3) e^{-(2k_j^2 + 2k_j + 4)\tau} \\ &=: U_{j,j}^+(\tau, x) + U_{j,j}^-(\tau, x). \end{aligned}$$

So we can decompose v_2 as $v_2 = v_{2,1} + v_{2,2} + v_{2,3}$, where

$$\begin{cases} v_{2,1}(t, x) := \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r \int_0^t e^{(t-\tau)\Delta} \mathcal{P}U_{j,j}^+(\tau, x) d\tau, \\ v_{2,2}(t, x) := \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r \int_0^t e^{(t-\tau)\Delta} \mathcal{P}U_{j,j}^-(\tau, x) d\tau, \\ v_{2,3}(t, x) := \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r \sum_{i < j} \int_0^t e^{(t-\tau)\Delta} \mathcal{P}(U_{i,j}(\tau, x) + U_{j,i}(\tau, x)) d\tau. \end{cases} \quad (2.3)$$

Note that $v_{2,1}$ will be the term producing the norm inflation.

Lemma 2.1. *We have the following key estimates:*

$$\|v_{2,1}(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \frac{\Gamma_1(r)}{\Gamma_2^2(r)} = \Gamma_1^{\frac{1}{3}}(r), \quad \text{for } t > 0 \quad (2.4)$$

and for each components of $v_{2,1}$ on the time interval $[T/320, T]$,

$$\|v_{2,1}^1(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-1}} = \|v_{2,1}^2(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-1}} = \|v_{2,1}^3(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-1}} \gtrsim \Gamma_1^{\frac{1}{3}}(r), \quad (2.5)$$

$$\|v_{2,1}^1(t, \cdot)\|_{L^3} = \|v_{2,1}^2(t, \cdot)\|_{L^3} = \|v_{2,1}^3(t, \cdot)\|_{L^3} \gtrsim \Gamma_1^{\frac{1}{3}}(r). \quad (2.6)$$

Moreover, for $t > 0$

$$\|v_{2,2}(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \frac{1}{\Gamma_2^2(r)} = \Gamma_1^{-\frac{2}{3}}(r), \quad (2.7)$$

$$\|v_{2,3}(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \frac{1}{\Gamma_2^2(r)} = \Gamma_1^{-\frac{2}{3}}(r), \quad (2.8)$$

$$\|u_1(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \frac{1}{\sqrt{t} \Gamma_2(r)} = \Gamma_1^{-\frac{1}{3}}(r) t^{-\frac{1}{2}}. \quad (2.9)$$

Proof of Lemma 2.1. First, a direct computation gives that

$$\mathcal{P}((\mathbf{v}' - \mathbf{v}) \sin(x_1 + 2x_2 - x_3)) = \frac{1}{3} \sin(x_1 + 2x_2 - x_3) (-1, 1, 1).$$

Then, by the definition of $v_{2,1}$ in (2.3) and the above equality,

$$v_{2,1}(t, x) = \frac{(-1, 1, 1)}{\Gamma_2^2(r)} \sum_{j=1}^r \frac{k_j^2}{6j} \int_0^t e^{(t-\tau)\Delta} \sin(x_1 + 2x_2 - x_3) e^{-(2k_j^2 + 2k_j + 4)\tau} d\tau.$$

Thus for $t > 0$

$$\|v_{2,1}(t, \cdot)\|_{L^\infty} \lesssim \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r \frac{k_j^2}{j} \int_0^t e^{-2k_j^2\tau} d\tau \lesssim \frac{\Gamma_1(r)}{\Gamma_2^2(r)}.$$

The vertical component of $v_{2,1}$ is given by

$$v_{2,1}^3(t, x) = \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r \frac{k_j^2}{6j} \int_0^t e^{(t-\tau)\Delta} \sin(x_1 + 2x_2 - x_3) e^{-(2k_j^2 + 2k_j + 4)\tau} d\tau. \quad (2.10)$$

Using this and that $k_1^2 T = 64$, we obtain

$$\begin{aligned} \|v_{2,1}^3(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-1}} &\gtrsim \frac{1}{\Gamma_2^2(r)} \sup_{s>0} \left(\sum_{j=1}^r \frac{k_j^2}{j} s^{\frac{1}{2}} \int_0^t e^{-6(t-\tau+s)} e^{-4k_j^2\tau} d\tau \right) \\ &\gtrsim \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r \frac{e^{-6t}}{j} \int_0^t k_j^2 e^{-4k_j^2\tau} d\tau \\ &\gtrsim \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r \frac{1}{j} (1 - e^{-4k_j^2 t}) \\ &\gtrsim \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r \frac{1}{j} (1 - e^{-1}) \gtrsim \frac{\Gamma_1(r)}{\Gamma_2^2(r)} \end{aligned} \quad (2.11)$$

for $t \in [T/256, T]$ with $T < 1$. Moreover, we see that the components of $v_{2,1}$ are comparable, and due to the embedding (1.19), we get (2.5) and (2.6) easily.

Next, let us estimate⁹ $v_{2,2}(t, x)$ and $u_1(t, x)$ for $t > 0$. We have by the definition of v_{22} in (2.3) that

$$\begin{aligned} \|v_{2,2}(t, \cdot)\|_{L^\infty} &\lesssim \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r \frac{k_j^2}{j} \int_0^t e^{-(1+(2k_j+1)^2)(t-\tau)} e^{-(2k_j^2 + 2k_j + 4)\tau} d\tau, \\ &\lesssim \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r k_j^2 t e^{-(2k_j^2 + 2k_j + 4)t} \frac{1 - e^{-(2k_j^2 + 2k_j - 2)t}}{(2k_j^2 + 2k_j - 2)t} \\ &\lesssim \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r k_j^2 t e^{-k_j^2 t} \lesssim \frac{1}{\Gamma_2^2(r)}, \end{aligned}$$

⁹In the computation follows, we drop the Helmholtz-Leray projector \mathcal{P} since its contribution is harmless.

where we used that $\frac{(1-e^{-(2k_j^2+2k_j-2)t})}{(2k_j^2+2k_j-2)t}$ is uniformly bounded for $t > 0$. Similarly, from (2.2) we have for $t > 0$,

$$\|\sqrt{t} u_1(t, \cdot)\|_{L^\infty} \lesssim \frac{1}{\Gamma_2(r)} \sum_{j=1}^r (k_j^2 t)^{\frac{1}{2}} e^{-k_j^2 t} \lesssim \frac{1}{\Gamma_2(r)}.$$

Thus, we have shown (2.7) and (2.9).

Finally, using $\frac{k_j}{2} \leq k_j - k_i - 1$ for $i < j$ and $\sum_{i < j} k_i \leq \frac{k_j}{4}$, it is easy to see that

$$-2(k_j - k_i)^2 = -2k_i^2 - 2(k_j - 2k_i)k_j \leq -(2k_i^2 + k_j^2).$$

Therefore we have for $v_{2,3}$

$$\begin{aligned} \|v_{2,3}(t, \cdot)\|_{L^\infty} &\lesssim \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r \sum_{i < j} \frac{k_i k_j}{\sqrt{ij}} \\ &\quad \times \int_0^t \left(e^{-(k_j - k_i - 1)^2(t-\tau)} + e^{-(k_j + k_i + 1)^2(t-\tau)} \right) e^{-(k_j^2 + k_i^2)\tau} d\tau \\ &\lesssim \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r k_j^2 \int_0^t e^{-\frac{1}{4}k_j^2(t-\tau)} e^{-k_j^2\tau} d\tau \\ &\lesssim \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^r k_j^2 t e^{-\frac{1}{4}k_j^2 t} \frac{1 - e^{-\frac{3}{4}k_j^2 t}}{k_j^2 t} \lesssim \frac{1}{\Gamma_2^2(r)}. \quad \square \end{aligned}$$

Step 3: error analysis. We will show that for appropriately chosen $0 < T < 1$, there exists a solution u on $[0, T] \times \mathbb{T}^3$. We will then analyze the remainder term w between u and the second iterate. Showing the existence of u is equivalent to finding w satisfying the integral equation

$$w = F_1 + F_2 + F_3 \tag{2.12}$$

with

$$\begin{aligned} F_1 &:= \mathcal{B}(w, u_1) + \mathcal{B}(u_1, w) + \mathcal{B}(w, v_2) + \mathcal{B}(v_2, w), \\ F_2 &:= \mathcal{B}(w, w), \\ F_3 &:= \mathcal{B}(u_1, v_2) + \mathcal{B}(v_2, u_1) + \mathcal{B}(v_2, v_2). \end{aligned}$$

Then u is given by $u = u_1 + v_2 + w$. From Lemma 2.1, we have for v_2 that

$$\begin{aligned} \|v_2(t, \cdot)\|_{L^\infty} &\lesssim \|v_{2,1}(t, \cdot)\|_{L^\infty} + \|v_{2,2}(t, \cdot)\|_{L^\infty} + \|v_{2,2}(t, \cdot)\|_{L^\infty} \\ &\lesssim \Gamma_1^{\frac{1}{3}}(r). \end{aligned} \tag{2.13}$$

From (2.9), we have for $t > 0$

$$\|u_1(t, \cdot)\|_{L^\infty} \lesssim \Gamma_1^{-\frac{1}{3}}(r) t^{-\frac{1}{2}}. \tag{2.14}$$

By the L^∞ bilinear estimate and estimates (2.13)-(2.14), we have for $0 < t \leq T < 1$,

$$\begin{aligned} \|\mathcal{B}(A, u_1)(t, \cdot)\|_{L^\infty} &\lesssim \Gamma_1^{-\frac{1}{3}}(r) \int_0^t (t-\tau)^{-\frac{1}{2}} \tau^{-\frac{1}{2}} d\tau \sup_{t \in [0, T]} \|A(t, \cdot)\|_{L^\infty} \\ &\lesssim \Gamma_1^{-\frac{1}{3}}(r) \int_0^1 (1-s)^{-\frac{1}{2}} s^{-\frac{1}{2}} ds \sup_{t \in [0, T]} \|A(t, \cdot)\|_{L^\infty} \\ &\lesssim \Gamma_1^{-\frac{1}{3}}(r) \sup_{t \in [0, T]} \|A(t, \cdot)\|_{L^\infty}, \end{aligned} \tag{2.15}$$

$$\begin{aligned}
\|\mathcal{B}(A, v_2)(t, \cdot)\|_{L^\infty} &\lesssim \Gamma_1^{\frac{1}{3}}(r) \int_0^t (t-\tau)^{-\frac{1}{2}} d\tau \sup_{t \in [0, T]} \|A(t, \cdot)\|_{L^\infty} \\
&\lesssim \Gamma_1^{\frac{1}{3}}(r) \sqrt{T} \sup_{t \in [0, T]} \|A(t, \cdot)\|_{L^\infty}, \tag{2.16}
\end{aligned}$$

$$\|\mathcal{B}(u_1, v_2)(t, \cdot)\|_{L^\infty} \lesssim \Gamma_1^{-\frac{1}{3}}(r) \Gamma_1^{\frac{1}{3}}(r) \int_0^t (t-\tau)^{-\frac{1}{2}} \tau^{-\frac{1}{2}} d\tau \lesssim 1, \tag{2.17}$$

$$\|\mathcal{B}(v_2, v_2)(t, \cdot)\|_{L^\infty} \lesssim \Gamma_1^{\frac{1}{3}}(r) \Gamma_1^{\frac{1}{3}}(r) \int_0^t (t-\tau)^{-\frac{1}{2}} d\tau \lesssim \Gamma_1^{\frac{2}{3}}(r) \sqrt{T} \tag{2.18}$$

and

$$\|\mathcal{B}(A, B)(t, \cdot)\|_{L^\infty} \lesssim \sqrt{T} \sup_{t \in [0, T]} \|A(t, \cdot)\|_{L^\infty} \sup_{t \in [0, T]} \|B(t, \cdot)\|_{L^\infty}. \tag{2.19}$$

We take $\delta = \Gamma_1^{-\frac{1}{3}}(r)$ and $T = \Gamma_1^{-1}(r)$. Notice that $T < \delta$ and $\sqrt{T}(1 + \Gamma_1^{\frac{2}{3}}(r)\sqrt{T}) \lesssim \Gamma_1^{-\frac{1}{3}}(r) \ll 1$ for large r . Using this and (2.15)-(2.19), we can apply [23, Lemma A.1]. This gives the existence of $w \in C([0, T] \times \mathbb{T}^3)$. We also infer that

$$\begin{aligned}
&\sup_{t \in [0, T]} \|w(t, \cdot)\|_{L^\infty} \\
&\lesssim \sup_{t \in [0, T]} (\|\mathcal{B}(w, u_1)\|_{L^\infty} + \|\mathcal{B}(w, v_2)\|_{L^\infty} + \|\mathcal{B}(w, w)\|_{L^\infty} + \|\mathcal{B}(u_1, v_2)\|_{L^\infty} \\
&\quad + \|\mathcal{B}(v_2, v_2)\|_{L^\infty}) \\
&\lesssim \left(\sup_{t \in [0, T]} \|w(t, \cdot)\|_{L^\infty} \right)^2 \sqrt{T} + \left(\Gamma_1^{-\frac{1}{3}}(r) + \Gamma_1^{\frac{1}{3}}(r) \sqrt{T} \right) \sup_{t \in [0, T]} \|w(t, \cdot)\|_{L^\infty} \\
&\quad + \left(1 + \Gamma_1^{\frac{2}{3}}(r) \sqrt{T} \right).
\end{aligned}$$

The choice of T made above allows us to apply an absorbing argument (see Lemma 1.2). Hence we have the following a priori bound for sufficiently large r

$$\sup_{t \in [0, T]} \|w(t, \cdot)\|_{L^\infty} \lesssim \Gamma_1^{\frac{1}{6}}(r). \tag{2.20}$$

We now prove the main theorem. Thanks to Lemma 2.1 and (2.20), (2.14), we conclude that for $t \in [T/256, T]$ and large enough r

$$\begin{aligned}
\|u^3(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-1}} &= \|u_1^3 + v_2^3 + w^3\|_{\dot{B}_{\infty, \infty}^{-1}} \\
&\geq \|v_2^3(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-1}} - \|u_1(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-1}} - \|w(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-1}} \\
&\geq \|v_{2,1}^3(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-1}} - \|v_{2,2}(t, \cdot)\|_{L^\infty} - \|v_{2,3}(t, \cdot)\|_{L^\infty} - \|u_1(t, \cdot)\|_{L^\infty} \\
&\quad - \|w(t, \cdot)\|_{L^\infty} \\
&\gtrsim \Gamma_1^{\frac{1}{3}}(r) - \Gamma_1^{-\frac{2}{3}}(r) - \Gamma_1^{\frac{1}{6}}(r) \gtrsim \Gamma_1^{\frac{1}{3}}(r) = \frac{1}{\delta},
\end{aligned}$$

where we used the embedding (1.19). We finally obtain (1.2) using (2.6), $u = u_1 + v_{2,1} + v_{2,2} + v_{2,3} + w$ and the fact that

$$\|u_1(t, \cdot)\|_{L^\infty} + \|v_{2,2}(t, \cdot)\|_{L^\infty} + \|v_{2,3}(t, \cdot)\|_{L^\infty} + \|w(t, \cdot)\|_{L^\infty} \ll \Gamma_1^{\frac{1}{3}}(r).$$

This completes the proof of Theorem A.

2.2. Proof of Theorem A'. The outcome of the previous proof is that the data (1.8) is well-designed to show the norm inflation on the third component. This data will serve as a first building block for constructing the initial data for Theorem A'. Two other blocks will be added in order to prove Isotropic Norm Inflation as stated in (1.3). The objective of this construction is to rule out the possibility of compensations between different components.

Step 1: choice of the initial data. Let Γ_1 and Γ_2 be defined as in (2.1). Let r be a large integer (to be specified later). We set initial data u_{in} as follows:

$$u_{\text{in}} = \frac{1}{\Gamma_2(r)} \sum_{j=1}^{r^3} \frac{k_j}{\sqrt{j}} (\mathbf{v}_j \cos(\mathbf{k}_j \cdot x) + \mathbf{v}'_j \cos(\mathbf{k}'_j \cdot x)), \quad (2.21)$$

where we define the sequence $k_j = 2^{3j} T^{-\frac{1}{2}}$ ($j = 1, \dots, r^3$)¹⁰ and where $\mathbf{v}_j, \mathbf{v}'_j, \mathbf{k}_j, \mathbf{k}'_j$ are vectors which (contrary to the construction in Theorem A) depend on j in the following way:

$$\mathbf{v}_j = \begin{cases} (1, -1, 0), & 1 \leq j \leq r, \\ (1, 0, 0), & r+1 \leq j \leq r^2, \\ (1, 1, 0), & r^2+1 \leq j \leq r^3, \end{cases}, \quad \mathbf{v}'_j = \begin{cases} (1, 0, 0), & 1 \leq j \leq r, \\ (1, 1, 0), & r+1 \leq j \leq r^2, \\ (0, 1, 0), & r^2+1 \leq j \leq r^3, \end{cases}$$

and

$$\mathbf{k}_j = \begin{cases} (1, 1, k_j), & 1 \leq j \leq r, \\ (0, 1, k_j), & r+1 \leq j \leq r^2, \\ (1, -1, k_j), & r^2+1 \leq j \leq r^3, \end{cases}, \quad \mathbf{k}'_j = \begin{cases} (0, -1, k_j+1), & 1 \leq j \leq r, \\ (-1, 1, k_j), & r+1 \leq j \leq r^2, \\ (1, 0, k_j), & r^2+1 \leq j \leq r^3. \end{cases}$$

Notice that we have the following relations

$$\mathbf{v}_j \cdot \mathbf{k}_j = 0 = \mathbf{v}'_j \cdot \mathbf{k}'_j, \quad \mathbf{v}_j \cdot \mathbf{e}_3 = 0 = \mathbf{v}'_j \cdot \mathbf{e}_3,$$

which guarantee that the data is incompressible and has vanishing third component. Moreover, we have a low frequency vector

$$\boldsymbol{\eta}_j := \mathbf{k}_j - \mathbf{k}'_j = \begin{cases} (1, 2, -1), & 1 \leq j \leq r, \\ (1, 0, 0), & r+1 \leq j \leq r^2, \\ (0, -1, 0), & r^2+1 \leq j \leq r^3 \end{cases} \quad (2.22)$$

that varies according to j . This is key to the Isotropic Norm Inflation mechanism. Notice that

$$\mathbf{v}_j \cdot \mathbf{k}'_j = \begin{cases} 1, & 1 \leq j \leq r, \\ -1, & r+1 \leq j \leq r^2, \\ 1, & r^2+1 \leq j \leq r^3, \end{cases}, \quad \mathbf{v}_j \cdot \boldsymbol{\eta}_j = \begin{cases} -1, & 1 \leq j \leq r, \\ 1, & r+1 \leq j \leq r^2, \\ -1, & r^2+1 \leq j \leq r^3. \end{cases}$$

Step 2: analysis of the second approximation. As above, we consider the second Duhamel iterate from which the norm inflation comes

$$v_2 := \mathcal{B}(u_1, u_1),$$

¹⁰As above, the existence time $0 < T < 1$ is to be determined in terms of r .

where $u_1 := e^{t\Delta} u_{\text{in}}$ is the first Duhamel iterate. We identify the inflation term $v_{2,1}$ by decomposing as above, cf. (2.3): $v_2 = v_{2,1} + v_{2,2} + v_{2,3}$, where

$$\begin{cases} v_{2,1}(t, x) := \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^{r^3} \int_0^t e^{(t-\tau)\Delta} \mathcal{P}U_{j,j}^+(\tau, x) d\tau, \\ v_{2,2}(t, x) := \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^{r^3} \int_0^t e^{(t-\tau)\Delta} \mathcal{P}U_{j,j}^-(\tau, x) d\tau, \\ v_{2,3}(t, x) := \frac{1}{\Gamma_2^2(r)} \sum_{j=1}^{r^3} \sum_{i < j} \int_0^t e^{(t-\tau)\Delta} \mathcal{P}(U_{i,j}(\tau, x) + U_{j,i}(\tau, x)) d\tau. \end{cases} \quad (2.23)$$

Here,

$$\begin{aligned} U_{i,j}(\tau, x) := & -\frac{k_i k_j}{2\sqrt{i_j}} \mathbf{v}_j \left((\mathbf{v}_i \cdot \mathbf{k}_j) \left(\sin((\mathbf{k}_j + \mathbf{k}_i) \cdot x) + \sin((\mathbf{k}_j - \mathbf{k}_i) \cdot x) \right) e^{-\tau(|\mathbf{k}_j|^2 + |\mathbf{k}_i|^2)} \right. \\ & + (\mathbf{v}_i' \cdot \mathbf{k}_j) \left(\sin((\mathbf{k}_j + \mathbf{k}_i') \cdot x) + \sin((\mathbf{k}_j - \mathbf{k}_i') \cdot x) \right) e^{-\tau(|\mathbf{k}_j|^2 + |\mathbf{k}_i'|^2)} \\ & - \frac{k_i k_j}{2\sqrt{i_j}} \mathbf{v}_j' \left((\mathbf{v}_i \cdot \mathbf{k}_j') \left(\sin((\mathbf{k}_j' + \mathbf{k}_i) \cdot x) + \sin((\mathbf{k}_j' - \mathbf{k}_i) \cdot x) \right) e^{-\tau(|\mathbf{k}_j'|^2 + |\mathbf{k}_i|^2)} \right. \\ & \left. \left. + (\mathbf{v}_i' \cdot \mathbf{k}_j') \left(\sin((\mathbf{k}_j' + \mathbf{k}_i') \cdot x) + \sin((\mathbf{k}_j' - \mathbf{k}_i') \cdot x) \right) e^{-\tau(|\mathbf{k}_j'|^2 + |\mathbf{k}_i'|^2)} \right) \right) \end{aligned}$$

and

$$U_{j,j}(\tau, x) = U_{j,j}^+(\tau, x) + U_{j,j}^-(\tau, x),$$

with

$$\begin{aligned} U_{j,j}^+(\tau, x) := & -\frac{k_j^2}{2j} \left(\mathbf{v}_j (\mathbf{v}_j' \cdot \mathbf{k}_j) \sin((\mathbf{k}_j + \mathbf{k}_j') \cdot x) e^{-\tau(|\mathbf{k}_j|^2 + |\mathbf{k}_j'|^2)} \right. \\ & \left. + \mathbf{v}_j' (\mathbf{v}_j \cdot \mathbf{k}_j') \sin((\mathbf{k}_j' + \mathbf{k}_j) \cdot x) e^{-\tau(|\mathbf{k}_j'|^2 + |\mathbf{k}_j|^2)} \right) \end{aligned}$$

Using the relation (2.22), it appears that

$$\begin{aligned} v_{2,1}(t, x) = & \frac{(-1, 1, 1)}{\Gamma_2^2(r)} \sum_{j=1}^r \frac{k_j^2}{6j} \int_0^t e^{-6(t-\tau)} e^{-(2k_j^2 + 2k_j + 4)\tau} \sin(x_1 + 2x_2 - x_3) d\tau \\ & + \frac{(0, -1, 0)}{\Gamma_2^2(r)} \sum_{j=r+1}^{r^2} \frac{k_j^2}{2j} \int_0^t e^{-(t-\tau)} e^{-(3+2k_j^2)\tau} \sin(x_1) d\tau \\ & + \frac{(1, 0, 0)}{\Gamma_2^2(r)} \sum_{j=r^2+1}^{r^3} \frac{k_j^2}{2j} \int_0^t e^{-(t-\tau)} e^{-(3+2k_j^2)\tau} \sin(-x_2) d\tau. \end{aligned} \quad (2.24)$$

Essentially the same arguments as in Lemma 2.1 yield that for all $t > 0$

$$\|v_{2,2}(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \frac{1}{\Gamma_2^2(r)} = \Gamma_1^{-\frac{2}{3}}(r), \quad (2.25)$$

$$\|v_{2,3}(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \frac{1}{\Gamma_2^2(r)} = \Gamma_1^{-\frac{2}{3}}(r), \quad (2.26)$$

$$\|u_1(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \frac{1}{\sqrt{t} \Gamma_2(r)} = \Gamma_1^{-\frac{1}{3}}(r) t^{-\frac{1}{2}}. \quad (2.27)$$

Let us focus on obtaining a lower bound in $\dot{B}_{\infty, \infty}^{-1}$ of the dot product of $v_{2,1}(t, \cdot)$ with any unit direction. This is the main difference with respect to the proof of

Theorem A. We claim that for all $t \in [T/320, T]$, for $r \geq 64$,

$$\inf_{\mathbf{e} \in \mathbb{R}^3: |\mathbf{e}|=1} \|v_{2,1}(t, \cdot) \cdot \mathbf{e}\|_{\dot{B}_{\infty, \infty}^{-1}} \gtrsim \Gamma_1^{\frac{1}{3}}(r). \quad (2.28)$$

To show this, we make use of the structure of the inflation term $v_{2,1}$ in (2.24) and we also utilize the following simple fact from algebra. Namely for any unit vector $(\alpha, \beta, \gamma) \in \mathbb{R}^3$,

$$\max \left(\left| \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} \right|, \left| \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} \cdot \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} \right|, \left| \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right| \right) \geq \frac{1}{4\sqrt{2}}. \quad (2.29)$$

According to (2.29), first suppose that the unit vector $\mathbf{e} = (\alpha, \beta, \gamma)$ satisfies

$$\left| \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} \right| \geq \frac{1}{4\sqrt{2}}. \quad (2.30)$$

Using this, the form of $v_{2,1}$ in (2.24) and the same arguments as in (2.10)-(2.11), we obtain that for $t \in [T/320, T]$

$$\begin{aligned} \sup_{s>0} s^{\frac{1}{2}} |e^{s\Delta} v_{2,1}(t, 0, 0, \frac{\pi}{2}) \cdot \mathbf{e}| &\gtrsim \frac{\sup_{s>0} s^{\frac{1}{2}} e^{-6s}}{\Gamma_2^2(r)} \sum_{j=1}^r \frac{k_j^2}{j} \int_0^t e^{-6(t-\tau)} e^{-\tau(2k_j^2+2k_j+4)} d\tau \\ &\gtrsim \Gamma_1^{\frac{1}{3}}(r). \end{aligned}$$

Hence, in the first case (2.30) we get that for all $t \in [T/320, T]$ and $r \geq 64$

$$\|v_{2,1}(t, \cdot) \cdot \mathbf{e}\|_{\dot{B}_{\infty, \infty}^{-1}} \gtrsim \Gamma_1^{\frac{1}{3}}(r).$$

For the second case according to (2.29), suppose that the unit vector $\mathbf{e} = (\alpha, \beta, \gamma)$ satisfies

$$\left| \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} \cdot \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} \right| \geq \frac{1}{4\sqrt{2}}. \quad (2.31)$$

From this, the form of $v_{2,1}$ in (2.24) and similar arguments as in the first case, we obtain that for $t \in [T/320, T]$ and $r \geq 64$

$$\begin{aligned} \sup_{s>0} s^{\frac{1}{2}} |e^{s\Delta} v_{2,1}(t, \frac{\pi}{2}, 0, \frac{\pi}{2}) \cdot \mathbf{e}| &\gtrsim \frac{\sup_{s>0} s^{\frac{1}{2}} e^{-s}}{\Gamma_2^2(r)} \sum_{j=r+1}^{r^2} \frac{k_j^2}{j} \int_0^t e^{-(t-\tau)} e^{-\tau(2k_j^2+3)} d\tau \\ &\gtrsim \frac{1}{\Gamma_2^2(r)} \sum_{j=r+1}^{r^2} \frac{1}{j} \gtrsim \frac{\Gamma_1(r)}{\Gamma_2^2(r)} \gtrsim \Gamma_1^{\frac{1}{3}}(r). \end{aligned}$$

Hence, in the second case (2.31) we get that for all $t \in [T/320, T]$ and $r \geq 64$

$$\|v_{2,1}(t, \cdot) \cdot \mathbf{e}\|_{\dot{B}_{\infty, \infty}^{-1}} \gtrsim \Gamma_1^{\frac{1}{3}}(r).$$

For the third and final case according to (2.29), suppose that the unit vector $\mathbf{e} = (\alpha, \beta, \gamma)$ satisfies

$$\left| \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right| \geq \frac{1}{4\sqrt{2}}. \quad (2.32)$$

From this, the form of $v_{2,1}$ in (2.24) and similar arguments as in the previous cases, we obtain that for $t \in [T/320, T]$ and $r \geq 64$

$$\begin{aligned} \sup_{s>0} s^{\frac{1}{2}} |e^{s\Delta} v_{2,1}(t, 0, \frac{\pi}{2}, 0) \cdot \mathbf{e}| &\gtrsim \frac{\sup_{s>0} s^{\frac{1}{2}} e^{-s}}{\Gamma_2^2(r)} \sum_{j=r^2+1}^{r^3} \frac{k_j^2}{j} \int_0^t e^{-(t-\tau)} e^{-\tau(2k_j^2+3)} d\tau \\ &\gtrsim \frac{1}{\Gamma_2^2(r)} \sum_{j=r^2+1}^{r^3} \frac{1}{j} \gtrsim \frac{\Gamma_1(r)}{\Gamma_2^2(r)} \gtrsim \Gamma_1^{\frac{1}{3}}(r). \end{aligned}$$

Hence, in the third case (2.32) we get that for all $t \in [T/320, T]$ and $r \geq 64$

$$\|v_{2,1}(t, \cdot) \cdot \mathbf{e}\|_{\dot{B}_{\infty, \infty}^{-1}} \gtrsim \Gamma_1^{\frac{1}{3}}(r).$$

Combing these three cases, we see that we have established (2.28).

Using (2.28) with (2.25)-(2.27), we see that the final error analysis is carried out as in **Step 3** of Theorem A above, choosing $\delta = \Gamma_1^{-\frac{1}{3}}(r)$ and $T = \Gamma_1^{-1}(r)$. This concludes the proof of Theorem A'.

3. SYMMETRY BREAKING IN THE PRESENCE OF FAVORABLE PRESSURE

The objective of this section is to prove Theorem B. In the following, we construct an initial data $(u_{\text{in}}^h, 0)$ satisfying condition (1.5) and such that the condition $u^3 \equiv 0$ is instantly broken for the Navier-Stokes problem (1.1). For further insights about the heuristics behind our construction, we refer to Subsection 1.4.

In this section, we use the data introduced in Theorem B:

$$u_{\text{in}} = \left(\cos x_2 \frac{N}{N + \sin x_3}, \cos x_1 \frac{N + \sin x_3}{N}, 0 \right).$$

First, let us explain why a unique solution u exists on $[0, 1] \times \mathbb{T}^3$ for N sufficiently large. Let

$$u_{\text{in}}^{2D} = (\cos x_2, \cos x_1, 0)$$

and let $u^{2D} \in L^\infty((0, 1) \times \mathbb{T}^3)$ be the two-dimensional global solution. Then,

$$u_{\text{in}} - u_{\text{in}}^{2D} = \left(-\cos x_2 \frac{\sin x_3}{N + \sin x_3}, \cos x_1 \frac{\sin x_3}{N}, 0 \right) \quad (3.1)$$

and we see finding u is equivalent to finding U on $[0, 1] \times \mathbb{T}^3$ satisfying

$$U = e^{t\Delta}(u_{\text{in}} - u_{\text{in}}^{2D}) + \mathcal{B}(U, u^{2D}) + \mathcal{B}(u^{2D}, U) + \mathcal{B}(U, U). \quad (3.2)$$

Using the previously discussed L^∞ -bilinear estimates and (3.1), we see that for sufficiently large N we can apply [23, Lemma A.1] on successive time intervals to get existence of $U \in L^\infty((0, 1) \times \mathbb{T}^3)$ satisfying (3.2).

We rewrite the Navier-Stokes equations (1.1) as

$$u = e^{t\Delta} u_{\text{in}} + \mathcal{B}(u, u).$$

Now, we define the first and the second approximate solutions in the following natural way: let $u_1 = e^{t\Delta} u_{\text{in}}$ and

$$u_2 := e^{t\Delta} u_{\text{in}} + v_2 \quad \text{with} \quad v_2 := \mathcal{B}(u_1, u_1).$$

We denote the difference between u and the second approximation u_2 by w . Then w satisfies the integral equation (2.12).

Step 1: analysis of v_2 . We show that for the initial data u_{in} the third component of the first approximate solution u_2^3 has a non-zero L^∞ norm for a short time interval. Notice that

$$u_1(t, x) = \begin{pmatrix} e^{-t} \cos x_2 e^{t\partial_3^2} \left(\frac{N}{N + \sin x_3} \right) \\ e^{-t} \cos x_1 \frac{N + e^{-t} \sin x_3}{N} \\ 0 \end{pmatrix}. \quad (3.3)$$

Recalling the definition

$$v_2(t, x) = - \int_0^t e^{(t-\tau)\Delta} \mathcal{P}(u_1 \cdot \nabla u_1)(\tau, x) d\tau,$$

we have

$$\begin{aligned} v_2(t, x) &= \int_0^t e^{(t-\tau)\Delta} \mathcal{P} \begin{pmatrix} e^{-2\tau} \cos x_1 \sin x_2 F(\tau, x_3) \\ e^{-2\tau} \sin x_1 \cos x_2 F(\tau, x_3) \\ 0 \end{pmatrix} d\tau \\ &= -2 \int_0^t e^{-2\tau} e^{(t-\tau)\Delta} \mathcal{P} \begin{pmatrix} 0 \\ 0 \\ \sin x_1 \sin x_2 \partial_3 F(\tau, x_3) \end{pmatrix} d\tau \\ &= - \int_0^t e^{-2\tau} e^{(t-\tau)\Delta} (-\Delta)^{-1} \begin{pmatrix} \cos x_1 \sin x_2 \partial_3^2 F(\tau, x_3) \\ \sin x_1 \cos x_2 \partial_3^2 F(\tau, x_3) \\ 2 \sin x_1 \sin x_2 \partial_3 F(\tau, x_3) \end{pmatrix} d\tau, \end{aligned} \quad (3.4)$$

where $F(\tau, x_3) := \frac{N + e^{-\tau} \sin x_3}{N} e^{\tau \partial_3^2} \left(\frac{N}{N + \sin x_3} \right)$.

Since we are not able to write an explicit formula for $\partial_3 F$, we need to determine the main contributions of $\partial_3 F$ while keeping control of the remainder parts. Unlike the case of [42], there is no way to use the Taylor series $e^{\tau \Delta} = \sum_{j \in \mathbb{N}} \frac{(\tau \Delta)^j}{j!}$ to single out the main parts of $\partial_3 F$. Indeed at each order of $e^{\tau \partial_3^2} \frac{\cos x_3}{(N + \sin x_3)^2}$ there is a remainder term $\frac{\cos x_3}{(N + \sin x_3)^2}$ and thus we are not able to control the tail, even for a short time. Therefore, our idea is to first write a Taylor expansion for $\frac{N}{N + \sin x_3}$ and then compute the associated heat flows.

Since $u_2^3 = v_2^3$ we need to consider

$$\begin{aligned} \partial_3 F(\tau, x_3) &= e^{-\tau} \left\{ N(1 - e^\tau) e^{\tau \partial_3^2} \left(\frac{\cos x_3}{(N + \sin x_3)^2} \right) + \cos x_3 (e^{\tau \partial_3^2} - 1) \left(\frac{1}{N + \sin x_3} \right) \right. \\ &\quad \left. - (N + \sin x_3) (e^{\tau \partial_3^2} - 1) \left(\frac{\cos x_3}{(N + \sin x_3)^2} \right) \right\}. \end{aligned}$$

It is clear that due to the structure of the initial data one has that $\partial_3 F(\tau, x_3)|_{\tau=0} = 0$ and $\partial_\tau \partial_3 F(\tau, x_3)|_{\tau=0} \sim \frac{1}{N^2}$. Thus, for a short time $\partial_3 F(\tau, x_3) \sim \frac{\tau}{N^2}$.

We write

$$\frac{1}{N + \sin x_3} = \frac{1}{N} - \frac{\sin x_3}{N^2} + R_1(x_3) \quad \text{with} \quad R_1(x_3) := \frac{\sin^2 x_3}{(N + \sin x_3)N^2}.$$

Then, we have

$$\begin{aligned} (e^{\tau \partial_3^2} - 1) \left(\frac{1}{N + \sin x_3} \right) &= (e^{\tau \partial_3^2} - 1) \left(\frac{1}{N} - \frac{\sin x_3}{N^2} + R_1(x_3) \right) \\ &= \frac{\sin x_3}{N^2} (1 - e^{-\tau}) + (e^{\tau \partial_3^2} - 1) R_1(x_3). \end{aligned} \quad (3.5)$$

Notice that $\frac{d}{dx_3} \frac{1}{N + \sin x_3} = -\frac{\cos x_3}{(N + \sin x_3)^2}$, so one has that

$$\frac{\cos x_3}{(N + \sin x_3)^2} = \frac{\cos x_3}{N^2} - \frac{\sin(2x_3)}{N^3} + R_2(x_3) \quad \text{with} \quad R_2 := \frac{(3N + 2 \sin x_3) \sin^2 x_3 \cos x_3}{(N + \sin x_3)^2 N^3}.$$

Furthermore,

$$\begin{aligned} (e^{\tau\partial_3^2} - 1) \left(\frac{\cos x_3}{(N + \sin x_3)^2} \right) &= (e^{\tau\partial_3^2} - 1) \left(\frac{\cos x_3}{N^2} - \frac{\sin(2x_3)}{N^3} + R_2(x_3) \right) \\ &= \frac{\cos x_3}{N^2} (e^{-\tau} - 1) + \frac{\sin(2x_3)}{N^3} (1 - e^{-4\tau}) + (e^{\tau\partial_3^2} - 1) R_2(x_3). \end{aligned} \quad (3.6)$$

For the first term in the formula of $\partial_3 F$, we note that

$$e^{\tau\partial_3^2} \left(\frac{\cos x_3}{(N + \sin x_3)^2} \right) = \frac{\cos x_3}{N^2} e^{-\tau} - \frac{\sin(2x_3)}{N^3} e^{-4\tau} + e^{\tau\partial_3^2} R_2(x_3). \quad (3.7)$$

Substituting (3.5)-(3.7) into the formula of $\partial_3 F$ and then using (3.4), we get

$$\begin{aligned} \Delta u_2^3(t, x) &= \frac{2}{N^2} \int_0^t e^{(t-\tau)\Delta} \left\{ \sin x_1 \sin x_2 \sin(2x_3) (e^{-6\tau} - e^{-4\tau}) \right\} d\tau \\ &\quad - \frac{2}{N^3} \int_0^t e^{(t-\tau)\Delta} \left\{ \sin x_1 \sin x_2 \sin x_3 \sin(2x_3) (e^{-3\tau} - e^{-7\tau}) \right\} d\tau \\ &\quad + \int_0^t e^{(t-\tau)\Delta} 2e^{-3\tau} \sin x_1 \sin x_2 \left\{ N(1 - e^\tau) e^{\tau\partial_3^2} R_2(x_3) \right. \\ &\quad \left. + \cos x_3 (e^{\tau\partial_3^2} - 1) R_1(x_3) - (N + \sin x_3) (e^{\tau\partial_3^2} - 1) R_2(x_3) \right\} d\tau \\ &=: S_1(t, x) + S_2(t, x) + S_3(t, x). \end{aligned}$$

Now, we are ready to estimate $\Delta u_2^3(t, x)$ in the space $\dot{B}_{\infty, \infty}^{-2}$ for a small time t ($0 < t \ll 1$). For $s > 0$, we find that

$$\begin{aligned} e^{s\Delta} S_1(t, x) &= \frac{2}{N^2} \sin x_1 \sin x_2 \sin(2x_3) \int_0^t e^{-6(t+s-\tau)} (e^{-6\tau} - e^{-4\tau}) d\tau \\ &= \frac{2}{N^2} \sin x_1 \sin x_2 \sin(2x_3) (te^{-6t} - \frac{1}{2}(e^{-4t} - e^{-6t})) e^{-6s}, \end{aligned}$$

thus

$$\begin{aligned} \|S_1(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-2}} &= \sup_{s>0} s \|e^{s\Delta} S_1(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \\ &= \frac{1}{N^2} (e^{-4t} - (2t+1)e^{-6t}) \sup_{s>0} s e^{-6s} \geq \frac{1}{6e} \frac{t^2}{N^2}. \end{aligned} \quad (3.8)$$

To estimate $e^{s\Delta} S_2$, let us recall the formula

$$\sin x_3 \sin(2x_3) = \frac{1}{2} \cos(x_3) - \frac{1}{2} \cos(3x_3),$$

then

$$\begin{aligned} e^{s\Delta} S_2(t, x) &= \frac{1}{N^3} \sin x_1 \sin x_2 e^{-2(t+s)} \int_0^t e^{(t+s-\tau)\partial_3^2} (\cos(3x_3) - \cos x_3) (e^{-\tau} - e^{-5\tau}) d\tau \\ &= \frac{1}{N^3} \sin x_1 \sin x_2 \cos(3x_3) \left(\frac{1}{8} e^{-3t} - \frac{1}{4} e^{-7t} + \frac{1}{8} e^{-11t} \right) e^{-11s} \\ &\quad - \frac{1}{N^3} \sin x_1 \sin x_2 \cos x_3 \left(te^{-3t} + \frac{1}{4} e^{-7t} - \frac{1}{4} e^{-3t} \right) e^{-3s}. \end{aligned}$$

Thus, we have

$$\begin{aligned} & \|e^{s\Delta}S_2(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \\ & \leq \frac{1}{N^3} \left(\frac{1}{8}e^{-3t} - \frac{1}{4}e^{-7t} + \frac{1}{8}e^{-11t} \right) e^{-11s} + \frac{1}{N^3} \left(te^{-3t} + \frac{1}{4}e^{-7t} - \frac{1}{4}e^{-3t} \right) e^{-3s} \end{aligned}$$

and

$$\begin{aligned} & \|S_2(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-2}} = \sup_{s>0} s \|e^{s\Delta}S_2(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \\ & \leq \frac{1}{N^3} \left(e^{-11t} \left(\frac{1}{8}(e^{8t} - 1) + \frac{1}{4}(1 - e^{4t}) \right) + e^{-3t} \left(t + \frac{1}{4}e^{-4t} - \frac{1}{4} \right) \right) \sup_{s>0} se^{-3s} \leq \frac{4}{3e} \frac{t^2}{N^3}. \end{aligned} \quad (3.9)$$

To estimate S_3 , we write

$$\begin{aligned} e^{s\Delta}S_3(t, x) &= 2e^{-2(t+s)} \sin x_1 \sin x_2 \int_0^t e^{-\tau} \left\{ N(1 - e^\tau)e^{(t+s)\partial_3^2} R_2(x_3) \right. \\ & \quad \left. + e^{(t+s-\tau)\partial_3^2} \left(\cos x_3(e^{\tau\partial_3^2} - 1)R_1(x_3) - (N + \sin x_3)(e^{\tau\partial_3^2} - 1)R_2(x_3) \right) \right\} d\tau, \end{aligned}$$

and by Lemma 1.1

$$\begin{aligned} & \|e^{s\Delta}S_3(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \\ & \leq 2e^{-2(t+s)} \int_0^t e^{-\tau} \left\{ (e^\tau - 1)N\|R_2\|_{L^\infty(\mathbb{T})} + \tau\|R_1''\|_{L^\infty(\mathbb{T})} + (N+1)\tau\|R_2''\|_{L^\infty(\mathbb{T})} \right\} d\tau \\ & \leq 2e^{-2(t+s)} \left((t-1 + e^{-t})N\|R_2\|_{L^\infty(\mathbb{T})} + (1 - (t+1)e^{-t})(\|R_1''\|_{L^\infty(\mathbb{T})} + (N+1)\|R_2''\|_{L^\infty(\mathbb{T})}) \right). \end{aligned}$$

It is easy to check that

$$\|R_2\|_{L^\infty(\mathbb{T})} \leq \frac{16}{N^4}$$

and

$$\|R_1''\|_{L^\infty(\mathbb{T})} \leq \frac{48}{N^3}, \quad \|R_2''\|_{L^\infty(\mathbb{T})} \leq \frac{928}{N^4}.$$

Thus

$$\|S_3(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-2}} = \sup_{s>0} s \|e^{s\Delta}S_3(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \leq \frac{496}{e} \frac{t^2}{N^3}. \quad (3.10)$$

Let $N_0 := 6000$. Combining (3.8)-(3.10), we have for all $N > N_0$,

$$\begin{aligned} & \|u_2^3(t, \cdot)\|_{L^\infty} \geq \|\Delta u_2^3(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-2}} \\ & \geq \|S_1(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-2}} - \|S_2(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-2}} - \|S_3(t, \cdot)\|_{\dot{B}_{\infty, \infty}^{-2}} \geq \frac{1}{12e} \frac{t^2}{N^2}. \end{aligned} \quad (3.11)$$

Step 2: further analysis of v_2 . Similar to previous computations, one obtains that

$$\|\partial_3 F(\tau, \cdot)\|_{L^\infty(\mathbb{T})} + \|\partial_3^2 F(\tau, \cdot)\|_{L^\infty(\mathbb{T})} \lesssim \frac{\tau}{N}. \quad (3.12)$$

Thus, from (3.4) and (3.12) we have

$$\|v_2(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \|\Delta v_2(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \int_0^t e^{-2\tau} \frac{\tau}{N} d\tau \lesssim \frac{t^2}{N}. \quad (3.13)$$

Meanwhile, from (3.3) we see that

$$\|u_1(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim e^{-t}. \quad (3.14)$$

Step 3: final error estimate. Now we analyze the remaining part of the solution, which we denote by w . We use L^∞ bilinear estimates for controlling the error. Recall equation (2.12), estimates (3.13), (3.14). Therefore using the equation for the perturbation (2.12) and the estimates (3.13), (3.14), we have for all $0 < T < 1$,

$$\begin{aligned} & \sup_{0 \leq t \leq T} \|w(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \\ & \leq \sup_{0 \leq t \leq T} (\|\mathcal{B}(w, u_1)\|_{L^\infty} + \|\mathcal{B}(u_1, w)\|_{L^\infty} + \|\mathcal{B}(w, v_2)\|_{L^\infty} + \|\mathcal{B}(v_2, w)\|_{L^\infty} \\ & \quad + \|\mathcal{B}(w, w)\|_{L^\infty} \|\mathcal{B}(u_1, v_2)\|_{L^\infty} + \|\mathcal{B}(v_2, u_1)\|_{L^\infty} + \|\mathcal{B}(v_2, v_2)\|_{L^\infty}) \\ & \lesssim \sup_{0 \leq t \leq T} (\|\mathcal{B}(w, u_1)\|_{L^\infty} + \|\mathcal{B}(w, v_2)\|_{L^\infty} + \|\mathcal{B}(w, w)\|_{L^\infty} + \|\mathcal{B}(u_1, v_2)\|_{L^\infty} \\ & \quad + \|\mathcal{B}(v_2, v_2)\|_{L^\infty}) \\ & \lesssim T^{\frac{1}{2}} \left(1 + \frac{T^2}{N} + \sup_{0 \leq t \leq T} \|w(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \right) \sup_{0 \leq t \leq T} \|w(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} + \left(\frac{T^{\frac{5}{2}}}{N} + \frac{T^{\frac{9}{2}}}{N^2} \right). \end{aligned}$$

By an absorbing argument (see Lemma 1.2) for $T \ll 1$ and $N \gg 1$, we obtain

$$\sup_{0 \leq t \leq T} \|w(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \frac{T^{\frac{5}{2}}}{N}. \quad (3.15)$$

Using the fact that $u^3 = u_2^3 + w^3$, estimates (3.11) and (3.15) imply that

$$\begin{aligned} \|u^3(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} & \geq \|u_2^3(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} - \|w(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \\ & \gtrsim t^2 \left(\frac{1}{N^2} - \frac{t^{\frac{1}{2}}}{N} \right) \gtrsim \frac{t^2}{N^2} \end{aligned}$$

and

$$\|u^3(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \leq \|u_2^3(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} + \|w(t, \cdot)\|_{L^\infty(\mathbb{T}^3)} \lesssim \frac{t^2}{N^2}$$

for any $0 \leq t \leq \frac{1}{2N^2} \ll 1$.

This completes the proof of Theorem B.

4. 2.75D SHEAR FLOWS

4.1. 2.75D shear flows and nonlinear inviscid damping. Let us now describe the aforementioned example of a 3D Euler solution that weakly converges but does not strongly converges to the Kolmogorov flow $u^K = (0, \sin x_3, 0)$ in $L^2(\mathbb{T}^3)$ as $t \rightarrow \infty$. In (1.11), take $f(y_1, y_2) = \sin y_1$ and $g(y_3) = -\sin y_3$. Then the following smooth initial data

$$(\sin(\lambda x_1 + x_2), -\lambda \sin(\lambda x_1 + x_2) + \sin x_3, 0), \quad x \in \mathbb{T}^3$$

gives a global-in-time solution to the 3D Euler equations of the form

$$u^E = \begin{pmatrix} \sin(\lambda x_1 + x_2 - t \sin x_3) \\ -\lambda \sin(\lambda x_1 + x_2 - t \sin x_3) + \sin x_3 \\ 0 \end{pmatrix}. \quad (4.1)$$

Next consider a continuous function $\eta : [0, \frac{\pi}{2}] \rightarrow \mathbb{R}$, which is compactly supported in $(0, \frac{\pi}{2})$. Moreover, consider

$$\int_0^{\frac{\pi}{2}} \sin(t \sin x_3) \eta(x_3) dx_3 = \int_0^1 \sin(ty) \frac{\eta(\arcsin(y))}{\sqrt{1-y^2}} dy \rightarrow 0, \quad \text{as } t \rightarrow \infty, \quad (4.2)$$

where we used the fact that $\{\sin(ty)\}_{t>0}$ converges weakly to zero in the space $L^2([0, 1])$. Using the same arguments used to establish (4.2), one can conclude that $\{\sin(\lambda x_1 + x_2 + t \cos x_3)\}_{t>0}$ weakly converges to zero in $L^2(\mathbb{T}^3)$ as $t \rightarrow \infty$. Hence, $u^E(\cdot, t)$ converges weakly to u^K in $L^2(\mathbb{T}^3)$ as $t \rightarrow \infty$. To see that $u^E(\cdot, t)$ does not converge strongly to u^K in $L^2(\mathbb{T}^3)$ as $t \rightarrow \infty$, note that the L^2 norm of $u^E(\cdot, t)$ is conserved in time and is initially not equal to the L^2 norm of u^K .

4.2. 2.75D shear flows and Onsager supercritical inviscid limits. The solution $u^\nu : \mathbb{R}_+ \times \mathbb{T}^3 \rightarrow \mathbb{R}^3$ defined by (1.9) satisfies the following energy equality

$$\|u^\nu(t, \cdot)\|_{L^2}^2 + 2\nu \int_0^t \int_{\mathbb{T}^3} |\nabla u^\nu(s, x)|^2 dx ds = \|u_{\text{in}}\|_{L^2}^2, \text{ for all } t \geq 0. \quad (4.3)$$

Such solutions are unique in the class of 2.75D flows sharing the same symmetry, yet may not necessarily be unique in the general class of weak Leray-Hopf solutions with the same initial data in $L^2(\mathbb{T}^3)$.¹¹ Recall from (1.11) that the corresponding Euler solution with initial data (1.8) is given by

$$u^E(t, x) = (f(\lambda x_1 + x_2 + tg(x_3), x_3), -\lambda f(\lambda x_1 + x_2 + tg(x_3), x_3) - g(x_3), 0). \quad (4.4)$$

In this section, we investigate properties of u^ν , u^E and the vanishing viscosity limit.

Proposition 4.1. *Let $f \in L^2(\mathbb{T}^2; \mathbb{R})$ and $g \in C^\infty(\mathbb{T})$. Consider initial data u_{in} of the form (1.8) with associated f, g . Suppose that u^ν is the global-in-time solution given by (1.9) to the problem (1.1) ($\nu > 0$) with initial data u_{in} , and u^E is the global-in-time solution given by (4.4) for the 3D Euler equations with initial data u_{in} .*

The above set up implies that the following holds true:

- (1) $\forall T > 0, u^\nu \rightarrow u^E$ in $L^2((0, T) \times \mathbb{T}^3)$ as $\nu \rightarrow 0$.
- (2) (pointwise convergence) $\forall t \geq 0, u^\nu(t, \cdot) \rightarrow u^E(t, \cdot)$ in $L^2(\mathbb{T}^3)$ as $\nu \rightarrow 0$.
- (3) There is an absence of anomalous dissipation in the vanishing viscosity limit. Namely, for all $t \geq 0$:

$$\lim_{\nu \rightarrow 0} \nu \int_0^t \int_{\mathbb{T}^3} |\nabla u^\nu(s, x)|^2 dx ds = 0. \quad (4.5)$$

- (4) Let $f \in L^3(\mathbb{T}^2; \mathbb{R})$ then u^E satisfies the local energy balance. Namely for all positive $\varphi \in C^\infty(\mathbb{R}_+ \times \mathbb{T}^3; \mathbb{R})$ and $t \geq 0$

$$\begin{aligned} \int_{\mathbb{T}^3} |u^E(t, x)|^2 \varphi(t, x) dx - \int_{\mathbb{T}^3} |u_{\text{in}}(x)|^2 \varphi(0, x) dx \\ = \int_0^t \int_{\mathbb{T}^3} \partial_s \varphi |u^E|^2 + |u^E|^2 u^E \cdot \nabla \varphi dx ds. \end{aligned} \quad (4.6)$$

Proof of Proposition 4.1. We first prove item (1). Following the same arguments as [5, Theorem 5], we see that

$$u^\nu \xrightarrow{*} u^E \text{ in } L^\infty(0, T; L^2(\mathbb{T}^3)) \text{ as } \nu \rightarrow 0. \quad (4.7)$$

Using arguments along the same lines as [4], we have that for all $t \geq 0$,

$$\|u^E(\cdot, t)\|_{L^2}^2 = \|u_{\text{in}}\|_{L^2}^2. \quad (4.8)$$

¹¹By weak-strong uniqueness, 2.75D shear flows (1.9) with $f \in L^p$, $p \geq 3$, and $g \in C^\infty$ are unique amongst the general class of weak Leray-Hopf solutions.

Thanks to (4.7), one has that

$$u^\nu \rightharpoonup u^E \text{ weakly in } L^2((0, T) \times \mathbb{T}^3), \text{ as } \nu \rightarrow 0. \quad (4.9)$$

Then it is enough to show

$$\int_0^T \int_{\mathbb{T}^3} |u^\nu(s, x)|^2 dx ds \rightarrow \int_0^T \int_{\mathbb{T}^3} |u^E(s, x)|^2 dx ds, \text{ as } \nu \rightarrow 0. \quad (4.10)$$

We have the integrated energy balances:

$$\begin{aligned} \int_0^T \int_{\mathbb{T}^3} |u^\nu(t, x)|^2 dx dt + 2\nu \int_0^T \int_0^t \int_{\mathbb{T}^3} |\nabla u^\nu(s, x)|^2 dx ds dt &= T \|u_{\text{in}}\|_{L^2}^2, \\ \int_0^T \int_{\mathbb{T}^3} |u^E(t, x)|^2 dx dt &= T \|u_{\text{in}}\|_{L^2}^2. \end{aligned}$$

From the first energy balance above,

$$\begin{aligned} &\limsup_{\nu \rightarrow 0} \left(\int_0^T \int_{\mathbb{T}^3} |u^\nu(t, x)|^2 dx dt \right) \\ &\leq \limsup_{\nu \rightarrow 0} \left(\int_0^T \int_{\mathbb{T}^3} |u^\nu(t, x)|^2 dx dt + 2\nu \int_0^T \int_0^t \int_{\mathbb{T}^3} |\nabla u^\nu(s, x)|^2 dx ds dt \right) = T \|u_{\text{in}}\|_{L^2}^2. \end{aligned}$$

From (4.9) and (4.8),

$$T \|u_{\text{in}}\|_{L^2}^2 = \int_0^T \int_{\mathbb{T}^3} |u^E(t, x)|^2 dx dt \leq \liminf_{\nu \rightarrow 0} \left(\int_0^T \int_{\mathbb{T}^3} |u^\nu(t, x)|^2 dx dt \right).$$

Thus (4.10) is satisfied and we get $u^\nu \rightarrow u^E$ in $L^2((0, T) \times \mathbb{T}^3)$ as $\nu \rightarrow 0$.

To prove item (2), we proceed similarly as for item (1). First, due to (1.9)-(1.10) and the energy equality (4.3), we have that

$$\sup_{\nu} \|\partial_t u^\nu\|_{L^2(0, T; W^{-1,2}(\mathbb{T}^3))} < \infty. \quad (4.11)$$

This, together¹² with a classical diagonalisation argument and (4.7), yields that

$$\forall t \geq 0 \quad u^\nu(\cdot, t) \rightharpoonup u^E(\cdot, t) \text{ in } L^2(\mathbb{T}^3). \quad (4.12)$$

Using (4.3) again, we write

$$\begin{aligned} &\limsup_{\nu \rightarrow 0} \int_{\mathbb{T}^3} |u^\nu(t, x)|^2 dx \\ &\leq \limsup_{\nu \rightarrow 0} \left(\int_{\mathbb{T}^3} |u^\nu(t, x)|^2 dx + 2\nu \int_0^t \int_{\mathbb{T}^3} |\nabla u^\nu(s, x)|^2 dx ds \right) = \|u_{\text{in}}\|_{L^2}^2. \end{aligned}$$

By virtue of (4.12) and the energy balance (4.8) for the associated Euler flow, we write

$$\|u_{\text{in}}\|_{L^2}^2 = \int_{\mathbb{T}^3} |u^E(t, x)|^2 dx \leq \liminf_{\nu \rightarrow 0} \int_{\mathbb{T}^3} |u^\nu(t, x)|^2 dx \leq \|u_{\text{in}}\|_{L^2}^2.$$

¹²We refer to [39, page 104] where a similar argument is used.

Thus

$$\int_{\mathbb{T}^3} |u^\nu(t, x)|^2 dx \rightarrow \int_{\mathbb{T}^3} |u^E(t, x)|^2 dx, \quad \text{as } \nu \rightarrow 0,$$

which together with (4.12) gives item (2).

For the proof of item (3), notice that for any $t \geq 0$,

$$\int_{\mathbb{T}^3} |u^\nu(t, x)|^2 dx + 2\nu \int_0^t \int_{\mathbb{T}^3} |\nabla u^\nu(s, x)|^2 dx ds = \|u_{\text{in}}\|_{L^2}^2 = \int_{\mathbb{T}^3} |u^E(t, x)|^2 dx.$$

It is then easy to find that

$$\lim_{\nu \rightarrow 0} \left(\int_{\mathbb{T}^3} |u^\nu(t, x)|^2 dx + 2\nu \int_0^t \int_{\mathbb{T}^3} |\nabla u^\nu(s, x)|^2 dx ds \right) = \int_{\mathbb{T}^3} |u^E(t, x)|^2 dx.$$

Combining with item (2) implies (4.5).

Let us now prove item (4). The proof includes two steps: at first, we mollify f to get a series of smooth solutions in the form of (1.11) and write down the local energy equalities for these smooth solutions, then we pass to the limit by using that $f \in L^3$ (which is sharp as an assumption in view of the nonlinear contribution to the right-hand-side of (4.6)).

Note that since $f \in L^3(\mathbb{T}^2; \mathbb{R})$, there exists sequence $\{f_k\}_{k \in \mathbb{N}} \in C^\infty(\mathbb{T}^2; \mathbb{R})$ such that $\|f_k - f\|_{L^3} \rightarrow 0$, $k \rightarrow \infty$. Define

$$u^{E,k}(t, x) = (f_k(\lambda x_1 + x_2 + tg(x_3), x_3), -\lambda f_k(\lambda x_1 + x_2 + tg(x_3), x_3) - g(x_3), 0).$$

By Fubini's theorem, one has

$$\begin{aligned} & \|u^{E,k}(t, \cdot) - u^E(t, \cdot)\|_{L^3}^3 \\ & \leq (|\lambda|^3 + 1) \int_{\mathbb{T}^3} |f_k(\lambda x_1 + x_2 + tg(x_3), x_3) - f(\lambda x_1 + x_2 + tg(x_3), x_3)|^3 dx \\ & = 2\pi(|\lambda|^3 + 1) \|f_k - f\|_{L^3}^3 \rightarrow 0, \quad \text{as } k \rightarrow \infty \end{aligned}$$

and

$$\lim_{k \rightarrow \infty} \|u^{E,k}(0, \cdot) - u_{\text{in}}(\cdot)\|_{L^2} = 0.$$

Now, since $u^{E,k}$ is smooth on $\mathbb{R}_+ \times \mathbb{T}^3$, we have that for any positive $\varphi \in C^\infty(\mathbb{R}_+ \times \mathbb{T}^3)$ and $t \geq 0$,

$$\begin{aligned} & \int_{\mathbb{T}^3} |u^{E,k}(t, x)|^2 \varphi(t, x) dx \\ & = \int_{\mathbb{T}^3} |u^{E,k}(0, x)|^2 \varphi(0, x) dx + \int_0^t \int_{\mathbb{T}^3} \partial_t \varphi |u^{E,k}|^2 + |u^{E,k}|^2 u^{E,k} \cdot \nabla \varphi dx ds. \end{aligned}$$

Using above convergence properties, we pass to the limit as $k \rightarrow \infty$, and obtain the energy balance (4.6). \square

Note that the third item in Proposition 4.1 only gives convergence of energy pointwise. Below we show that if the Euler solution has some additional Onsager *supercritical* Sobolev regularity, then one obtains a uniform convergence of the energy (i.e. $u^\nu \rightarrow u^E$ in $L^\infty(\mathbb{R}_+; L^2(\mathbb{T}^3))$) and a rate of vanishing of the anomalous dissipation. Proposition 4.1 also implies that the sufficient conditions in [18] and [36] on the Euler flow for the inviscid limit to hold in $L^\infty(\mathbb{R}_+; L^2(\mathbb{T}^3))$ are not

necessary conditions. For 2D results on the vanishing viscosity and conservation of energy in Onsager supercritical regimes we refer to [17, 34].

In particular, we have the following result.

Proposition 4.2. *Let $f \in H^\ell(\mathbb{T}^2; \mathbb{R})$ with $\ell \in (0, \frac{5}{6})$. Consider initial data u_{in} in the form of (1.8) with associated f and $g \in C^\infty(\mathbb{T})$. Suppose that u^ν is the global-in-time solution given by (1.9) for the problem (1.1) ($\nu > 0$) with initial data u_{in} , and u^E is the global-in-time solution given by (4.4) for the 3D Euler equations with initial data u_{in} .*

The above set up implies that the following holds true:

(1) *For $t \geq 0$ and $\nu \in (0, 1)$,*

$$\nu \int_0^t \int_{\mathbb{T}^3} |\nabla u^\nu| dx ds \leq \nu^\ell C(\lambda, \ell, t, \|g'\|_{L^\infty}, \|f\|_{H^\ell}).$$

(2) *If $f \in H^\ell(\mathbb{T}^2; \mathbb{R})$ with $\ell \in [\frac{1}{2}, \frac{5}{6})$, then*

$$u^\nu \rightarrow u^E \quad \text{in } C([0, T]; L^2(\mathbb{T}^3)), \quad \forall T > 0$$

and for any $\ell' \in [0, \ell)$ and $\nu \in (0, 1)$,

$$\sup_{t \in [0, T]} \|u^\nu(t, \cdot) - u^E(t, \cdot)\|_{\dot{H}^{\ell'}} \leq \nu^{\frac{1}{4}(1-\frac{\ell'}{\ell})} C(\lambda, \ell, \ell', T, \|g\|_{W^{3, \infty}}, \|f\|_{H^\ell}). \quad (4.13)$$

Proof. First let us establish item (1). Fix $\nu > 0$. For $f \in L^2(\mathbb{T}^2; \mathbb{R})$, let $T_\nu(f) := F_\nu$ be such that F_ν satisfies (1.10). Then

$$T_\nu : L^2(\mathbb{T}^2) \rightarrow L^2(0, \infty; \dot{H}^1(\mathbb{T}^2)) \cap C([0, \infty; L^2(\mathbb{T}^2))$$

is a well-defined linear operator, due to the uniqueness of solutions to (1.10) in the class $L^2(0, \infty; \dot{H}^1(\mathbb{T}^2)) \cap C([0, \infty; L^2(\mathbb{T}^2))$ with L^2 initial data. Furthermore, an L^2 energy estimate on (1.10) yields

$$\|T_\nu(f)\|_{L^\infty(0, \infty; L^2)}, \nu^{\frac{1}{2}} \|T_\nu(f)\|_{L^2(0, \infty; \dot{H}^1)} \leq \|f\|_{L^2}. \quad (4.14)$$

When $f \in H^1(\mathbb{T}^2)$, applying an H^1 energy estimate to (1.10) and then Gronwall's inequality yields

$$\|T_\nu(f)\|_{L^\infty(0, \infty; \dot{H}^1)}, \nu^{\frac{1}{2}} \|T_\nu(f)\|_{L^2(0, \infty; \dot{H}^2)} \lesssim e^{ct\|g'\|_{L^\infty}} \|f\|_{H^1}. \quad (4.15)$$

Here, $c > 0$ is a universal constant. Using (4.14)-(4.15), we can apply [28, Theorem 2.2.10], [6, Theorem 6.45] and the interpolation theory for linear operators [1, Theorem 7.23]. This yields that for every $f \in H^\ell(\mathbb{T}^2)$ ($\ell \in (0, \frac{5}{6})$) and $t \geq 0$

$$\nu^{\frac{1}{2}} \|F_\nu\|_{L^2(0, t; \dot{H}^{1+\ell})} \lesssim e^{c\ell t\|g'\|_{L^\infty}} \|f\|_{H^\ell}. \quad (4.16)$$

Applying similar arguments pointwise in time to $T_\nu(f)$ also yields that for every $f \in H^\ell(\mathbb{T}^2)$ ($\ell \in (0, \frac{5}{6})$) and $t \geq 0$

$$\|F_\nu\|_{L^\infty(0, t; \dot{H}^\ell)} \lesssim e^{c\ell t\|g'\|_{L^\infty}} \|f\|_{H^\ell}. \quad (4.17)$$

Using the interpolation inequality for Sobolev spaces, Hölder's inequality and (4.16)-(4.17) gives that for any $f \in H^\ell(\mathbb{T}^2)$ ($\ell \in (0, \frac{5}{6})$) and $t \geq 0$:

$$\begin{aligned} \nu \int_0^t \int_{\mathbb{T}^2} |\nabla F_\nu|^2 dx ds &\leq \nu \int_0^t \|F_\nu(s, \cdot)\|_{\dot{H}^\ell}^{2\ell} \|F_\nu(s, \cdot)\|_{\dot{H}^{1+\ell}}^{2(1-\ell)} ds \\ &\leq (\nu t)^\ell \|F_\nu\|_{L^\infty(0, t; \dot{H}^\ell)}^{2\ell} \left(\nu \int_0^t \|F_\nu(s, \cdot)\|_{\dot{H}^{1+\ell}}^2 ds \right)^{1-\ell} \end{aligned}$$

$$\lesssim_\ell (\nu t)^\ell e^{2c\ell t \|g'\|_{L^\infty}} \|f\|_{H^\ell}^2. \quad (4.18)$$

Recall that u^ν is given by (1.9), where $\lambda \in \mathbb{Z} \setminus \{0\}$. Together with (4.18), this gives

$$\begin{aligned} \nu \int_0^t \int_{\mathbb{T}^3} |\nabla u^\nu|^2 dx ds &\lesssim \lambda^4 \nu \int_0^t \int_{\mathbb{T}^3} |\nabla F_\nu|^2 dx ds + \nu \int_0^t \int_{\mathbb{T}} |\nabla e^{\nu s \partial_3^2} g|^2 dx ds \\ &\lesssim_\ell (\nu t)^\ell e^{2c\ell t \|g'\|_{L^\infty}} \|f\|_{H^\ell}^2 + t\nu \|g'\|_{L^2}^2. \end{aligned}$$

This establishes item (1).

Let us now prove item (2). Define

$$F(t, x_1, x_2) := f(x_1 + tg(x_2), x_2),$$

which is a distributional solution to

$$\begin{cases} \partial_t F - g(y_2) \partial_1 F = 0 & \text{in } \mathbb{T}^2 \times \mathbb{R}_+ \\ F(0, y_1, y_2) = f(y_1, y_2). \end{cases} \quad (4.19)$$

Furthermore, by Fubini's theorem

$$\|F(t, \cdot)\|_{L^2} = \|f\|_{L^2} \quad \forall t \geq 0. \quad (4.20)$$

Similar arguments as those used to establish (4.17) give that for all $f \in H^\ell$ and $t \geq 0$

$$\|F\|_{L^\infty(0,t; \dot{H}^\ell)} \lesssim \max(1, t \|g'\|_{L^\infty})^\ell \|f\|_{H^\ell}. \quad (4.21)$$

Hence, using this and (4.17) gives

$$\|F_\nu - F\|_{L^\infty(0,t; \dot{H}^\ell)} \lesssim (\max(1, t \|g'\|_{L^\infty})^\ell + e^{c\ell t \|g'\|_{L^\infty}}) \|f\|_{H^\ell} \quad (4.22)$$

Using (4.21), (4.20) and the interpolation inequality for Sobolev spaces, one deduces that

$$\|F\|_{L^\infty(0,t; \dot{H}^{\frac{1}{2}})} \lesssim_\ell \max(1, t \|g'\|_{L^\infty})^{\frac{1}{2}} \|f\|_{H^\ell}. \quad (4.23)$$

Similarly, (4.14) and (4.17) imply

$$\|F_\nu\|_{L^\infty(0,t; \dot{H}^{\frac{1}{2}})} \lesssim_\ell e^{\frac{c\ell t \|g'\|_{L^\infty}}{2}} \|f\|_{H^\ell}. \quad (4.24)$$

Using the interpolation inequality for Sobolev spaces and Hölder's inequality, we have

$$\begin{aligned} \nu \int_0^t \|F_\nu\|_{\dot{H}^{\frac{3}{2}}}^2 ds &\leq \nu \int_0^t \|F_\nu\|_{\dot{H}^\ell}^{2(\ell-\frac{1}{2})} \|F_\nu\|_{\dot{H}^{1+\ell}}^{2(\frac{3}{2}-\ell)} ds \\ &\leq \|F_\nu\|_{L^\infty(0,t; \dot{H}^\ell)}^{2(\ell-\frac{1}{2})} \left(\nu \int_0^t \|F_\nu\|_{\dot{H}^{1+\ell}}^2 ds \right)^{\frac{3}{2}-\ell} (t\nu)^{\ell-\frac{1}{2}}. \end{aligned}$$

Using this and (4.16)-(4.17) gives

$$\nu \int_0^t \|F_\nu\|_{\dot{H}^{\frac{3}{2}}}^2 ds \leq e^{c\ell t \|g'\|_{L^\infty}} \|f\|_{H^\ell}^2 (t\nu)^{\ell-\frac{1}{2}}. \quad (4.25)$$

Next, we consider the equation satisfied by $F_\nu - F$:

$$\begin{cases} \partial_t (F_\nu - F) - g(y_2) \partial_1 (F_\nu - F) = (e^{\nu \partial_2^2} g - g(y_2)) \partial_1 F_\nu \\ \quad + \nu ((\lambda^2 + 1) \partial_1^2 + \partial_2^2) F_\nu & \text{in } \mathbb{T}^2 \times \mathbb{R}_+ \\ (F_\nu - F)(0, y_1, y_2) = 0. \end{cases} \quad (4.26)$$

Performing an L^2 energy estimate¹³ on (4.26) gives

$$\begin{aligned} \|(F_\nu - F)(t, \cdot)\|_{L^2}^2 &= 2 \int_0^t \int_{\mathbb{T}^2} (e^{s\nu\partial_{y_2}^2} g - g(y_2)) \partial_1 F_\nu (F_\nu - F) dy_1 dy_2 ds \\ &\quad + 2 \int_0^t \int_{\mathbb{T}^2} \nu ((\lambda^2 + 1) \partial_1^2 + \partial_2^2) F_\nu (F_\nu - F) dy_1 dy_2 ds := I + II. \end{aligned} \quad (4.27)$$

First, let us estimate I :

$$|I| \leq 2t^{\frac{1}{2}} \|e^{s\nu\partial_{y_2}^2} g - g\|_{L^\infty(\mathbb{T} \times (0, t))} \|F_\nu - F\|_{L^\infty(0, t; L^2)} \left(\int_0^t \|\partial_1 F_\nu\|_{L^2}^2 ds \right)^{\frac{1}{2}}.$$

This, Lemma 1.1, (4.14) and (4.20) imply that

$$|I| \lesssim t^{\frac{3}{2}} \nu^{\frac{1}{2}} \|f\|_{L^2} \|g''\|_{L^\infty}. \quad (4.28)$$

Now we estimate II in (4.27):

$$|II| \leq 2(\nu t)^{\frac{1}{2}} (\lambda^2 + 1) \|F_\nu - F\|_{L^\infty(0, t; \dot{H}^{\frac{1}{2}})} \left(\nu \int_0^t \|F_\nu\|_{\dot{H}^{\frac{3}{2}}}^2 ds \right)^{\frac{1}{2}}.$$

Using (4.23)-(4.24) and (4.25) gives

$$|II| \lesssim_\ell (\nu t)^{\frac{1}{4} + \frac{\ell}{2}} (\lambda^2 + 1) e^{\frac{c\ell t \|g'\|_{L^\infty}}{2}} \left(\max(1, t \|g'\|_{L^\infty})^{\frac{1}{2}} + e^{\frac{c t \|g'\|_{L^\infty}}{2}} \right) \|f\|_{H^\ell}. \quad (4.29)$$

Combining (4.28)-(4.29) gives

$$\begin{aligned} &\sup_{t \in [0, T]} \|F_\nu(t, \cdot) - F(t, \cdot)\|_{L^2}^2 \\ &\lesssim_\ell (\nu T)^{\frac{1}{4} + \frac{\ell}{2}} (\lambda^2 + 1) e^{\frac{c\ell T \|g'\|_{L^\infty}}{2}} \left(\max(1, T \|g'\|_{L^\infty})^{\frac{1}{2}} \right. \\ &\quad \left. + e^{\frac{c T \|g'\|_{L^\infty}}{2}} \right) \|f\|_{H^\ell} + T^{\frac{3}{2}} \nu^{\frac{1}{2}} \|f\|_{L^2} \|g''\|_{L^\infty}. \end{aligned} \quad (4.30)$$

Using this, the fact that u^ν and u^E are given by (1.9) and (4.4) and Lemma 1.1, we get the following. Namely,

$$\begin{aligned} &\sup_{t \in [0, T]} \|u^\nu(t, \cdot) - u^E(t, \cdot)\|_{L^2}^2 \\ &\lesssim_\ell \sup_{t \in [0, T]} \|e^{t\nu\partial_{y_2}^2} g - g\|_{L^2}^2 + (\nu T)^{\frac{1}{4} + \frac{\ell}{2}} (\lambda^2 + 1) e^{\frac{c\ell T \|g'\|_{L^\infty}}{2}} \left(\max(1, T \|g'\|_{L^\infty})^{\frac{1}{2}} \right. \\ &\quad \left. + e^{\frac{c T \|g'\|_{L^\infty}}{2}} \right) \|f\|_{H^\ell} + T^{\frac{3}{2}} \nu^{\frac{1}{2}} \|f\|_{L^2} \|g''\|_{L^\infty} \\ &\lesssim (\nu T)^2 \|g''\|_{L^\infty} + (\nu T)^{\frac{1}{4} + \frac{\ell}{2}} (\lambda^2 + 1) e^{\frac{c\ell T \|g'\|_{L^\infty}}{2}} \left(\max(1, T \|g'\|_{L^\infty})^{\frac{1}{2}} \right. \\ &\quad \left. + e^{\frac{c T \|g'\|_{L^\infty}}{2}} \right) \|f\|_{H^\ell} + T^{\frac{3}{2}} \nu^{\frac{1}{2}} \|f\|_{L^2} \|g''\|_{L^\infty}. \end{aligned} \quad (4.31)$$

This gives (4.13) for $\ell' = 0$ as required. To get (4.13) for $\ell' \in (0, \ell)$ we interpolate (4.30) with (4.22) to get

$$\sup_{t \in [0, T]} \|F_\nu(t, \cdot) - F(t, \cdot)\|_{\dot{H}^{\ell'}} \leq \nu^{\frac{1}{4}(1 - \frac{\ell'}{\ell})} C(\ell, \ell', T, \|g\|_{W^{2, \infty}}, \|f\|_{H^\ell}). \quad (4.32)$$

Furthermore, using Lemma 1.1 we see that

$$\sup_{t \in [0, T]} \|e^{t\nu\partial_{y_2}^2} g - g\|_{\dot{H}^{\ell'}} \lesssim (\nu T) \|g\|_{W^{3, \infty}}. \quad (4.33)$$

¹³All subsequent estimates can be rigorously justified by approximating $f \in H^\ell$ by smooth $f_k \rightarrow f$ in H^ℓ .

By similar reasoning as the $\ell' = 0$ case we then get

$$\begin{aligned} & \sup_{t \in [0, T]} \|u^\nu(t, \cdot) - u^E(t, \cdot)\|_{\dot{H}^{\ell'}} \\ & \leq |\lambda|^{1+\ell'} \sup_{t \in [0, T]} \|F_\nu(t, \cdot) - F(t, \cdot)\|_{\dot{H}^{\ell'}} + \sup_{t \in [0, T]} \|e^{t\nu\partial_y^2} g - g\|_{\dot{H}^{\ell'}}. \end{aligned} \quad (4.34)$$

Combining this with (4.32)-(4.33) gives (4.13) for all $\ell' \in [0, \ell)$ as required. \square

4.3. Strong ill-posedness for 3D Euler equations in anisotropic spaces.

Let $f \in W^{1,p}(\mathbb{T}; W^{1,q}(\mathbb{T}))$, $g \in W^{1,q}(\mathbb{T})$ and $p, q \in [2, \infty)$. For $\lambda \in \mathbb{Z} \setminus \{0\}$, consider the following initial data

$$(f(\lambda x_1 + x_2, x_3), -\lambda f(\lambda x_1 + x_2, x_3) - g(x_3), 0) \in W^{1,p}(\mathbb{T}^2; W^{1,q}(\mathbb{T})),$$

which generates an explicit solution¹⁴ u^E in the form of (1.11) to the 3D Euler equations on the torus. Using identical reasoning as in [3], it is clear that the roughness of g means that $u^E(t, x)$ will not lie in the expected solution space $W^{1,p}(\mathbb{T}^2; W^{1,q}(\mathbb{T}))$ for any positive time t , which shows strong ill-posedness in the sense of Hadamard (non-existence) in the anisotropic Sobolev space. We also anticipate that it is possible to show illposedness of the 3D Euler equations for initial data that has dependence on three spatial dimensions and belongs to other anisotropic spaces (analogous to the isotropic spaces considered in [4]).

APPENDIX A. HEURISTICS FOR THE STRUCTURE OF 2.75D SHEAR FLOWS

In this appendix we give some heuristics about the derivation of 2.75D shear flows. As mentioned earlier, they are rotated versions of the parallel flows introduced by Wang [44]. Here we outline another derivation based on the analysis of the following reduced Navier-Stokes system

$$\begin{cases} \partial_t u^h + u^h \cdot \nabla_h u^h + \nabla_h P = \Delta u^h, \\ \partial_3 P = 0, \\ \operatorname{div}_h u^h = 0, \\ u^h|_{t=0} = u_{\text{in}}^h. \end{cases} \quad (\text{A.1})$$

We dub that system the ‘2.75D Navier-Stokes equations’.¹⁵ It is well-known that solutions to this system with H^1 data are smooth.¹⁶

Consider initial data u_{in}^h of the form

$$u_{\text{in}}^h = (\partial_2 \phi(x), -\partial_1 \phi(x)). \quad (\text{A.2})$$

Let us look for solutions of system (A.1) under the form

$$u^h = (\partial_2 \Phi(t, x), -\partial_1 \Phi(t, x)) = \nabla_h^\perp \Phi, \quad (\text{A.3})$$

¹⁴The fact that this is a weak solution to the 3D Euler equations uses the same arguments as in [4, Theorem 2]

¹⁵Once we get a solution for system (A.1), then it also satisfies the so-called primitive equations, see for example the works of Cao and Titi [9] and Hieber and Kashiwabara [25] on primitive equations.

¹⁶A byproduct of Theorem B is that system (A.1) is ill-posed for generic data in H^1 . The proof is by contradiction. In fact, if for any initial data u_{in}^h satisfying condition (1.5), there always exists a solution u^h to the problem (A.1) on some time interval $[0, T]$, then one can extend u^h to a solution $(u^h, 0)$ for the 3D Navier-Stokes problem (1.1). By local well-posedness theory for (1.1), regularity results in [37] and weak-strong uniqueness, one confirms that $u = (u^h, 0)$ is the unique solution on $[0, T]$ supplemented with initial data $(u_{\text{in}}^h, 0)$. In particular, it implies that $u^3 \equiv 0$ will be preserved.

where $\nabla_h^\perp := (\partial_2, -\partial_1)$. Notice that the pressure is now given by

$$\begin{cases} \Delta_h P = -\operatorname{div}_h(u^h \cdot \nabla_h u^h) = 2 \det(\operatorname{Hessian}_h \Phi) \\ \partial_3 P = 0, \end{cases}$$

where we used the vector identities

$$\begin{aligned} \operatorname{div}_h\left((\nabla_h^\perp \Phi) \cdot \nabla_h(\nabla_h^\perp \Phi)\right) &= \operatorname{div}_h \operatorname{div}_h\left((\nabla_h^\perp \Phi) \otimes (\nabla_h^\perp \Phi)\right) \\ &= (\nabla_h \nabla_h^\perp \Phi) : (\nabla_h \nabla_h^\perp \Phi)^\top = -2 \det(\operatorname{Hessian}_h \Phi), \end{aligned}$$

where

$$\operatorname{Hessian}_h := \begin{pmatrix} \partial_1^2 & \partial_1 \partial_2 \\ \partial_1 \partial_2 & \partial_2^2 \end{pmatrix}.$$

In order to have $\partial_3 P = 0$, one has to satisfy

$$\partial_3 \det(\operatorname{Hessian}_h \Phi) = 0.$$

If there exist a function $\Psi : \mathbb{T} \times \mathbb{R}_+ \rightarrow \mathbb{R}$ and a constant $\lambda \in \mathbb{Z}$ such that

$$\mathcal{L}_\lambda \Phi(t, x) = \Psi(t, x_3) \quad \text{with} \quad \mathcal{L}_\lambda := \partial_1 - \lambda \partial_2, \quad (\text{A.4})$$

then we have

$$\partial_1 \mathcal{L}_\lambda \Phi = \partial_2 \mathcal{L}_\lambda \Phi = 0 \quad \text{i.e.} \quad \begin{pmatrix} \partial_1^2 \Phi \\ \partial_1 \partial_2 \Phi \end{pmatrix} = \lambda \begin{pmatrix} \partial_1 \partial_2 \Phi \\ \partial_2^2 \Phi \end{pmatrix},$$

and thus

$$\det(\operatorname{Hessian}_h \Phi) = 0. \quad (\text{A.5})$$

In the following, we will focus on the case (A.4) for the Cauchy problem (A.1). Concerning the initial data, we also need to look for a function $\psi : \mathbb{T} \rightarrow \mathbb{R}$ such that

$$\mathcal{L}_\lambda \phi(x) - \psi(x_3) = 0. \quad (\text{A.6})$$

Recalling that the velocity $u^h = \nabla_h^\perp \Phi$ and taking into consideration (A.4), one has

$$u^h \cdot \nabla_h u^h = (-1, \lambda) \Psi \partial_2^2 \Phi$$

and P is a constant. Finally, we are lead to considering the following system

$$\begin{cases} \partial_t \partial_2 \Phi - \Psi(t, x_3) \partial_2^2 \Phi = \Delta \partial_2 \Phi & \text{in } \mathbb{R}_+ \times \mathbb{T}^3, \\ \partial_t (\lambda \partial_2 \Phi + \Psi(t, x_3)) - \lambda \Psi(t, x_3) \partial_2^2 \Phi = \Delta (\lambda \partial_2 \Phi + \Psi(t, x_3)) & \text{in } \mathbb{R}_+ \times \mathbb{T}^3, \\ (\Phi, \Psi)|_{t=0} = (\phi, \psi) \quad \text{with} \quad \mathcal{L}_\lambda \phi - \psi(x_3) = 0, \end{cases}$$

which can be simplified as

$$\begin{cases} \partial_t \partial_2 \Phi - \Psi(t, x_3) \partial_2^2 \Phi = \Delta \partial_2 \Phi & \text{in } \mathbb{R}_+ \times \mathbb{T}^3, \\ \partial_t \Psi(t, x_3) = \partial_3^2 \Psi(t, x_3) & \text{in } \mathbb{R}_+ \times \mathbb{T}, \\ (\Phi, \Psi)|_{t=0} = (\phi, \psi) \quad \text{with} \quad \mathcal{L}_\lambda \phi - \psi(x_3) = 0. \end{cases} \quad (\text{A.7})$$

In conclusion, $\Psi(t, x_3) = (\mathcal{K} \star \psi)(t, x_3)$, where \mathcal{K} is the one-dimensional heat kernel see (1.20), and $\partial_2 \Phi$ satisfies the *linear* transport-heat equation

$$\begin{cases} \partial_t v + V \cdot \nabla v = \Delta v & \text{in } \mathbb{R}_+ \times \mathbb{T}^3 \quad \text{with} \quad V = (0, -\Psi(t, x_3), 0), \\ v_{\text{in}} = \partial_2 \phi. \end{cases} \quad (\text{A.8})$$

Take $\psi(x_3) = g(x_3)$ and

$$\phi(x) = \phi(\lambda x_1 + x_2, x_3) = \int_0^{\lambda x_1 + x_2} f(y_1, x_3) dy_1 + x_1 \psi(x_3).$$

Obviously, $\phi(x)$ and $\psi(x_3)$ satisfy (A.6), so the associated solution $\partial_2 \Phi(t, x)$ of (A.8) for which u^h given by (A.3) solves problem (A.1).

ACKNOWLEDGMENT

The authors thank Hao Jia for stimulating discussions concerning the inviscid damping for the 2.75D shear flows, as well as Jean-Yves Chemin for a discussion relating to Theorem A. We also thank Pierre Gilles Lemarié-Rieusset and Helena J. Nussenzweig Lopes for their comments on an earlier version of this paper.

CP and JT are partially supported by the Agence Nationale de la Recherche, project BORDS, grant ANR-16-CE40-0027-01. CP is also partially supported by the Agence Nationale de la Recherche, project SINGFLOWS, grant ANR-18-CE40-0027-01, project CRISIS, grant ANR-20-CE40-0020-01, by the CY Initiative of Excellence, project CYNA (CY Nonlinear Analysis) and project CYFI (CYngular Fluids and Interfaces). JT is also supported by the Labex MME-DII. TB and CP thank the Institute of Advanced Studies of Cergy Paris University for their hospitality.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

- [1] R. A. Adams and J. J. F. Fournier. *Sobolev spaces*, volume 140 of *Pure and Applied Mathematics (Amsterdam)*. Elsevier/Academic Press, Amsterdam, second edition, 2003.
- [2] C. Bardos, M. C. Lopes Filho, D. Niu, H. Nussenzweig Lopes, and E. S. Titi. Stability of two-dimensional viscous incompressible flows under three-dimensional perturbations and inviscid symmetry breaking. *SIAM Journal on Mathematical Analysis*, 45(3):1871–1885, 2013.
- [3] C. Bardos and E. S. Titi. Euler equations for an ideal incompressible fluid. *Uspekhi Mat. Nauk*, 62(3(375)):5–46, 2007.
- [4] C. Bardos and E. S. Titi. Loss of smoothness and energy conserving rough weak solutions for the 3d Euler equations. *Discrete Contin. Dyn. Syst. Ser. S*, 3(2):185–197, 2010.
- [5] C. Bardos, E. S. Titi, and E. Wiedemann. The vanishing viscosity as a selection principle for the Euler equations: the case of 3D shear flow. *C. R. Math. Acad. Sci. Paris*, 350(15-16):757–760, 2012.
- [6] J. Bergh and J. Löfström. *Interpolation spaces. An introduction*. Grundlehren der Mathematischen Wissenschaften, No. 223. Springer-Verlag, Berlin-New York, 1976.
- [7] J. Bourgain and N. Pavlović. Ill-posedness of the Navier-Stokes equations in a critical space in 3D. *J. Funct. Anal.*, 255(9):2233–2247, 2008.
- [8] T. Buckmaster and V. Vicol. Nonuniqueness of weak solutions to the Navier-Stokes equation. *Ann. of Math. (2)*, 189(1):101–144, 2019.
- [9] C. Cao and E. S. Titi. Global well-posedness of the three-dimensional viscous primitive equations of large scale ocean and atmosphere dynamics. *Ann. of Math. (2)*, 166(1):245–267, 2007.
- [10] D. Chae and J. Wolf. On the Serrin-type condition on one velocity component for the Navier-Stokes equations. *Arch. Ration. Mech. Anal.*, 240(3):1323–1347, 2021.
- [11] J.-Y. Chemin, I. Gallagher, and P. Zhang. Some remarks about the possible blow-up for the Navier-Stokes equations. *Comm. Partial Differential Equations*, 44(12):1387–1405, 2019.
- [12] J.-Y. Chemin and P. Zhang. On the critical one component regularity for 3-D Navier-Stokes systems. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(1):131–167, 2016.
- [13] J.-Y. Chemin, P. Zhang, and Z. Zhang. On the critical one component regularity for 3-D Navier-Stokes system: general case. *Arch. Ration. Mech. Anal.*, 224(3):871–905, 2017.
- [14] J. Chen and T. Y. Hou. Stable nearly self-similar blowup of the 2D Boussinesq and 3D Euler equations with smooth data. *arXiv e-prints*, page arXiv:2210.07191, Oct. 2022.
- [15] A. Cheskidov, P. Constantin, S. Friedlander, and R. Shvydkoy. Energy conservation and Onsager’s conjecture for the Euler equations. *Nonlinearity*, 21(6):1233, 2008.
- [16] A. Cheskidov and M. Dai. Norm inflation for generalized Navier-Stokes equations. *Indiana Univ. Math. J.*, 63(3):869–884, 2014.

- [17] A. Cheskidov, M. C. L. Filho, H. J. N. Lopes, and R. Shvydkoy. Energy conservation in two-dimensional incompressible ideal fluids. *Comm. Math. Phys.*, 348(1):129–143, 2016.
- [18] P. Constantin. Note on loss of regularity for solutions of the 3-D incompressible Euler and related equations. *Comm. Math. Phys.*, 104(2):311–326, 1986.
- [19] P. Constantin, E. Weinan, and E. S. Titi. Onsager’s conjecture on the energy conservation for solutions of euler’s equation. *Communications in Mathematical Physics*, 165(1):207–209, 1994.
- [20] M. Coti Zelati, T. M. Elgindi, and K. Widmayer. Stationary Structures near the Kolmogorov and Poiseuille Flows in the 2d Euler Equations. *arXiv e-prints*, page arXiv:2007.11547, July 2020.
- [21] C. De Lellis and L. Székelyhidi, Jr. The Euler equations as a differential inclusion. *Ann. of Math. (2)*, 170(3):1417–1436, 2009.
- [22] T. Elgindi. Finite-time singularity formation for $C^{1,\alpha}$ solutions to the incompressible Euler equations on \mathbb{R}^3 . *Ann. of Math. (2)*, 194(3):647–727, 2021.
- [23] I. Gallagher, D. Iftimie, and F. Planchon. Asymptotics and stability for global solutions to the Navier-Stokes equations. *Ann. Inst. Fourier (Grenoble)*, 53(5):1387–1424, 2003.
- [24] J. Guillod and V. Šverák. Numerical investigations of non-uniqueness for the Navier-Stokes initial value problem in borderline spaces. *arXiv e-prints*, page arXiv:1704.00560, Apr. 2017.
- [25] M. Hieber and T. Kashiwabara. Global strong well-posedness of the three dimensional primitive equations in L^p -spaces. *Arch. Ration. Mech. Anal.*, 221(3):1077–1115, 2016.
- [26] T. Y. Hou. Potential singularity of the 3D Euler equations in the interior domain. *Foundations of Computational Mathematics*, pages 1–47, 2022.
- [27] T. Y. Hou. Potentially singular behavior of the 3D Navier–Stokes equations. *Foundations of Computational Mathematics*, pages 1–49, 2022.
- [28] T. Hytönen, J. van Neerven, M. Veraar, and L. Weis. *Analysis in Banach spaces. Vol. I. Martingales and Littlewood-Paley theory*, volume 63 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer, Cham, 2016.
- [29] P. Isett. A proof of onsager’s conjecture. *Annals of Mathematics*, 188(3):871–963, 2018.
- [30] A. N. Kolmogoroff. Dissipation of energy in the locally isotropic turbulence. *C. R. (Doklady) Acad. Sci. URSS (N.S.)*, 32:16–18, 1941.
- [31] A. Kolmogorov. The local structure of turbulence in incompressible viscous fluid for very large Reynold’s numbers. *C. R. (Doklady) Acad. Sci. URSS (N.S.)*, 30:301–305, 1941.
- [32] A. N. Kolmogorov. On degeneration of isotropic turbulence in an incompressible viscous liquid. *C. R. (Doklady) Acad. Sci. URSS (N. S.)*, 31:538–540, 1941.
- [33] O. A. Ladyženskaja. Unique global solvability of the three-dimensional Cauchy problem for the Navier-Stokes equations in the presence of axial symmetry. *Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 7:155–177, 1968.
- [34] S. Lanthaler, S. Mishra, and C. Parés-Pulido. On the conservation of energy in two-dimensional incompressible flows. *Nonlinearity*, 34(2):1084, 2021.
- [35] Y. Maekawa and Y. Terasawa. The Navier-Stokes equations with initial data in uniformly local L^p spaces. *Differential Integral Equations*, 19(4):369–400, 2006.
- [36] N. Masmoudi. Remarks about the inviscid limit of the Navier-Stokes system. *Comm. Math. Phys.*, 270(3):777–788, 2007.
- [37] J. Neustupa and P. Penel. Regularity of a suitable weak solution to the Navier-Stokes equations as a consequence of regularity of one velocity component. In *Applied nonlinear analysis*, pages 391–402. Kluwer/Plenum, New York, 1999.
- [38] V. Scheffer. An inviscid flow with compact support in space-time. *J. Geom. Anal.*, 3(4):343–401, 1993.
- [39] G. Seregin. *Lecture notes on regularity theory for the Navier-Stokes equations*. World Scientific, 2014.
- [40] A. Shnirelman. On the nonuniqueness of weak solution of the Euler equation. *Comm. Pure Appl. Math.*, 50(12):1261–1286, 1997.
- [41] M. R. Ukhovskii and V. I. Iudovich. Axially symmetric flows of ideal and viscous fluids filling the whole space. *J. Appl. Math. Mech.*, 32:52–61, 1968.
- [42] B. Wang. Ill-posedness for the Navier-Stokes equations in critical Besov spaces $\dot{B}_{\infty,q}^{-1}$. *Adv. Math.*, 268:350–372, 2015.
- [43] W. Wang, D. Wu, and Z. Zhang. Scaling invariant Serrin criterion via one velocity component for the Navier-Stokes equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 2023.
- [44] X. Wang. A Kato type theorem on zero viscosity limit of Navier-Stokes flows. *Indiana University Mathematics Journal*, pages 223–241, 2001.

- [45] T. Yoneda. Ill-posedness of the 3D-Navier-Stokes equations in a generalized Besov space near BMO^{-1} . *J. Funct. Anal.*, 258(10):3376–3387, 2010.

(T. Barker)

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF BATH, BATH BA2 7AY. UK
Email address: `tobiasbarker5@gmail.com`

(C. Prange)

LABORATOIRE DE MATHÉMATIQUES AGM, UMR CNRS 8088, CERGY PARIS UNIVERSITÉ, 2 AVENUE ADOLPHE CHAUVIN, 95302 CERGY-PONTOISE CEDEX, FRANCE
Email address: `christophe.prange@cyu.fr`

(J. Tan)

LABORATOIRE DE MATHÉMATIQUES AGM, UMR CNRS 8088, CERGY PARIS UNIVERSITÉ, 2 AVENUE ADOLPHE CHAUVIN, 95302 CERGY-PONTOISE CEDEX, FRANCE
Email address: `jin.tan@cyu.fr`